




Lecture Notes

Geometrical Anatomy of Theoretical Physics

Dr. Fredric P Schuller*

PIYUSH KUMAR SINGH †

December 2024

*YouTube [playlist](#)

†iamPiyushKrSingh.github.io

Structure of this Course

In theoretical physics we mainly deal with three big part of mathematics, namely, ‘Analysis’, ‘Algebra’ and ‘Geometry’. And at the mutual intersection of these three, we have different branches of physics like ‘Quantum Mechanics’, ‘General Relativity’, ‘Statistical Mechanics’, etc. see [fig. 0.1](#).




Figure 0.1: Structure of Theoretical Physics

This course mainly focuses on differential geometry and topology, and their applications in theoretical physics. And we will start with the basic propositional logic and set theory, and then move to the topology and geometry of manifolds, and then we will see the applications of these in physics. Following is the structure of this course:

- Logic
- Set Theory
- Topology
- Topological Manifolds
- Differential Manifolds
- Bundles
- Geometry: Symplectic Geometry, Metric Geometry, etc.
- Physics: Classical Mechanics, Electrodynamics, Quantum Mechanics, Statistical Mechanics, Special and General Relativity, etc.

Contents

Structure of this Course	ii
Contents	iv
1 Axiomatic Set Theory	1
Lecture 1	1
1.1 Propositional Logic	1
1.1.1 Logical Operators	1
1.2 Predicate Logic	2
1.3 Axiomatic Systems and Theory of Proofs	3
Lecture 2	4
1.4 The $\in$ -Relation	4
1.5 Zermelo-Fraenkel Axioms of Set Theory	5
Lecture 3	11
1.6 Classifications of Sets	11
1.7 Equivalence Relations	12
1.8 Construction of Naturals, Integers, Rationals and Reals	14
1.8.1 Natural Numbers	14
1.8.2 Integers	15
1.8.3 Rational Numbers	16
1.8.4 Real Numbers	17
2 Topological Spaces	19
Lecture 4	19
2.1 Topological Spaces	19
2.2 Construction of new Topologies from given topologies	21
2.2.1 Induced Topology	21
2.2.2 Quotient Topology	22
2.2.3 Product Topology	22
2.3 Convergence	23
2.4 Continuity	24
2.4.1 Homeomorphism	25
Lecture 5	26
2.5 Topological Properties I: Separation Axioms	26
2.6 Compactness and Paracompactness	27
2.6.1 Compactness	27
2.6.2 Paracompactness	28
2.7 Connectedness and Path-Connectedness	30
2.7.1 Connectedness	30
2.7.2 Path-Connectedness	31
2.8 Homotopic Curves and Fundamental Group	32
2.8.1 Fundamental Group	33
3 Topological Manifolds and Bundles	37
Lecture 6	37
3.1 Definition and Construction of Topological Manifolds	37
3.1.1 Submanifolds	37
3.1.2 Product Manifolds	38
3.2 Bundles	38
3.2.1 Fiber Bundles	39
3.2.2 Constructing Bundles	40
3.3 Bundle Morphisms	41

3.4	Viewing Manifolds from Atlases	43
4	Differentiable Manifolds	45
	Lecture 7	45
4.1	Adding Structure by refining the (maximal) $\mathcal{C}^0$ -atlas	45
	Lecture 8	48
4.2	Review of Vector Spaces	48
4.2.1	Tensors	52
4.2.2	Dimension of a Vector Space	53
4.2.3	Change of Basis	56
4.2.4	Determinants	59
4.2.4.1	Permutation Group	59
	Lecture 9	61
4.3	Tangent Spaces to a Manifold	62
4.3.1	Algebras and Derivations	65
4.3.2	Basis and Dimension of Tangent Space	67
4.3.3	Change of Basis and Coordinate Transformation	69
4.3.3.1	Position Vector in Physics	71
	Lecture 10: Construction of Tangent Bundle	71
4.4	Cotangent Spaces and Gradient	71
4.4.1	Basis of Cotangent Space	72
4.5	Push-Forward and Pull-Back	72
4.6	Immersions and Embeddings	74
4.7	Tangent Bundle and Vector Fields	75
4.7.1	Smooth Structure on Tangent Bundle	76
	Lecture 11	80
4.8	Tensor Fields and Modules	81
4.8.1	Zorn's Lemma	83
4.8.2	Proof of Module Basis Theorem	84
4.8.3	Module Construction and Important Terms	84
	Lecture 12	87
4.9	Differential Forms and de Rham Cohomology	87

1 Axiomatic Set Theory

LECTURE 1

§1.1 Propositional Logic

Definition 1.1 (Proposition):

A proposition p is a variable that can take the values “true” or “false”. No other values are allowed.

It is not the task of ‘propositional logic’ to determine whether a proposition is true or false. It is only concerned with the logical relationship between propositions.

Note: We can build new propositions from existing ones with the help of logical operators.

§1.1.1 Logical Operators

- (a) Unary Operators: These operators operate on a single proposition. There are four unary operators:

p	$\neg p$ Negation	$\text{id } p$ Identity	$\top p$ Tautology	$\perp p$ Contradiction
T	F	T	T	F
F	T	F	T	F

Table 1.1: Unary Operators

- (b) Binary Operators: These operators operate on two propositions. There are sixteen binary operators. Some important ones are:

p	q	$p \wedge q$ Conjunction (AND)	$p \vee q$ Disjunction (OR)	$p \vee\vee q$ Exclusive Or	$p \Rightarrow q$ Implication	$p \Leftrightarrow q$ Equivalence
T	T	T	T	F	T	T
T	F	F	T	T	F	F
F	T	F	T	T	T	F
F	F	F	F	F	T	T

Table 1.2: Binary Operators

Remark 1.2 (*ex falso quodlibet*). The definition of implication has two not so obvious cases. The first one is when the hypothesis is false, and the conclusion is true. The second one is when both the hypothesis and the conclusion are false. In both cases, the implication is true.

In other words, this says that we can conclude anything from a false assumption.

Theorem 1.3

$$(p \Rightarrow q) \Leftrightarrow ((\neg q) \Rightarrow (\neg p))$$

Proof:

We can prove this by truth table.

p	q	$\neg p$	$\neg q$	$p \Rightarrow q$	$(\neg q) \Rightarrow (\neg p)$
T	T	F	F	T	T
T	F	F	T	F	F
F	T	T	F	T	T
F	F	T	T	T	T

Since the columns for $p \Rightarrow q$ and $(\neg q) \Rightarrow (\neg p)$ are identical, we have shown that $(p \Rightarrow q) \Leftrightarrow ((\neg q) \Rightarrow (\neg p))$. Q.E.D.

Corollary 1.4

We can prove assertions by way of contradiction.

Remark 1.5 (Binding Order). We agree on the decreasing binding strength of the logical operators as follows:

$$\neg, \wedge, \vee, \Rightarrow, \Leftrightarrow$$

Remark 1.6 (Higher Order Operators). All higher order operators (e.g. $\heartsuit(p_1, p_2, \dots, p_N)$) can be constructed from one single binary operator i.e. NAND ($\uparrow$).

p	q	$p \uparrow q$
T	T	F
T	F	T
F	T	T
F	F	T

Table 1.3: NAND Operator

§1.2 Predicate Logic

Definition 1.7 (Predicate):

A predicate is a proposition-valued function of some variable(s).

Example 1.8

At this point, we don't know how to construct a predicate. But we can only talk about its value for a given value of the variable. In general, predicates are denoted as follows:

1. $P(x)$: true or false depending on the value of x .
2. $Q(x, y)$: true or false depending on the values of x and y .

We can construct new predicates from existing ones.

- (a) Let $P(x)$ and $R(y, z)$ are two given predicates then we can define a third predicate $Q(x, y, z) :\Leftrightarrow P(x) \wedge R(y, z)$ ¹.
- (b) Convert predicate P of one variable into a proposition:

¹The symbol $(:\Leftrightarrow)$ means that the left-hand side is defined to be equivalent to the right-hand side.

Definition 1.9 ('Universal' quantifier):

$$\boxed{\forall x : P(x)} \quad \text{reads as "for all } x, P(x) \text{ is true."} \quad (1.1)$$

defined to be true, if $P(x)$ is true independently of x .

Using for all quantifier, we can define a new proposition from a given predicate *i.e.* 'existence' quantifier.

Definition 1.10 ('Existential' quantifier):

$$\boxed{\exists x : P(x)} \quad \text{reads as "there exists an } x \text{ such that } P(x) \text{ is true."} \quad (1.2)$$

defined as $\exists x : P(x) :\Leftrightarrow \neg(\forall x : \neg P(x))$

Corollary 1.11

$$\forall x : \neg P(x) \Leftrightarrow \neg(\exists x : P(x)) \quad (1.3)$$

(c) Quantification for predicates of more than one variable:

$$Q(y) :\Leftrightarrow \forall x : P(x, y)$$

here, x is a *bound variable* and y is a *free variable*.

Remark 1.12 (Order of Quantifiers). The order of quantifiers is important. For example,

$$\underbrace{\forall x : \exists y : P(x, y)}_{\text{used for definition of inverse}} \quad \text{generically different proposition than} \quad \underbrace{\exists y : \forall x : P(x, y)}_{\text{used for definition of identity}}$$

§1.3 Axiomatic Systems and Theory of Proofs

Definition 1.13 (Axiomatic System):

An *axiomatic system* is a finite sequence of propositions $a_1, a_2, \dots, a_N$ called *axioms*.

Definition 1.14 (Proof):

A proof of a proposition p is within an axiomatic system $a_1, a_2, \dots, a_N$ is a finite sequence of propositions $q_1, q_2, \dots, (q_M = p)$ such that for any $1 \leq i \leq M$ in the sequence, either

(A) q_i is a proposition from the list of axioms, or

(T) q_i is a tautology, or

(M) "modus ponens"

$$\exists 1 \leq m, n < i : (q_m \wedge q_n \Rightarrow q_i) \text{ is true.}$$

This definition allows to easily recognize a proof by checking the sequence of propositions.

An altogether different matter is to actually find a proof.

Remark 1.15. If proposition p can be proven from an axiomatic system $a_1, a_2, \dots, a_N$, we often write this as

$$a_1, a_2, \dots, a_N \vdash p$$

and say that axiomatic system proves proposition p .

Remark 1.16 (Redundant Axioms). Any tautology, should it occur in the axioms, can be removed from the list of axioms without impairing the power of the axiomatic system.

Extreme case of this is: axiomatic system for propositional logic is ‘empty sequence.’

Definition 1.17 (Consistency):

An axiomatic system is consistent if there exists a proposition q which cannot be proven from the axioms.

$$\exists q : \neg(a_1, a_2, \dots, a_N \vdash q) \quad (1.4)$$

Idea behind consistency: Consider an axiomatic system containing contradicting propositions:

$$a_1, a_2, \dots, s, \dots, \neg s, \dots, a_N$$

Then by *modus ponens* (M), we can prove any proposition p as

$$s \wedge \neg s \Rightarrow p \text{ is a tautology}$$

This can be used as a marker for inconsistency *i.e.* if an axiomatic system can prove every proposition, then it is inconsistent.

Theorem 1.18

Propositional logic is consistent.

Proof:

Suffices to show that there exists a proposition which cannot be proven within propositional logic.

Propositional logic has an empty sequence of axioms. Only (T) and (M) must carry any proof $\Rightarrow$ only tautologies can be proven; *i.e.* for a proposition p , we can’t prove $p \wedge \neg p$. Q.E.D.

Theorem 1.19 (Gödel’s Incompleteness Theorem)

Any axiomatic system that is powerful enough to encode the elementary arithmetic of natural numbers is either inconsistent or contains a proposition that can neither be proven nor disproven.

Proof:

The proof of this theorem is complicated; but the basic idea is as follows:

- 1) assign a number to each (meta-)mathematical statement, now called *Gödel number*.
- 2) Use a “The barber shaves all man in his village who do not shave themselves”- type of argument to identify a proposition that is neither provable nor disprovable.

Q.E.D.

LECTURE 2

§1.4 The $\in$ -Relation

Set theory built on the postulate that there is a fundamental relation² called $\in$. There will be no definition of what $\in$ is, or what a set is.

Instead of defining what a set is, we have 9 axioms that speak of $\in$ and sets. Overviews of these axioms are as follows:

²A predicate of two variables

E	basic existence axiom
E	
P	Construction axiom
U	
R	
P	
I	further existence and construction
C	
F	Non-existence axiom

Table 1.4: Mnemonic for Axioms of Set Theory

Using the $\in$ -relation, we can immediately define following relations:

$$\text{Not an element: } x \notin y :\Leftrightarrow \neg(x \in y) \quad (1.5)$$

$$\text{Subset: } x \subseteq y :\Leftrightarrow \forall z : (z \in x \Rightarrow z \in y) \quad (1.6)$$

$$\text{Equality: } x = y :\Leftrightarrow x \subseteq y \wedge y \subseteq x \quad (1.7)$$

§1.5 Zermelo-Fraenkel Axioms of Set Theory

Axiom 1.20 (Axiom on $\in$ -relation): $x \in y$ is a proposition if and only if x and y are both sets.

$$\forall x : \forall y : (x \in y) \vee \neg(x \in y) \quad (1.8)$$

Counter example (Russell's Paradox): Assume there is some u that contains all sets that do not contain themselves as an element. To be precise,

$$\exists u : \forall z : (z \in u \Leftrightarrow z \notin z) \quad (1.9)$$

Problem: Is u a set?

If u was a set, then one must be able to determine whether $u \in u$ is true or false. As [axiom 1.20](#) states, $u \in u$ is a proposition if and only if u is a set.

Case 1: Assume $u \in u$ is true: $\xRightarrow[\text{def of } u]{} u \notin u. \quad \downarrow$

Case 2: Assume $u \in u$ is false: $\xLeftrightarrow[\text{def } \notin]{} u \notin u \xRightarrow[\text{def } u]{} u \in u. \quad \downarrow$

Conclusion: u is not a set. □

Axiom 1.21 (Existence of an empty set): *There exists a set that contains no elements.*

$$\exists x : \forall y : y \notin x \quad (1.10)$$

Theorem 1.22 (Uniqueness of the empty set)

There is only one empty set. And it is denoted by $\emptyset$.

Proof (Standard textbook style):

Assume x and x' are both empty sets. But then

since hypothesis is always false the implication is always true

$$\forall y : \underbrace{(y \notin x) \Rightarrow (y \in x')}_{\text{always false}}$$

But this just means that $x \subseteq x'$ from [eq. \(1.6\)](#). Conversely,

$$\forall y : (y \notin x') \Rightarrow (y \in x)$$

But thus $x' \subseteq x$. Hence, $x = x'$.

Q.E.D.

The above proof is a bit informal. Now we will prove it using the definition of proof in an axiomatic system.

Proof (Formal Version):

Suffices to show that if x and x' are both empty sets, then $x = x'$.

$$\begin{aligned} a_1 &\Leftrightarrow \forall y : y \notin x \\ a_2 &\Leftrightarrow \forall y : y \notin x' \end{aligned} \left. \vphantom{\begin{aligned} a_1 &\Leftrightarrow \forall y : y \notin x \\ a_2 &\Leftrightarrow \forall y : y \notin x' \end{aligned}} \right\} \text{assumptions or axioms}$$

$$q_1 \xLeftrightarrow[(T)] \forall y : y \notin x \Rightarrow \forall y : (y \in x \Rightarrow y \in x')$$

$$q_2 \xLeftrightarrow[(A)1] \forall y : y \notin x$$

$$q_3 \xLeftrightarrow[(M)1,2] \forall y : (y \in x \Rightarrow y \in x') \quad \boxed{\Leftrightarrow x \subseteq x'}$$

$$q_4 \xLeftrightarrow[(T)] \forall y : y \notin x' \Rightarrow \forall y : (y \in x' \Rightarrow y \in x)$$

$$q_5 \xLeftrightarrow[(A)2] \forall y : y \notin x'$$

$$q_6 \xLeftrightarrow[(M)4,5] \forall y : (y \in x' \Rightarrow y \in x) \quad \boxed{\Leftrightarrow x' \subseteq x}$$

$$q_7 \xLeftrightarrow[(M)3,6] x = x'$$

Q.E.D.

Axiom 1.23 (Axiom on Pair Sets): *Let x and y be sets. Then there exists a set that contains as its elements precisely the sets x and y .*

$$\forall x : \forall y : \exists m : \forall u : (u \in m \Leftrightarrow u = x \vee u = y) \quad (1.11)$$

Notation: denote this set m by $\{x, y\}$.

Theorem 1.24 (Ordering in Pair Sets)

Let x and y be sets. Then $\{x, y\} = \{y, x\}$.

Proof:

$$\begin{aligned} a &\in \{x, y\} \Rightarrow a \in \{y, x\} \Rightarrow \{x, y\} \subseteq \{y, x\} \\ a &\in \{y, x\} \Rightarrow a \in \{x, y\} \Rightarrow \{y, x\} \subseteq \{x, y\} \end{aligned}$$

Thus, $\{x, y\} = \{y, x\}$.

Q.E.D.

Definition 1.25 (Set with one element):

Let x be a set. Then there exists a set that contains as its element precisely the set x .

$$\{x\} := \{x, x\} \quad (1.12)$$

Axiom 1.26 (Axiom on Union Sets): Let x be a set. Then there exists a set whose elements are precisely the elements of the elements of x .

$$\forall x : \exists u : \forall y : (y \in u \Leftrightarrow \exists z : (z \in x \wedge y \in z)) \quad (1.13)$$

Notation: denote this set u by $\bigcup x$.

Example 1.27

Let a, b be sets. Then by [axiom 1.23](#), $\{a\}$ and $\{b\}$ are sets, and hence $x := \{\{a\}, \{b\}\}$ is a set. Then $\bigcup x = \{a, b\}$.

Observe that, since a and b are sets, then by [axiom 1.23](#), $\{a, b\}$ is a set. So it gives us a false impression that [axiom 1.26](#) is redundant. But it is not. Consider the following example:

Example 1.28

Let a, b, c be sets. Then by [axiom 1.23](#), $\{a, b\}$ and $\{c\}$ are sets, and hence $x := \{\{a, b\}, \{c\}\}$ is a set. Then

$$\bigcup x =: \{a, b, c\}. \quad (1.14)$$

With this example, we can generalize the definition of a finite set.

Definition 1.29 (Finite Set):

Let $a_1, a_2, \dots, a_N$ be sets. Define, recursively for all $N \geq 3$,

$$\{a_1, a_2, \dots, a_N\} := \bigcup \{\{a_1, a_2, \dots, a_{N-1}\}, \{a_N\}\} \quad (1.15)$$

Axiom 1.30 (Axiom of Replacement): Let R be a functional relation. Let m be a set. Then the image of m under R $[\text{im}_R(m)]$ is a set.

For this axiom to make sense, we need to define what a functional relation is.

Definition 1.31 (Functional Relation):

A relation R is functional if and only if

$$\forall x : \exists! y : R(x, y)^3 \quad (1.16)$$

We define the image of a set under a functional relation keeping in mind that [axiom 1.30](#) guarantees this object to be a set.

Definition 1.32 (Image of a Set under a Functional Relation):

Let R be a functional relation. Let m be a set. Then the image of m under R consists of all those elements y for which there is an element $x \in m$ such that $R(x, y)$.

The axiom of replacement is a very powerful axiom. It implies, but is not implied by, the “*principle of restricted comprehension*”.

³The symbol $\exists!$ is ‘Uniqueness’ quantifier and read as “there exists a unique” and is a shorthand for $\exists y : R(x, y) \wedge \forall y' : (R(x, y') \Rightarrow y = y')$.

Theorem 1.33 (Principle of Restricted Comprehension (PRC))

Let P be a predicate of one variable and let m be a set. Then those elements $y \in m$ for which $P(y)$ is true constitute a set.

Notation: this set is denoted by $\{y \in m \mid P(y)\}$.

PRC is not to be confused with the inconsistent “*principle of unrestricted comprehension*” (PUC) which allows for paradoxes like ‘Self-Reference Paradox.’

Consider the predicate $P(x)$ is true if x is a set that doesn’t contain themselves as an element. Thus, from PUC we have $\{y \mid P(y)\}$ is a set (sick), but this give rise to Russell’s Paradox in ‘*naive set theory*’

Intuitively PRC makes sure that the image set doesn’t grow bigger than a set itself.

Proof:

We can prove PRC using the axiom of replacement in following cases:

Case 1: $\neg \exists y \in m : P(y)$ ⁴ in this case,

$$\{y \in m \mid P(y)\} := \emptyset$$

Case 2: $\exists \hat{y} \in m : P(\hat{y})$, then define a relation R as follows:

$$R(x, y) := (P(x) \wedge x = y) \vee (\neg P(x) \wedge \hat{y} = y)$$

Claim: R is a functional relation.

Pf: Let x be a set. If $P(x)$ is true, then $R(x, y)$ is true if and only if $y = x$. If $P(x)$ is false, then $R(x, y)$ is true if and only if $y = \hat{y}$.

Thus, $\forall x : \exists! y : R(x, y)$. Hence, R is a functional relation. $\square$

Then by [axiom 1.30](#), $\text{im}_R(m)$ is a set. And by definition of R , $\forall y \in \text{im}_R(m) : P(y)$ ⁵, thus we can define $\{y \in m \mid P(y)\} := \text{im}_R(m)$.

Q.E.D.

Definition 1.34 (Set Difference):

Let u and m be sets such that $u \subseteq m$. Then the set difference of m and u is defined as

$$m \setminus u := \{x \in m \mid x \notin u\}. \quad (1.17)$$

The object $m \setminus u$ is a set due to PRC *i.e.* ultimately due to the axiom of replacement.

Definition 1.35 (Intersection of Sets):

Let x be a set. Then the intersection of x is defined as

$$\bigcap x := \{y \in \bigcup x \mid \forall z \in x : y \in z\} \quad (1.18)$$

Historically, in naive set theory, PUC was thought to be needed in order to define, for any set m ,

$$\mathcal{P}(m) \underset{\text{inconsistently}}{:=} \{u \mid u \subseteq m\}$$

This definition is circular, as we need to know a priori, from which bigger set the elements of $\mathcal{P}(m)$ are taken.

⁴The quantifier $\exists y \in m : P(y)$ is logically equivalent to $\neg(\forall y \in m : \neg P(y))$.

⁵The quantifier $\forall y \in m : P(y)$ is logically equivalent to $\forall y : (y \in m \Rightarrow P(y))$.

Axiom 1.36 (Axiom on existence of power sets): *Let m be a set. Then there exists a set, denoted by $\mathcal{P}(m)$, whose elements are precisely the subsets of m .*

$$\forall m : \exists p : \forall u : (u \in p \Leftrightarrow u \subseteq m) \quad (1.19)$$

Example 1.37

Let $m = \{a, b\}$, then $\mathcal{P}(m) = \{\emptyset, \{a\}, \{b\}, \{a, b\}\}$.

At this point, we can define what an ordered pair is and what a Cartesian product is.

Definition 1.38 (Ordered Pair):

Let X and Y be non-empty sets. For any $x \in X$ and $y \in Y$, define the ordered pair

$$(x, y) := \{\{x\}, \{x, y\}\} \quad (1.20)$$

From this definition, we can prove the following theorem which justifies the name ‘ordered pair.’

Theorem 1.39 (Ordering in Ordered Pairs)

Let X and Y be non-empty sets. Then $(x, y) = (x', y')$ if and only if $x = x'$ and $y = y'$.

Proof:

[$\Rightarrow$]

We will prove this theorem in two cases:

Case 1: Assume $x = y$:

$$\begin{aligned} (x, y) &= \{\{x\}, \{x, y\}\} = \{\{x\}, \{x, x\}\} = \{\{x\}, \{x\}\} = \{\{x\}\} \\ \{x', y'\} &\in \{\{x'\}, \{x', y'\}\} = (x', y') = (x, y) = \{\{x\}\} \Rightarrow \{x', y'\} = \{x\} \end{aligned}$$

Thus by [axiom 1.23](#), $x' = y' = x$.

Case 2: Assume $x \neq y$:

Observe that $x' \neq y'$ if not then $\{x, y\} \in (x, y) = (x', y') = \{\{x'\}\} \Rightarrow x = y = x'$. $\nmid$

As $\{x'\} \in (x', y') = (x, y) = \{\{x\}, \{x, y\}\}$, we have $\{x'\} = \{x\}$ as $\{x'\} \neq \{x, y\}$ since $x \neq y$. Hence, $x' = x$.

Now, $\{x', y'\} \in (x', y') = (x, y) = \{\{x\}, \{x, y\}\}$, then $\{x', y'\} = \{x\}$ or $\{x', y'\} = \{x, y\}$. Observe $\{x', y'\} \neq \{x\}$ if not then $x' = y' = x$. $\nmid$

Hence $\{x', y'\} = \{x, y\}$ and thus $y' = y$ as $y' \neq x$ as $x' \neq y'$. Thus, $x' = x$ and $y' = y$.

[$\Leftarrow$]

Trivial.

Q.E.D.

Is collection of all ordered pairs a set? Answer to this question leads us to the definition of Cartesian product.

Theorem 1.40 (Existence of Cartesian Product)

Let X and Y be non-empty sets. Then the collection of all ordered pairs of elements of X and Y is a set *i.e.*

$$X \times Y := \{(x, y) \mid x \in X \wedge y \in Y\} \quad (1.21)$$

is a set.

Proof:

Suffice to show that for any $x \in X$ and $y \in Y$, (x, y) exists in “bigger” set that we know exists. Then by PRC, $X \times Y$ is a set.

Fix $x \in X$ and $y \in Y$. Then by [axiom 1.23](#), $\{x\}$ and $\{y\}$ are sets. Then by [axiom 1.23](#), $\{x, y\}$ is a set. So by [eq. \(1.6\)](#), $\{x\} \subseteq X \Rightarrow \{x\} \subseteq \bigcup\{X, Y\}$ and $\{x, y\} \subseteq \bigcup\{X, Y\}$. So by [axiom 1.36](#) we know $\mathcal{P}(\bigcup\{X, Y\})$ exists. Thus,

$$\{x\} \in \mathcal{P}\left(\bigcup\{X, Y\}\right) \quad \text{and} \quad \{x, y\} \in \mathcal{P}\left(\bigcup\{X, Y\}\right) \quad (1.22)$$

With this we can write $(x, y) = \{\{x\}, \{x, y\}\} \subseteq \mathcal{P}(\bigcup\{X, Y\})$. Hence, $(x, y) \in \mathcal{P}(\mathcal{P}(\bigcup\{X, Y\}))$. Thus, by PRC,

$$X \times Y = \left\{u \in \mathcal{P}\left(\mathcal{P}\left(\bigcup\{X, Y\}\right)\right) \mid \exists x \in X : \exists y \in Y : u = (x, y)\right\} \quad (1.23)$$

is a set.

Q.E.D.

Axiom 1.41 (Axiom of Infinity): *There exists a set that contains the empty set as an element and with every of its elements y it also contains the set $\{y\}$ as an element.*

$$\exists x : \emptyset \in x \wedge \forall y : (y \in x \Rightarrow \{y\} \in x). \quad (1.24)$$

Remark 1.42 (Smallest Infinite Set). Let x be the set guaranteed by [axiom 1.41](#). Then $\emptyset \in x \Rightarrow \{\emptyset\} \in x \Rightarrow \dots$,

$$x = \{\emptyset, \{\emptyset\}, \{\{\emptyset\}\}, \{\{\{\emptyset\}\}\} \dots\} \quad (1.25)$$

Notationally, writing down the set x is cumbersome as there are too many braces. So we denote each element of x by a natural number as follows:

$$0 := \emptyset, \quad 1 := \{\emptyset\}, \quad 2 := \{\{\emptyset\}\}, \quad 3 := \{\{\{\emptyset\}\}\}, \quad \dots \quad (1.26)$$

Thus, the set x can be written as

$$x = \{0, 1, 2, 3, \dots\} \quad (1.27)$$

Corollary 1.43 (Existence of Natural Numbers)

$\mathbb{N}$ is a set.

Remark 1.44 (Set of Real Numbers). As a set, $\mathbb{R} = \mathcal{P}(\mathbb{N})$.

Axiom 1.45 (Axiom of Choice): *Let x be a set whose elements are non-empty and mutually disjoint sets, then there exists a set y which contains precisely one element from each element of x .*

$$\forall x : P(x) \Rightarrow \exists y : \forall w \in x : \exists! z \in w : z \in y \quad (1.28)$$

where, $P(x) :\Leftrightarrow (\exists u : u \in x) \wedge (\forall u \in x : \forall v \in x : u \neq v \Rightarrow \bigcap\{u, v\} = \emptyset)$.

The set y is called “dark” set. Say we have a set x that contains pair of shoes, then we can algorithmically choose the left shoe from each pair of shoes to form the set y . But if we have a set x that contains a pair of socks, then we can’t algorithmically choose one sock from each pair of socks to form the set y . This is where the axiom of choice comes into play.

The axiom of choice is independent of the other 8 axioms, which means that one could have set theory with or without the axiom of choice. However, standard mathematics uses the axiom of choice and hence so will we.

There is a number of theorems that can only be proved by using the axiom of choice. Amongst these we have:

- Proof that every vector space has a basis needs the axiom of choice.
- Proof that there exists a complete system of representatives for the equivalence classes of a set under an equivalence relation needs the axiom of choice.

Axiom 1.46 (Axiom of foundation): *Every non-empty set x contains an element y that has none of its elements in common with x .*

$$\forall x : (\exists y : y \in x) \Rightarrow \exists y \in x : \bigcap\{x, y\} = \emptyset \quad (1.29)$$

Corollary 1.47

There is no set that contains itself as an element.

$$x \in x \quad \text{is false for all sets } x \quad (1.30)$$

The totality of these 9 axioms is called ZFC (Zermelo-Fraenkel with the axiom of choice) set theory.

LECTURE 3

§1.6 Classifications of Sets

A recurrent theme in mathematics is the study/classification of spaces by means of structure-preserving maps between those spaces.

A space is usually meant to be some set equipped with some additional structure. In this context, we are interested in the classification of sets which is a space without any additional structure.

Definition 1.48 (Map):

A map $\phi : A \rightarrow B$ is a relation such that for every $a \in A$, there exists exactly one $b \in B$ such that $\phi(a, b)$.

Notation: $\phi : A \rightarrow B$ or $A \xrightarrow{\phi} B$. Since there is a unique b for every a , we have a notational abuse as $a \mapsto b =: \phi(a)$,

Some basic terminologies:

- A is called the *domain* of ϕ .
- B is called the *co-domain* of ϕ .
- The set $\phi(A) \equiv \text{im}_\phi(A) := \{\phi(a) \in B \mid a \in A\}$ is called the *image* of A under ϕ .

Definition 1.49:

Let A and B be sets. A map $\phi : A \rightarrow B$ is called

- Surjective (or onto) if and only if $\text{im}_\phi(A) = B$.
- Injective (or one-to-one) if and only if for all $a_1, a_2 \in A$, $\phi(a_1) = \phi(a_2) \Rightarrow a_1 = a_2$.
- Bijective if and only if it is both surjective and injective.

Definition 1.50 (Iso-morphism):

Let A and B be sets. We say that A and B are (set-theoretic) isomorphic if there exists a bijection $\phi : A \rightarrow B$. In this case, we write $A \cong_{\text{set}} B$.

For two sets to be isomorphic, we only need to prove the existence of a bijection between them. We do not need to construct the bijection explicitly.

Remark 1.51 (Number of isomorphisms). If there is any bijection between two sets, then generically there are many bijections between them.

Intuition: pair the elements of the two sets in any way you like.

In case of set theory, the structure-preserving maps are bijections.

Definition 1.52:

Let A be a set. The set A is

- Infinite if there exists a proper subset $B \subsetneq A$ such that $B \cong_{\text{set}} A$.
 - * A is called countable infinite if and only if $A \cong_{\text{set}} \mathbb{N}$.
 - * A is called uncountable infinite if and only if it is infinite but not countably infinite.
- Finite if it is not infinite.

In this case, we have $A \cong_{\text{set}} \{1, 2, 3, \dots, N\}$ for some $N \in \mathbb{N}$. Then we write $|A| = N$ and call N the cardinality of A .

Composition of maps: Given two maps $\phi : A \rightarrow B$ and $\psi : B \rightarrow C$, we can construct a new map known as the composition of ϕ and ψ , denoted by $\psi \circ \phi$, defined as

$$\begin{aligned} \psi \circ \phi : A &\rightarrow C \\ a &\mapsto \psi(\phi(a)) \end{aligned}$$

Diagrammatically, we can represent the composition of maps as follows:

$$\begin{array}{ccc} & B & \\ \phi \nearrow & & \searrow \psi \\ A & \xrightarrow{\psi \circ \phi} & C \end{array} \quad [\text{Commutes}]$$

And the composition of maps is associative, i.e., $\xi \circ (\psi \circ \phi) = (\xi \circ \psi) \circ \phi$.

Identity map: For any set A , there exists a unique map $\text{id}_A : A \rightarrow A$ such that

$$\forall a \in A : \quad a \xrightarrow{\text{id}_A} a$$

Definition 1.53 (Inverse of a map):

Let $\phi : A \rightarrow B$ be a bijection. Then the inverse of ϕ , is the map $\phi^{-1} : B \rightarrow A$ defined uniquely by

$$\begin{array}{ccc} \text{id}_A \hookrightarrow A & \begin{array}{c} \xrightarrow{\phi} \\ \xleftarrow{\phi^{-1}} \end{array} & B \hookrightarrow \text{id}_B \end{array} \quad \begin{aligned} \phi^{-1} \circ \phi &= \text{id}_A \\ \phi \circ \phi^{-1} &= \text{id}_B \end{aligned} \quad (1.31)$$

Definition 1.54 (Pre-image):

Let $\phi : A \rightarrow B$ be a map and let $B' \subseteq B$. Then define the set

$$\text{preim}_\phi(B') := \{a \in A \mid \phi(a) \in B'\}. \quad (1.32)$$

$\text{preim}_\phi(B')$ is called the pre-image of B' under ϕ .

§1.7 Equivalence Relations

Definition 1.55 (Equivalence relation):

Let M be a set and let $\sim$ be a relation such that:

- Reflexivity: $\forall m \in M : m \sim m$.
- Symmetry: $\forall m, n \in M : m \sim n \Rightarrow n \sim m$.
- Transitivity: $\forall m, n, p \in M : m \sim n \wedge n \sim p \Rightarrow m \sim p$.

Then $\sim$ is called an equivalence relation on M .

Example 1.56

Consider the following wordy examples.

- (a) $p \sim q :\Leftrightarrow p$ is of the same opinion as q . This relation is reflexive, symmetric and transitive. Hence, it is an equivalence relation.
- (b) $p \sim q :\Leftrightarrow p$ is a sibling of q . This relation is symmetric and transitive but not reflexive and hence, it is not an equivalence relation.
- (c) $p \sim q :\Leftrightarrow p$ is taller q . This relation is transitive, but neither reflexive nor symmetric and hence, it is not an equivalence relation.
- (d) $p \sim q :\Leftrightarrow p$ is in love with q . This relation is generally not reflexive. People don't like themselves very much. It is certainly not normally symmetric, which is the basis of much drama in literature. It is also not transitive, except in some French films.

Definition 1.57 (Equivalence class):

Let M be a set and let $\sim$ be an equivalence relation on M . For any $m \in M$, the set

$$[m] := \{n \in M \mid n \sim m\} \quad (1.33)$$

is called the equivalence class of m under $\sim$.

Two key properties of equivalence classes are:

Proposition 1.58

Let M be a set and let $\sim$ be an equivalence relation on M . Then:

- (a) $a \in [m] \Rightarrow [a] = [m]$.

In other words, this means that “any element of an equivalence class can act as a representative of the equivalence class”.

- (b) Let $m, n \in M$, then either $[m] = [n]$ or $\bigcap \{[m], [n]\} = \emptyset$.

Proof: (a) Let $a \in [m]$. Then by definition, $a \sim m$. Let $b \in [a]$. Then $b \sim a$. By transitivity, $b \sim m$. Hence, $b \in [m]$. Therefore, $[a] \subseteq [m]$. Now, let $b \in [m]$. Then $b \sim m$. By transitivity, $b \sim a$. Hence, $b \in [a]$. Therefore, $[m] \subseteq [a]$. Hence, $[a] = [m]$.

- (b) For two elements, either $m \sim n$ or $m \not\sim n$. If $m \sim n$, then $n \in [m]$ and hence, $[m] = [n]$. If $m \not\sim n$, then $\forall p \in M : (p \in [m]) \vee (p \in [n])$. Hence, $\bigcap \{[m], [n]\} = \emptyset$.

Q.E.D.

Definition 1.59 (Quotient set):

Let M be a set and let $\sim$ be an equivalence relation on M . Then define the quotient set

$$M/\sim := \{[m] \in \mathcal{P}(M) \mid m \in M\} \quad (1.34)$$

Notation: $M/\sim$ is read as “ M modulo $\sim$ ”. The elements of $M/\sim$ are the equivalence classes of M under $\sim$.

Intuition: The quotient set is the set of all equivalence classes of M under $\sim$.

Remark 1.60 (Set of representatives). Due to the [Axiom of Choice](#), there exists a complete system of representatives for $\sim$ i.e. a set R such that $\forall m \in M : \exists! r \in R : m \sim r$. Then we can write

$$R \cong_{\text{set}} M/\sim. \quad (1.35)$$

This essentially means that we can choose a representative from each equivalence class and put them in a set namely R . Then this provides us a natural bijection between the set of representatives R and the quotient set $M/\sim$.

Remark 1.61. Care must be taken while defining maps whose domain or co-domain is a quotient set and if one uses representatives to define the map. The map must be well-defined, *i.e.* the map must not depend on the choice of representatives.

Example 1.62

Let $M = \mathbb{Z}$ and let $\sim$ be a relation defined by $m \sim n :\Leftrightarrow m - n \in 2\mathbb{Z}$. Then $\sim$ is an equivalence relation. The equivalence classes are

$$\begin{aligned} [0] &= [2] = [4] = \dots = [-2] = [-4] = \dots = 2\mathbb{Z}, \\ [1] &= [3] = [5] = \dots = [-1] = [-3] = \dots = 2\mathbb{Z} + 1. \end{aligned}$$

Therefore, the quotient set is

$$\mathbb{Z}/\sim = \{[0], [1]\}.$$

Idea: on $\mathbb{Z}$ we have addition $+: \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$. We wish to inherit an addition on $\mathbb{Z}/\sim$ from the addition on $\mathbb{Z}$ *i.e.* $\oplus: \mathbb{Z}/\sim \times \mathbb{Z}/\sim \rightarrow \mathbb{Z}/\sim$. We can define this map as follows:

$$[a] \oplus [b] \mapsto [a + b]$$

Care needs to be taken precisely because this could be inconsistent (not well-defined). Check whether choice of representatives matters:

Let $a', b' \in \mathbb{Z}$ such that $[a] = [a']$ and $[b] = [b']$. Then we need to check whether $[a'] \oplus [b'] = [a] \oplus [b]$. So by assumptions we have $a \sim a' \Rightarrow a - a' = 2n \in 2\mathbb{Z}$ and $b \sim b' \Rightarrow b - b' = 2m \in 2\mathbb{Z}$. Then

$$\begin{aligned} [a'] \oplus [b'] &\stackrel{\text{def}}{=} [a' + b'] = [a - 2n + b - 2m] \\ &= [a + b - 2(n + m)] \\ &= [a + b] \stackrel{\text{def}}{=} [a] \oplus [b]. \end{aligned}$$

Therefore, the addition is well-defined.

§1.8 Construction of $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$, $\mathbb{R}$

We are only going through the outline of the construction of these sets.

§1.8.1 Natural Numbers $\mathbb{N}$

Recall, from the [Axiom of Infinity](#), that there exists a set $\mathbb{N}$ such that

$$\begin{aligned} \mathbb{N} &= \{0, 1, 2, 3, \dots\} \\ 0 &:= \emptyset, \quad 1 := \{\emptyset\}, \quad 2 := \{\{\emptyset\}\}, \quad 3 := \{\{\{\emptyset\}\}\}, \quad \dots \end{aligned} \tag{1.36}$$

We wish to establish addition on $\mathbb{N}$. For that we need to define the *successor map* as

$$\begin{aligned} S: \mathbb{N} &\rightarrow \mathbb{N} \\ n &\mapsto \{n\} \end{aligned} \tag{1.37}$$

This works as, $S(2) = S(\{\{\emptyset\}\}) = \{\{\{\emptyset\}\}\} = 3$.

We need to define the *predecessor map* as

$$\begin{aligned} P: \mathbb{N}^* &\rightarrow \mathbb{N} \\ n &\mapsto m \quad \text{such that} \quad m \in n \end{aligned} \tag{1.38}$$

Here, $\mathbb{N}^* = \mathbb{N} \setminus \{0\}$. In this case, $P(2) = P(\{\{\emptyset\}\}) = \{\emptyset\} = 1$ as there is only one element in 2. Now, with this we can define $(n \in \mathbb{N})^{\text{th}}$ power of S :

$$\begin{aligned} S^n &= S \circ S^{P(n)}, \quad \text{if } n \in \mathbb{N}^* \\ S^0 &= \text{id}_{\mathbb{N}} \end{aligned} \quad (1.39)$$

At this point, we can define addition on $\mathbb{N}$ as

Definition 1.63 (Addition on $\mathbb{N}$):

$$\begin{aligned} + : \mathbb{N} \times \mathbb{N} &\rightarrow \mathbb{N} \\ (m, n) &\mapsto m + n := S^n(m) \end{aligned} \quad (1.40)$$

Example 1.64 ($2 + 1 = 3$)

With this definition, we can verify our intuition of addition on $\mathbb{N}$. For example, $2 + 1 = 3$.

$$2 + 1 = S^1(2) = S(S^0(2)) = S(\text{id}_{\mathbb{N}}(2)) = S(2) = 3$$

And with this, we should be able to check whether the addition is commutative or not.

$$1 + 2 = S^2(1) = S(S^1(1)) = S(S(S^0(1))) = S(S(1)) = S(2) = 3$$

This same calculation generalizes to any two natural numbers and hence, the addition is commutative.

Remark 1.65 (Identity element of addition). We can see that 0 is the identity element of addition on $\mathbb{N}$. This is because

$$\begin{aligned} \forall n \in \mathbb{N} : 0 + n &= S^n(0) = \dots = n, \\ \forall n \in \mathbb{N} : n + 0 &= S^0(n) = n. \end{aligned}$$

What about the inverse of addition? We can see that there is no inverse of addition on $\mathbb{N}$. This is because, for any $n \in \mathbb{N}$, there is no $m \in \mathbb{N}$ such that $n + m = 0$. In language of group theory, $\mathbb{N}$ is not a commutative group under addition.

Now this hurdle motivates us to define the set of integers $\mathbb{Z}$.

§1.8.2 Integers $\mathbb{Z}$

Let's start with defining an equivalence relation $\sim_{\mathbb{Z}}$ on $\mathbb{N} \times \mathbb{N}$ as

$$(m, n) \sim_{\mathbb{Z}} (p, q) :\Leftrightarrow m + q = n + p \quad (1.41)$$

Check that this is an equivalence relation.

1. *Reflexivity:* $(m, n) \sim_{\mathbb{Z}} (m, n) \Leftrightarrow m + n = n + m$.
2. *Symmetry:* $(m, n) \sim_{\mathbb{Z}} (p, q) \Leftrightarrow m + q = n + p \Leftrightarrow n + p = m + q \Leftrightarrow (p, q) \sim_{\mathbb{Z}} (m, n)$.
3. *Transitivity:* $(m, n) \sim_{\mathbb{Z}} (p, q) \wedge (p, q) \sim_{\mathbb{Z}} (r, s) \Rightarrow m + q = n + p \wedge p + s = q + r \Rightarrow m + q + p + s = n + p + q + r \Rightarrow m + s = n + r \Rightarrow (m, n) \sim_{\mathbb{Z}} (r, s)$.

Thus, $\sim_{\mathbb{Z}}$ is an equivalence relation on $\mathbb{N} \times \mathbb{N}$.

Idea: in this case (m, n) corresponds to $m - n$. So, we can define the set of integers as

$$\mathbb{Z} := (\mathbb{N} \times \mathbb{N}) / \sim_{\mathbb{Z}} \quad (1.42)$$

Intuitively, we want $\mathbb{N} \subseteq \mathbb{Z}$ but at this stage this is nonsense as both the sets have different structures. We can resolve this by embedding $\mathbb{N}$ into $\mathbb{Z}$ with the help of an *inclusion map*:

$$\begin{aligned}\iota : \mathbb{N} &\hookrightarrow \mathbb{Z} \\ n &\mapsto [(n, 0)]\end{aligned}\tag{1.43}$$

With this new definition of $\mathbb{N}$ in $\mathbb{Z}$, we can say that $\mathbb{N} \subseteq \mathbb{Z}$. Now, in similar fashion we can define negative integers as well.

Definition 1.66 (Negative integers):

Let $n \in \mathbb{N}$, then we define the negative integer as

$$-n := [(0, n)]\tag{1.44}$$

With this, let's define addition on $\mathbb{Z}$.

Definition 1.67 (Addition on $\mathbb{Z}$):

$$\begin{aligned}+_{\mathbb{Z}} : \mathbb{Z} \times \mathbb{Z} &\rightarrow \mathbb{Z} \\ [(m, n)] +_{\mathbb{Z}} [(p, q)] &:= [(m + p, n + q)]\end{aligned}\tag{1.45}$$

It is easy to see that this addition is well-defined.

Example 1.68 ($2 + (-3) = -1$)

Using our definition of $2 = [(2, 0)]$ and $-3 = [(0, 3)]$, we can calculate $2 +_{\mathbb{Z}} (-3)$ as:

$$2 +_{\mathbb{Z}} (-3) = [(2, 0)] +_{\mathbb{Z}} [(0, 3)] = [(2 + 0, 0 + 3)] = [(2, 3)] = [(0, 1)] = -1$$

We will now assume that we have constructed the multiplication on $\mathbb{Z}^6$, and then we can define the set of rational numbers $\mathbb{Q}$.

§1.8.3 Rational Numbers $\mathbb{Q}$

With similar construction as above, we define an equivalence relation $\sim_{\mathbb{Q}}$ on $\mathbb{Z} \times \mathbb{Z}^7$ as

$$(m, n) \sim_{\mathbb{Q}} (p, q) :\Leftrightarrow mq = np\tag{1.46}$$

Check that this is an equivalence relation.

1. *Reflexivity*: $(m, n) \sim_{\mathbb{Q}} (m, n) \Leftrightarrow mn = nm$.
2. *Symmetry*: $(m, n) \sim_{\mathbb{Q}} (p, q) \Leftrightarrow mq = np \Leftrightarrow np = mq \Leftrightarrow (p, q) \sim_{\mathbb{Q}} (m, n)$.
3. *Transitivity*: $(m, n) \sim_{\mathbb{Q}} (p, q) \wedge (p, q) \sim_{\mathbb{Q}} (r, s) \Rightarrow mq = np \wedge ps = qr \Rightarrow mqps = npqr \Rightarrow ms = nr \Rightarrow (m, n) \sim_{\mathbb{Q}} (r, s)$.

Thus, $\sim_{\mathbb{Q}}$ is an equivalence relation on $\mathbb{Z} \times \mathbb{Z}^*$.

Idea: in this case (m, n) corresponds to $\frac{m}{n}$. So, we can define the set of rational numbers as

$$\mathbb{Q} := (\mathbb{Z} \times \mathbb{Z}^*) / \sim_{\mathbb{Q}}\tag{1.47}$$

Intuitively, we want $\mathbb{Z} \subseteq \mathbb{Q}$ but at this stage this is nonsense as both the sets have different structures. We can resolve this by embedding $\mathbb{Z}$ into $\mathbb{Q}$ with the help of an *inclusion map*:

$$\begin{aligned}\iota : \mathbb{Z} &\hookrightarrow \mathbb{Q} \\ n &\mapsto [(n, 1)]\end{aligned}\tag{1.48}$$

With this new definition of $\mathbb{Z}$ in $\mathbb{Q}$, we can say that $\mathbb{Z} \subseteq \mathbb{Q}$.

⁶Multiplication $\cdot_{\mathbb{Z}} : \mathbb{Z} \times \mathbb{Z} \rightarrow \mathbb{Z}$, defined as $[(m, n)] \cdot_{\mathbb{Z}} [(p, q)] = [(mp + nq, mq + np)]$.

⁷ $\mathbb{Z}^* := \mathbb{Z} \setminus \{0\}$

Definition 1.69 (Addition on $\mathbb{Q}$):

$$\begin{aligned} +_{\mathbb{Q}} : \mathbb{Q} \times \mathbb{Q} &\rightarrow \mathbb{Q} \\ [(m, n)] +_{\mathbb{Q}} [(p, q)] &:= [(mq + np, nq)] \end{aligned} \quad (1.49)$$

Similarly, we can define the multiplication on $\mathbb{Q}$

Definition 1.70 (Multiplication on $\mathbb{Q}$):

$$\begin{aligned} \cdot_{\mathbb{Q}} : \mathbb{Q} \times \mathbb{Q} &\rightarrow \mathbb{Q} \\ [(m, n)] \cdot_{\mathbb{Q}} [(p, q)] &:= [(mp, nq)] \end{aligned} \quad (1.51)$$

§1.8.4 Real Numbers $\mathbb{R}$

There are many ways to construct the reals from the rationals. One is to define a set $\mathcal{A}$ of *almost homomorphism* on $\mathbb{Z}$ and hence define:

$$\mathbb{R} := \mathcal{A} / \sim, \quad (1.51)$$

where $\sim$ is a “suitable” equivalence relation on $\mathcal{A}$.

We will not go into the details of this construction. We will just assume that the reals are constructed, and we have all the usual operations defined on them. Now define the set of positive reals as

$$\mathbb{R}^+ := \{r \in \mathbb{R} \mid r > 0\} \quad (1.52)$$

With this construction, we can define $\mathbb{R}^d$ for any $d \in \mathbb{N}$ as

$$\mathbb{R}^d := \underbrace{\mathbb{R} \times \mathbb{R} \times \cdots \times \mathbb{R}}_{d \text{ times}} \quad (1.53)$$

For $x \in \mathbb{R}^d$, we write $x = (x^1, x^2, \dots, x^d)$ where $x^i \in \mathbb{R}$. For further convenience, we need to define the *norm* on $\mathbb{R}^d$.

Definition 1.71 (norm):

Let $x = (x^1, x^2, \dots, x^d) \in \mathbb{R}^d$. Then the norm of x is defined as

$$\begin{aligned} \|\cdot\| : \mathbb{R}^d &\rightarrow \mathbb{R}^+ \\ x &\mapsto \sqrt{\sum_{i=1}^d (x^i)^2} \end{aligned} \quad (1.54)$$

We can extend this definition of norm as

$$x \mapsto \sqrt[2n]{\sum_{i=1}^d (x^i)^{2n}} \quad \text{for any } n \in \mathbb{N} \quad (1.55)$$

In order to avoid confusion, we will denote the norm of x as $\|x\|_2$ for $n = 1$ and in general as $\|x\|_{2n}$ for $n \in \mathbb{N}$. For convenience, we will use $n = 1$ most of the time.

With this, one can define a ball in $\mathbb{R}^d$ as

Definition 1.72 (Ball in $\mathbb{R}^d$):

Let $x \in \mathbb{R}^d$ and $r \in \mathbb{R}^+$. Then define the set $B_r(x)$ as

$$B_r(x) := \left\{ y \in \mathbb{R}^d \mid \sqrt{\sum_{i=1}^d (x^i - y^i)^2} < r \right\} \quad (1.56)$$

is called the ball of radius r centered at x .

Later on, we will see that we denote these sets as “open balls”.

2 Topological Spaces

LECTURE 4

We will now discuss topological spaces based on our previous development of set theory. As we will see, a topology on a set provides the weakest structure in order to define the two very important notions of convergence of sequences to points in a set, and of continuity of maps between two sets.

§2.1 Topological Spaces

Definition 2.1 (Topology & Topological Space):

Let M be a set. Then a choice of subsets $\mathcal{O} \subseteq \mathcal{P}(M)$ is called a topology on M if

- (i) $\emptyset \in \mathcal{O}$ and $M \in \mathcal{O}$,
- (ii) $U, V \in \mathcal{O} \Rightarrow \bigcap \{U, V\} \in \mathcal{O}$,
- (iii) $C \subseteq \mathcal{O} \Rightarrow \bigcup C \in \mathcal{O}$.

The pair $(M, \mathcal{O})$ is then called a topological space.

Remark 2.2 (Open & Closed Sets). Let $(M, \mathcal{O})$ be a topological space. Then

- (a) A set $U \in \mathcal{O}$ is called an *open set* in $(M, \mathcal{O})$.
- (b) A set $C \subseteq M$ is called a *closed set* in $(M, \mathcal{O})$ if $M \setminus C \in \mathcal{O}$ i.e. $M \setminus C$ is open in $(M, \mathcal{O})$.

Remark 2.3 (choice of topology). Unless $|M| = 1$, there are different topologies $\mathcal{O}$ one can choose on one and the same set M . Based on the choice of topology $\mathcal{O}$, the notion of convergence of sequences and continuity of maps will change.

For finite set, we can count the number of topologies on it.

$ M $	Number of Topologies
1	1
2	4
3	29
4	355
5	6,942
6	209,527
7	9,535,241
$\vdots$	$\vdots$

Example 2.4 (Basic Topologies)

Let M be any set, then we can define following “trivial” topologies on M :

- (a) $\mathcal{O} = \{\emptyset, M\}$. It is easy to see that, $\mathcal{O}$ is a topology on M . This is called the *chaotic topology*.
- (b) $\mathcal{O} = \mathcal{P}(M)$. It is easy to see that, $\mathcal{O}$ is a topology on M . This is called the *discrete topology*.

For a less abstract example, consider $M = \{1, 2, 3\}$, then we can define 29 different topologies on M . For example, $\mathcal{O} = \{\emptyset, \{1\}, \{2\}, \{1, 2\}, \{1, 2, 3\}\}$ is a topology on M .

Example 2.5 (Standard Topology on $\mathbb{R}^d$)

This is the most important and heavily used topology in mathematics and physics.

Consider $M = \mathbb{R}^d$, where $d \in \mathbb{N}$. Then the *standard topology* on $\mathcal{O}_{\text{std.}}$ is constructed as follows:

$$U \in \mathcal{O}_{\text{std.}} \Leftrightarrow \forall p \in U : \exists r \in \mathbb{R}^+ : B_r(p) \subseteq U$$

where $B_r(p) = \{x \in \mathbb{R}^d \mid \|x - p\| < r\}$ is the open ball of radius r around p . In more involved terms, we can write the definition of the standard topology on $\mathbb{R}^d$ as

$$\mathcal{O}_{\text{std.}} = \left\{ U \in \mathcal{P}(\mathbb{R}^d) \mid \forall p \in U : \exists r \in \mathbb{R}^+ : B_r(p) \subseteq U \right\}$$

Then by [theorem 1.33](#), $\mathcal{O}_{\text{std.}}$ is a set. Now we have to prove that $\mathcal{O}_{\text{std.}}$ is indeed a topology on $\mathbb{R}^d$.

(i) First, we need to check whether $\emptyset \in \mathcal{O}_{\text{std.}}$, *i.e.* whether:

$$\forall p \in \emptyset : \exists r \in \mathbb{R}^+ : B_r(p) \subseteq \emptyset$$

is true. This proposition is of the form $\forall p \in \emptyset : Q(p)$, which was defined as being equivalent to:

$$\forall p : p \in \emptyset \Rightarrow Q(p).$$

However, since $p \in \emptyset$ is false, the implication is true independent of p . Hence, the initial proposition is true and thus $\emptyset \in \mathcal{O}_{\text{std.}}$.

Second, by definition, we have $B_r(x) \subseteq \mathbb{R}^d$ independent of x and r , hence:

$$\forall p \in \mathbb{R}^d : \exists r \in \mathbb{R}^+ : B_r(p) \subseteq \mathbb{R}^d$$

is true and thus $\mathbb{R}^d \in \mathcal{O}_{\text{std.}}$.

(ii) Let $U, V \in \mathcal{O}_{\text{std.}}$, then we have to show that $U \cap V \in \mathcal{O}_{\text{std.}}$. Let $p \in U \cap V$, then $p \in U$ and $p \in V$. Since $U, V \in \mathcal{O}_{\text{std.}}$, we have:

$$\begin{aligned} \exists r_1 \in \mathbb{R}^+ : B_{r_1}(p) &\subseteq U, \\ \exists r_2 \in \mathbb{R}^+ : B_{r_2}(p) &\subseteq V. \end{aligned}$$

Now, let $r = \min\{r_1, r_2\}$, then we have $B_r(p) \subseteq B_{r_1}(p) \subseteq U$ and $B_r(p) \subseteq B_{r_2}(p) \subseteq V$. Hence, $B_r(p) \subseteq U \cap V$ and thus $U \cap V \in \mathcal{O}_{\text{std.}}$.

(iii) Let $C \subseteq \mathcal{O}_{\text{std.}}$, then we have to show that $\bigcup C \in \mathcal{O}_{\text{std.}}$. Let $p \in \bigcup C$, then $p \in U$ for some $U \in C$. Since $U \in \mathcal{O}_{\text{std.}}$, we have:

$$\exists r \in \mathbb{R}^+ : B_r(p) \subseteq U \subseteq \bigcup C.$$

Hence, $\bigcup C \in \mathcal{O}_{\text{std.}}$.

Thus, $\mathcal{O}_{\text{std.}}$ is a topology on $\mathbb{R}^d$.

Definition 2.6 (Comparison of Topologies):

Let M be a set and $\mathcal{O}_1, \mathcal{O}_2$ be two topologies on M . If $\mathcal{O}_1 \subseteq \mathcal{O}_2$, then we say that $\mathcal{O}_2$ is finer (or stronger) topology than $\mathcal{O}_1$, and $\mathcal{O}_1$ is coarser (or weaker) topology than $\mathcal{O}_2$.

From this definition, we can see that the chaotic topology is the coarsest topology on a set, and the discrete topology is the finest topology on a set.

§2.2 Construction of new Topologies from given topologies

§2.2.1 Induced Topology

Theorem 2.7 (Induced Topology)

Let $(M, \mathcal{O})$ be a topological space and $N \subset M^1$. Then the following set

$$\mathcal{O}|_N = \{U \cap N \mid U \in \mathcal{O}\} \subseteq \mathcal{P}(N) \quad (2.1)$$

is a topology on N , called the *induced topology* on N .

Proof:

We have to show that $\mathcal{O}|_N$ is a topology on N .

(i) First, as $\emptyset \in \mathcal{O}$, we have $\emptyset \cap N = \emptyset \in \mathcal{O}|_N$.

Second, since $M \in \mathcal{O}$, we have $M \cap N = N \in \mathcal{O}|_N$.

(ii) $U, V \in \mathcal{O}|_N \stackrel{?}{\Rightarrow} U \cap V \in \mathcal{O}|_N$. From (2.1), $\exists U_1, V_1 \in \mathcal{O} : U = U_1 \cap N \wedge V = V_1 \cap N$. Then we have:

$$U \cap V = (U_1 \cap N) \cap (V_1 \cap N) = (U_1 \cap V_1) \cap N \in \mathcal{O}|_N$$

as $U_1 \cap V_1 \in \mathcal{O}$.

(iii) $C \subseteq \mathcal{O}|_N \stackrel{?}{\Rightarrow} \bigcup C \in \mathcal{O}|_N$. Since $C \subseteq \mathcal{O}|_N$, we have $C = \{U_i \cap N \mid U_i \in \mathcal{O}\}$. Then we have:

$$\bigcup C = \bigcup \{U_i \cap N \mid U_i \in \mathcal{O}\} = \left\{ \bigcup U_i \right\} \cap N \in \mathcal{O}|_N.$$

Thus, $\mathcal{O}|_N$ is a topology on N .

Q.E.D.

Example 2.8

Consider $(\mathbb{R}, \mathcal{O}_{\text{std.}})$ and $N = [0, 1] := \{x \in \mathbb{R} \mid 0 \leq x \leq 1\}$. Then from theorem 2.7, $(N, \mathcal{O}_{\text{std.}}|_N)$ is a topological space.

It is easy to see that $(0, 1] \notin \mathcal{O}_{\text{std.}}$ so $(0, 1]$ is not open in $(\mathbb{R}, \mathcal{O}_{\text{std.}})$, but $(0, 1] = (0, 2) \cap N \in \mathcal{O}_{\text{std.}}|_N$. Hence, $(0, 1]$ is an open set in $(N, \mathcal{O}_{\text{std.}}|_N)$.

At this point we can talk about openness and closedness of sets in a topological space.

Remark 2.9. Let $(M, \mathcal{O})$ be a topological space and $N \subset M$. Then N is either

- open, or
- closed, or
- open and closed, or
- open and not closed, or
- not open and closed, or
- not open and not closed.

For this very unintuitive idea of a set to be open and closed at the same time, see the following example.

Example 2.10

For any topological space $(M, \mathcal{O})$, we know by definition of topology that $\emptyset, M \in \mathcal{O}$. Hence, $\emptyset$ and M are open in $(M, \mathcal{O})$.

- Now, let $N = M$, then $M \setminus N = \emptyset \in \mathcal{O}$. Hence, M is closed in $(M, \mathcal{O})$.

¹We will use either $N \subset M$ or $N \subsetneq M$ or $N \subsetneqq M$ to denote that N is a proper subset of M .

- Let $N = \emptyset$, then $M \setminus N = M \in \mathcal{O}$. Hence, $\emptyset$ is closed in $(M, \mathcal{O})$.

Thus, $\emptyset$ and M are both open and closed in $(M, \mathcal{O})$.

§2.2.2 Quotient Topology

Definition 2.11 (Quotient Topology):

Let $(M, \mathcal{O})$ be a topological space and $\sim$ be an equivalence relation on M . Then the quotient set $M/\sim$ can be equipped with the quotient topology $\mathcal{O}_{M/\sim}$ defined as follows:

$$\mathcal{O}_{M/\sim} = \left\{ U \subseteq M/\sim \mid \bigcup U = \bigcup_{[u] \in U} [u] \in \mathcal{O} \right\} \quad (2.2)$$

An equivalent definition of the quotient topology is as follows: Let $\pi : M \rightarrow M/\sim$ be the map defined as $\pi(m) = [m]$. Then the quotient topology is defined as

$$\mathcal{O}_{M/\sim} = \left\{ U \subseteq M/\sim \mid \text{preim}_\pi(U) \in \mathcal{O} \right\} \quad (2.3)$$

Example 2.12 (Topology on 1-sphere)

Consider $M = \mathbb{R}^2$. The *circle* (or *1-sphere*) is defined as a set $S^1 := \{(x, y) \in \mathbb{R}^2 \mid x^2 + y^2 = 1\}$. We have two ways to construct a new topological space:

- (a) Induced Topology: As $S^1 \subset \mathbb{R}^2$, we can construct a topology $\mathcal{O}$ on S^1 by [theorem 2.7](#) using $(\mathbb{R}^2, \mathcal{O}_{\text{std.}})$ topological space. This is the *standard topology* on S^1 .




Figure 2.1: 1-sphere induced topology

From [fig. 2.1](#), we can see that U_i 's are open in $\mathbb{R}^2$ and $U_i \cap S^1$ are open in S^1 . See, $U_3 \in \mathcal{O}_{\text{std.}} \Rightarrow \emptyset \in \mathcal{O}$ and $U_1 \in \mathcal{O}_{\text{std.}} \Rightarrow S_1 \in \mathcal{O}$. Also, the set S_2 is not open in S^1 as $\forall U \in \mathcal{P}(\mathbb{R}) : U \cap S^1 \neq S_2$. Hence, $S_2 \notin \mathcal{O}$.

- (b) Quotient Topology: Let $\sim$ be the equivalence relation on $\mathbb{R}$ defined by:

$$x \sim y \Leftrightarrow \exists n \in \mathbb{Z} : x = y + 2\pi n.$$

Then the circle can be defined as the set $S^1 := \mathbb{R}/\sim$ equipped with the quotient topology.

§2.2.3 Product Topology

Definition 2.13 (Product Topology):

Let $(A, \mathcal{O}_A)$ and $(B, \mathcal{O}_B)$ be two topological spaces. Then the set $\mathcal{O}_{A \times B}$ defined implicitly as


$$U \in \mathcal{O}_{A \times B} \Leftrightarrow \forall (a, b) \in U : \exists U_A \in \mathcal{O}_A, U_B \in \mathcal{O}_B : a \in U_A \wedge b \in U_B \wedge U_A \times U_B \subseteq U \quad (2.4)$$

is a topology on $A \times B$, called the product topology on $A \times B$.

Here we used the existence of U_A and U_B such that it contains a and b respectively. And we will use this kind of argument throughout our discussion.

Let $(M, \mathcal{O})$ be a topological space. Let $a \in M$ be a point. Then we say that a set $U \in \mathcal{O}$ is an *open neighbourhood* of a if $a \in U$.

Intuitively, the definition can be understood with the help of following diagram:



Remark 2.14 (Extending product topology to finite cartesian product). Let $A_1, A_2, \dots, A_N$ be topological spaces. Then in same spirit, we can define the product topology

$$\mathcal{O}_{A_1 \times A_2 \times \dots \times A_N}$$

With this, we can check that the standard topology on $\mathbb{R}^d$ is a product topology with open ball defined using $\|\cdot\|_\infty$

$$\mathcal{O}_{\text{std. } \mathbb{R}^d} = \mathcal{O}_{\underbrace{\mathbb{R} \times \mathbb{R} \times \dots \times \mathbb{R}}_{d \text{ times}}}$$

Till this point, we have seen topology and topological spaces from geometric view-point. But one can use these things beyond the geometric (or manifold) applications. For example, a guy named Furstenberg introduced a simple topology on the set of integers to prove the infinitude of primes.

§2.3 Convergence

Definition 2.15 (Sequence):

A sequence (of points) in a set M is a map from $\mathbb{N}$ onto M , i.e. $q : \mathbb{N} \rightarrow M$.

Definition 2.16 (Convergence and Limit Point):

Let $(M, \mathcal{O})$ be a topological space. A sequence q in M is said to converge against a limit point $a \in M$ if

$$\forall U \in \mathcal{O} : a \in U \Rightarrow \exists N \in \mathbb{N} : \forall n > N : q(n) \in U \quad (2.5)$$

Example 2.17 (Justifying different topologies)

With the notion of convergence, we can justify the different topologies on a set.

- (a) Chaotic Topology: Consider the chaotic topology $\mathcal{O} = \{\emptyset, M\}$ on M . Then any sequence in M converges against any point in M . Let $a \in M$ and q be a sequence in M , then we have only two choice for $U \in \mathcal{O}$, i.e. $U = \emptyset$ or $U = M$. In case $U = \emptyset$, the definition of convergence is vacuously true. In case $U = M$, the definition of convergence is true as $q(n) \in M$ for all $n \in \mathbb{N}$. Hence, (arbitrary) sequence q converges against (arbitrary) point a in $(M, \mathcal{O})$.

This weird behavior is the reason for the name “chaotic topology”.

- (b) Discrete Topology: Consider the discrete topology $\mathcal{O} = \mathcal{P}(M)$ on M . Then only almost constant sequences converge against a (unique) point in M . Let $a \in M$ and q be a sequence in M , then

we have $U = \{a\} \in \mathcal{O}$. Then the definition of convergence is true if and only if $q(n) = a$ for all $n > N$. Hence, only almost constant sequences converge against a point in $(M, \mathcal{O})$.

Here we can see that every sequence is convergent in chaotic topology (the coarsest topology) and only a specific kind of sequence is convergent in discrete topology (the finest topology). So we deduce the following remark.

Remark 2.18. Let M be a set and $\mathcal{O}_1, \mathcal{O}_2$ be two topologies on M such that $\mathcal{O}_1 \subseteq \mathcal{O}_2$ and let q be a sequence in M . Then if q converges in $(M, \mathcal{O}_2)$, then q converges in $(M, \mathcal{O}_1)$ but not necessarily vice-versa.

Theorem 2.19 (Convergence in standard topology on $\mathbb{R}^d$)

Let q be a sequence in $\mathbb{R}^d$ and $a \in \mathbb{R}^d$. Then q converges against a in $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ if and only if

$$\forall \varepsilon > 0 : \exists N \in \mathbb{N} : \forall n > N : \|q(n) - a\| < \varepsilon \quad (2.6)$$

Proof:

Let q be a sequence in $\mathbb{R}^d$ and $a \in \mathbb{R}^d$. Then q converges against a in $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ if and only if

$$\forall U \in \mathcal{O}_{\text{std.}} : a \in U \Rightarrow \exists N \in \mathbb{N} : \forall n > N : q(n) \in U \quad (2.7)$$

From the definition of standard topology on $\mathbb{R}^d$, we have $U \in \mathcal{O}_{\text{std.}} \Rightarrow \exists r \in \mathbb{R}^+ : B_r(a) \subseteq U$. Hence, (2.7) is equivalent to

$$\forall r > 0 : \exists N \in \mathbb{N} : \forall n > N : q(n) \in B_r(a) \quad (2.8)$$

which is equivalent to (2.6). Q.E.D.

Example 2.20 (Convergence in $\mathbb{R}$)

Let $q(n) = 1 - \frac{1}{n+1}$ be a sequence in $\mathbb{R}$. Then q converges against 1 in $(\mathbb{R}, \mathcal{O}_{\text{std.}})$ as

$$\forall \varepsilon \in \mathbb{R}^+ : \exists N \in \mathbb{N} : \forall n > N : |q(n) - 1| = \left| 1 - \frac{1}{n+1} - 1 \right| = \frac{1}{n+1} < \varepsilon$$

But observe q is not almost constant sequence and hence does not converge in $(\mathbb{R}, \mathcal{P}(\mathbb{R}))$.

§2.4 Continuity

Definition 2.21 (Continuity):

Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two topological spaces and $\phi : M \rightarrow N$ be a map. Then ϕ is said to be continuous if

$$\forall V \in \mathcal{O}_N : \text{preim}_\phi(V) \in \mathcal{O}_M \quad (2.9)$$

In other words, ϕ is continuous iff the pre-image of every open set in N is open in M .

Example 2.22 ('Trivial' Continuity)

Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two topological spaces and $\phi : M \rightarrow N$ be a map. Then

- (a) Let $\mathcal{O}_M = \mathcal{P}(M)$ and $\mathcal{O}_N$ be any topology on N . Then every map is continuous.

Let $V \in \mathcal{O}_N$ be any open set in N . Then $\text{preim}_\phi(V) \subseteq M$ and hence $\text{preim}_\phi(V) \in \mathcal{O}_M$. Hence, ϕ is continuous.

- (b) Let $\mathcal{O}_M$ be any topology on M and $\mathcal{O}_N = \{\emptyset, N\}$. Then every map is continuous.

Here, either $V = \emptyset$ or $V = N$. In case $V = \emptyset$, we have $\text{preim}_\phi(\emptyset) = \emptyset \in \mathcal{O}_M$. In case $V = N$, we have $\text{preim}_\phi(N) = M \in \mathcal{O}_M$. Hence, ϕ is continuous.

Recovering the definition of $\varepsilon - \delta$ definition of continuity on ‘Euclidean’ spaces from the definition of continuity in topological spaces.

Theorem 2.23 (Continuity in Euclidean spaces)

Let $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a map. Then f is continuous if and only if

$$\forall a \in \mathbb{R}^m : \forall \varepsilon > 0 : \exists \delta > 0 : \|x - a\| < \delta \Rightarrow \|f(x) - f(a)\| < \varepsilon \quad (2.10)$$

Proof:

Let $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a map. Then f is continuous if and only if

$$\forall U \in \mathcal{O}_{\mathbb{R}^n} : \text{preim}_f(U) \in \mathcal{O}_{\mathbb{R}^m} \quad (2.11)$$

From the definition of standard topology on $\mathbb{R}^n$, we have $U \in \mathcal{O}_{\mathbb{R}^n} \Rightarrow \exists \varepsilon > 0 : B_\varepsilon(f(a)) \subseteq U$. Hence, (2.11) is equivalent to

$$\forall \varepsilon > 0 : \exists \delta > 0 : \text{preim}_f(B_\varepsilon(f(a))) \in \mathcal{O}_{\mathbb{R}^m} \quad (2.12)$$

which is equivalent to (2.10).

Q.E.D.

§2.4.1 Homeomorphism

Definition 2.24 (Homeomorphism):

Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two topological spaces. Let $\phi : M \rightarrow N$ be a bijection. Then ϕ is said to be a homeomorphism if

- (a) ϕ is continuous, and
- (b) ϕ^{-1} is continuous.

Remark 2.25 (Structure preserving maps). Homeomorphisms are structure preserving maps in topology.

Remark 2.26 (One-to-one pairing of open sets). Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two topological spaces and let $\phi : M \rightarrow N$ be a homeomorphism.

$$\begin{array}{ccc} M & \xrightarrow{\phi} & N \\ & \xleftarrow{\phi^{-1}} & \end{array}$$

Then ϕ provides a *one-to-one* pairing of the open sets of M with those of N .

Definition 2.27 (Homeomorphic):

Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two topological spaces. Then M and N are said to be homeomorphic if there exists a homeomorphism between them. And we write

$$M \cong_{\text{top.}} N \quad (2.13)$$

Remark 2.28 (Homeomorphic & Isomorphic). Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two homeomorphic topological spaces. Then they are isomorphic as sets. But the converse is not true.

$$M \cong_{\text{top.}} N \Rightarrow M \cong_{\text{set}} N \quad (2.14)$$

LECTURE 5

In case of set theory, we had a notion of ‘cardinality’ of a set. With the help of this notion, we can say that if two sets are isomorphic (set theoretically) then they have the same cardinality. But in case of topological spaces, we don’t have any such quantity *i.e.* we don’t have any property which can tell us that two topological spaces are ‘homeomorphic’ or not.

Building on this idea, we had classified sets into different types based on their cardinality like finite, countable, uncountable, etc. But in case of topology, classification of topological spaces is an open problem.

§2.5 Topological Properties I: Separation Axioms

Definition 2.29 (T1 Space):

A topological space $(M, \mathcal{O})$ is called T1 if for any two distinct points $p, q \in M$ (*i.e.* $p \neq q$), then

$$\exists U_p \in \mathcal{O} : q \notin U_p \quad (2.15)$$

here U_p is an open neighbourhood of p .

Definition 2.30 (T2 Space):

A topological space $(M, \mathcal{O})$ is called T2 or Hausdorff if for any two distinct points $p, q \in M$ (*i.e.* $p \neq q$), then

$$\exists U_p, V_q \in \mathcal{O} : U_p \cap V_q = \emptyset \quad (2.16)$$

here U_p and V_q are open neighbourhoods of p and q respectively.

Intuitively, we can see why [definition 2.29](#) and [definition 2.30](#) are called ‘separation axioms’. Consider standard topology on $\mathbb{R}^2$, see [fig. 2.2](#).




Figure 2.2: Separation Axioms

Remark 2.31 (T1 vs T2). Every T2 space is a T1 space, but the converse is not true.

Example 2.32

Some examples of topological spaces based on separation axioms are:

- (a) $\mathbb{R}^d$ with standard topology is a T2 space.
- (b) Zariski topology (algebraic geometry) is T1 but not T2.
- (c) Chaotic topology is neither T1 nor T2.
- (d) Discrete topology is always T1 and T2.

There are many other “T” properties, like T3, T4, etc. All of them have use cases in some area of mathematics or physics.

Remark 2.33 ($T2\frac{1}{2}$ Space). A topological space $(M, \mathcal{O})$ is called $T2\frac{1}{2}$ if for any two distinct points $p, q \in M$ (i.e. $p \neq q$), then there exists a closed neighbourhood U_p of p such that $q \notin U_p$.

Theorem 2.34 (Uniqueness of Limit Point)

Let $(M, \mathcal{O})$ be a Hausdorff space. Then every convergent sequence in M has a unique limit point.

§2.6 Compactness and Paracompactness

Before we define compactness, we need to understand the concept of ‘open cover’.

Definition 2.35 (Cover and Open Cover):

Let $(M, \mathcal{O})$ be a topological space. A set $C \subseteq \mathcal{P}(M)$ is called a cover of M if

$$\bigcup C = M. \quad (2.17)$$

If $C \subseteq \mathcal{O}$ then it is called an open cover.

Definition 2.36 (Subcover and Finite Subcover):

Let C be a cover of M . A subset $\tilde{C} \subseteq C$ is called a subcover of C if

$$\bigcup \tilde{C} = M. \quad (2.18)$$

If $\tilde{C}$ is finite then it is called a finite subcover.

§2.6.1 Compactness

Definition 2.37 (Compact Space):

A topological space $(M, \mathcal{O})$ is called compact if every open cover of M has a finite subcover.

It is clear from the definition that compactness is a topological property. So we have,

Remark 2.38. Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two homeomorphic topological spaces. Then M is compact if and only if N is compact.

Definition 2.39 (Compact Subset):

Let $(M, \mathcal{O})$ be a topological space. A subset $N \subseteq M$ is called compact if the induced topology $(N, \mathcal{O}|_N)$ is compact.

Determining whether a topological space is compact or not, is not an easy task. With the help of a suitable homeomorphism we can simplify this task.

Theorem 2.40 (Heine-Borel Theorem)

Let $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ be a topological space. Then a subset $K \subseteq \mathbb{R}^d$ is compact if and only if it is closed and bounded.

Remark 2.41 (Bounded Set in $\mathbb{R}^d$). A set $K \subseteq \mathbb{R}^d$ is called *bounded* if

$$\exists r > 0 : K \subseteq B_r(0) \quad (2.19)$$

We can extend [Heine-Borel Theorem](#) to any metric space. But first we need to define the concept of ‘metric space’.

Definition 2.42 (Metric Space):

Let M be a set. A function $d : M \times M \rightarrow \mathbb{R}$ is called a metric on M if it satisfies the following properties for all $p, q, r \in M$:

- Symmetry: $d(p, q) = d(q, p)$.
- Positivity: $d(p, q) \geq 0$ and $d(p, q) = 0 \Leftrightarrow p = q$.
- Triangle Inequality: $d(p, q) + d(q, r) \geq d(p, r)$.

A set M equipped with a metric d is called a metric space.

We can define ‘metric-induced topology’ on a metric space as we did for $\mathbb{R}^d$.

Remark 2.43 (Metric-Induced Topology). Let (M, d) be a metric space. Then the set $B_r(p)$ is an open ball of radius $r \in \mathbb{R}^+$ centered at $p \in M$ defined as

$$B_r(p) = \{q \in M : d(p, q) < r\}. \quad (2.20)$$

With this, we can define the metric-induced topology $\mathcal{O}_d$ on M as

$$U \in \mathcal{O}_d :\Leftrightarrow \forall p \in U, \exists r \in \mathbb{R}^+ : B_r(p) \subseteq U. \quad (2.21)$$

Theorem 2.44 (Generalized Heine-Borel Theorem)

Let (M, d) be a metric space with metric-induced topology $\mathcal{O}_d$. Then a subset $K \subseteq M$ is compact if and only if it is *complete* and *totally bounded*.

Example 2.45

Example of compact subsets in $\mathbb{R}$ are:

- $[0, 1]$ is compact in $(\mathbb{R}, \mathcal{O}_{\text{std}})$.
- $\mathbb{R}$ is not compact in $(\mathbb{R}, \mathcal{O}_{\text{std}})$. It suffices to show that there exists an open cover of $\mathbb{R}$ which does not have a finite subcover. Consider the open cover

$$C := \{(n, n+1) \mid n \in \mathbb{Z}\} \cup \{(n+1/2, n+3/2) \mid n \in \mathbb{Z}\}. \quad (2.22)$$

We can prove that the product of two compact spaces is compact for product topology.

Theorem 2.46

Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be two compact topological spaces. Then the product space $(M \times N, \mathcal{O}_{M \times N})$ is compact.

Now extending this to finite product spaces is easy.

Compactness is a very strong property, and sometimes it doesn’t hold. So we need a weaker property which can be used in such cases. This is where the concept of ‘paracompactness’ comes in.

§2.6.2 Paracompactness

Definition 2.47 (Refinement of a Cover):

Let C be a cover of M . A cover R of M is called a refinement of C if

$$\forall \tilde{U} \in R : \exists U \in C : \tilde{U} \subseteq U. \quad (2.23)$$

A refinement R is said to be:

- open if $R \subseteq \mathcal{O}$.
- Locally finite if for every $p \in M$, there exists an open neighbourhood U_p of p such that

$$\left| \left\{ \tilde{U} \in R \mid \tilde{U} \cap U_p \neq \emptyset \right\} \right| < \infty. \quad (2.24)$$

Remark 2.48. Let C be a cover of M and let $\tilde{C}$ be a subcover of C . Then $\tilde{C}$ is a refinement of C . But the converse is not true.

Definition 2.49 (Paracompact Space):

A topological space $(M, \mathcal{O})$ is called paracompact if every open cover of M has a locally finite open refinement.

Corollary 2.50

Every compact space is paracompact.

Definition 2.51 (Metrisable Space):

A topological space $(M, \mathcal{O})$ is called metrizable if there exists a metric d on M such that the metric-induced topology $\mathcal{O}_d$ is same as $\mathcal{O}$.

Theorem 2.52 (Stone's Theorem)

Every metrizable space is paracompact.

Example 2.53

Let's see some topological spaces which are paracompact or not:

- The topological space $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ is metrizable since $\mathcal{O}_{\text{std.}} = \mathcal{O}_d$ where $d = \|\cdot\|_2$. Hence, $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ is paracompact.
- Alexandroff Long Line is not paracompact. To construct it, we first observe that we could “build” $\mathbb{R}$ by taking the interval $[0, 1)$ and stacking countably many copies of it one after the other. Hence, in a sense, $\mathbb{R}$ is equivalent to $\mathbb{Z} \times [0, 1)$. The long line L is defined analogously as $L : \omega_1 \times [0, 1)$, where ω_1 is an uncountable infinite set. The resulting space L is not paracompact.

Theorem 2.54

Let $(M, \mathcal{O})$ be a paracompact space and $(N, \mathcal{O}_N)$ be a compact space. Then the product space $(M \times N, \mathcal{O}_{M \times N})$ is paracompact.

Corollary 2.55

Let $(M, \mathcal{O})$ be a paracompact space and $(N_i, \mathcal{O}_{N_i})$ be compact spaces for all $i \in \{1, 2, \dots, n\}$. Then the product space $M \times N_1 \times N_2 \times \dots \times N_n$ is paracompact.

At this point, we can discuss the sufficient and necessary criteria for paracompactness which are used as definitions in some books. For that we need to define the concept of ‘*partition of unity*.’

Definition 2.56 (Partition of Unity):

Let $(M, \mathcal{O})$ be a topological space. A set $\mathcal{F}$ of continuous functions $\phi : M \rightarrow [0, 1]$ is called a partition of unity if for each $p \in M$,

- (a) There exists an open neighbourhood U_p such that $\phi(p) \neq 0$ for finitely many $\phi \in \mathcal{F}$.
- (b) $\sum_{\phi \in \mathcal{F}} \phi(p) = 1$.

Definition 2.57 (Partition of Unity Subordinate to a Cover):

Let $(M, \mathcal{O})$ be a topological space and let C be a cover of M . A partition of unity $\mathcal{F}$ is said to be subordinate to C if:

$$\forall \phi \in \mathcal{F} : \exists U \in C : \phi(p) \neq 0 \Rightarrow p \in U. \quad (2.25)$$

Theorem 2.58

Let $(M, \mathcal{O})$ be a Hausdorff space. Then M is paracompact if and only if every open cover C admits a partition of unity subordinate to C .

Example 2.59

Let $(\mathbb{R}, \mathcal{O}_{\text{std.}})$ be a topological space. By [theorem 2.52](#), $(\mathbb{R}, \mathcal{O}_{\text{std.}})$ is paracompact. Consider the open cover $C := \{U_1 = (-\infty, 1), U_2 = (0, \infty)\}$. We can construct a partition of unity $\mathcal{F} = \{\phi_1, \phi_2\}$ subordinate to C as follows:

$$\phi_1(x) = \begin{cases} 1 & \text{if } x \leq 0 \\ 1 - x^2 & \text{if } 0 \leq x \leq 1 \\ 0 & \text{if } x \geq 1 \end{cases} \quad \text{and} \quad \phi_2(x) = \begin{cases} 0 & \text{if } x \leq 0 \\ x^2 & \text{if } 0 \leq x \leq 1 \\ 1 & \text{if } x \geq 1 \end{cases}$$

For ϕ_1 , we have $U_1 = (-\infty, 1)$ for which $\phi_1(x) \neq 0$ and for ϕ_2 , we have $U_2 = (0, \infty)$ for which $\phi_2(x) \neq 0$. Also, $\phi_1(x) + \phi_2(x) = 1$ for all $x \in \mathbb{R}$.

§2.7 Connectedness and Path-Connectedness

§2.7.1 Connectedness

Definition 2.60 (Connected Space):

A topological space $(M, \mathcal{O})$ is called connected unless there exists two non-empty open sets A, B such that

$$A \cup B = M \quad \text{and} \quad A \cap B = \emptyset. \quad (2.26)$$

Example 2.61

Let $(M := \mathbb{R} \setminus \{0\}, \mathcal{O} := \mathcal{O}_{\text{std.}}|_{\mathbb{R} \setminus \{0\}})$ be a topological space. Then M is not connected since we can write $M = A \cup B$ where $A = (-\infty, 0)$ and $B = (0, \infty)$ with $A \cap B = \emptyset$.

Theorem 2.62

The closed interval $[0, 1]$ with induced topology from $(\mathbb{R}, \mathcal{O}_{\text{std.}})$ is connected.

Theorem 2.63

Let $(M, \mathcal{O})$ be a topological space. Then M is connected if and only if $\emptyset$ and M are the only subsets of M that are both open and closed.

Proof:[$\Rightarrow$]:

Suppose not there exists a set $U \subseteq M$ that is both open and closed and $U \neq \emptyset, M$. Then $M = U \dot{\cup} (M \setminus U)$. Since U is open and $M \setminus U$ is open (as U is closed). Hence, M is not connected. $\nmid$

[$\Leftarrow$]:

Suppose not M is not connected. Then there exists two non-empty open sets A, B such that $A \cup B = M$ and $A \cap B = \emptyset$. Clearly, $A \neq M$, if not then $B = \emptyset$. Since A is open, $M \setminus A$ is closed. But $M \setminus A = B$ is also open. Hence, $B \neq \emptyset, M$ is open and closed. $\nmid$ Q.E.D.

§2.7.2 Path-Connectedness**Definition 2.64 (Path-Connected Space):**

A topological space $(M, \mathcal{O})$ is called path-connected if for every pair of points $p, q \in M$, there exists a continuous curve $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = p$ and $\gamma(1) = q$.

Example 2.65


Let $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ be a topological space. Then $(\mathbb{R}^d, \mathcal{O}_{\text{std.}})$ is path-connected.

Proof:

Let $p, q \in \mathbb{R}^d$. Then the curve $\gamma(\lambda) := (1 - \lambda)p + \lambda q$ is continuous and $\gamma(0) = p$ and $\gamma(1) = q$. Q.E.D.

Example 2.66

Let $S := \left\{ \left(x, \sin\left(\frac{1}{x}\right) \right) \mid x \in (0, 1] \right\} \cup \{(0, 0)\} \subseteq \mathbb{R}^2$. Then $(S, \mathcal{O}_{\text{std.}}|_S)$ is not path-connected but connected.

**Theorem 2.67**

Every path-connected space is connected.

Proof:

Let $(M, \mathcal{O})$ be a path-connected space. Suppose not M is not connected. Then there exists two non-empty open sets A, B such that $A \dot{\cup} B = M$. Since, A and B are non-empty, we can choose $a \in A$ and $b \in B$. Since M is path-connected, there exists a continuous curve $\gamma : [0, 1] \rightarrow M$ such that $\gamma(0) = a$

and $\gamma(1) = b$. Moreover, γ is continuous and A, B are open, disjoint sets, $\text{preim}_\gamma(A)$ and $\text{preim}_\gamma(B)$ are open, disjoint sets. We know

$$[0, 1] = \text{preim}_\gamma(M) = \text{preim}_\gamma(A \dot{\cup} B) = \text{preim}_\gamma(A) \dot{\cup} \text{preim}_\gamma(B).$$

Thus, existence of $\text{preim}_\gamma(A)$ and $\text{preim}_\gamma(B)$ implies that $[0, 1]$ is not connected. $\nexists$

Q.E.D.

§2.8 Homotopic Curves and Fundamental Group

Definition 2.68 (Homotopic Curves):

Let $(M, \mathcal{O})$ be a topological space. Two continuous curves $\gamma_1, \gamma_2 : [0, 1] \rightarrow M$ such that

$$\gamma_1(0) = \gamma_2(0) \quad \text{and} \quad \gamma_1(1) = \gamma_2(1)$$

are called homotopic if there exists a continuous function $h : [0, 1] \times [0, 1] \rightarrow M$ such that for all $\lambda \in [0, 1]$,

$$h(0, \lambda) = \gamma_1(\lambda) \quad \text{and} \quad h(1, \lambda) = \gamma_2(\lambda). \quad (2.27)$$

Intuitively, two curves are homotopic if one can be continuously deformed into the other. Pictorially, see fig. 2.3.




Figure 2.3: Homotopic Curves

Proposition 2.69 (Homotopy is an Equivalence Relation)

Let $(M, \mathcal{O})$ be a topological space. Define a relation $\sim$ on the set of continuous curves $\{\gamma : [0, 1] \rightarrow M\}$ by

$$\gamma_1 \sim \gamma_2 :\Leftrightarrow \gamma_1 \text{ and } \gamma_2 \text{ are homotopic.} \quad (2.28)$$

Then $\sim$ is an equivalence relation.

Proof:

Let $\gamma_1, \gamma_2, \gamma_3 : [0, 1] \rightarrow M$ be continuous curves.

- Reflexivity:** Define a continuous function $h : [0, 1] \times [0, 1] \rightarrow M$ as $h(t, \lambda) := \gamma_1(\lambda)$. Then $h(0, \lambda) = \gamma_1(\lambda)$ and $h(1, \lambda) = \gamma_1(\lambda)$ for all $\lambda \in [0, 1]$. Hence, $\gamma_1 \sim \gamma_1$.
- Symmetry:** Let $\gamma_1 \sim \gamma_2$. Then there exists a continuous function $h : [0, 1] \times [0, 1] \rightarrow M$ such that $h(0, \lambda) = \gamma_1(\lambda)$ and $h(1, \lambda) = \gamma_2(\lambda)$ for all $\lambda \in [0, 1]$. Define a continuous function $h' : [0, 1] \times [0, 1] \rightarrow M$ as $h'(t, \lambda) := h(1 - t, \lambda)$. Then $h'(0, \lambda) = \gamma_2(\lambda)$ and $h'(1, \lambda) = \gamma_1(\lambda)$ for all $\lambda \in [0, 1]$. Hence, $\gamma_2 \sim \gamma_1$.

- (c) Transitivity: Let $\gamma_1 \sim \gamma_2$ and $\gamma_2 \sim \gamma_3$. Then there exists continuous functions $h_1, h_2 : [0, 1] \times [0, 1] \rightarrow M$ such that $h_1(0, \lambda) = \gamma_1(\lambda)$, $h_1(1, \lambda) = \gamma_2(\lambda)$ and $h_2(0, \lambda) = \gamma_2(\lambda)$, $h_2(1, \lambda) = \gamma_3(\lambda)$ for all $\lambda \in [0, 1]$. Define a continuous function $h : [0, 1] \times [0, 1] \rightarrow M$ as

$$h(t, \lambda) := \begin{cases} h_1(2t, \lambda) & \text{if } 0 \leq t \leq \frac{1}{2} \\ h_2(2t - 1, \lambda) & \text{if } \frac{1}{2} \leq t \leq 1 \end{cases} \quad (2.29)$$

Then $h(0, \lambda) = \gamma_1(\lambda)$ and $h(1, \lambda) = \gamma_3(\lambda)$ for all $\lambda \in [0, 1]$. Hence, $\gamma_1 \sim \gamma_3$.

Q.E.D.

Definition 2.70 (Space of Loops):

Let $(M, \mathcal{O})$ be a topological space. Then, for every point $p \in M$, the set $\mathcal{L}_p$ defined as

$$\mathcal{L}_p := \{\gamma : [0, 1] \rightarrow M \mid \gamma \text{ is continuous and } \gamma(0) = \gamma(1) = p\} \quad (2.30)$$

is called the space of loops at p .

Definition 2.71 (Concatenation of Loops):

Let $(M, \mathcal{O})$ be a topological space. Fix a point $p \in M$. We define the concatenation operation $*_p : \mathcal{L}_p \times \mathcal{L}_p \rightarrow \mathcal{L}_p$ as

$$(\gamma_1 *_p \gamma_2)(\lambda) := \begin{cases} \gamma_1(2\lambda) & \text{if } 0 \leq \lambda < \frac{1}{2} \\ \gamma_2(2\lambda - 1) & \text{if } \frac{1}{2} \leq \lambda \leq 1 \end{cases} \quad (2.31)$$

§2.8.1 Fundamental Group

We will first recall the definition of a group.

Definition 2.72 (Group):

A set G equipped with a binary operation $*$: $G \times G \rightarrow G$ is called a group if it satisfies the following properties:

- (a) Associativity: $\forall a, b, c \in G : (a * b) * c = a * (b * c)$.
- (b) Identity Element: $\exists e \in G : \forall a \in G : a * e = e * a = a$.
- (c) Inverse Element: $\forall a \in G : \exists a^{-1} \in G : a * a^{-1} = a^{-1} * a = e$.

A group is called abelian (or commutative) if $\forall a, b \in G : a * b = b * a$.

For future discussion, we need the notion of isomorphism between groups.

Definition 2.73 (Group Isomorphism):

Let $(G, *)$ and $(H, \star)$ be two groups. A function $\varphi : G \rightarrow H$ is called a group isomorphism if it is bijective and satisfies

$$\forall a, b \in G : \varphi(a * b) = \varphi(a) \star \varphi(b). \quad (2.32)$$

Notation: If there exists an isomorphism between groups G and H , we say that G and H are isomorphic and write $G \cong_{\text{grp}} H$.

Now we can define the fundamental group.

Definition 2.74 (Fundamental Group):

Let $(M, \mathcal{O})$ be a topological space and let $p \in M$. Define the set $\pi_1(p)$ as

$$\pi_1(p) := \mathcal{L}_p / \sim = \{[\gamma] \mid \gamma \in \mathcal{L}_p\} \quad (2.33)$$

where $\sim$ is the equivalence relation defined using homotopy. Define a binary operation $\bullet_p : \pi_1(p) \times \pi_1(p) \rightarrow \pi_1(p)$ as

$$[\gamma_1] \bullet_p [\gamma_2] := [\gamma_1 *_p \gamma_2]. \quad (2.34)$$

Then $\pi_1(p)$ equipped with the operation $\bullet_p$ is called the fundamental group of M at p .

To prove that $(\pi_1(p), \bullet_p)$ is a group, we need to show following properties:

1. $\bullet_p$ is well-defined.
2. $\bullet_p$ is associative.
3. There exists an identity element.
4. There exists an inverse element for every element.

Proof:

Using the steps mentioned above, we can prove that $(\pi_1(p), \bullet_p)$ is a group.

(a) Well-Defined:

Let $\gamma_1, \gamma'_1, \gamma_2, \gamma'_2 \in \mathcal{L}_p$ such that $\gamma_1 \sim \gamma'_1$ and $\gamma_2 \sim \gamma'_2$. Then there exists continuous functions $h_1, h_2 : [0, 1] \times [0, 1] \rightarrow M$ such that $h_1(0, \lambda) = \gamma_1(\lambda)$, $h_1(1, \lambda) = \gamma'_1(\lambda)$ and $h_2(0, \lambda) = \gamma_2(\lambda)$, $h_2(1, \lambda) = \gamma'_2(\lambda)$ for all $\lambda \in [0, 1]$. Define a continuous function $h : [0, 1] \times [0, 1] \rightarrow M$ as

$$h(t, \lambda) := \begin{cases} h_1(2t, \lambda) & \text{if } 0 \leq t \leq \frac{1}{2} \\ h_2(2t - 1, \lambda) & \text{if } \frac{1}{2} \leq t \leq 1 \end{cases} \quad (2.35)$$

Then $h(0, \lambda) = \gamma_1(\lambda) *_p \gamma_2(\lambda)$ and $h(1, \lambda) = \gamma'_1(\lambda) *_p \gamma'_2(\lambda)$. Hence, $\gamma_1 *_p \gamma_2 \sim \gamma'_1 *_p \gamma'_2$.

(b) Associativity:

Let $[\gamma_1], [\gamma_2], [\gamma_3] \in \pi_1(p)$. We need to show that $([\gamma_1] \bullet_p [\gamma_2]) \bullet_p [\gamma_3] = [\gamma_1] \bullet_p ([\gamma_2] \bullet_p [\gamma_3])$ which is equivalent to showing $(\gamma_1 *_p \gamma_2) *_p \gamma_3$ and $\gamma_1 *_p (\gamma_2 *_p \gamma_3)$ are homotopic. From definition of concatenation,

$$(\gamma_1 *_p \gamma_2) *_p \gamma_3(\lambda) = \begin{cases} \gamma_1(4\lambda) & \text{if } 0 \leq \lambda \leq \frac{1}{4} \\ \gamma_2(4\lambda - 2) & \text{if } \frac{1}{4} \leq \lambda \leq \frac{1}{2} \\ \gamma_3(2\lambda - 1) & \text{if } \frac{1}{2} \leq \lambda \leq 1 \end{cases}; \quad \gamma_1 *_p (\gamma_2 *_p \gamma_3)(\lambda) = \begin{cases} \gamma_1(2\lambda) & \text{if } 0 \leq \lambda \leq \frac{1}{2} \\ \gamma_2(4\lambda - 2) & \text{if } \frac{1}{2} \leq \lambda \leq \frac{3}{4} \\ \gamma_3(4\lambda - 3) & \text{if } \frac{3}{4} \leq \lambda \leq 1 \end{cases}$$

Now define $h : [0, 1] \times [0, 1] \rightarrow [0, 1]$ as

$$h(t, \lambda) := \begin{cases} \gamma_1((2\lambda)t + (4\lambda)(1-t)) & \text{if } 0 \leq \lambda \leq \frac{t}{2} + \frac{1-t}{4} \\ \gamma_2(4\lambda - 2) & \text{if } \frac{t}{2} + \frac{1-t}{4} \leq \lambda \leq \frac{3t}{4} + \frac{1-t}{2} \\ \gamma_3((4\lambda - 3)t + (2\lambda - 1)(1-t)) & \text{if } \frac{3t}{4} + \frac{1-t}{2} \leq \lambda \leq 1 \end{cases}$$

Then $h(0, \lambda) = (\gamma_1 *_p \gamma_2) *_p \gamma_3(\lambda)$ and $h(1, \lambda) = \gamma_1 *_p (\gamma_2 *_p \gamma_3)(\lambda)$. Hence, $(\gamma_1 *_p \gamma_2) *_p \gamma_3 \sim \gamma_1 *_p (\gamma_2 *_p \gamma_3)$.

(c) Identity Element:

Observe that the constant curve $\gamma_{e,p} : [0, 1] \rightarrow M$ defined as $\gamma_{e,p}(\lambda) = p$ for all $\lambda \in [0, 1]$ is the identity element.

(d) Inverse Element:

Let $[\gamma] \in \pi_1(p)$. Then the curve $\gamma^{-1} : [0, 1] \rightarrow M$ defined as $\gamma^{-1}(\lambda) = \gamma(1 - \lambda)$ is the inverse element.

This completes the proof.

Q.E.D.

Remark 2.75 (Notion of Topological Invariance). A “topological property” (*i.e.* a property that only depends upon the topological space) is said to be *topologically invariant* if it is shared between homeomorphic spaces.

For example, connectedness, compactness, path-connectedness, etc., are topologically invariant properties.

Example 2.76 (2-sphere S^2)

The 2-sphere S^2 is defined as

$$S^2 := \{(x, y, z) \in \mathbb{R}^3 \mid x^2 + y^2 + z^2 = 1\}. \quad (2.36)$$

We define the topological space $(S^2, \mathcal{O})$ where $\mathcal{O}$ is the induced topology from $(\mathbb{R}^3, \mathcal{O}_{\text{std.}})$.




Figure 2.4: A homeomorphic representation of the 2-sphere

The sphere has the property that all the loops at any point are homotopic, hence the fundamental group (at every point) of the sphere is the trivial group:

$$\forall p \in S^2 : \pi_1(p) = \{[\gamma_{e,p}]\}. \quad (2.37)$$

Example 2.77 (Cylinder $S^1 \times \mathbb{R}$)

The cylinder C is defined as

$$C := \mathbb{R} \times S^1 \quad (2.38)$$

equipped with product topology.




Figure 2.5: A homeomorphic representation of the cylinder

A loop in C can either go around the cylinder (*i.e.* around its central axis) or not. If it does not, then it can be continuously deformed to a point (the identity loop). If it does, then it cannot be deformed to the identity loop (intuitively because the cylinder is infinitely long) and hence it is a homotopically different loop. The number of times a loop winds around the cylinder is called the *winding number*. Loops with different winding numbers are not homotopic. Moreover, loops with different orientations are also not homotopic. Hence, the fundamental group of the cylinder at any point is isomorphic to the additive group of integers:

$$\forall p \in C : (\pi_1(p), \bullet_p) \cong_{\text{grp}} (\mathbb{Z}, +). \quad (2.39)$$

Example 2.78 (2-Torus $S^1 \times S^1$)

The 2-torus T^2 is defined as

$$T^2 := S^1 \times S^1 \quad (2.40)$$

equipped with product topology.




Figure 2.6: A homeomorphic representation of the 2-torus

For the 2-torus, the fundamental group at any point is isomorphic to the additive group of integers squared:

$$\forall p \in T^2 : (\pi_1(p), \bullet_p) \cong_{\text{grp}} (\mathbb{Z} \times \mathbb{Z}, +). \quad (2.41)$$

Here the group operation is component-wise addition.

But still, the question remains: does a complete list of topological invariants exist *i.e.* can we conclude that two spaces are homeomorphic if and only if they have the same topological invariants?

This problem is still open in topology and is known as the *classification of topological spaces*.

3 Topological Manifolds and Bundles

LECTURE 6

Roughly speaking, a topological manifold is a topological space that locally looks like $\mathbb{R}^d$ for some fixed $d \in \mathbb{N}$. For example, 2-sphere is a topological manifold as well as a 2-torus and a pretzel with $d = 2$ in each case.

§3.1 Definition and Construction of Topological Manifolds

Definition 3.1 (Topological Manifold):

A paracompact Hausdorff topological space $(M, \mathcal{O})$ is called a d -dimensional (topological) manifold if for every $p \in M$ there exists an open neighborhood $U_p \in \mathcal{O}$ of p and a homeomorphism $x : U_p \rightarrow x(U_p) \subseteq \mathbb{R}^d$. And we write $\dim M = d$.

Example 3.2

Let's go through some known topological spaces to see if they are topological manifolds.

1. Trivially, $\mathbb{R}^d$ is a d -dimensional topological manifold for all $d \geq 1$.
2. The 1-sphere S^1 is a 1-dimensional topological manifold.
3. The topological spaces S^2 , C and T^2 are 2-dimensional topological manifolds.

Remark 3.3 (Local Homeomorphisms vs. Homeomorphisms). The definition of a topological manifold requires only local homeomorphisms. It is not necessary for the whole space to be homeomorphic to $\mathbb{R}^d$. For example, the 1-sphere S^1 is a 1-dimensional topological manifold, but it is not topologically isomorphic to $\mathbb{R}$.

§3.1.1 Submanifolds


Definition 3.4 (Submanifold):

Let $(M, \mathcal{O})$ be a topological manifold and $N \subseteq M$ be a subset. Then $(N, \mathcal{O}|_N)$ is called a submanifold of $(M, \mathcal{O})$ if it is a topological manifold in its own right.


Example 3.5

The manifold S^1 is a submanifold of $\mathbb{R}^2$ while the manifolds S^2 , C and T^2 are submanifolds of $\mathbb{R}^3$.

Counterexample: The figure-eight curve is not a submanifold of $\mathbb{R}^2$. As at the intersection point, we can't define dimension.



(a) Figure-Eight in $\mathbb{R}^2$



(b) Light-Cone in Minkowski Spacetime

Figure 3.1: Non-Submanifolds

We can see this in case of light-cone in Minkowski spacetime as well. The light-cone is also not a submanifold of Minkowski spacetime.

§3.1.2 Product Manifolds

Definition 3.6 (Product Manifold):

Let $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ be topological manifolds. Then $(M \times N, \mathcal{O}_{M \times N})$ is a topological manifold called the product manifold with $\dim(M \times N) = \dim M + \dim N$.

Example 3.7

This shows that $T^2 = S^1 \times S^1$ is a topological manifold of dimension 2. And we can generalize this to define n -torus as

$$T^n := \underbrace{S^1 \times \cdots \times S^1}_{n \text{ times}} \quad (3.1)$$

which is a topological manifold of dimension n .

Products are very useful. Very often in physics one intuitively thinks of the product of two manifolds as attaching a copy of the second manifold to each point of the first.

Counterexample: Möbius strip is not a product manifold.




Figure 3.2: Non-Product Manifolds

§3.2 Bundles

Definition 3.8 (Bundle):

A bundle (of a topological space) is a triple (E, π, M) where E and M are topological spaces called the total space and the base space respectively, and $\pi : E \rightarrow M$ is a continuous surjective map called the projection map.

We often denote the bundle as

$$E \xrightarrow{\pi} M \quad (3.2)$$

Definition 3.9 (Fiber):

Let (E, π, M) be a bundle and $p \in M$. Then the set $F_p := \text{preim}_\pi(\{p\})$ is called the fiber over p .

Example 3.10 (Product Bundle)

Let M and F be topological spaces. Define the product space $E := M \times F$ and the projection map $\pi : E \rightarrow M$ as $\pi(m, f) = m$. Then (E, π, M) is a bundle called the *product bundle*.

Define $\pi' : E \rightarrow F$ as $\pi'(m, f) = f$. Then (E, π', F) is also a product bundle.

Example 3.11 (Möbius Strip as a Bundle)

Consider the Möbius strip as a bundle. Define the total space E as the Möbius strip and the projection map $\pi : E \rightarrow S^1$ as the projection of the Möbius strip onto the circle. Then (E, π, S^1) is a bundle.

It is easy to see that the fiber over any point $p \in S^1$ is a line segment. See [fig. 3.3](#).

$$\forall p \in S^1 : F_p = [-1, 1] \quad (3.3)$$





Figure 3.3: Möbius Strip as a Bundle

In both the examples, the fiber over any point in the base space is the same. But this is not necessary. The fiber can be different over different points in the base space.

Example 3.12

Consider the base space $M = \mathbb{R}$ and the total space as defined in fig. 3.4.

Figure 3.4: Total Space E

Here, the fiber over every point in the base space is not the same.

$$\forall p \in \mathbb{R} : F_p \cong_{\text{top.}} \begin{cases} S^1 & \text{if } p < 0 \\ \{0\} & \text{if } p = 0 \\ (1, -1) & \text{if } p > 0 \end{cases} \quad (3.4)$$

§3.2.1 Fiber Bundles

Definition 3.13 (Fiber Bundle):

Let $E \xrightarrow{\pi} M$ be a bundle. Then (E, π, M) is called a fiber bundle if

$$\exists F (= \text{topological space}) : \forall p \in M : F_p \cong_{\text{top.}} F \quad (3.5)$$

We can think of a fiber bundle as an attachment of a copy of the fiber F to each point of the base space M . Notationally, we write

$$\begin{array}{ccc} F & \longrightarrow & E \\ & & \downarrow \pi \\ & & M \end{array} \quad (3.6)$$

It is easy to see that the product bundle is a fiber bundle. But the converse is not true.

Example 3.14 (Möbius Strip as a Fiber Bundle)

Let Möbius strip be our total space and 1-sphere S^1 be our base space. Then the Möbius strip is a fiber bundle over the 1-sphere with the typical fiber $[-1, 1]$. See fig. 3.3.

Keep in mind that $E \neq S^1 \times [-1, 1]$ i.e. Möbius strip is not a product bundle.

Example 3.15 ($\mathbb{C}^1$ -line bundle)

A $\mathbb{C}$ -line bundle over M is the fiber bundle (E, π, M) with fiber $\mathbb{C}$.

Note that $M \times \mathbb{C} \xrightarrow{\pi} M$ is a $\mathbb{C}$ -line bundle over M , but a $\mathbb{C}$ -line bundle over M need not be a product bundle.

Definition 3.16 (Section):

Let $E \xrightarrow{\pi} M$ be a bundle. A map $\sigma : M \rightarrow E$ is called a (cross-)section of the bundle if $\pi \circ \sigma = \text{id}_M$.

Intuitively, a section is a map σ which sends each point $p \in M$ to some point $\sigma(p)$ in its fiber F_p , so that the projection map π takes $\sigma(p) \in F_p \subseteq E$ back to the point $p \in M$.

Example 3.17 (Special Case for Product Bundles)

Let $M \times F \xrightarrow{\pi} M$ be a product bundle. Let $s : M \rightarrow F$ be a map. Then the map

$$\begin{aligned}\sigma : M &\rightarrow M \times F \\ p &\mapsto (p, s(p))\end{aligned}\tag{3.7}$$

is a section of the product bundle.

Thus, for product bundles, sections are in one-to-one correspondence with maps $M \rightarrow F$ i.e. by choosing a map $s : M \rightarrow F$, we get a section $\sigma : M \rightarrow M \times F$.

In case of product bundles, we can analyze ‘section’ of the bundle by looking at the corresponding map $s : M \rightarrow F$.

With this we can order the bundles as follows:

$$\text{Product Bundles} \subset \text{Fiber Bundles} \subset \text{Bundles}\tag{3.8}$$

At this point, we can have a physics example (from quantum mechanics) but to understand it fully, we require more maturity.

Example 3.18 (Wave functions)

Consider our base space M to be our physical space (for case $\mathbb{R}^3$). The mathematical structure, we are analyzing in quantum mechanics is $\mathbb{C}^1$ -line bundle over physical space,

$$\text{wave function} := \text{section of } (E, \pi, M).\tag{3.9}$$

In case, E is a product manifold of M and $\mathbb{C}$, then wave function is a function.

§3.2.2 Constructing Bundles**Definition 3.19 (Sub-Bundle):**

Let $E \xrightarrow{\pi} M$ and $E' \xrightarrow{\pi'} M'$ be bundles. Then (E', π', M') is called a sub-bundle of (E, π, M) if

1. $E' \subseteq E$ be a submanifold,
2. $M' \subseteq M$ be a submanifold, and
3. $\pi' = \pi|_{E'}$.

Definition 3.20 (Restricted Bundle):

Let $E \xrightarrow{\pi} M$ be a bundle and $N \subseteq M$ be a submanifold. Then the bundle

$$\text{preim}_{\pi}(N) \xrightarrow{\pi|_{\text{preim}_{\pi}(N)}} N\tag{3.10}$$

is called the restricted bundle of (E, π, M) to N .

§3.3 Bundle Morphisms

Definition 3.21 (Bundle Morphism):

Let $E \xrightarrow{\pi} M$ and $E' \xrightarrow{\pi'} M'$ be bundles and $u : E \rightarrow E'$ and $v : M \rightarrow M'$ be continuous maps. Then (u, v) is called a bundle morphism if

$$\begin{array}{ccc} E & \xrightarrow{u} & E' \\ \pi \downarrow & & \downarrow \pi' \\ M & \xrightarrow{v} & M' \end{array} \quad (3.11)$$

commutes i.e. $\pi' \circ u = v \circ \pi$.

Definition 3.22 (Bundle Isomorphism):

Let $E \xrightarrow{\pi} M$ and $E' \xrightarrow{\pi'} M'$ be bundles. They are called isomorphic (as bundles) if there exist bundle morphisms (u, v) such that

$$\begin{array}{ccc} E & \begin{array}{c} \xrightarrow{u} \\ \xleftarrow{u^{-1}} \end{array} & E' \\ \pi \downarrow & & \downarrow \pi' \\ M & \begin{array}{c} \xrightarrow{v} \\ \xleftarrow{v^{-1}} \end{array} & M' \end{array} \quad (3.12)$$

is a commutative diagram. Such a bundle morphism (u, v) is called a bundle isomorphism. And we write $E \xrightarrow{\pi} M \cong_{\text{bdl}} E' \xrightarrow{\pi'} M'$.

Bundle isomorphisms are structure-preserving maps between bundles.

Definition 3.23 (Local Isomorphism):

Let $E \xrightarrow{\pi} M$ and $E' \xrightarrow{\pi'} M'$ be bundles. Then $E \xrightarrow{\pi} M$ is called locally isomorphic to $E' \xrightarrow{\pi'} M'$ if

$$\forall p \in M : \exists U_p \in \mathcal{O}_M : \text{preim}_{\pi}(U_p) \xrightarrow{\pi|_{\text{preim}_{\pi}(U_p)}} U_p \cong_{\text{bdl}} E' \xrightarrow{\pi'} M'. \quad (3.13)$$

Some useful terminologies:

Definition 3.24:

A bundle $E \xrightarrow{\pi} M$ is called

- (a) trivial if it is isomorphic to a product bundle $M \times F \xrightarrow{\pi_1} M$,
- (b) locally trivial if it is locally isomorphic to some product bundle.

It is obvious that every trivial bundle is locally trivial. But the converse is not true.

Example 3.25

Let's see some insightful examples:

1. A cylinder C is trivial $\Rightarrow$ locally trivial.
2. Möbius Strip is not trivial, but it is locally trivial.
3. A very specific construction, such that the bundle is not even locally trivial.




Figure 3.5: Non Locally Trivial bundle

From now on, we will be focusing on locally trivial bundles unless mentioned otherwise.

Remark 3.26. Since we have restricted ourselves to locally trivial bundles. Thus, any section can be thought of as a map $M \rightarrow F$ locally.

Example 3.27 (Wave functions (revisited))

Assume that our $\mathbb{C}^1$ -line bundle over $M(=\mathbb{R}^3)$ is locally trivial. Then the wave function is a function $M \rightarrow \mathbb{C}$ locally.

Definition 3.28 (Pull-Back Bundle):

Let $E \xrightarrow{\pi} M$ be a bundle and $f : M' \rightarrow M$ be a map from another manifold M' to M . Define the manifold E' as

$$E' := \{(m', e) \in M' \times E \mid f(m') = \pi(e)\}. \quad (3.14)$$

Then the bundle (E', π', M') with $\pi'(m', e) = m'$ is called the pull-back bundle of $E \xrightarrow{\pi} M$ induced by $f : M' \rightarrow M$.

Remark 3.29 (Bundle Morphism on Pull-Back Bundle). Let $E' \xrightarrow{\pi'} M'$ be a pull-back bundle of $E \xrightarrow{\pi} M$ induced by $v : M' \rightarrow M$. Define the map

$$\begin{aligned} u : E' &\rightarrow E \\ (m', e) &\mapsto e. \end{aligned} \quad (3.15)$$

Then (u, v) is a bundle morphism. This corresponds to the commutative diagram

$$\begin{array}{ccc} E' & \xrightarrow{u} & E \\ \pi' \downarrow & & \downarrow \pi \\ M' & \xrightarrow{v} & M \end{array}$$

Remark 3.30 (Pull-back of Sections). Let $E' \xrightarrow{\pi'} M'$ be a pull-back bundle of $E \xrightarrow{\pi} M$ induced by $f : M' \rightarrow M$. Let σ be a section of $E \xrightarrow{\pi} M$.

$$\begin{array}{ccc} E' & & E \\ \uparrow \sigma' & \nearrow \sigma \circ f & \uparrow \sigma \\ M' & \xrightarrow{f} & M \end{array}$$

$\pi' \downarrow$ and $\downarrow \pi$ are also indicated in the original diagram.

Observe that $\sigma \circ f$ is a map $M' \rightarrow E$. Using the fact that σ is a section, we have,

$$(\pi \circ (\sigma \circ f))(m') = ((\pi \circ \sigma) \circ f)(m') = (\text{id}_M \circ f)(m') = f(m')$$

This can be interpreted as $\pi((\sigma \circ f)(m')) = f(m')$. Thus, $(m', (\sigma \circ f)(m')) \in E'$. Define the map σ' as

$$\begin{aligned} \sigma' : M' &\rightarrow E' \\ m' &\mapsto (m', (\sigma \circ f)(m')). \end{aligned} \quad (3.16)$$

Furthermore, $\pi' \circ \sigma' = \text{id}_{M'}$ as $\pi'(m', (\sigma \circ f)(m')) = m'$. Thus, σ' is a section of the pull-back bundle.

§3.4 Viewing Manifolds from Atlases

Definition 3.31 (Chart):

Let $(M, \mathcal{O})$ be a d -dimensional manifold. Then the pair (U, x) , where $U \in \mathcal{O}$ and $x : U \rightarrow x(U) \subseteq \mathbb{R}^d$ is a homeomorphism, is called a chart on M .

Remark 3.32 (Component Functions). Let (U, x) be a chart on M . Then the *component functions* of x are the functions

$$\begin{aligned} x^i : U &\rightarrow \mathbb{R} \\ p &\mapsto \text{proj}^i(x(p)) \end{aligned} \quad (3.17)$$

for $1 \leq i \leq d$, where proj^i is the i -component of $x(p) \in \mathbb{R}^d$. The $x^i(p)$'s are called *co-ordinate* of the point $p \in U$ with respect to the chart (U, x) .

Definition 3.33 (Atlas):

Let $(M, \mathcal{O})$ be a d -dimensional manifold. Then the collection of all charts $\mathcal{A} := \{(U_\alpha, x_\alpha) \mid \alpha \in I\}$ is called an atlas of the manifold M if,

$$\bigcup_{\alpha \in I} U_\alpha = M, \quad (3.18)$$

for some index set I .

This is obvious to see that this collection is a set, as from the definition of manifolds, we have an open set U_p at each point with an appropriate homeomorphism x_p . Thus, this provides a trivial atlas of M ,

$$\mathcal{A} = \{(U_p, x_p) \mid p \in M\} \quad (3.19)$$

Definition 3.34 ($\mathcal{C}^0$ -Compatibility):

Let (U_α, x_α) and (U_β, x_β) be two charts of a manifold M . They are called $\mathcal{C}^0$ -compatible either

- (a) $U_\alpha \cap U_\beta = \emptyset$, or
- (b) the map $x_\beta \circ x_\alpha^{-1} : x_\alpha(U_\alpha \cap U_\beta) \rightarrow x_\beta(U_\alpha \cap U_\beta)$ is continuous.

$x_\beta \circ x_\alpha^{-1}$ (and its inverse $x_\alpha \circ x_\beta^{-1}$) is a map from a subset of $\mathbb{R}^d$ to another subset of $\mathbb{R}^d$.

$$\begin{array}{ccc} & U_\alpha \cap U_\beta & \\ x_\alpha \swarrow & & \searrow x_\beta \\ \mathbb{R}^d \supseteq x_\alpha(U_\alpha \cap U_\beta) & \xrightleftharpoons[x_\alpha \circ x_\beta^{-1}]{x_\beta \circ x_\alpha^{-1}} & x_\beta(U_\alpha \cap U_\beta) \subseteq \mathbb{R}^d \end{array}$$

Here it may seem like a redundant definition, since x_α and x_β are homeomorphisms, the composition map $x_\beta \circ x_\alpha^{-1}$ (or its inverse $x_\alpha \circ x_\beta^{-1}$) is also a homeomorphism thus continuous. Therefore, any two charts on a topological manifold are $\mathcal{C}^0$ -compatible.

This notion will be useful, when we want to talk about differentiable manifolds. In that case, we require the map $x_\beta \circ x_\alpha^{-1}$ to be continuously differentiable or k -smooth for $\mathcal{C}^1$ or $\mathcal{C}^k$ -compatible respectively using the notion of differentiability on $\mathbb{R}^d$.

Remark 3.35 (Co-ordinate Change). The map $\tau_{\alpha,\beta} := x_\beta \circ x_\alpha^{-1}$ (and its inverse $\tau_{\beta,\alpha} := x_\alpha \circ x_\beta^{-1}$) is called the *co-ordinate change map* or *chart transition map*.

This also express that, all the physics that we have done so far lives in chart. For example, the motion of a particle is a subset of our physical manifold, and we use different co-ordinates (or charts) for our understanding.

Definition 3.36 ($\mathcal{C}^0$ -Atlas):

An atlas $\mathcal{A}$ is a $\mathcal{C}^0$ -atlas if all the charts in $\mathcal{A}$ are pairwise $\mathcal{C}^0$ -compatible.

Any atlas is a $\mathcal{C}^0$ -atlas.

Definition 3.37 (Maximal Atlas):

A $\mathcal{C}^0$ -atlas $\mathcal{A}$ is called a maximal if it contains every chart that is $\mathcal{C}^0$ -compatible with every chart in $\mathcal{A}$.

Example 3.38

Let $(\mathbb{R}, \mathcal{O}_{\text{std.}})$ be our topological manifold. Then we have a trivial atlas $\mathcal{A} = \{(\mathbb{R}, \text{id}_{\mathbb{R}})\}$, but it is not maximal. Since chart $((0, 1), \text{id}_{(0,1)})$ is $\mathcal{C}^0$ -compatible with the chart $(\mathbb{R}, \text{id}_{\mathbb{R}})$, but it is not in the atlas $\mathcal{A}$.

We can look at “objects on” topological manifolds from two point of view. Consider a curve γ in a d -dimensional manifold, *i.e.* $\gamma : \mathbb{R} \rightarrow M$. From our previous knowledge, we know this curve has to be continuous if it describes the trajectory of a classical particle.

One way to verify this, let an open set in M and check whether its pre-image is open or not (topological definition).

Another way is through the concept of charts. Let (U, x) be a chart, rather than studying γ directly, we will focus on $x \circ \gamma : \mathbb{R} \supseteq \text{preim}_\gamma(U) \rightarrow x(U) \subseteq \mathbb{R}^d$. Checking continuity of this composition map is easier as well as guarantees the continuity of γ .

Let the problem is very complicated in this co-ordinate system, let (U, y) be another chart. And this change of co-ordinate is facilitated by transition maps. We can summarize all this in the following commutative diagram:




Figure 3.6: Looking at Continuity of a curve in a manifold using charts

Intuitively, this description allows us to forget about the inner structure (*i.e.* U and the maps γ , x and y) which, in a sense, is the real world, and only consider $\text{preim}_\gamma(U) \subseteq \mathbb{R}$ and $x(U), y(U) \subseteq \mathbb{R}^d$ together with the maps between them, which is our representation of the real world.

4 Differentiable Manifolds

§4.1 Adding Structure by refining the (maximal) $\mathcal{C}^0$ -atlas

Definition 4.1 ($\mathfrak{A}$ -Atlas):

Let $(M, \mathcal{O})$ be a d -dimensional manifold. An atlas $\mathcal{A}$ is called $\mathfrak{A}$ -atlas, if any two charts $(U, x), (V, y) \in \mathcal{A}$ are $\mathfrak{A}$ -compatible.

In other words, either $U \cap V = \emptyset$ or if $U \cap V \neq \emptyset$, then the transition map $y \circ x^{-1}$ is $\mathfrak{A}$ as a map from $\mathbb{R}^d$ to $\mathbb{R}^d$.

$$\begin{array}{ccc} & U \cap V & \\ x \swarrow & & \searrow y \\ \mathbb{R}^d \supseteq x(U \cap V) & \xrightleftharpoons[x \circ y^{-1}]{y \circ x^{-1}} & y(U \cap V) \subseteq \mathbb{R}^d \end{array}$$

Now, we can define the placeholder symbol $\mathfrak{A}$ as:

- $\mathfrak{A} = \mathcal{C}^0$: see [definition 3.36](#).
- $\mathfrak{A} = \mathcal{C}^k$: the transition map is k -times continuously differentiable as maps $\mathbb{R}^d \rightarrow \mathbb{R}^d$.
- $\mathfrak{A} = \mathcal{C}^\infty$: the transition map is smooth (infinitely many times differentiable); i.e., k -times continuously differentiable for all $k \in \mathbb{N}$.
- $\mathfrak{A} = \mathcal{C}^\omega$: the transition map is real-analytic; i.e., it can be locally represented by a convergent power series.
- $\mathfrak{A} = \mathcal{C}_{\mathbb{C}}^\omega$: the transition map is complex-analytic; equivalently, it satisfies the Cauchy-Riemann conditions.

Here for completeness, we need to define what are the Cauchy-Riemann conditions:

Set theoretical we know that $\mathbb{C} \cong_{\text{set}} \mathbb{R}^2$. Let $f : \mathbb{C} \rightarrow \mathbb{C}$ be a complex function defined as

$$\begin{aligned} f : \mathbb{C} &\rightarrow \mathbb{C} \\ x + iy &\mapsto u(x, y) + iv(x, y) \end{aligned} \tag{4.1}$$

where $u, v : \mathbb{R}^2 \rightarrow \mathbb{R}$ are real-valued functions. Then the Cauchy-Riemann conditions says that f is complex-analytic at $x_0 + iy_0$ if and only if the following two conditions are satisfied:

1. All the partial derivatives of u and v exist at (x_0, y_0) and are continuous in a neighborhood of (x_0, y_0) .
2. The following two equations are satisfied:

$$\frac{\partial u}{\partial x}(x_0, y_0) = \frac{\partial v}{\partial y}(x_0, y_0) \quad \text{and} \quad \frac{\partial u}{\partial y}(x_0, y_0) = -\frac{\partial v}{\partial x}(x_0, y_0). \tag{4.2}$$

Theorem 4.2 (Whitney)

Any maximal $\mathcal{C}^k$ -atlas (for any $k \geq 1$) contains a $\mathcal{C}^\infty$ -atlas. Moreover, any two maximal $\mathcal{C}^k$ -atlases that contains the same $\mathcal{C}^\infty$ -atlas are identical.

In other words, we refine (or remove) all the charts in a maximal $\mathcal{C}^k$ -atlas which are not $\mathcal{C}^\infty$ -compatible, and we get a maximal $\mathcal{C}^\infty$ -atlas. This is the reason why we can always work with $\mathcal{C}^\infty$ -atlases given that we are working with a differentiable manifold. Immediate consequence of this theorem is that if any result is true for a $\mathcal{C}^k$ -atlas for any $k \geq 1$, then it is also true for a $\mathcal{C}^\infty$ -atlas.

But in the case of $\mathcal{C}^0$ -atlas, it may happen that it doesn't admit a $\mathcal{C}^1$ -atlas, and hence we cannot refine it to a $\mathcal{C}^\infty$ -atlas.

Hence, we will not make any distinction between $\mathcal{C}^k$ -manifolds and $\mathcal{C}^\infty$ -manifolds, and we will always work with $\mathcal{C}^\infty$ -manifolds.

Definition 4.3 ($\mathcal{C}^k$ -manifold):

A triple $(M, \mathcal{O}, \mathcal{A})$ is called a $\mathcal{C}^k$ -manifold where

- $(M, \mathcal{O})$ is a topological manifold.
- $\mathcal{A}$ is a maximal $\mathcal{C}^k$ -atlas on M .

Definition 4.4 (Incompatible Atlases):

Let two $\mathfrak{B}$ -compactible atlases $\mathcal{A}_1$ and $\mathcal{A}_2$ on a topological manifold $(M, \mathcal{O})$ be called compatible if $\mathcal{A}_1 \cup \mathcal{A}_2$ is a $\mathfrak{B}$ -atlas on M . Otherwise, they are called incompatible.

Remark 4.5. A given topological manifold $(M, \mathcal{O})$ can have different incompatible atlases.

A simple example of incompatible atlases,

Example 4.6

Let $M = \mathbb{R}$ with the standard topology, and let $\mathcal{A}_1 = \{(\mathbb{R}, \text{id}_{\mathbb{R}})\}$ and $\mathcal{A}_2 = \{(\mathbb{R}, a \mapsto \sqrt[3]{a})\}$. Since each atlas contains only one chart, they are trivially $\mathcal{C}^\infty$ -compatible as the transition map is the identity map in both cases. But $\mathcal{A}_1 \cup \mathcal{A}_2$ is not a $\mathcal{C}^\infty$ -atlas on M because the transition maps $\text{id}_{\mathbb{R}} \circ x^{-1} \equiv a \mapsto a^3$ which is a smooth map, but the transition map $x \circ \text{id}_{\mathbb{R}}^{-1} \equiv a \mapsto \sqrt[3]{a}$ is not smooth as it is not differentiable at $a = 0$. Hence, $\mathcal{A}_1$ and $\mathcal{A}_2$ are incompatible atlases on M .

This example shows that we can equip the real line $\mathbb{R}$ with different incompatible $\mathcal{C}^\infty$ -structures. This sounds bad as we want to do physics on $\mathbb{R}$, and we want to have a unique $\mathcal{C}^\infty$ -structure on it. But this is not a problem, as we are given an atlas by the definition of differentiable manifold.

Definition 4.7 (Differentiability):

Let $(M, \mathcal{O}_M, \mathcal{A}_M)$ and $(N, \mathcal{O}_N, \mathcal{A}_N)$ be two $\mathcal{C}^k$ -manifolds of dimension m and n respectively. A map $f : M \rightarrow N$ is called $\mathcal{C}^k$ -differentiable at a point $p \in M$ if there exists a chart $(U, x) \in \mathcal{A}_M$ around p and a chart $(V, y) \in \mathcal{A}_N$ around $f(p)$ such that the map $(y \circ f \circ x^{-1}) : x(U) \rightarrow y(V)$ is $\mathcal{C}^k$ -differentiable at $x(p)$ as a map from $\mathbb{R}^m$ to $\mathbb{R}^n$.

$$\begin{array}{ccc}
 M \supseteq U & \xrightarrow{f} & N \subseteq V \\
 \downarrow x & & \downarrow y \\
 \mathbb{R}^m \supseteq x(U) & \xrightarrow{y \circ f \circ x^{-1}} & y(V) \subseteq \mathbb{R}^n
 \end{array}$$

If f is $\mathcal{C}^k$ -differentiable at every point $p \in M$, then we say that f is a $\mathcal{C}^k$ -differentiable map from M to N .

Proposition 4.8

The definition of $\mathcal{C}^k$ -differentiability is independent of the choice of charts $(U, x) \in \mathcal{A}_M$ and $(V, y) \in \mathcal{A}_N$ i.e. the definition is well-defined.

Proof:

Consider two charts $(U, x), (\tilde{U}, \tilde{x}) \in \mathcal{A}_M$ around p and two charts $(V, y), (\tilde{V}, \tilde{y}) \in \mathcal{A}_N$ around $f(p)$. We need to show that if f is $\mathcal{C}^k$ -differentiable at p with respect to the charts (U, x) and (V, y) , then it should also be $\mathcal{C}^k$ -differentiable with respect to the charts $(\tilde{U}, \tilde{x})$ and $(\tilde{V}, \tilde{y})$. Consider the following commutative diagram:

$$\begin{array}{ccc}
 \mathbb{R}^m \supseteq \tilde{x}(U \cap \tilde{U}) & \xrightarrow{\tilde{y} \circ f \circ \tilde{x}^{-1}} & \tilde{y}(V \cap \tilde{V}) \subseteq \mathbb{R}^n \\
 \uparrow \tilde{x} & & \uparrow \tilde{y} \\
 M \supseteq U \cap \tilde{U} \ni p & \xrightarrow{f} & f(p) \in V \cap \tilde{V} \subseteq N \\
 \downarrow x & & \downarrow y \\
 \mathbb{R}^m \supseteq x(U \cap \tilde{U}) & \xrightarrow{y \circ f \circ x^{-1}} & y(V \cap \tilde{V}) \subseteq \mathbb{R}^n \\
 \swarrow x \circ \tilde{x}^{-1} & & \searrow \tilde{y} \circ y^{-1}
 \end{array}$$

We know that the transition maps $x \circ \tilde{x}^{-1}$ and $\tilde{y} \circ y^{-1}$ are $\mathcal{C}^k$ -differentiable as they are transition maps between charts in the same atlas. So the composition of the maps

$$\tilde{y} \circ f \circ \tilde{x}^{-1} = (\tilde{y} \circ y^{-1}) \circ (y \circ f \circ x^{-1}) \circ (x \circ \tilde{x}^{-1}) \quad (4.3)$$

is also $\mathcal{C}^k$ -differentiable as a composition of $\mathcal{C}^k$ -differentiable maps. Hence, f is $\mathcal{C}^k$ -differentiable at p with respect to the charts $(\tilde{U}, \tilde{x})$ and $(\tilde{V}, \tilde{y})$. Q.E.D.

Definition 4.9 (Diffeomorphism):

Let $(M, \mathcal{O}_M, \mathcal{A}_M)$ and $(N, \mathcal{O}_N, \mathcal{A}_N)$ be two $\mathcal{C}^k$ -manifolds. A map $f : M \rightarrow N$ is called a $\mathcal{C}^k$ -diffeomorphism if it is a bijection and both f and its inverse $f^{-1} : N \rightarrow M$ are $\mathcal{C}^k$ -differentiable.

Definition 4.10 (Diffeomorphic):

Two $\mathcal{C}^k$ -manifolds $(M, \mathcal{O}_M, \mathcal{A}_M)$ and $(N, \mathcal{O}_N, \mathcal{A}_N)$ are called diffeomorphic if there exists a $\mathcal{C}^k$ -diffeomorphism $f : M \rightarrow N$. Then we write

$$M \cong_{\text{diff}} N \quad \text{or} \quad (M, \mathcal{O}_M, \mathcal{A}_M) \cong_{\text{diff}} (N, \mathcal{O}_N, \mathcal{A}_N). \quad (4.4)$$

With this new notation, we want to finally answer the question: whether, for instance

$$(\mathbb{R}, \mathcal{O}_{\text{std.}}, \mathcal{A}_{1, \text{max}}) \cong_{\text{diff}} (\mathbb{R}, \mathcal{O}_{\text{std.}}, \mathcal{A}_{2, \text{max}}) \quad (4.5)$$

where $\mathcal{A}_{1, \text{max}}$ and $\mathcal{A}_{2, \text{max}}$ are the maximal $\mathcal{C}^\infty$ -atlases on $\mathbb{R}$ defined in the previous example.

In principle, we want to know, how many different differentiable structures are there on a given topological manifold $(M, \mathcal{O})$ – up to diffeomorphism? The answer to this question is not known in general, but we know that it depends on the dimension of the manifold M .

Theorem 4.11 (Radon-Moise)

For $d \leq 3$, any two $\mathcal{C}^\infty$ -manifolds of dimension d are diffeomorphic if and only if they are homeomorphic. In other words, let $(M, \mathcal{O}_M, \mathcal{A}_M)$ and $(N, \mathcal{O}_N, \mathcal{A}_N)$ be two $\mathcal{C}^\infty$ -manifolds of dimension d and $d \leq 3$. Then

$$(M, \mathcal{O}_M, \mathcal{A}_M) \cong_{\text{diff}} (N, \mathcal{O}_N, \mathcal{A}_N) \Leftrightarrow (M, \mathcal{O}_M) \cong_{\text{top.}} (N, \mathcal{O}_N). \quad (4.6)$$

So in particular, if $(M, \mathcal{O}_M)$ and $(N, \mathcal{O}_N)$ are homeomorphic, then we have a unique $\mathcal{C}^\infty$ -structure on M and N up to diffeomorphism.

From the above theorem, we can say that given a topological manifold $(M, \mathcal{O}_M)$, there is a unique $\mathcal{C}^\infty$ -structure on it up to diffeomorphism if $\dim M \leq 3$.

The answer for $d > 4$ (specifically for $d \neq 4$) is that there are finitely many different differentiable structures on a given *compact* topological manifold $(M, \mathcal{O}_M)$ up to diffeomorphism. This answer is provided by *surgery theory* (or obstruction theory). This is a collection of tools and techniques in topology with which one obtains a new manifold from given ones by performing surgery on them, *i.e.* by cutting, replacing and gluing parts in such a way as to control topological invariants like the fundamental group. The idea is to understand all manifolds in dimensions higher than 4 by performing surgery systematically.

Remark 4.12 (Good News for String Theorists). According to many string theorists, our space-time is a 10-dimensional manifold. Since we don't have a unique differentiable structure on a 10-dimensional manifold, so in principle, different differentiable structures can lead to different predictions in physics, which is not what we want. But the good news is that for $d = 10$, there are only finitely many different differentiable structures, so we can decide which one is the correct for our space-time by performing finite number of experiments.

For $d = 4$, the situation is very different. In fact, the problem of classifying all smooth differentiable structures is still open. But we know following partial results:

- Non-compact 4-manifolds:

There are uncountably many non-diffeomorphic $\mathcal{C}^\infty$ -structures, specifically on $\mathbb{R}^4$.

- Compact 4-manifolds:

The classification is not yet complete, but one of the most interesting results is that there are countably many non-diffeomorphic $\mathcal{C}^\infty$ -structures on a given compact 4-manifold with $b_2 > 18$ (where b_2 is the second Betti number, which is a topological invariant of the manifold).

Remark 4.13 (Betti Numbers). Betti numbers are topological invariants defined using homology groups (notion of algebraic topology). But, intuitively, the k -th Betti number b_k of a topological space is the number of k -dimensional holes in it.

- b_0 is the number of connected components;
- b_1 is the number of 1-dimensional or “circular” holes;
- b_2 is the number of 2-dimensional “voids” or “cavities”.
- And so on.

For example, the 2-sphere S^2 has $b_0 = 1$, $b_1 = 0$ and $b_2 = 1$ as it has one connected component, no circular holes and one 2-dimensional cavity. And the 2-torus T^2 has $b_0 = 1$, $b_1 = 2$ and $b_2 = 1$ as it has one connected component, two circular holes (one equatorial and one meridional) and one 2-dimensional cavity.

Key feature of a differentiable manifold is that there exists a “*tangent space*” at each point of the manifold.

LECTURE 8

§4.2 Review of Vector Spaces

Definition 4.14 ((Algebraic) Field):

Let K be a non-empty set with two binary operations $+$ and $\cdot$ (addition and multiplication) such that

- $(K, +)$ is an abelian group with identity element 0.
- $(K \setminus \{0\}, \cdot)$ is an Abelian group with identity element 1.
- Multiplication is distributive over addition, *i.e.* for all $a, b, c \in K$,

$$a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{and} \quad (a + b) \cdot c = a \cdot c + b \cdot c. \quad (4.7)$$

Then K is called a field.

Later we will use a set R equipped with two binary operations $+$ and $\cdot$ but fewer axioms than a field, and we will call it a *ring*.

Definition 4.15 (Ring):

A ring is a set R equipped with two binary operations $+$ and $\cdot$ such that

- $(R, +)$ is an Abelian group with identity element 0 .
- $(R, \cdot)$ is a semigroup, i.e. multiplication is associative: for all $a, b, c \in R$, $(a \cdot b) \cdot c = a \cdot (b \cdot c)$.
- Multiplication is distributive over addition, i.e. for all $a, b, c \in R$,

$$a \cdot (b + c) = a \cdot b + a \cdot c \quad \text{and} \quad (a + b) \cdot c = a \cdot c + b \cdot c. \quad (4.8)$$

Some examples which we are using from our school days:

- $(\mathbb{Z}, +, \cdot)$ is a commutative ring but not a field as it has no multiplicative inverses for all non-zero elements.
- $(\mathbb{Q}, +, \cdot)$ and $(\mathbb{R}, +, \cdot)$ are fields.
- $(\mathcal{M}_n(\mathbb{R}), +, \circ)$ is a ring, where $\mathcal{M}_n(\mathbb{R})$ is the set of $n \times n$ real matrices with the usual matrix addition and multiplication. Matrix multiplication is associative but not commutative, and it has an identity element I_n (the identity matrix). But it does not have multiplicative inverses for all non-zero elements, so it is not a field.

Definition 4.16 (Vector Space over a Field K):

A vector space V over a field $(K, +, \cdot)$ is a set equipped with two operations $\oplus : V \times V \rightarrow V$ (vector addition) and $\odot : K \times V \rightarrow V$ (scalar multiplication) such that

- $(V, \oplus)$ is an abelian group with identity element $\mathbf{0}$ (the zero vector).
- The scalar multiplication satisfies the following properties
 - (i) Let 1 be the multiplicative identity of the field K . Then for all $\mathbf{v} \in V : 1 \odot \mathbf{v} = \mathbf{v}$.
 - (ii) $\forall a, b \in K : \forall \mathbf{v} \in V : a \odot (b \odot \mathbf{v}) = (a \cdot b) \odot \mathbf{v}$.
 - (iii) $\forall a \in K : \forall \mathbf{u}, \mathbf{v} \in V : a \odot (\mathbf{u} \oplus \mathbf{v}) = a \odot \mathbf{u} \oplus a \odot \mathbf{v}$.
 - (iv) $\forall a, b \in K : \forall \mathbf{v} \in V : (a + b) \odot \mathbf{v} = a \odot \mathbf{v} \oplus b \odot \mathbf{v}$.

Definition 4.17 ((Vector) Subspace):

A subset $U \subseteq V$ of a vector space V over a field K is called a subspace if it is closed under the vector addition and scalar multiplication, i.e.

$$\forall \mathbf{u}, \mathbf{v} \in U : \mathbf{u} \oplus \mathbf{v} \in U \quad \text{and} \quad \forall a \in K : \forall \mathbf{u} \in U : a \odot \mathbf{u} \in U. \quad (4.9)$$

Since we are already comfortable with vector spaces, we drop the special notation for vector addition and scalar multiplication and use the usual $+$ and $\cdot$ for these operations.

Now, continuing with our usual structure, now we will define structure preserving maps between vector spaces i.e. *linear maps*.

Definition 4.18 (Linear Map):

Let $(V, \oplus, \odot)$ and $(W, \boxplus, \boxdot)$ be two vector spaces over the same field K . A map $L : V \rightarrow W$ is called a linear map if it satisfies the following properties:

- (i) $\forall \mathbf{u}, \mathbf{v} \in V : L(\mathbf{u} \oplus \mathbf{v}) = L(\mathbf{u}) \boxplus L(\mathbf{v})$ (preserves vector addition).
- (ii) $\forall a \in K : \forall \mathbf{v} \in V : L(a \odot \mathbf{v}) = a \boxdot L(\mathbf{v})$ (preserves scalar multiplication).

If a linear map $L : V \rightarrow W$ is bijective, it is called a linear isomorphism. We write $V \cong_{\text{vec}} W$ if there exists a linear isomorphism between V and W .

We can compress the definition of a linear map into a single equation:

$$\forall \mathbf{v}_1, \mathbf{v}_2 \in V, \forall a, b \in K : L(a \odot \mathbf{v}_1 \oplus b \odot \mathbf{v}_2) = a \boxplus L(\mathbf{v}_1) \boxplus b \boxplus L(\mathbf{v}_2). \quad (4.10)$$

Remark 4.19 (Inverse being a linear map). As in case of topological spaces, we need to check that the inverse of a continuous map is continuous, here linearity of the inverse map follows from the linearity of the original map. Thus, it is enough to check that a linear map $L : V \rightarrow W$ is bijective to conclude that its inverse $L^{-1} : W \rightarrow V$ is also a linear map.

Pf: Let $L : V \rightarrow W$ be a linear map which is bijective. We need to show that $L^{-1} : W \rightarrow V$ is linear.

(i) Since L is bijective, there exists a unique $\mathbf{v} \in V$ such that $L(\mathbf{v}) = \mathbf{w}$ for any $\mathbf{w} \in W$. Thus, $L^{-1}(\mathbf{w}) = \mathbf{v}$.

(ii) For any $\mathbf{w}_1, \mathbf{w}_2 \in W$, we have

$$L^{-1}(\mathbf{w}_1 + \mathbf{w}_2) = L^{-1}(L(\mathbf{v}_1) + L(\mathbf{v}_2)) = L^{-1}(L(\mathbf{v}_1 + \mathbf{v}_2)) = \mathbf{v}_1 + \mathbf{v}_2 = L^{-1}(\mathbf{w}_1) + L^{-1}(\mathbf{w}_2),$$

where we used the linearity of L and the fact that L^{-1} is the inverse of L .

(iii) For any $a \in K$ and $\mathbf{w} \in W$, we have

$$L^{-1}(a \cdot \mathbf{w}) = L^{-1}(a \cdot L(\mathbf{v})) = L^{-1}(L(a \cdot \mathbf{v})) = a \cdot \mathbf{v} = a \cdot L^{-1}(\mathbf{w}),$$

again using the linearity of L .

Thus, L^{-1} is a linear map. □

Let's consider the set of all linear maps from V to W , denoted by $\text{Hom}(V, W)$.

$$\text{Hom}(V, W) := \{L : V \rightarrow W \mid L \text{ is a linear map}\} \equiv \{L : V \xrightarrow{\sim} W\} \quad (4.11)$$

here $\xrightarrow{\sim}$ denotes that the map is a linear map.

Proposition 4.20

$\text{Hom}(V, W)$ is a vector space over the field K with the following operations:

- $\boxplus : \text{Hom}(V, W) \times \text{Hom}(V, W) \rightarrow \text{Hom}(V, W)$ defined by

$$(L_1, L_2) \mapsto L_1 \boxplus L_2 \quad (4.12)$$

where

$$L_1 \boxplus L_2 : V \xrightarrow{\sim} W, \quad \mathbf{v} \mapsto (L_1 \boxplus L_2)(\mathbf{v}) := L_1(\mathbf{v}) \boxplus L_2(\mathbf{v}). \quad (4.13)$$

- $\boxtimes : K \times \text{Hom}(V, W) \rightarrow \text{Hom}(V, W)$ defined by

$$(a, L) \mapsto a \boxtimes L \quad (4.14)$$

where

$$a \boxtimes L : V \xrightarrow{\sim} W, \quad \mathbf{v} \mapsto (a \boxtimes L)(\mathbf{v}) := a \boxplus L(\mathbf{v}). \quad (4.15)$$

To establish this, we need to verify that the operations defined above satisfy the linear map properties:

Proof:

We need to show that $\text{Hom}(V, W)$ is a vector space over K with the operations $\boxplus$ and $\boxtimes$.

(i) Closure under addition: Let $L_1, L_2 \in \text{Hom}(V, W)$. Then $L_1 \boxplus L_2$ is defined as

$$(L_1 \boxplus L_2)(\mathbf{v}) = L_1(\mathbf{v}) \boxplus L_2(\mathbf{v}). \quad (4.16)$$

So for all $\mathbf{v}_1, \mathbf{v}_2 \in V$ and $a, b \in K$,

$$(L_1 \boxplus L_2)(a \odot \mathbf{v}_1 \oplus b \odot \mathbf{v}_2) = L_1(a \odot \mathbf{v}_1 \oplus b \odot \mathbf{v}_2) \boxplus L_2(a \odot \mathbf{v}_1 \oplus b \odot \mathbf{v}_2)$$

$$\begin{aligned}
&= (a \boxplus L_1(\mathbf{v}_1) \boxplus b \boxplus L_1(\mathbf{v}_2)) \boxplus (a \boxplus L_2(\mathbf{v}_1) \boxplus b \boxplus L_2(\mathbf{v}_2)) \\
&= a \boxplus (L_1(\mathbf{v}_1) \boxplus L_2(\mathbf{v}_1)) \boxplus b \boxplus (L_1(\mathbf{v}_2) \boxplus L_2(\mathbf{v}_2)) \\
&= a \boxplus (L_1 \boxplus L_2)(\mathbf{v}_1) \boxplus b \boxplus (L_1 \boxplus L_2)(\mathbf{v}_2).
\end{aligned}$$

Thus, $L_1 \boxplus L_2$ is a linear map.

(ii) Closure under scalar multiplication: Let $a \in K$ and $L \in \text{Hom}(V, W)$. Then $a \boxtimes L$ is defined as

$$(a \boxtimes L)(\mathbf{v}) = a \boxplus L(\mathbf{v}). \quad (4.17)$$

So for all $\mathbf{v}_1, \mathbf{v}_2 \in V$ and $b \in K$,

$$\begin{aligned}
(a \boxtimes L)(b \odot \mathbf{v}_1 \oplus c \odot \mathbf{v}_2) &= a \boxplus L(b \odot \mathbf{v}_1 \oplus c \odot \mathbf{v}_2) \\
&= a \boxplus (b \boxplus L(\mathbf{v}_1) \boxplus c \boxplus L(\mathbf{v}_2)) \\
&= (a \cdot b) \boxplus L(\mathbf{v}_1) \boxplus (a \cdot c) \boxplus L(\mathbf{v}_2) \\
&= (b \cdot a) \boxplus L(\mathbf{v}_1) \boxplus (c \cdot a) \boxplus L(\mathbf{v}_2) \\
&= b \boxplus (a \boxplus L(\mathbf{v}_1)) \boxplus c \boxplus (a \boxplus L(\mathbf{v}_2)) \\
&= b \boxplus (a \boxtimes L)(\mathbf{v}_1) \boxplus c \boxplus (a \boxtimes L)(\mathbf{v}_2).
\end{aligned}$$

Thus, $a \boxtimes L$ is a linear map.

We have shown that $\text{Hom}(V, W)$ is closed under both operations $\boxplus$ and $\boxtimes$. Thus, $\text{Hom}(V, W)$ is a vector space over the field K . Q.E.D.

Till now, we haven't used the inverse element of the field K in the definition of a vector space. So, we can define a similar structure on a set M over a (unital) ring R with two operations $\oplus : M \times M \rightarrow M$ (addition) and $\odot : R \times M \rightarrow M$ (scalar multiplication), we call this a *module* over the ring R .

Remark 4.21 (Case of $\text{Hom}(V, W)$ as a module). In case of modules, we can still define $\text{Hom}(V, W)$ as a set of all linear maps from V to W over a ring R . But as in general, ring multiplication is not commutative, $\text{Hom}(V, W)$ is not a module over R .

Some useful terminology:

- **Endomorphism** is a linear map $L : V \rightarrow V$ from a vector space V to itself.

$$\text{End}(V) := \text{Hom}(V, V) = \{L : V \xrightarrow{\sim} V\} \quad (4.18)$$

- **Automorphism** is a linear isomorphism $L : V \rightarrow V$ from a vector space V to itself.

$$\text{Aut}(V) := \{L : V \xrightarrow{\sim} V \mid L \text{ is a linear isomorphism}\}. \quad (4.19)$$

It is easy to see that $\text{Aut}(V)$ is a subspace of $\text{End}(V)$.

- A field K can be considered as a vector space over itself, *i.e.* K is a vector space over K with the usual addition and multiplication operations. With this view, we can define the notion of *linear functionals* as linear maps from V to K :

$$V^* := \text{Hom}(V, K) \quad (4.20)$$

This set V^* is called the *dual space* of V .

§4.2.1 Tensors

Definition 4.22 (Tensor):

A type (p, q) -tensor is a multilinear map (linear in each argument) of the form

$$T : \underbrace{V^* \times \cdots \times V^*}_{p \text{ times}} \times \underbrace{V \times \cdots \times V}_{q \text{ times}} \rightarrow K \quad (4.21)$$

where V is a vector space over a field K and V^* is its dual space.

$$\mathbb{T}_q^p V := \underbrace{V \otimes \cdots \otimes V}_{p \text{ times}} \otimes \underbrace{V^* \otimes \cdots \otimes V^*}_{q \text{ times}} = \{T \mid T \text{ is type } (p, q)\text{-tensor}\} \quad (4.22)$$

Let's define the operations on tensors similar to the operations we defined on the set of linear maps $\text{Hom}(V, W)$:

1. Addition of tensors:

$$\oplus : \mathbb{T}_q^p V \times \mathbb{T}_q^p V \rightarrow \mathbb{T}_q^p V \quad (4.23)$$

Let $T_1, T_2 \in \mathbb{T}_q^p V$ be two tensors. Then their sum $T_1 \oplus T_2$ is defined as

$$(T_1 \oplus T_2)(\omega_1, \dots, \omega_p, \mathbf{v}_1, \dots, \mathbf{v}_q) = T_1(\omega_1, \dots, \omega_p, \mathbf{v}_1, \dots, \mathbf{v}_q) + T_2(\omega_1, \dots, \omega_p, \mathbf{v}_1, \dots, \mathbf{v}_q) \quad (4.24)$$

2. Scalar multiplication of tensors:

$$\odot : K \times \mathbb{T}_q^p V \rightarrow \mathbb{T}_q^p V \quad (4.25)$$

Let $a \in K$ and $T \in \mathbb{T}_q^p V$ be a tensor. Then the scalar multiplication $a \odot T$ is defined as

$$(a \odot T)(\omega_1, \dots, \omega_p, \mathbf{v}_1, \dots, \mathbf{v}_q) = a \cdot T(\omega_1, \dots, \omega_p, \mathbf{v}_1, \dots, \mathbf{v}_q) \quad (4.26)$$

Proposition 4.23

$\mathbb{T}_q^p V$ is a vector space over the field K with the operations defined above.

Sometimes, we refer (p, q) as the valence of the tensor, and $p + q$ as the rank of the tensor.

Now, we define a binary operation on tensors, called the *tensor product* of two tensors.

Definition 4.24 (Tensor Product):

Let $T_1 \in \mathbb{T}_q^p V$ and $T_2 \in \mathbb{T}_s^r V$ be two tensors. The tensor product $T_1 \otimes T_2$ is defined as a tensor of type $(p + r, q + s)$ given by

$$(T_1 \otimes T_2)(\omega_1, \dots, \omega_{p+r}, \mathbf{v}_1, \dots, \mathbf{v}_{q+s}) = T_1(\omega_1, \dots, \omega_p, \mathbf{v}_1, \dots, \mathbf{v}_q) \cdot T_2(\omega_{p+1}, \dots, \omega_{p+r}, \mathbf{v}_{q+1}, \dots, \mathbf{v}_{q+s}). \quad (4.27)$$

Example 4.25

Now, let's look at some examples and interesting results involving tensors:

- (a) The set $\mathbb{T}_0^0 V = K$ is the set of all scalar tensors, which is just the field K itself.
- (b) The set $\mathbb{T}_1^0 V = V^* := \text{Hom}(V, K)$.
- (c) The set $\mathbb{T}_1^1 V \equiv V \otimes V^* = \{T \mid T : V^* \times V \rightarrow K \text{ is a linear in both argument}\} \cong_{\text{vec}} \text{End}(V^*)$.

Pf: Let $T \in \mathbb{T}_1^1 V$ be a tensor. We need to construct a linear map $L \in \text{End}(V^*)$ using T . For each $\omega \in V^*$, we define a linear map $L_\omega : V \rightarrow K \in V^*$ by

$$\forall \mathbf{v} \in V : L_\omega(\mathbf{v}) = T(\omega, \mathbf{v}). \quad (4.28)$$

Now, we can define a linear map $L : V^* \rightarrow V^*$ by

$$L(\omega) = L_\omega \quad \Rightarrow \quad \omega \mapsto \underbrace{(\mathbf{v} \mapsto T(\omega, \mathbf{v}))}_{L_\omega}. \quad (4.29)$$

Now, let $L \in \text{End}(V^*)$ be a linear map. We need to show that there exists a tensor $T \in \mathbb{T}_1^1 V$ corresponding to L . Define $T : V^* \times V \rightarrow K$ by

$$T(\omega, \mathbf{v}) = L(\omega)(\mathbf{v}) \quad (4.30)$$

as $L(\omega) \in V^*$ is a linear map from V to K . $\square$

- (d) From previous example, we expect that $\mathbb{T}_0^1 V \cong_{\text{vec}} V$; but this is not true in general. However, if V is finite-dimensional, then $\mathbb{T}_0^1 V \cong_{\text{vec}} V$.
- (e) Similarly, $\mathbb{T}_1^1 V \cong_{\text{vec}} \text{End}(V)$ is also not true in general, but if V is finite-dimensional, then $\mathbb{T}_1^1 V \cong_{\text{vec}} \text{End}(V)$.
- (f) All these examples which are not true in general, but true for finite-dimensional vector spaces, can be summarized as

$$(V^*)^* \not\cong_{\text{vec}} V. \quad (4.31)$$

This is true only for finite-dimensional vector spaces.

§4.2.2 Dimension of a Vector Space

We have mentioned dimension of a vector space multiple times, but we have not defined it properly yet. To define the dimension of a vector space, we first need to define *basis* of a vector space, and a natural choice is to use *Hamel basis* as it doesn't require any additional structure on the vector space.

Definition 4.26 (Hamel Basis):

Let $(V, +, \cdot)$ be a vector space over a field K . A subset $\mathcal{B} \subseteq V$ is called a Hamel basis if it satisfies the following properties:

- (i) $\mathcal{B}$ is a linearly independent set, i.e. for any finite subset $\{\mathbf{b}_1, \dots, \mathbf{b}_N\} \subseteq \mathcal{B}$, the only solution to the following equation is the trivial solution $\lambda^i = 0$ for all $i = 1, \dots, N$:

$$\sum_{i=1}^N \lambda^i \cdot \mathbf{b}_i = \mathbf{0} \quad \text{for } \lambda^i \in K \quad (4.32)$$

- (ii) $\mathcal{B}$ is a spanning set, i.e. every element $\mathbf{v} \in V$ can be expressed as a finite linear combination of elements from $\mathcal{B}$, i.e.

$$\forall \mathbf{v} \in V : \exists M \in \mathbb{N} : \exists v^1, \dots, v^M \in K : \exists \mathbf{b}_1, \dots, \mathbf{b}_M \in \mathcal{B} : \mathbf{v} = \sum_{i=1}^M v^i \cdot \mathbf{b}_i. \quad (4.33)$$

Say we all have decided to use a single Hamel basis $\mathcal{B}$ for our vector space V . Then it is more convenient to talk about the elements of V as an array of coefficients i.e. $(v^1, \dots, v^M)$ also called the *coordinates* of the vector with respect to the Hamel basis $\mathcal{B}$.

With this definition, we can define the dimension of a vector space. But first, we need to ensure that the dimension is well-defined, i.e. it does not depend on the choice of Hamel basis. This is guaranteed by the following proposition.

Proposition 4.27

Let $(V, +, \cdot)$ be a vector space over a field K and let $\mathcal{B}$ be a Hamel basis of V . Then $\mathcal{B}$ is the minimal spanning set of V , and maximal linearly independent set of V . In other words, let $S \subseteq V$.

- (i) If $\text{span}(S) = V$, then $|S| \geq |\mathcal{B}|$.

(ii) If S is linearly independent, then $|S| \leq |\mathcal{B}|$.

Definition 4.28 (Dimension):

The dimension of a vector space V over a field K is defined as the cardinality of any Hamel basis of V . We denote the dimension of V by $\dim V$.

$$\dim V := |\mathcal{B}| \quad \text{for any Hamel basis } \mathcal{B} \subseteq V. \quad (4.34)$$

If V has a finite Hamel basis, we say that V is finite-dimensional and $\dim V < \infty$. If V does not have a finite Hamel basis, we say that V is infinite-dimensional and $\dim V = \infty$.

Now, we have proper tools to prove following theorem.

Theorem 4.29

Let V be a finite-dimensional vector space. Then

$$(V^*)^* \cong_{\text{vec}} V. \quad (4.35)$$

Proof:

Let V be an n -dimensional vector space over a field K . Define a linear map $L : V \rightarrow (V^*)^*$ by

$$\begin{aligned} L : V &\rightarrow (V^*)^* \\ \mathbf{v} &\mapsto L_{\mathbf{v}} \end{aligned} \quad (4.36)$$

where $L_{\mathbf{v}} : V^* \rightarrow K \in (V^*)^*$ is defined by

$$\begin{aligned} L_{\mathbf{v}} : V^* &\rightarrow K \\ \omega &\mapsto L_{\mathbf{v}}(\omega) := \omega(\mathbf{v}) \end{aligned} \quad (4.37)$$

We need to show that L is a linear isomorphism.

(i) Linearity of L :

For any $\mathbf{v}_1, \mathbf{v}_2 \in V$ and $a, b \in K$,

$$\begin{aligned} L(a\mathbf{v}_1 + b\mathbf{v}_2)(\omega) &= L_{\{a\mathbf{v}_1 + b\mathbf{v}_2\}}(\omega) \\ &= \omega(a\mathbf{v}_1 + b\mathbf{v}_2) \\ &= a \cdot \omega(\mathbf{v}_1) + b \cdot \omega(\mathbf{v}_2) \\ &= a \cdot L_{\mathbf{v}_1}(\omega) + b \cdot L_{\mathbf{v}_2}(\omega) \\ &= (aL_{\mathbf{v}_1} + bL_{\mathbf{v}_2})(\omega). \end{aligned}$$

Thus, $L(a\mathbf{v}_1 + b\mathbf{v}_2) = aL_{\mathbf{v}_1} + bL_{\mathbf{v}_2}$, which shows that L is linear.

(ii) Injectivity of L :

Let $\mathbf{v} \neq \mathbf{0} \in V$. We need to show that $L(\mathbf{v}) \neq 0 \in (V^*)^*$. Since $\mathbf{v} \neq \mathbf{0}$, so $\text{span}(\mathbf{v}) \subseteq V$ is a proper subspace of V , so we can extend this to a Hamel basis $\mathcal{B} = \{\mathbf{v}, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ of V . Thus, any $\mathbf{u} \in V$ can be expressed as

$$\mathbf{u} = b \cdot \mathbf{v} + \sum_{i=2}^n b^i \cdot \mathbf{e}_i \quad (4.38)$$

for some $b, b^2, \dots, b^n \in K$. Now, we can define a linear functional $\omega \in V^*$ as follows:

$$\mathbf{u} \mapsto \omega(\mathbf{u}) := b. \quad (4.39)$$

Thus, we have

$$L(\mathbf{v})(\omega) = \omega(\mathbf{v}) = 1 \neq 0. \quad (4.40)$$

This shows that $L(\mathbf{v}) \neq 0$, hence L is injective.

Claim (Dual Basis): For a finite-dimensional vector space V , with a Hamel basis $\mathcal{B} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$, there exists a unique dual basis $\mathcal{B}^* = \{\mathbf{f}^1, \dots, \mathbf{f}^n\}$ such that

$$\forall i, j \in \{1, \dots, n\} : \mathbf{f}^i(\mathbf{e}_j) = \delta_j^i := \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases} \quad (4.41)$$

(iii) Surjectivity of L :

Let $F \in (V^*)^*$ be any linear functional. We need to show that there exists a $\mathbf{v} \in V$ such that $L(\mathbf{v}) = F$. Since V is finite-dimensional, we can choose a Hamel basis $\mathcal{B} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ of V . Now, we can define a vector $\mathbf{v} \in V$ as follows:

$$\mathbf{v} = \sum_{i=1}^n F(\mathbf{f}^i) \cdot \mathbf{e}_i \quad (4.42)$$

where $\mathbf{f}^i$ is the i -th element of the dual basis $\mathcal{B}^* = \{\mathbf{f}^1, \dots, \mathbf{f}^n\}$ corresponding to the Hamel basis $\mathcal{B}$. Now, we can check that

$$\begin{aligned} L(\mathbf{v})(\omega) &= \sum_{i=1}^n F(\mathbf{f}^i) \cdot \omega(\mathbf{e}_i) \\ &= \sum_{i=1}^n F(\mathbf{f}^i) \cdot \delta_i^j \\ &= F(\omega). \end{aligned}$$

Thus, $L(\mathbf{v}) = F$, which shows that L is surjective.

Q.E.D.

Corollary 4.30

If V is a finite-dimensional vector space, then

$$(i) \quad \mathbb{T}_1^1 V \cong_{\text{vec}} \text{End}(V).$$

$$(ii) \quad \mathbb{T}_0^1 V \cong_{\text{vec}} V.$$

Till now, we have only used basis to classify vector spaces, either finite-dimensional or infinite-dimensional. In vector theory, we don't use basis to construct objects, but once we have the definition of the object without using basis, we can use basis to study the object in a more convenient way.

We have seen the components of a vector with respect to a basis, now we can define the components of a tensor with respect to a basis.

Definition 4.31 (Components of a Tensor):

Let V be a finite-dimensional vector space over a field K with a Hamel basis $\mathcal{B} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ and the corresponding dual basis $\mathcal{B}^* = \{\epsilon^1, \dots, \epsilon^n\}$. The components of a tensor $T \in \mathbb{T}_q^p V$ with respect to the basis $\mathcal{B}$ and $\mathcal{B}^*$ are defined as follows:

$$T^{i_1 \dots i_p}_{j_1 \dots j_q} := T(\epsilon^{i_1}, \dots, \epsilon^{i_p}, \mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_q}) \quad \text{for } i_k, j_l \in \{1, \dots, n\}. \quad (4.43)$$

With these tensor components, we can express the tensor T as a sum of its components multiplied by the basis 'tensors':

$$T = \underbrace{\sum_{i_1=1}^n \dots \sum_{j_q=1}^n}_{p+q \text{ sums}} T^{i_1 \dots i_p}_{j_1 \dots j_q} \cdot \mathbf{e}_{i_1} \otimes \dots \otimes \mathbf{e}_{i_p} \otimes \epsilon^{j_1} \otimes \dots \otimes \epsilon^{j_q}. \quad (4.44)$$

Here $\otimes$ denotes the tensor product of tensors, as $\mathbf{e}_{i_k} \in V \cong_{\text{vec}} T_0^1 V$ and $\epsilon^{j_l} \in V^* \cong_{\text{vec}} T_1^0 V$. The components are field elements, so the implicit multiplication is the scalar multiplication of tensors of type (p, q) .

From now on, we will do this big sums multiple times, so to reduce the clutter, we will use the Einstein summation convention. In this convention, we will omit the summation symbol and assume that repeated indices (one upper and one lower) are summed over. For example, the above expression can be written as

$$T = \underbrace{\sum_{i_1=1}^n \cdots \sum_{j_q=1}^n}_{p+q \text{ sums}} T^{i_1 \dots i_p}_{j_1 \dots j_q} \cdot \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_p} \otimes \epsilon^{j_1} \otimes \cdots \otimes \epsilon^{j_q} \equiv T^{i_1 \dots i_p}_{j_1 \dots j_q} \cdot \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_p} \otimes \epsilon^{j_1} \otimes \cdots \otimes \epsilon^{j_q}. \quad (4.45)$$

For Up-Down convection, we only need to remember the fundamental convention *i.e.*

- Basis vectors of V are denoted by lower indices, $\mathbf{e}_i$.
- Basis vectors of V^* are denoted by upper indices, ϵ^i .

So the indices of the components of a tensor are due to the basis vectors of V and V^* used to define it.

$$T^{i_1 \dots i_p}_{j_1 \dots j_q} := T(\epsilon^{i_1}, \dots, \epsilon^{i_p}, \mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_q}). \quad (4.46)$$

We can recover the indices for the coordinates of a vector and a covector as follows (given a choice of basis and dual basis)

$$\mathbf{v} = v^i \cdot \mathbf{e}_i \quad \text{where } v^i = \epsilon^i(\mathbf{v}) \quad (4.47)$$

$$\omega = \omega_j \cdot \epsilon^j \quad \text{where } \omega_j = \mathbf{e}_j(\omega). \quad (4.48)$$

Remark 4.32 (Caution). But we need to be careful about this convention, as it is only works for linear spaces and linear maps. For example, if we have a tensor $T : V^* \times V \rightarrow K$, then for any $\omega \in V^*$ and $\mathbf{v} \in V$, we have

$$\begin{aligned} T(\omega, \mathbf{v}) &= T\left(\sum_{i=1}^n \omega_i \epsilon^i, \sum_{j=1}^n v^j \mathbf{e}_j\right) \\ &= \sum_{i=1}^n \sum_{j=1}^n T(\omega_i \epsilon^i, v^j \mathbf{e}_j) \\ &= \sum_{i=1}^n \sum_{j=1}^n \omega_i v^j T(\epsilon^i, \mathbf{e}_j) \\ &= \sum_{i=1}^n \sum_{j=1}^n \omega_i v^j T^i_j. \end{aligned}$$

So, if we omit the summation symbol, then there is no way to distinguish between the first and second equation. But the second equation is only true for bilinear maps, and not for non-bilinear maps. So, we need to be careful about the context in which we are using the Einstein summation convention.

It is important to know that, how to go from one basis to another basis. This is called the *change of basis*.

§4.2.3 Change of Basis

Consider two bases $\mathcal{B} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ and $\tilde{\mathcal{B}} = \{\tilde{\mathbf{e}}_1, \dots, \tilde{\mathbf{e}}_n\}$ of a vector space V . Since each vector in $\tilde{\mathcal{B}}$ is a vector in V , we can express each $\tilde{\mathbf{e}}_i$ as a linear combination of the vectors in $\mathcal{B}$: (using the Einstein summation convention)

$$\tilde{\mathbf{e}}_i = A^j_i \mathbf{e}_j \quad (4.49)$$

where $A^j_i \in K$. Similarly, we can express each $\mathbf{e}_i$ as a linear combination of the vectors in $\tilde{\mathcal{B}}$:

$$\mathbf{e}_i = B^j_i \tilde{\mathbf{e}}_j \quad (4.50)$$

where $B^j_i \in K$. In linear algebra, we can define two matrices A and B with entries A^j_i and B^j_i respectively, such that

$$A^{-1} = B \quad \text{and} \quad B^{-1} = A. \quad (4.51)$$

But till now, we haven't explored the vector spaces using arrays, so we define following notation for using usual notion of matrices and (row and column) vectors from our linear algebra course.

Remark 4.33 (Matrix Notation). Fix a basis $\mathcal{B} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ of a vector space V and a dual basis $\mathcal{B}^* = \{\epsilon^1, \dots, \epsilon^n\}$. Then we will express

$$\mathbf{v} = v^i \mathbf{e}_i \quad \rightsquigarrow \quad \mathbf{v} \hat{=} \begin{pmatrix} v^1 \\ \vdots \\ v^n \end{pmatrix} \quad (4.52)$$

and

$$\omega = \omega_j \epsilon^j \quad \rightsquigarrow \quad \omega^\top \hat{=} (\omega_1 \quad \dots \quad \omega_n). \quad (4.53)$$

Now let $\phi \in \text{End}(V) \cong_{\text{vec}} \mathbb{T}_1^1 V$ be a linear map. Then we can express ϕ as

$$\phi = \phi^i_j \mathbf{e}_i \otimes \epsilon^j \quad \rightsquigarrow \quad \Phi \hat{=} \begin{pmatrix} \phi^1_1 & \dots & \phi^1_n \\ \vdots & \ddots & \vdots \\ \phi^n_1 & \dots & \phi^n_n \end{pmatrix}. \quad (4.54)$$

So in general, the convention is that we will use the upper indices for the rows and lower indices for the columns of a matrix.

We know for finite-dimensional vector space V , we have $\text{End}(V) \cong_{\text{vec}} \mathbb{T}_1^1 V$. Let $\phi \in \text{End}(V)$, so we can rethink ϕ as a type $(1, 1)$ -tensor in following sense:

$$\phi(\omega, \mathbf{v}) := \omega(\phi(\mathbf{v})). \quad (4.55)$$

So the components of tensor ϕ is defined as,

$$\phi^i_j := \phi(\epsilon^i, \mathbf{e}_j) = \epsilon^i(\phi(\mathbf{e}_j)). \quad (4.56)$$

Let $\phi, \psi \in \text{End}(V)$. Let's compute the components of linear map $\phi \circ \psi$

$$\begin{aligned} (\phi \circ \psi)^i_j &= \epsilon^i((\phi \circ \psi)(\mathbf{e}_j)) \\ &= \epsilon^i(\phi(\psi(\mathbf{e}_j))) \\ &= \epsilon^i(\phi(\psi^a_j \mathbf{e}_a)) \\ &= \epsilon^i(\psi^a_j \phi(\mathbf{e}_a)) \\ &= \epsilon^i(\psi^a_j \phi^b_a \mathbf{e}_b) \\ &= \psi^a_j \cdot \phi^b_a \epsilon^i(\mathbf{e}_b) \\ &= \psi^a_j \cdot \phi^b_a \cdot \delta^i_b \\ &= \psi^a_j \cdot \phi^i_a = \phi^i_a \cdot \psi^a_j. \end{aligned}$$

With this, we have a definition for matrix multiplication *i.e.* say Φ, Ψ are matrices corresponding to linear maps ϕ, ψ , so the matrix for the composed linear map $\phi \circ \psi$ is given as

$$\Phi \cdot \Psi \hat{=} \left(\phi^i_a \cdot \psi^a_j \right)_{i,j=1}^n \quad (4.57)$$

We call this rule of producing matrices “*matrix-multiplication*.”

Similarly, the action of a covector $\omega(\mathbf{v}) = \omega_m v^m$, and this can be thought of as matrix multiplication as $\omega^\top \cdot \mathbf{v}$. This notation gives us a false picture that these are basis independent, but they are not. And in the similar spirit we also write,

$$\phi(\omega, \mathbf{v}) = \omega_i \phi^i_j v^j \quad \rightsquigarrow \quad \omega^\top \cdot \Phi \cdot \mathbf{v} \quad (4.58)$$

There is another issue with these representations, *i.e.* for tensor of type other than these three, we don't have nice picture to draw. So it is better to think them as multilinear maps instead of matrices.

Now, let's talk about the effect of change of basis on the vectors, covectors, and tensors. Recall we have two basis, $\mathcal{B} = \{\mathbf{e}_1, \dots, \mathbf{e}_n\}$ and $\tilde{\mathcal{B}} = \{\tilde{\mathbf{e}}_1, \dots, \tilde{\mathbf{e}}_n\}$ of a vector space V . Then the change of basis for vectors is given by

$$\tilde{\mathbf{e}}_i = A^j_i \mathbf{e}_j \quad \mathbf{e}_j = B^i_j \tilde{\mathbf{e}}_i. \quad (4.59)$$

Since these relations are invertible, we have

$$A^i_m B^m_j = B^i_m A^m_j = \delta^i_j \quad (4.60)$$

For both the bases, we have the dual bases $\mathcal{B}^* = \{\epsilon^1, \dots, \epsilon^n\}$ and $\tilde{\mathcal{B}}^* = \{\tilde{\epsilon}^1, \dots, \tilde{\epsilon}^n\}$.

1. Covectors:

Let $\omega = \omega_j \epsilon^j$ be a covector, where $\omega_j = \mathbf{e}_j(\omega)$. Here $\mathbf{e}_j$ is an element of $T_0^1 V$. Then we have

$$\omega_j = \mathbf{e}_j(\omega) = \omega(\mathbf{e}_j) = \omega(B^i_j \tilde{\mathbf{e}}_i) = B^i_j \omega(\tilde{\mathbf{e}}_i) = B^i_j \tilde{\omega}_i(\omega) = B^i_j \tilde{\omega}_i. \quad (4.61)$$

2. Vectors:

Let $\mathbf{v} = v^i \mathbf{e}_i$ be a vector, where $v^i = \epsilon^i(\mathbf{v})$. Then we have

$$v^i = \epsilon^i(\mathbf{v}) = \epsilon^i(\tilde{v}^j \tilde{\mathbf{e}}_j) = \tilde{v}^j \epsilon^i(\tilde{\mathbf{e}}_j) = \tilde{v}^j \epsilon^i(A^k_j \mathbf{e}_k) = \tilde{v}^j A^k_j \epsilon^i(\mathbf{e}_k) = \tilde{v}^j A^k_j \delta^i_k = A^i_j \tilde{v}^j. \quad (4.62)$$

Summarizing the above two equations, we have

$$v^i = A^i_j \tilde{v}^j \quad \omega_j = B^i_j \tilde{\omega}_i \quad (4.63)$$

$$\tilde{v}^i = B^i_j v^j \quad \tilde{\omega}_j = A^i_j \omega_i. \quad (4.64)$$

Now, let's see how the basis of dual space changes with respect to the change of basis of the vector space. Recall that the dual basis is defined as follows

$$\epsilon^i(\mathbf{e}_j) = \delta^i_j \quad \tilde{\epsilon}^i(\tilde{\mathbf{e}}_j) = \delta^i_j. \quad (4.65)$$

Let $\tilde{\epsilon}^i = C^i_j \epsilon^j$, where $C^i_j \in K$. Then we have

$$\tilde{\epsilon}^i(\tilde{\mathbf{e}}_j) = C^i_k \epsilon^k(A^l_j \mathbf{e}_l) = C^i_k A^l_j \epsilon^k(\mathbf{e}_l) = C^i_k A^l_j \delta^k_l = C^i_k A^k_j = \delta^i_j. \quad (4.66)$$

Thus, using unique existence of the inverse, we have

$$C^i_j = B^i_j. \quad (4.67)$$

So we have the following relations for the dual basis:

$$\tilde{\epsilon}^i = B^i_j \epsilon^j \quad \epsilon^j = A^j_i \tilde{\epsilon}^i. \quad (4.68)$$

3. Tensors:

Let $T \in T_q^p V$ be a tensor, then the components of T with respect to the basis $\mathcal{B}$ and $\mathcal{B}^*$ are given by

$$T^{i_1 \dots i_p}_{j_1 \dots j_q} = T(\epsilon^{i_1}, \dots, \epsilon^{i_p}, \mathbf{e}_{j_1}, \dots, \mathbf{e}_{j_q}). \quad (4.69)$$

Now, we can express the components of T with respect to the basis $\tilde{\mathcal{B}}$ and $\tilde{\mathcal{B}}^*$ as follows:

$$\tilde{T}^{i_1 \dots i_p}_{j_1 \dots j_q} = T(\tilde{\epsilon}^{i_1}, \dots, \tilde{\epsilon}^{i_p}, \tilde{\mathbf{e}}_{j_1}, \dots, \tilde{\mathbf{e}}_{j_q}) \quad (4.70)$$

$$= T(B^{i_1}_{a_1} \epsilon^{a_1}, \dots, B^{i_p}_{a_p} \epsilon^{a_p}, A^{b_1}_{j_1} \mathbf{e}_{b_1}, \dots, A^{b_q}_{j_q} \mathbf{e}_{b_q}) \quad (4.71)$$

$$= B^{i_1}_{a_1} \dots B^{i_p}_{a_p} \cdot A^{b_1}_{j_1} \dots A^{b_q}_{j_q} \cdot T(\epsilon^{a_1}, \dots, \epsilon^{a_p}, \mathbf{e}_{b_1}, \dots, \mathbf{e}_{b_q}) \quad (4.72)$$

$$= B^{i_1}_{a_1} \dots B^{i_p}_{a_p} \cdot A^{b_1}_{j_1} \dots A^{b_q}_{j_q} \cdot T^{a_1 \dots a_p}_{b_1 \dots b_q}. \quad (4.73)$$

Thus, the reversal of the basis is given by

$$T^{i_1 \dots i_p}_{j_1 \dots j_q} = A^{i_1}_{a_1} \dots A^{i_p}_{a_p} \cdot B^{b_1}_{j_1} \dots B^{b_q}_{j_q} \cdot \tilde{T}^{a_1 \dots a_p}_{b_1 \dots b_q}. \quad (4.74)$$

§4.2.4 Determinants

From our previous knowledge of linear algebra, we know that the determinant is a scalar-valued function that takes a square matrix and returns a scalar. But since we know that the square matrix is just a convention for a linear map, we need to define the determinant in a more general way, independent of the basis. But first, look at this weird result which is purely due to the fact that we are using a witchcraft called *matrices* to represent linear maps.

Remark 4.34. Let $\phi \in T_1^1 V$ be an endomorphism and $g \in T_2^0 V$ be a bilinear form. We have

$$\phi = \phi_j^i \mathbf{e}_i \otimes \epsilon^j \quad \text{and} \quad g = g_{ij} \epsilon^i \otimes \epsilon^j. \quad (4.75)$$

We can arrange the components of ϕ and g in a matrix form as follows:

$$\Phi = \begin{pmatrix} \phi_1^1 & \cdots & \phi_1^n \\ \vdots & \ddots & \vdots \\ \phi_n^1 & \cdots & \phi_n^n \end{pmatrix} \quad \text{and} \quad G = \begin{pmatrix} g_{11} & \cdots & g_{1n} \\ \vdots & \ddots & \vdots \\ g_{n1} & \cdots & g_{nn} \end{pmatrix}. \quad (4.76)$$

But say, we change the basis of V to $\tilde{\mathcal{B}}$, then the components of ϕ and g change as follows:

$$\tilde{\phi}_j^i = B^i_k \cdot \phi_l^k \cdot A^l_j \quad \text{and} \quad \tilde{g}_{ij} = A^k_i \cdot g_{kl} \cdot A^l_j. \quad (4.77)$$

Recall $(A^{-1})^i_j = B^i_j$, so we can write the above equations as (remember that $A^i_j = A_{ij}$, which give rise to transpose in the expression)

$$\tilde{\Phi} = A^{-1} \cdot \Phi \cdot A \quad \text{and} \quad \tilde{G} = A^\top \cdot G \cdot A. \quad (4.78)$$

Now, we keep the old notion of determinant, so we can find determinant of G . But that is not true, as determinant is only defined for endomorphisms. And to see, why this the case, we have to look at basis free definition of determinants.

§4.2.4.1 Permutation Group

Before we define the determinant, we need a few definitions. We will use the *permutation group* to define the determinant.

Definition 4.35 (Permutation and Permutation Group):

A permutation of a non-empty set M is a bijective function $\sigma : M \rightarrow M$.

Let the set M be finite with n elements, for purpose of this course, we will assume $M = \{1, 2, \dots, n\}$. The set of all permutations of M is denoted by S_n and is called the permutation group of order n (or symmetric group of degree n). Binary operation on S_n is defined as the composition of two permutations.

There are some special type of permutations, which are called *transpositions*. A transposition is a permutation that swaps two elements of the set and leaves the rest unchanged. For example, in S_3 , the permutation $(1, 2, 3) \mapsto (2, 1, 3)$ is a transposition, usually denoted as $(1, 2)$.

There is a result from group theory, which states that every permutation can be expressed as a product of transpositions (precisely composition of transpositions). The number of transpositions in this product is called the *signature* or *sign* or *parity* of the permutation. Let $\sigma \in S_n$ be a permutation, then we define the sign of σ as follows:

$$\text{sgn}(\sigma) := \begin{cases} +1 & \text{if } \sigma \text{ is the product of an even number of transpositions} \\ -1 & \text{if } \sigma \text{ is the product of an odd number of transpositions} \end{cases} \quad (4.79)$$

But there is an issue with this definition, *i.e.* the decomposition of a permutation into transpositions is not unique. For example, the permutation $(1, 2, 3)$ can be expressed as $(1, 2)(2, 3)$ or $(1, 3)(1, 2)$. But we can still show that the sign of a permutation is well-defined, *i.e.* it does not depend on the decomposition

of the permutation into transpositions. Since the number of transpositions in the decomposition is either even or odd, and this property is preserved under composition of permutations.

Now we can define n -form for a vector space V of dimension d .

Definition 4.36 (n -form):

Let V be a finite-dimensional vector space of dimension d . An n -form on V (where $0 \leq n \leq d$) is a type $(0, n)$ -tensor $\omega \in \mathbf{T}_n^0 V$ such that it is totally antisymmetric, i.e. for any n vectors $\mathbf{v}_1, \dots, \mathbf{v}_n \in V$ and any permutation $\sigma \in S_n$, we have

$$\omega(\mathbf{v}_1, \dots, \mathbf{v}_n) = \text{sgn}(\sigma) \cdot \omega(\mathbf{v}_{\sigma(1)}, \dots, \mathbf{v}_{\sigma(n)}). \quad (4.80)$$

For $n = 0$, we define the 0-form to be the scalar field K itself, which is trivially totally antisymmetric.

Special case of n -form is the *top form* on V , which is for $n = d$.

Proposition 4.37

A $(0, n)$ -tensor ω is an n -form if and only if, $\omega(\mathbf{v}_1, \dots, \mathbf{v}_n) = 0$ whenever any $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is linearly dependent.

Since any collection of $n > d$ vectors in V is linearly dependent, we can conclude that any n -form (for $n > d$) is identically zero.

We have a special symbol for the space of n -forms on V , which is denoted by $\Lambda^n V$. With this we have following important result

Theorem 4.38 (Dimension of n -forms)

Let V be a finite-dimensional vector space of dimension d . Then the dimension of the space of n -forms on V is given by

$$\dim \Lambda^n V = \begin{cases} \binom{d}{n} & \text{if } 0 \leq n \leq d \\ 0 & \text{if } n > d \end{cases} \quad (4.81)$$

In particular, the dimension of the space of top forms on V is 1, i.e. $\dim \Lambda^d V = 1$. This means that for any two top forms $\omega_1, \omega_2 \in \Lambda^d V$, there exists a scalar $\lambda \in K$ such that $\omega_1 = \lambda \cdot \omega_2$.

With this, we define *volume form* on V as a non-zero top form $\omega \in \Lambda^d V$. And a vector space V with a specified volume form is called a *vector space with volume form*.

Definition 4.39 (Volume Span):

Let V be a d -dimensional vector space with a volume form ω . Given a set of d vectors $\{\mathbf{v}_1, \dots, \mathbf{v}_d\} \subseteq V$, the volume span of these vectors is defined as

$$\text{vol}(\mathbf{v}_1, \dots, \mathbf{v}_d) := \omega(\mathbf{v}_1, \dots, \mathbf{v}_d). \quad (4.82)$$

Due to the antisymmetry of the volume form, we have $\text{vol}(\mathbf{v}_1, \dots, \mathbf{v}_d) = 0$ if and only if the vectors are linearly dependent. Indeed, in this case, these vectors at most span a $(d-1)$ -dimensional hyperplane in V , which should have zero volume.

Now, we can define the determinant of a linear map $\phi \in \text{End}(V)$ as follows:

Definition 4.40 (Determinant):

Let V be a d -dimensional vector space. The determinant of a linear map $\phi \in \text{End}(V)$ is defined as

$$\det(\phi) := \frac{\omega(\phi(\mathbf{e}_1), \dots, \phi(\mathbf{e}_d))}{\omega(\mathbf{e}_1, \dots, \mathbf{e}_d)} \quad (4.83)$$

for some volume form ω on V and a basis $\{\mathbf{e}_1, \dots, \mathbf{e}_d\}$ of V .

First and foremost, we need to check that this definition is independent of the choice of volume form and basis. For any other volume form $\tilde{\omega}$ on V , we can write

$$\tilde{\omega} = \lambda \cdot \omega \quad \Rightarrow \quad \det(\phi) = \frac{\tilde{\omega}(\phi(\mathbf{e}_1), \dots, \phi(\mathbf{e}_d))}{\tilde{\omega}(\mathbf{e}_1, \dots, \mathbf{e}_d)} = \frac{\lambda \cdot \omega(\phi(\mathbf{e}_1), \dots, \phi(\mathbf{e}_d))}{\lambda \cdot \omega(\mathbf{e}_1, \dots, \mathbf{e}_d)}. \quad (4.84)$$

So, the determinant is independent of the choice of volume form. Now, let $\{\tilde{\mathbf{e}}_1, \dots, \tilde{\mathbf{e}}_d\}$ be another basis of V . Then we can write

$$\begin{aligned} \det(\phi) &= \frac{\omega(\phi(\tilde{\mathbf{e}}_1), \dots, \phi(\tilde{\mathbf{e}}_d))}{\omega(\tilde{\mathbf{e}}_1, \dots, \tilde{\mathbf{e}}_d)} \\ &= \frac{\omega(\phi(A^{i_1}_1 \mathbf{e}_{i_1}), \dots, \phi(A^{i_d}_d \mathbf{e}_{i_d}))}{\omega(A^{i_1}_1 \mathbf{e}_{i_1}, \dots, A^{i_d}_d \mathbf{e}_{i_d})} \\ &= \frac{A^{i_1}_1 \cdots A^{i_d}_d \cdot \omega(\phi(\mathbf{e}_{i_1}), \dots, \phi(\mathbf{e}_{i_d}))}{A^{i_1}_1 \cdots A^{i_d}_d \cdot \omega(\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_d})} \end{aligned}$$

say there is a permutation $\sigma_{i_1 \dots i_d}$ such that $\sigma_{i_1 \dots i_d}(i_1, \dots, i_d) = (1, \dots, d)$, then we can write

$$\begin{aligned} &= \frac{A^{i_1}_1 \cdots A^{i_d}_d \cdot \omega(\phi(\mathbf{e}_{\sigma_{i_1 \dots i_d}(1)}), \dots, \phi(\mathbf{e}_{\sigma_{i_1 \dots i_d}(d)}))}{A^{i_1}_1 \cdots A^{i_d}_d \cdot \omega(\mathbf{e}_{\sigma_{i_1 \dots i_d}(1)}, \dots, \mathbf{e}_{\sigma_{i_1 \dots i_d}(d)})} \\ &= \frac{\text{sgn}(\sigma_{i_1 \dots i_d}) \cdot A^{i_1}_1 \cdots A^{i_d}_d \cdot \omega(\phi(\mathbf{e}_1), \dots, \phi(\mathbf{e}_d))}{\text{sgn}(\sigma_{i_1 \dots i_d}) \cdot A^{i_1}_1 \cdots A^{i_d}_d \cdot \omega(\mathbf{e}_1, \dots, \mathbf{e}_d)} \end{aligned}$$

call $\text{sgn}(\sigma_{i_1 \dots i_d}) \cdot A^{i_1}_1 \cdots A^{i_d}_d = \lambda$, as all the indices are summed over, so we can write

$$= \frac{\lambda \cdot \omega(\phi(\mathbf{e}_1), \dots, \phi(\mathbf{e}_d))}{\lambda \cdot \omega(\mathbf{e}_1, \dots, \mathbf{e}_d)} = \det(\phi).$$

With this, we can conclude that the determinant is independent of the choice of basis. So we have

$$\det(\tilde{\Phi}) = \det(A^{-1} \cdot \Phi \cdot A) = \det(A^{-1}) \cdot \det(\Phi) \cdot \det(A) = \det(\Phi). \quad (4.85)$$

Recall the transformation of the bilinear form g under change of basis, we have

$$g \rightarrow A^\top g A. \quad (4.86)$$

So the determinant of the bilinear form g transforms as follows:

$$\det(A^\top g A) = \det(A^\top) \cdot \det(g) \cdot \det(A) = \det(A)^2 \cdot \det(g). \quad (4.87)$$

i.e. the determinant of a bilinear form transforms is not invariant under change of basis. But say there is another quantity, which transforms as follows:

$$X \rightarrow \frac{1}{\det(A)^2} X. \quad (4.88)$$

Then the quantity $\det(g) \cdot X$ is invariant under change of basis, *i.e.* it transforms as

$$\det(g) \cdot X \rightarrow \det(A)^2 \cdot \det(g) \cdot \frac{1}{\det(A)^2} X = \det(g) \cdot X. \quad (4.89)$$

Here, it seems like two wrongs make a right, but this is not the case. In order to make this mathematically precise, we will have to introduce principal fibre bundles. Using them, we will be able to give a bundle definition of tensor and of tensor densities which are, loosely speaking, quantities that transform with powers of $\det A$ under a change of basis.

§4.3 Tangent Spaces to a Manifold

Let M be a smooth manifold (from now on, whenever we say a smooth manifold, the associated topology and atlas are always implied). Then we can construct the following vector space over $\mathbb{R}$:

$$(\mathcal{C}^\infty(M), +, \cdot) \quad (4.90)$$

where $\mathcal{C}^\infty(M) := \{f : M \rightarrow \mathbb{R} \mid f \text{ is smooth}\}$ is the set of all smooth real-valued functions on M . The notion of smoothness is via smooth charts in the atlas of M . The addition and scalar multiplication are defined pointwise, for all $f, g \in \mathcal{C}^\infty(M)$, $\lambda \in \mathbb{R}$, and $p \in M$, as

$$(f + g)(p) := f(p) + g(p) \quad (4.91)$$

$$(\lambda \cdot f)(p) := \lambda \cdot f(p). \quad (4.92)$$

It is easy to check that $(\mathcal{C}^\infty(M), +, \cdot)$ is indeed a vector space over $\mathbb{R}$.

Definition 4.41 (Directional Derivative):

Let $\gamma : \mathbb{R} \rightarrow M$ be a smooth curve¹ through a point $p \in M$, and WLOG let $\gamma(0) = p$. Then the directional derivative operator along γ at p is a map

$$X_{\gamma,p} : \mathcal{C}^\infty(M) \rightarrow \mathbb{R} \quad (4.93)$$

defined as

$$\mathcal{C}^\infty(M) \ni f \mapsto X_{\gamma,p}(f) := (f \circ \gamma)'(0) \in \mathbb{R}. \quad (4.94)$$

Here the notion of smoothness is via charts in the atlas of M , i.e., for all charts (U, x) of M such that $p \in U$, the composition $x \circ \gamma : \mathbb{R} \rightarrow \mathbb{R}^d$ is a smooth map in the usual sense.

Note the composition $f \circ \gamma$ is a map from $\mathbb{R}$ to $\mathbb{R}$, and hence the derivative is the usual derivative of real-valued functions of a real variable.

Proposition 4.42

The directional derivative operator $X_{\gamma,p}$ along a smooth curve γ through p is a linear map, i.e., for all $f, g \in \mathcal{C}^\infty(M)$ and $\lambda, \mu \in \mathbb{R}$,

$$X_{\gamma,p}(\lambda f + \mu g) = \lambda X_{\gamma,p}(f) + \mu X_{\gamma,p}(g). \quad (4.95)$$

Proof:

This follows from the linearity of the usual derivative of real-valued functions of a real variable.

$$\begin{aligned} X_{\gamma,p}(\lambda f + \mu g) &= ((\lambda f + \mu g) \circ \gamma)'(0) \\ &= (\lambda(f \circ \gamma) + \mu(g \circ \gamma))'(0) \\ &= \lambda(f \circ \gamma)'(0) + \mu(g \circ \gamma)'(0) \\ &= \lambda X_{\gamma,p}(f) + \mu X_{\gamma,p}(g). \end{aligned}$$

Q.E.D.

In differential geometry, $X_{\gamma,p}$ is usually called a *tangent vector* to curve γ at point p . Physically, we can think of $X_{\gamma,p}$ as the velocity vector of a particle moving along the curve γ at point p . To see this, let $\gamma_1, \gamma_2 : \mathbb{R} \rightarrow M$ be two smooth curves through p such that $\gamma_1(0) = \gamma_2(0) = p$, and $\gamma_1(t) = \gamma_2(2t)$ for all $t \in \mathbb{R}$. Let $f \in \mathcal{C}^\infty(M)$ be a smooth function. Then

$$X_{\gamma_1,p}(f) = (f \circ \gamma_1)'(0) = (f \circ \gamma_2)'(0) \cdot 2 = 2(f \circ \gamma_2)'(0) = 2X_{\gamma_2,p}(f). \quad (4.96)$$

This means that $X_{\gamma_1,p} = 2X_{\gamma_2,p}$. So, if we think of γ_1 and γ_2 as the trajectories of two particles moving through point p on manifold M , then the velocity vector of the first particle at p is twice that of the second particle at p , which is consistent with our physical intuition.

Definition 4.43 (Tangent Vector Space):

Let M be a smooth manifold and $p \in M$. The tangent vector space to M at p , denoted by $\mathcal{T}_p M$, is defined as

$$\mathcal{T}_p M := \{X_{\gamma,p} \mid \gamma : \mathbb{R} \rightarrow M \text{ is a smooth curve with } \gamma(0) = p\}. \quad (4.97)$$

Equipped with following operations:

$$\oplus : \mathcal{T}_p M \times \mathcal{T}_p M \rightarrow \mathcal{T}_p M,$$

$$\odot : \mathbb{R} \times \mathcal{T}_p M \rightarrow \mathcal{T}_p M,$$

defined pointwise as

$$\begin{aligned} (X_{\gamma_1,p} \oplus X_{\gamma_2,p})(f) &:= X_{\gamma_1,p}(f) + X_{\gamma_2,p}(f), \quad \forall f \in \mathcal{C}^\infty(M), \\ (\lambda \odot X_{\gamma,p})(f) &:= \lambda \cdot X_{\gamma,p}(f), \quad \forall f \in \mathcal{C}^\infty(M) \text{ and } \lambda \in \mathbb{R}, \end{aligned}$$

$(\mathcal{T}_p M, \oplus, \odot)$ is a vector space over $\mathbb{R}$.

This definition is still incomplete, as the pointwise addition and scalar multiplication doesn't guarantee that the results are still in $\mathcal{T}_p M$. So, we have to prove the following proposition.

Proposition 4.44

The operations $\oplus$ and $\odot$ defined above are closed in $\mathcal{T}_p M$, i.e., for all $X_{\gamma_1,p}, X_{\gamma_2,p} \in \mathcal{T}_p M$ and $\lambda \in \mathbb{R}$,

$$X_{\gamma_1,p} \oplus X_{\gamma_2,p} \in \mathcal{T}_p M, \quad (4.98)$$

$$\lambda \odot X_{\gamma,p} \in \mathcal{T}_p M. \quad (4.99)$$

So we need to show that there exist smooth curves $\gamma_3, \gamma_4 : \mathbb{R} \rightarrow M$ such that $\gamma_3(0) = \gamma_4(0) = p$ and

$$\begin{aligned} X_{\gamma_3,p} &= X_{\gamma_1,p} \oplus X_{\gamma_2,p}, \\ X_{\gamma_4,p} &= \lambda \odot X_{\gamma,p}. \end{aligned}$$

Since the notion of derivative is local, so if two curves agree on a neighborhood of $0 \in \mathbb{R}$, then they have the same derivative at 0 i.e., if $\gamma_1(t) = \gamma_2(t)$ for all t in some open interval containing 0, then $X_{\gamma_1,p} = X_{\gamma_2,p}$. So, it is sufficient to construct γ_3 and γ_4 on some open interval containing 0.

Proof:

Let (U, x) be a chart of M around p , i.e., $p \in U$ and $x : U \rightarrow x(U) \subseteq \mathbb{R}^d$ is a homeomorphism.

Let $I \subseteq \mathbb{R}$ be an open interval containing 0 such that $\gamma(t), \gamma_1(t), \gamma_2(t) \in U$ for all $t \in I$. Such an interval exists since $\gamma, \gamma_1, \gamma_2$ are continuous and $\gamma(0) = \gamma_1(0) = \gamma_2(0) = p \in U$, and U is open in M .

1. Construct a curve $\gamma_3 : I \rightarrow M$ using the chart (U, x) as follows:

$$\gamma_3(t) := x^{-1} \circ (x \circ \gamma_1(t) + x \circ \gamma_2(t) - x(p)), \quad \forall t \in I. \quad (4.100)$$

Note that $x \circ \gamma_1(t), x \circ \gamma_2(t) \in \mathbb{R}^d$ and $x(p) \in \mathbb{R}^d$, so the addition and subtraction are well-defined. Also, since x is a diffeomorphism, $x \circ \gamma_1$ and $x \circ \gamma_2$ are smooth maps from I to $\mathbb{R}^d$, and hence their sum is also a smooth map from I to $\mathbb{R}^d$. Therefore, the composition γ_3 is a smooth map from I to M . Moreover, $\gamma_3(0) = x^{-1}(x(p) + x(p) - x(p)) = p$.

Now, for all $f \in \mathcal{C}^\infty(M)$,

$$\begin{aligned} X_{\gamma_3,p}(f) &= (f \circ \gamma_3)'(0) \\ &= \left(f \circ x^{-1} \circ (x \circ \gamma_1 + x \circ \gamma_2 - x(p)) \right)'(0) \end{aligned}$$

map $f \circ x^{-1} : x(U) \subseteq \mathbb{R}^d \rightarrow \mathbb{R}$, and $x \circ \gamma_1 + x \circ \gamma_2 - x(p) : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^d$, so we can apply the multivariate chain rule, taking derivative in j -th coordinate direction, $j = 1, \dots, d$:

$$= \left[\partial_j (f \circ x^{-1})(x(p)) \right] \cdot \left(x^j \circ \gamma_1 + x^j \circ \gamma_2 - x^j(p) \right)'(0)$$

where x^j is the j -th coordinate function of chart x . Here sum over j from 1 to d is implied. Using linearity of the usual derivative, we have:

$$\begin{aligned} &= \left[\partial_j (f \circ x^{-1})(x(p)) \right] \cdot \left((x^j \circ \gamma_1)'(0) + (x^j \circ \gamma_2)'(0) \right) \\ &= [\partial_j (f \circ x^{-1})(x(p)) \cdot (x^j \circ \gamma_1)'(0)] + [\partial_j (f \circ x^{-1})(x(p)) \cdot (x^j \circ \gamma_2)'(0)] \end{aligned}$$

combining the term in the square brackets, we get:

$$\begin{aligned} &= (f \circ \gamma_1)'(0) + (f \circ \gamma_2)'(0) \\ &= X_{\gamma_1,p}(f) + X_{\gamma_2,p}(f) \\ &= (X_{\gamma_1,p} \oplus X_{\gamma_2,p})(f). \end{aligned}$$

Since this is true for all $f \in \mathcal{C}^\infty(M)$, we have

$$X_{\gamma_3,p} = X_{\gamma_1,p} \oplus X_{\gamma_2,p}. \quad (4.101)$$

2. Construct a curve $\gamma_4 : I \rightarrow M$ using the chart (U, x) as follows:

$$\gamma_4(t) := x^{-1} \circ (x \circ \gamma(\lambda t)), \quad \forall t \in I. \quad (4.102)$$

Here it is tempting rewrite $\gamma_4(t) = \gamma(\lambda t)$, but this lead to a problem that how to define $f'(p)$ when we find $X_{\gamma_4,p}(f)$.

Note that since x is a diffeomorphism, $x \circ \gamma$ is a smooth map from I to $\mathbb{R}^d$, and hence the composition γ_4 is also a smooth map from I to M . Moreover, $\gamma_4(0) = x^{-1}(x(p)) = p$.

Now, for all $f \in \mathcal{C}^\infty(M)$,

$$\begin{aligned} X_{\gamma_4,p}(f) &= (f \circ \gamma_4)'(0) \\ &= \left(f \circ x^{-1} \circ (x \circ \gamma(\lambda t)) \right)'(0) \end{aligned}$$

map $f \circ x^{-1} : x(U) \subseteq \mathbb{R}^d \rightarrow \mathbb{R}$, and $x \circ \gamma(\lambda t) : I \subseteq \mathbb{R} \rightarrow \mathbb{R}^d$, so we can apply the multivariate chain rule, taking derivative in j -th coordinate direction, $j = 1, \dots, d$:

$$= \left[\partial_j (f \circ x^{-1})(x(p)) \right] \cdot \left(x^j \circ \gamma(\lambda t) \right)'(0)$$

where x^j is the j -th coordinate function of chart x . Using the chain rule for the usual derivative, we have:

$$\begin{aligned} &= \left[\partial_j (f \circ x^{-1})(x(p)) \right] \cdot \left[(x^j \circ \gamma)'(0) \cdot \lambda \right] \\ &= \lambda [\partial_j (f \circ x^{-1})(x(p)) \cdot (x^j \circ \gamma)'(0)] \end{aligned}$$

combining the term in the square brackets, we get:

$$\begin{aligned} &= \lambda (f \circ \gamma)'(0) \\ &= \lambda X_{\gamma,p}(f) \\ &= (\lambda \odot X_{\gamma,p})(f). \end{aligned}$$

Since this is true for all $f \in \mathcal{C}^\infty(M)$, we have

$$X_{\gamma_4,p} = \lambda \odot X_{\gamma,p}. \quad (4.103)$$

Q.E.D.

Remark 4.45 (Independence of Chart Choice). The construction of γ_3 and γ_4 depends on the choice of chart (U, x) . However, the resulting tangent vectors $X_{\gamma_3,p}$ and $X_{\gamma_4,p}$ do not depend on the choice of chart. This is because if we choose another chart (V, y) around p , then the transition map $y \circ x^{-1} : x(U \cap V) \rightarrow y(U \cap V)$ is a diffeomorphism between open subsets of $\mathbb{R}^d$, and hence the construction of γ_3 and γ_4 using chart (V, y) will yield the same tangent vectors $X_{\gamma_3,p}$ and $X_{\gamma_4,p}$.

§4.3.1 Algebras and Derivations

Definition 4.46 (Algebra over a Field):

Let $(V, +, \cdot)$ be a vector space over a field K equipped with a “product” operation,

$$\bullet : V \times V \rightarrow V, \quad (4.104)$$

such that $\bullet$ is bilinear. Then $(V, +, \cdot, \bullet)$ is called an algebra over field K .

In the future, we will impose more conditions on the product operation $\bullet$, such as anti-symmetry to get something called a Lie algebra. A typical example for that is the cross product in $\mathbb{R}^3$.

Example 4.47 (Algebra of Smooth Functions)

We have already seen that $(\mathcal{C}^\infty(M), +, \cdot)$ is a vector space over $\mathbb{R}$. Now, we can define a product operation $\bullet$ on $\mathcal{C}^\infty(M)$ as follows:

$$\begin{aligned} \bullet : \mathcal{C}^\infty(M) \times \mathcal{C}^\infty(M) &\rightarrow \mathcal{C}^\infty(M), \\ (f, g) &\mapsto f \bullet g \end{aligned} \quad (4.105)$$

where $(f \bullet g)(p) := f(p) \cdot g(p)$ for all $p \in M$. It is easy to check that $\bullet$ is bilinear, and hence $(\mathcal{C}^\infty(M), +, \cdot, \bullet)$ is an algebra over $\mathbb{R}$.

Note the difference between $\cdot$ and $\bullet$: the former is scalar multiplication, while the latter is function multiplication, both at heart uses the field multiplication in $\mathbb{R}$.

Let's look at some special algebras where the product operation satisfies some special properties.

Definition 4.48:

Let $(V, +, \cdot, \bullet)$ be an algebra over a field K . The algebra is called:

- Associative if for all $u, v, w \in V$,

$$(u \bullet v) \bullet w = u \bullet (v \bullet w). \quad (4.106)$$

- Commutative if for all $u, v \in V$,

$$u \bullet v = v \bullet u. \quad (4.107)$$

- Unital if there exists an element $\mathbf{1} \in V$ such that

$$\mathbf{1} \bullet v = v \bullet \mathbf{1} = v, \quad \forall v \in V. \quad (4.108)$$

Now let's look at more important class of algebras, which are not necessarily associative or commutative.

Definition 4.49 (Lie Algebra):

A Lie algebra over a field K is an algebra $(V, +, \cdot, [\star, \star])$ such that the product operation $[\star, \star]$, called the Lie bracket, satisfies the following properties:

- Antisymmetry: for all $u, v \in V$,

$$[u, v] = -[v, u]. \quad (4.109)$$

- Jacobi Identity: for all $u, v, w \in V$,

$$[u, [v, w]] + [v, [w, u]] + [w, [u, v]] = 0. \quad (4.110)$$

Note that the 0 here is the additive identity of the vector space $(V, +, \cdot)$.

It is easy to see that for a non-trivial Lie bracket, the algebra cannot be unital.

Definition 4.50 (Derivation):

Let $(V, +, \cdot, \bullet)$ be an algebra over a field K . A derivation on V is a linear map $D : V \rightarrow V$ such that it satisfies the Leibniz rule:

$$D(u \bullet v) = D(u) \bullet v + u \bullet D(v), \quad \forall u, v \in V. \quad (4.111)$$

Example 4.51 (Derivation on Smooth Functions)

We have already seen that $(\mathcal{C}^\infty(M), +, \cdot, \bullet)$ is an algebra over $\mathbb{R}$. Fix a point $p \in M$, take any tangent vector $X_{\gamma,p} \in \mathcal{T}_p M$. We know from the definition that $X_{\gamma,p} : \mathcal{C}^\infty(M) \rightarrow \mathbb{R}$ is a linear map. Now let's check if it satisfies the Leibniz rule, for all $f, g \in \mathcal{C}^\infty(M)$,

$$\begin{aligned} X_{\gamma,p}(f \bullet g) &= ((f \bullet g) \circ \gamma)'(0) \\ &= ((f \circ \gamma) \cdot (g \circ \gamma))'(0) \\ &= (f \circ \gamma)'(0) \cdot (g \circ \gamma)(0) + (f \circ \gamma)(0) \cdot (g \circ \gamma)'(0) \\ &= X_{\gamma,p}(f) \cdot g(p) + f(p) \cdot X_{\gamma,p}(g). \end{aligned}$$

So, $X_{\gamma,p}$ satisfies the Leibniz rule. However, note that $X_{\gamma,p}(f)$ is a real number, not a smooth function on M . So, $X_{\gamma,p}$ is not a derivation on the algebra $(\mathcal{C}^\infty(M), +, \cdot, \bullet)$, usually called a derivation at point p .

Now define a map

$$\begin{aligned} D : \mathcal{C}^\infty(M) &\rightarrow \mathcal{C}^\infty(M), \\ f &\mapsto D(f) \end{aligned} \quad (4.112)$$

where $D(f)(p) := X_{\gamma,p}(f)$ for all $p \in M$. Then D is a derivation on the algebra $(\mathcal{C}^\infty(M), +, \cdot, \bullet)$.

Example 4.52

Let V be the vector space over $\mathbb{R}$, define $A := \text{End}(V)$, we know that $(A, +, \cdot)$ is a vector space over $\mathbb{R}$. Now define a product operation on A as follows:

$$\begin{aligned} [\star, \star] : A \times A &\rightarrow A, \\ (\phi, \psi) &\mapsto [\phi, \psi] := \phi \circ \psi - \psi \circ \phi, \end{aligned} \quad (4.113)$$

where $\circ$ is the composition of linear maps. It is easy to see that $[\star, \star]$ is bilinear, and hence $(A, +, \cdot, [\star, \star])$ is an algebra over $\mathbb{R}$. Moreover, $[\star, \star]$ is antisymmetric, and for all $\phi, \psi, \rho \in A$,

$$[\phi, [\psi, \rho]] + [\psi, [\rho, \phi]] + [\rho, [\phi, \psi]] = 0, \quad (4.114)$$

which is called the *Jacobi identity*. So, $(A, +, \cdot, [\star, \star])$ is a Lie algebra over $\mathbb{R}$.

Now fix $H \in A$, define a map

$$\begin{aligned} D_H : A &\rightarrow A, \\ \phi &\mapsto D_H(\phi) := [H, \phi]. \end{aligned} \quad (4.115)$$

Let's check if D_H is a derivation, for all $\phi, \psi \in A$,

$$D_H([\phi, \psi]) = [H, [\phi, \psi]]$$

using the Jacobi identity, we have:

$$= -[\phi, [\psi, H]] - [\psi, [H, \phi]]$$

rearranging the terms and use antisymmetry, we get:

$$\begin{aligned} &= [[H, \phi], \psi] + [\phi, [H, \psi]] \\ &= [D_H(\phi), \psi] + [\phi, D_H(\psi)]. \end{aligned}$$

So, D_H is a derivation on the Lie algebra $(A, +, \cdot, [\star, \star])$.

With this example, we can see the algebraic structure of Poisson brackets in classical mechanics, and the commutator in quantum mechanics.

Remark 4.53 (Poisson Bracket). In classical mechanics, the state of a system is represented by a point in phase space (which is a symplectic manifold), and observables are represented by smooth functions on the phase space. The Poisson bracket defines a Lie algebra structure on the space of observables. If we fix an observable H (the Hamiltonian), then the map $D_H(f) := \{H, f\}$ is a derivation on the Lie algebra of observables, which generates the time evolution of the system according to Hamilton's equations.

Remark 4.54 (Commutator in Quantum Mechanics). Similarly, in quantum mechanics, the state of a system is represented by a vector in a Hilbert space, and observables are represented by self-adjoint operators on that space. The commutator defines a Lie algebra structure on the space of observables. If we fix an observable H (the Hamiltonian operator), then the map $D_H(\phi) := [H, \phi]$ is a derivation on the Lie algebra of observables, which generates the time evolution of the system according to the Heisenberg equation of motion.

§4.3.2 Basis and Dimension of Tangent Space

We have shown that for a smooth manifold M and a point $p \in M$, the set of tangent vectors $\mathcal{T}_p M$ is a vector space over $\mathbb{R}$. Now we will prove a very crucial theorem in differential geometry, which states that the dimension of the tangent space $\mathcal{T}_p M$ is equal to the dimension of the manifold M .

Theorem 4.55 (Dimension of Tangent Space)

Let M be a smooth manifold of dimension d , then for all $p \in M$, the tangent vector space $\mathcal{T}_p M$ is a vector space over $\mathbb{R}$ of dimension d .

$$\dim(\mathcal{T}_p M) = \dim(M) = d. \quad (4.116)$$

Note that we have used the same symbol $\dim$ for the dimension of a manifold and the dimension of a vector space, but they are different concepts. The dimension of a manifold is defined as the dimension of the Euclidean space that it locally resembles, while the dimension of a vector space is defined as the cardinality of its basis.

Proof:

Fix a point $p \in M$, and fix a chart (U, x) of M around p .

To prove this theorem, we will construct a basis of $\mathcal{T}_p M$ consisting of d tangent vectors.

Define d curves $\gamma_j : \mathbb{R} \rightarrow U$, $j = 1, \dots, d$, such that

$$\gamma_j(0) = p; \quad x^i \circ \gamma_j(t) = \delta_j^i t, \quad \forall t \in \mathbb{R}, \quad (4.117)$$

where x^i is the i -th coordinate function of chart x , and δ_j^i is the Kronecker delta. So pictorially, γ_j is a curve that moves along the j -th coordinate axis in the Euclidean space $\mathbb{R}^d$ under the chart x .

Name the corresponding tangent vectors at p as

$$\mathbf{e}_j := X_{\gamma_j, p}, \quad j = 1, \dots, d. \quad (4.118)$$

Let's look at how $\mathbf{e}_j$ acts on a smooth function $f \in \mathcal{C}^\infty(M)$:

$$\mathbf{e}_j(f) = (f \circ \gamma_j)'(0) = (f \circ \text{id}_U \circ \gamma_j)'(0)$$

insert the identity map $\text{id}_U = x^{-1} \circ x$ on U :

$$\begin{aligned} &= (f \circ x^{-1} \circ (x \circ \gamma_j))'(0) \\ &= [\partial_i (f \circ x^{-1})(x(p))] \cdot (x^i \circ \gamma_j)'(0) \\ &= [\partial_i (f \circ x^{-1})(x(p))] \cdot \delta_j^i \end{aligned}$$

$$= \partial_j(f \circ x^{-1})(x(p)).$$

Define a formal symbol as

$$\left(\frac{\partial}{\partial x^j}\right)_p(f) := \partial_j(f \circ x^{-1})(x(p)), \quad \forall f \in \mathcal{C}^\infty(M). \quad (4.119)$$

Don't confuse this notation with the usual partial derivative of a function of several real variables.

$$\begin{aligned} (\partial_j)_p &: \mathcal{C}^\infty(\mathbb{R}^d, \mathbb{R}) \rightarrow \mathbb{R}, \\ \left(\frac{\partial}{\partial x^j}\right)_p &: \mathcal{C}^\infty(M) \rightarrow \mathbb{R}. \end{aligned}$$

So we have

$$\mathbf{e}_j = \left(\frac{\partial}{\partial x^j}\right)_p, \quad j = 1, \dots, d. \quad (4.120)$$

Define the set

$$\mathcal{B} := \{\mathbf{e}_1, \dots, \mathbf{e}_d\} = \left\{ \left(\frac{\partial}{\partial x^1}\right)_p, \dots, \left(\frac{\partial}{\partial x^d}\right)_p \right\}. \quad (4.121)$$

We will show that $\mathcal{B}$ is a basis of $\mathcal{T}_p M$, *i.e.*, for any $X \in \mathcal{T}_p M$, there exist unique real numbers $X^1, \dots, X^d$ such that

$$X = X^j \mathbf{e}_j = X^j \left(\frac{\partial}{\partial x^j}\right)_p. \quad (\text{sum over } j \text{ from } 1 \text{ to } d \text{ is implied}) \quad (4.122)$$

1. **Spanning:** We know $\exists \gamma : \mathbb{R} \rightarrow M$ such that $X = X_{\gamma, p}$. For all $f \in \mathcal{C}^\infty(M)$,

$$X(f) = (f \circ \gamma)'(0) = (f \circ \text{id}_U \circ \gamma)'(0)$$

insert the identity map $\text{id}_U = x^{-1} \circ x$ on U :

$$\begin{aligned} &= (f \circ x^{-1} \circ (x \circ \gamma))'(0) \\ &= [\partial_i(f \circ x^{-1})(x(p))] \cdot (x^i \circ \gamma)'(0) \\ &= (x^i \circ \gamma)'(0) \cdot \left(\frac{\partial}{\partial x^i}\right)_p(f). \end{aligned}$$

Note that $x^i \circ \gamma : \mathbb{R} \rightarrow \mathbb{R}$ is a smooth function of a real variable, so $(x^i \circ \gamma)'(0) \in \mathbb{R}$. Define

$$X^i := (x^i \circ \gamma)'(0) \in \mathbb{R}, \quad i = 1, \dots, d. \quad (4.123)$$

So we have

$$X(f) = X^i \left(\frac{\partial}{\partial x^i}\right)_p(f) = X^i \mathbf{e}_i(f), \quad \forall f \in \mathcal{C}^\infty(M). \quad (4.124)$$

Since this is true for all $f \in \mathcal{C}^\infty(M)$, we have

$$X = X^i \mathbf{e}_i. \quad (4.125)$$

Thus, $\mathcal{T}_p M = \text{span}(\mathcal{B})$.

Remark 4.56 (Smoothness of Chart map and co-ordinate functions). In general to talk about smooth of any function $f : M \rightarrow \mathbb{R}$, we have used charts such that f is smooth if and only if $f \circ x^{-1} : x(U) \subseteq \mathbb{R}^d \rightarrow \mathbb{R}$.

So by this definition, the chart map $x : U \rightarrow x(U) \subseteq \mathbb{R}^d$ is trivially smooth, since $x \circ x^{-1} = \text{id}_{x(U)}$ is smooth. Similarly, the coordinate functions $x^i : U \rightarrow \mathbb{R}$ are also smooth, since $x^i \circ x^{-1} : x(U) \subseteq \mathbb{R}^d \rightarrow \mathbb{R}$ is just the projection onto the i -th coordinate, which is a linear map and hence smooth.

2. **Linear Independence:** Suppose that $\mathcal{B}$ is linearly dependent, then there exist real numbers $X^1, \dots, X^d$, not all zero, such that

$$X^j \mathbf{e}_j = 0. \quad (4.126)$$

So for all $f \in \mathcal{C}^\infty(M)$, we have $X^j \mathbf{e}_j(f) = 0$. In particular, take $f = x^i$, the i -th coordinate function of chart x , then

$$\begin{aligned} 0 &= X^j \mathbf{e}_j(x^i) = X^j \left(\frac{\partial}{\partial x^j} \right)_p (x^i) = X^j \partial_j (x^i \circ x^{-1})(x(p)) \\ &= X^j \partial_j (\text{proj}^i)(x(p)) = X^j \delta_j^i = X^i. \end{aligned}$$

Since this is true for all $i = 1, \dots, d$, we have $X^1 = X^2 = \dots = X^d = 0$, which contradicts our assumption. Hence, $\mathcal{B}$ is linearly independent.

Therefore, $\mathcal{B}$ is a basis of $\mathcal{T}_p M$, and $\dim(\mathcal{T}_p M) = d$. Q.E.D.

Terminology: Let $X \in \mathcal{T}_p M$, then we have

$$X = X^j \left(\frac{\partial}{\partial x^j} \right)_p, \quad (4.127)$$

where $X^j = X(x^j) = (x^j \circ \gamma)'(0)$ are called the *components* of X with respect to the basis $\mathcal{B}$ induced by the chart (U, x) .

§4.3.3 Change of Basis and Coordinate Transformation

In the construction of the basis $\mathcal{B}$ of $\mathcal{T}_p M$, we have used a chart (U, x) around p . Now suppose we have another chart (V', y) around p , we can similarly construct another basis of $\mathcal{T}_p M$ as

$$\tilde{\mathcal{B}} := \{\tilde{\mathbf{e}}_1, \dots, \tilde{\mathbf{e}}_d\} = \left\{ \left(\frac{\partial}{\partial y^1} \right)_p, \dots, \left(\frac{\partial}{\partial y^d} \right)_p \right\}. \quad (4.128)$$

For simplicity, define $V := U \cap V'$, which is also a chart neighborhood of p , now we have two charts (V, x) and (V, y) around p .

These two charts induce two different bases $\mathcal{B}$ and $\tilde{\mathcal{B}}$ of the same vector space $\mathcal{T}_p M$, so there must exist a change of basis map between them $A, B \in \text{End}(\mathcal{T}_p M)$ such that

$$\tilde{\mathbf{e}}_j = A_j^i \mathbf{e}_i, \quad \mathbf{e}_j = B_j^i \tilde{\mathbf{e}}_i, \quad (4.129)$$

where $A_j^i, B_j^i \in \mathbb{R}$ are the components of the linear maps A and B with respect to the basis $\mathcal{B}$. Since change of basis maps are invertible, we have the following relation

$$A_j^i B_i^k = \delta_j^k = B_j^i A_i^k. \quad (4.130)$$

Now let's find the explicit form of the components A_j^i and B_j^i . Take x^k , the k -th coordinate function of chart x , and apply $\tilde{\mathbf{e}}_j$ on it, we have

$$\begin{aligned} \tilde{\mathbf{e}}_j(x^k) &= A_j^i \mathbf{e}_i(x^k) \\ \left(\frac{\partial}{\partial y^j} \right)_p (x^k) &= A_j^i \left(\frac{\partial}{\partial x^i} \right)_p (x^k) \\ \partial_j (x^k \circ y^{-1})(y(p)) &= A_j^i \partial_i (x^k \circ x^{-1})(x(p)) \\ \partial_j (x^k \circ y^{-1})(y(p)) &= A_j^i \delta_i^k \\ \partial_j (x^k \circ y^{-1})(y(p)) &= A_j^k. \end{aligned}$$

So we have

$$A_j^k = \partial_j (x^k \circ y^{-1})(y(p)). \quad (4.131)$$

Similarly, take y^k , the k -th coordinate function of chart y , and apply $\mathbf{e}_j$ on it, we have

$$B_j^k = \partial_j (y^k \circ x^{-1})(x(p)). \quad (4.132)$$

With this let's look at the transformation of components of a tangent vector $X \in \mathcal{T}_p M$ under change of basis. We have

$$X = X^j \mathbf{e}_j = \tilde{X}^j \tilde{\mathbf{e}}_j, \quad (4.133)$$

where $X^j = X(x^j)$ and $\tilde{X}^j = X(y^j)$ are the components of X with respect to the bases $\mathcal{B}$ and $\tilde{\mathcal{B}}$ respectively. Using the change of basis relation (4.129), we have

$$X = X^j \mathbf{e}_j = X^j B^i_j \tilde{\mathbf{e}}_i = \tilde{X}^i \tilde{\mathbf{e}}_i.$$

Since $\tilde{\mathcal{B}}$ is a basis, we have

$$\tilde{X}^i = X^j B^i_j = X^j \partial_j (y^i \circ x^{-1})(x(p)). \quad (4.134)$$

Now, let's write this component of change of basis in a more familiar form, *i.e.*, the Jacobian matrix of the coordinate transformation. Note that the transition map between the two charts (V, x) and (V, y) is given by

$$y^i \circ x^{-1} : x(V) \subseteq \mathbb{R}^d \rightarrow \mathbb{R}. \quad (4.135)$$

Abusing the notation a bit, for now call $y^i \circ x^{-1}$ as y^i , so y^i is a real-valued smooth function of d real variables. Since we are only interested in a single point $p \in V$, we can think of x^j as independent variables for the function y^i . So we have

$$y^i = y^i(x^1, \dots, x^d). \quad (4.136)$$

With this new interpretation, we have

$$\partial_j (y^i \circ x^{-1})(x(p)) = \frac{\partial y^i}{\partial x^j}(x(p)). \quad (4.137)$$

Similarly, we can also write

$$\partial_j (x^i \circ y^{-1})(y(p)) = \frac{\partial x^i}{\partial y^j}(y(p)). \quad (4.138)$$

So the change of basis relation (4.129) can be rewritten as

$$\tilde{\mathbf{e}}_j = \frac{\partial y^i}{\partial x^j}(x(p)) \mathbf{e}_i, \quad \mathbf{e}_j = \frac{\partial x^i}{\partial y^j}(y(p)) \tilde{\mathbf{e}}_i, \quad (4.139)$$

and the transformation of components of a tangent vector $X \in \mathcal{T}_p M$ under change of basis can be rewritten as

$$\tilde{X}^i = X^j \frac{\partial y^i}{\partial x^j}(x(p)). \quad (4.140)$$

Remark 4.57 (Jacobian Matrix). Let $f : \mathbb{R}^m \rightarrow \mathbb{R}^n$ be a smooth map, then the Jacobian matrix of f at a point $a \in \mathbb{R}^m$ is defined as the $n \times m$ matrix whose (i, j) -th entry is given by

$$\frac{\partial f^i}{\partial x^j}(a), \quad (4.141)$$

where f^i is the i -th component function of f , and x^j is the j -th coordinate of $\mathbb{R}^m$.

In older literature, you may see that the transformation law (see (4.140)) is being used to define a 'vector'. It goes like this: an array of d real numbers $(X^1, \dots, X^d)$ is called a vector if under a change of coordinates $x \rightarrow y = y(x)$, the components transform as

$$\tilde{X}^i = X^j \frac{\partial y^i}{\partial x^j}. \quad (4.142)$$

This definition is not wrong, but it is not very useful, since it does not tell you what a vector actually is.

§4.3.3.1 Position Vector in Physics

In physics, we often talk about the position vector of a point in space. From all this discussion, only coordinate functions x^i looks like a good candidate for a position vector. However, this does not make sense, as we have seen that vectors have a specific transformation law under change of coordinates. Let's consider a chart (U, y) such that y is highly non-linear, for example, polar coordinates in $\mathbb{R}^2$. So the transformation of components of y^i under change of basis will not be linear, and hence y^i cannot be a vector.

This shows that the concept of position vector in physics is not very well-defined in the context of differential geometry. In fact, the position vector is not a vector in the sense of differential geometry, but rather an image of a point on a manifold under a chart map.

LECTURE 10

§4.4 Cotangent Spaces and Gradient

Definition 4.58 (Cotangent Space):

Let M be a smooth manifold and $p \in M$. The cotangent space at p , denoted $\mathcal{T}_p^*M$, is the dual space of the tangent space $\mathcal{T}_pM$:

$$\mathcal{T}_p^*M := (\mathcal{T}_pM)^*. \quad (4.143)$$

Since the manifold is finite-dimensional², the tangent space $\mathcal{T}_pM$ is a finite-dimensional vector space. Therefore, its dual space $\mathcal{T}_p^*M$ is also a finite-dimensional vector space of the same dimension, moreover, we have the isomorphism

$$\mathcal{T}_pM \cong_{\text{vec}} \mathcal{T}_p^*M. \quad (4.144)$$

However, this isomorphism is not canonical, *i.e.*, there is no natural way to identify vectors in $\mathcal{T}_pM$ with covectors in $\mathcal{T}_p^*M$ without introducing some additional structure on these tangent and cotangent spaces.

Now, using these vector and dual vector spaces, we can define tensor spaces at a point $p \in M$,

$$\mathcal{T}_s^r(\mathcal{T}_pM) := \left\{ t : \underbrace{\mathcal{T}_p^*M \times \cdots \times \mathcal{T}_p^*M}_{r \text{ times}} \times \underbrace{\mathcal{T}_pM \times \cdots \times \mathcal{T}_pM}_{s \text{ times}} \rightarrow \mathbb{R} \mid t \text{ is multilinear} \right\}. \quad (4.145)$$

This is set (even vector space) of all (r, s) -type tensors at the point p .

Definition 4.59 (Gradient):

Let $f \in \mathcal{C}^\infty(M)$. Then at each point $p \in M$, we have a linear map

$$\begin{aligned} d_p : \mathcal{C}^\infty(M) &\rightarrow \mathcal{T}_p^*M \\ f &\mapsto d_p f, \end{aligned} \quad (4.146)$$

where $d_p f$ is defined by

$$d_p f(X) := X(f), \quad \forall X \in \mathcal{T}_pM. \quad (4.147)$$

The linear map d_p is called the gradient operator at the point p , and $d_p f$ is called the gradient of the function f at the point p .

Observe that the gradient $d_p f$ is a covector at the point p not a vector. Also, the gradient of a function f at a point p is “orthogonal” to the level set of f passing through the point p , *i.e.*, let a tangent vector $X \in \mathcal{T}_pM$ be tangent to the level set³ of f at the point p , then we have

$$d_p f(X) = X(f) = 0,$$

²The way we have defined manifolds in this course, they are finite-dimensional by default as we always map them to $\mathbb{R}^d$. For infinite-dimensional manifolds, we need more sophisticated tools like Banach and Hilbert spaces, which are beyond the scope of this course.

since the function f is constant on its level set. Thus, the gradient $d_p f$ annihilates all tangent vectors to the level set of f at the point p .

§4.4.1 Basis of Cotangent Space

We have defined the gradient operator at a point $p \in M$, using which we can construct a basis for the cotangent space $\mathcal{T}_p^* M$.

Theorem 4.60 (Basis of Cotangent Space)

Let M be a smooth manifold and $p \in M$. Let (U, x) be a chart around the point p . Then the set

$$\{d_p x^1, d_p x^2, \dots, d_p x^{\dim M}\} \quad (4.148)$$

where x^i is the i -th component function of the chart x , forms a basis of the cotangent space $\mathcal{T}_p^* M$.

Proof:

Since the dimension of the cotangent space $\mathcal{T}_p^* M$ is equal to the dimension of the manifold M , it is sufficient to show that the set $\{d_p x^i\}_{i=1}^{\dim M}$ is linearly independent. Suppose not the set is linearly dependent, then there exist real numbers $\omega_i \in \mathbb{R}$, not all zero, such that

$$\omega_i d_p x^i = 0. \quad (\text{sum over } i \text{ from } 1 \text{ to } \dim M \text{ is implied}).$$

So for any tangent vector $X \in \mathcal{T}_p M$, we have $\omega_i d_p x^i(X) = 0$. In particular, let us choose $X = \left(\frac{\partial}{\partial x^j}\right)_p$, the j -th basis vector of the tangent space $\mathcal{T}_p M$ induced by the chart (U, x) . Then we have

$$\begin{aligned} 0 &= \omega_i d_p x^i \left(\left(\frac{\partial}{\partial x^j} \right)_p \right) = \omega_i \left(\frac{\partial}{\partial x^j} \right)_p (x^i) = \omega_i \partial_j (x^i \circ x^{-1})(x(p)) \\ &= \omega_i \partial_j (\text{proj}^i)(x(p)) = \omega_i \delta_j^i = \omega_j. \end{aligned}$$

Here, $\text{proj}^i : \mathbb{R}^{\dim M} \rightarrow \mathbb{R}$ is the projection map onto the i -th coordinate. Since j was arbitrary, we have $\omega_j = 0$ for all $j = 1, 2, \dots, \dim M$, which contradicts our assumption that not all ω_i are zero. Thus, the set $\{d_p x^i\}_{i=1}^{\dim M}$ is linearly independent and hence forms a basis of the cotangent space $\mathcal{T}_p^* M$. Q.E.D.

Moreover, we have the following duality relation between the basis of the tangent space and the basis of the cotangent space induced by the same chart (U, x) :

$$d_p x^i \left(\left(\frac{\partial}{\partial x^j} \right)_p \right) = \delta_j^i. \quad (4.149)$$

Thus, the basis $\{d_p x^i\}_{i=1}^{\dim M}$ is *dual basis* to the basis $\left\{ \left(\frac{\partial}{\partial x^i} \right)_p \right\}_{i=1}^{\dim M}$.

§4.5 Push-Forward and Pull-Back

Definition 4.61 (Push-Forward):

Let M and N be two smooth manifolds and $\phi : M \rightarrow N$ be a smooth map between them. Then the push-forward of ϕ at a point $p \in M$, denoted by ϕ_{*p} , is the linear map

$$\begin{aligned} \phi_{*p} : \mathcal{T}_p M &\rightarrow \mathcal{T}_{\phi(p)} N \\ X &\mapsto \phi_{*p}(X), \end{aligned} \quad (4.150)$$

where $\phi_{*p}(X)$ is defined by its action on smooth functions $f \in C^\infty(N)$ as

$$\phi_{*p}(X)(f) := X(f \circ \phi). \quad (4.151)$$

³The level set of a function $f : M \rightarrow \mathbb{R}$ corresponding to a value $c \in \mathbb{R}$ is defined as the set $\mathcal{L}_c(f) := \{p \in M \mid f(p) = c\}$.

It is easy to reconstruct the definition of push-forward by observing that the composition $f \circ \phi$ is a smooth function on the manifold M , and hence the tangent vector $X \in \mathcal{T}_p M$ can act on it. The result of this action is a real number, which is exactly what we want from the tangent vector $\phi_{*p}(X) \in \mathcal{T}_{\phi(p)} N$ when it acts on the smooth function $f \in \mathcal{C}^\infty(N)$.

The push-forward of the smooth map $\phi : M \rightarrow N$ is also called the *differential* of ϕ at the point p , and is often denoted by $d_p \phi$.

Proposition 4.62

Let $X_{\gamma,p} \in \mathcal{T}_p M$ be the tangent vector at the curve γ at the point p (with $\gamma(0) = p$). Then the push-forward of $X_{\gamma,p}$ by the smooth map $\phi : M \rightarrow N$ is the tangent vector at the curve $\phi \circ \gamma$ at the point $\phi(p)$, that is,

$$\phi_{*p}(X_{\gamma,p}) = X_{\phi \circ \gamma, \phi(p)}. \quad (4.152)$$

Proof:

For any smooth function $f \in \mathcal{C}^\infty(N)$, we have

$$\begin{aligned} \phi_{*p}(X_{\gamma,p})(f) &= X_{\gamma,p}(f \circ \phi) && \text{(by definition of push-forward)} \\ &= ((f \circ \phi) \circ \gamma)'(0) && \text{(by definition of } X_{\gamma,p}) \\ &= (f \circ (\phi \circ \gamma))'(0) && \text{(by associativity of function composition } \circ) \\ &= X_{\phi \circ \gamma, \phi(p)}(f). && \text{(by definition of } X_{\phi \circ \gamma, \phi(p)}) \end{aligned}$$

Since this is true for all smooth functions $f \in \mathcal{C}^\infty(N)$, we have $\phi_{*p}(X_{\gamma,p}) = X_{\phi \circ \gamma, \phi(p)}$. Q.E.D.

Definition 4.63 (Pull-Back):

Let M and N be two smooth manifolds and $\phi : M \rightarrow N$ be a smooth map between them. Then the pull-back of ϕ at a point $p \in M$, denoted by ϕ_p^* , is the linear map

$$\begin{aligned} \phi_p^* : \mathcal{T}_{\phi(p)}^* N &\rightarrow \mathcal{T}_p^* M \\ \omega &\mapsto \phi_p^*(\omega), \end{aligned} \quad (4.153)$$

where $\phi_p^*(\omega)$ is defined by its action on tangent vectors $X \in \mathcal{T}_p M$ as

$$\phi_p^*(\omega)(X) := \omega(\phi_{*p}(X)). \quad (4.154)$$

This definition can also be reconstructed by observing that the push-forward $\phi_{*p}(X)$ is a tangent vector at the point $\phi(p) \in N$, and hence the covector $\omega \in \mathcal{T}_{\phi(p)}^* N$ can act on it. The result of this action is a real number, which is exactly what we want from the covector $\phi_p^*(\omega) \in \mathcal{T}_p^* M$ when it acts on the tangent vector $X \in \mathcal{T}_p M$.

Proposition 4.64

Let $d_{\phi(p)} f \in \mathcal{T}_{\phi(p)}^* N$ be the gradient of the function $f \in \mathcal{C}^\infty(N)$ at the point $\phi(p) \in N$. Then the pull-back of $d_{\phi(p)} f$ by the smooth map $\phi : M \rightarrow N$ is the gradient of the function $f \circ \phi \in \mathcal{C}^\infty(M)$ at the point $p \in M$, that is,

$$\phi_p^*(d_{\phi(p)} f) = d_p(f \circ \phi). \quad (4.155)$$

Proof:

For any tangent vector $X \in \mathcal{T}_p M$, we have

$$\begin{aligned} \phi_p^*(d_{\phi(p)} f)(X) &= d_{\phi(p)} f(\phi_{*p}(X)) && \text{(by definition of pull-back)} \\ &= \phi_{*p}(X)(f) && \text{(by definition of } d_{\phi(p)} f) \\ &= X(f \circ \phi) && \text{(by definition of push-forward } \phi_{*p}) \\ &= d_p(f \circ \phi)(X). && \text{(by definition of } d_p(f \circ \phi)) \end{aligned}$$

Since this is true for all tangent vectors $X \in \mathcal{T}_p M$, we have $\phi_p^*(d_{\phi(p)} f) = d_p(f \circ \phi)$. Q.E.D.

So given a smooth map $\phi : M \rightarrow N$ between two smooth manifolds M and N ,

*vectors are pushed forward from M to N , and
covectors are pulled back from N to M .*

However, if the smooth map $\phi : M \rightarrow N$ is a diffeomorphism, then we can also pull back vectors from N to M and push forward covectors from M to N using the inverse map $\phi^{-1} : N \rightarrow M$.

$$\begin{aligned} \phi_p^* : \mathcal{T}_{\phi(p)}N &\rightarrow \mathcal{T}_pM & \phi_p^* : \mathcal{T}_p^*M &\rightarrow \mathcal{T}_{\phi(p)}^*N \\ \phi_p^*(Y) &:= (\phi^{-1})_{*\phi(p)}(Y), \quad \forall Y \in \mathcal{T}_{\phi(p)}N, & \phi_p^*(\eta) &:= (\phi^{-1})_{\phi(p)}^*(\eta), \quad \forall \eta \in \mathcal{T}_p^*M. \end{aligned} \quad (4.156)$$

§4.6 Immersions and Embeddings


Now we want to go back to usual intuitive notion of tangent vectors as arrows attached to points on the manifold. For that, we need to think about the manifold being “inside” some higher-dimensional space like $\mathbb{R}^n$ (for some n). We want to decide under which conditions some smooth manifold M can ‘sit’ inside some $\mathbb{R}^n$. Here we are using the term ‘sit inside’ in intuitive sense, as there are two different notions of sitting inside another manifold, namely, immersion and embedding. Embedding is a stronger notion than immersion, however, both of these notions have their use cases sometimes we need only immersion, sometimes we need embedding.

Definition 4.65 (Immersion):

Let M and N be two smooth manifolds and $\phi : M \rightarrow N$ be a smooth map between them. The map ϕ is called an immersion if for every point $p \in M$, the differential $d_p\phi : \mathcal{T}_pM \rightarrow \mathcal{T}_{\phi(p)}N$ is injective.

Example 4.66 (1-Sphere S^1 immersed in $\mathbb{R}^2$)

Consider the smooth map $\phi : S^1 \rightarrow \mathbb{R}^2$ such that



Notice that both points p_1 and p_2 on the circle S^1 are mapped to the same point $\phi(p_1) = \phi(p_2)$ in $\mathbb{R}^2$. Thus, the tangent spaces at these points are same, $\mathcal{T}_{\phi(p_1)}\mathbb{R}^2 = \mathcal{T}_{\phi(p_2)}\mathbb{R}^2$. However, the push-forwards of the tangent vectors at these points are different, represented by the red and blue arrows respectively. Thus, the differential $d_p\phi$ is injective at both points p_1 and p_2 . Hence, the map $\phi : S^1 \rightarrow \mathbb{R}^2$ is an immersion.

Definition 4.67 (Embedding):

Let M and N be two smooth manifolds and $\phi : M \rightarrow N$ be a smooth map between them. The map ϕ is called an embedding if

- (i) ϕ is an immersion, and
- (ii) $M \cong_{top.} \phi(M) \subseteq N$ via the subspace topology induced from N .

Using the condition of embedding being an immersion, we can show that the image is also diffeomorphic to the original manifold.

Proposition 4.68

Let M and N be two smooth manifolds and $\phi : M \rightarrow N$ be an embedding. Then the image $\phi(M)$ is a smooth manifold and $M \cong_{\text{diff}} \phi(M)$.

It is easy to see that the [example 4.66](#) is not an embedding, since the map ϕ itself is not injective.

With these notions of immersion and embedding, now we can comment on the two different ways of thinking about tangent vectors:

- **Abstract Viewpoint:** In this viewpoint, we think of tangent vectors as derivations acting on smooth functions at a point on the manifold.
- **Concrete Viewpoint:** In this viewpoint, we think of tangent vectors as arrows attached to points on the manifold, which can be visualized when the manifold is immersed or embedded in a higher-dimensional space like $\mathbb{R}^n$.

Whitney's embedding theorem states that both these viewpoints are equivalent,

Theorem 4.69 (Whitney's Embedding Theorem)

Any smooth manifold M can be

- embedded in $\mathbb{R}^{2 \dim M}$, and
- immersed in $\mathbb{R}^{2 \dim M - 1}$.

This theorem doesn't say that we only can embed or immerse a manifold in these dimensions, rather it says that these dimensions are sufficient to embed or immerse any smooth manifold. For example, we can embed the 2-sphere S^2 in $\mathbb{R}^3$ itself, even though Whitney's embedding theorem states that we can embed it in $\mathbb{R}^4$. Similarly, we have another extreme example of the Klein bottle, which cannot be embedded in $\mathbb{R}^3$ but can be immersed in $\mathbb{R}^3$.

This version of Whitney's embedding theorem is called the *strong Whitney embedding theorem*. There is also a *weak Whitney embedding theorem*, but there are even stronger versions of the strong Whitney embedding theorem.

Theorem 4.70 (Strong Whitney Embedding Theorem (Improved Version))

Any smooth manifold M can be immersed in $\mathbb{R}^{2 \dim M - \alpha(M)}$, where $\alpha(M)$ is the number of 1's in the binary representation of $\dim M$.

For example, for $\dim M = 3$, the binary representation is 11_2 , which has two 1's. Thus, any 3-dimensional smooth manifold can be immersed in $\mathbb{R}^{2 \times 3 - 2} = \mathbb{R}^4$, which is an improvement over the previous version of the strong Whitney embedding theorem that stated that any 3-dimensional smooth manifold can be immersed in $\mathbb{R}^5$.

§4.7 Tangent Bundle and Vector Fields

So far we have defined the tangent space at a point p on a smooth manifold M . Now say we want to study all tangent spaces at all points on the manifold M together.

Definition 4.71 (Tangent Bundle):

Let M be a smooth manifold. The tangent bundle of M , denoted by $\mathcal{T}M$, is the disjoint union of all tangent spaces at all points on the manifold M :

$$\mathcal{T}M := \bigsqcup_{p \in M} \mathcal{T}_p M \quad (4.157)$$

There is a canonical projection map $\pi : \mathcal{T}M \rightarrow M$, called the *bundle projection*, defined by

$$\begin{aligned}\pi : \mathcal{T}M &\rightarrow M \\ X &\mapsto p,\end{aligned}\tag{4.158}$$

where $X \in \mathcal{T}_p M$ is a tangent vector at the point $p \in M$.

Remark 4.72 (Implications of disjoint union). Note that the tangent bundle $\mathcal{T}M$ is defined as a disjoint union of all tangent spaces at all points on the manifold M . Since the tangent spaces are finite-dimensional real vector spaces, thus, each tangent space is isomorphic to $\mathbb{R}^{\dim M}$, so each tangent space is isomorphic to each other as vector spaces. However, since the tangent bundle is defined as a disjoint union, the elements of different tangent spaces are considered distinct even if they are isomorphic as vector spaces.

A consequence of this definition is that zero vector in each tangent space are distinct elements in the tangent bundle. That is, if $0_p \in \mathcal{T}_p M$ and $0_q \in \mathcal{T}_q M$ are the zero vectors in the tangent spaces at points p and q respectively, then 0_p and 0_q are distinct elements in the tangent bundle $\mathcal{T}M$ even though they are both zero vectors.

From the definition of the bundle projection π , it is clear that for each point $p \in M$, the preimage of p under the bundle projection π is the tangent space at the point p , thus, the map π is a surjection. That is the triple $(\mathcal{T}M, M, \pi)$ forms a *set bundle* over the manifold M , for it to be a (topological) bundle we need to define a suitable topology on the tangent bundle $\mathcal{T}M$ such that the bundle projection $\pi : \mathcal{T}M \rightarrow M$ is continuous.

§4.7.1 Smooth Structure on Tangent Bundle

Let's start with defining a suitable topology on the tangent bundle $\mathcal{T}M$. To make this set a topological bundle we need to define topology on it such that the bundle projection $\pi : \mathcal{T}M \rightarrow M$ is continuous and furthermore, to make it a manifold we need to define charts on it.

Consider a chart (U, x) on the manifold M . Define the coordinate map ξ for the tangent bundle $\mathcal{T}M$ as

$$\begin{aligned}\xi : \text{preim}_\pi(U) &\rightarrow \xi(\text{preim}_\pi(U)) = x(U) \times \mathbb{R}^{\dim M} \subseteq \mathbb{R}^{2 \dim M} \\ X &\mapsto \left(x^1(\pi(X)), \dots, x^{\dim M}(\pi(X)), X^1, \dots, X^{\dim M} \right),\end{aligned}\tag{4.159}$$

where $X \in \mathcal{T}_p M$ is a tangent vector at the point $p = \pi(X) \in U$, and X^i are the components of the tangent vector X in the basis induced by the chart (U, x) , that is

$$X = X^i \left(\frac{\partial}{\partial x^i} \right)_p.$$

Proposition 4.73

The map $\xi : \text{preim}_\pi(U) \rightarrow \mathbb{R}^{2 \dim M}$ is a bijection onto its image.

Proof:

To show that the map ξ is injective, let $X, Y \in \text{preim}_\pi(U)$ such that $\xi(X) = \xi(Y)$. Then we have

$$\left(x^1(p), \dots, x^{\dim M}(p), X^1, \dots, X^{\dim M} \right) = \left(x^1(q), \dots, x^{\dim M}(q), Y^1, \dots, Y^{\dim M} \right),$$

where $X \in \mathcal{T}_p M$ and $Y \in \mathcal{T}_q M$. From the equality of the first $\dim M$ components, we have $x(p) = x(q)$. Since the chart (U, x) is a bijection onto its image, we have $p = q$. Now, from the equality of the last $\dim M$ components, we have $X^i = Y^i$ for all $i = 1, 2, \dots, \dim M$. Thus, we have

$$X = X^i \left(\frac{\partial}{\partial x^i} \right)_p = Y^i \left(\frac{\partial}{\partial x^i} \right)_p = Y.$$

Hence, the map ξ is injective.

To show that the map ξ is surjective onto its image, let $(a^1, \dots, a^{\dim M}, v^1, \dots, v^{\dim M}) \in \xi(\text{preim}_\pi(U))$. Then by definition of the image, there exists a tangent vector $X \in \text{preim}_\pi(U)$ such that

$$\xi(X) = (a^1, \dots, a^{\dim M}, v^1, \dots, v^{\dim M}).$$

Thus, the map ξ is surjective onto its image. Hence, the map ξ is a bijection onto its image. Q.E.D.

Using this coordinate map ξ , we can define a topology on the tangent bundle $\mathcal{T}M$ as follows:

Proposition 4.74 (Topology on Tangent Bundle)

A subset $W \subseteq \mathcal{T}M$ is defined to be open if and only if for every chart (U, x) on the manifold M , the set $\xi(W \cap \text{preim}_\pi(U))$ is open in $\mathbb{R}^{2 \dim M}$.

Proof:

We need to show that this definition satisfies the axioms of a topology.

(i) Empty Set and Whole Set are Open: The empty set $\emptyset \subseteq \mathcal{T}M$ is open since for any chart (U, x) on the manifold M , we have

$$\xi(\emptyset \cap \text{preim}_\pi(U)) = \xi(\emptyset) = \emptyset,$$

which is open in $\mathbb{R}^{2 \dim M}$. Similarly, the whole set $\mathcal{T}M$ is open since for any chart (U, x) on the manifold M , we have

$$\xi(\mathcal{T}M \cap \text{preim}_\pi(U)) = \xi(\text{preim}_\pi(U)) = x(U) \times \mathbb{R}^{\dim M}$$

which is open in $\mathbb{R}^{2 \dim M}$ as $x(U)$ is open in $\mathbb{R}^{\dim M}$ by the definition of charts on the manifold M and $\mathbb{R}^{\dim M}$ is open in itself, thus their product $x(U) \times \mathbb{R}^{\dim M}$ is also open in $\mathbb{R}^{2 \dim M}$.

(ii) Arbitrary Unions are Open: Let $\{W_\alpha\}_{\alpha \in I}$ be a collection of open sets in $\mathcal{T}M$, where I is an arbitrary index set. Then for any chart (U, x) on the manifold M , we have

$$\xi\left(\bigcup_{\alpha \in I} W_\alpha \cap \text{preim}_\pi(U)\right) = \bigcup_{\alpha \in I} \xi(W_\alpha \cap \text{preim}_\pi(U)),$$

which is open in $\mathbb{R}^{2 \dim M}$ since each set $\xi(W_\alpha \cap \text{preim}_\pi(U))$ is open in $\mathbb{R}^{2 \dim M}$ by the definition of open sets in $\mathcal{T}M$. Thus, the arbitrary union $\bigcup_{\alpha \in I} W_\alpha$ is open in $\mathcal{T}M$.

(iii) Finite Intersections are Open: Let $W_1, W_2, \dots, W_n$ be a finite collection of open sets in $\mathcal{T}M$. Then for any chart (U, x) on the manifold M , we have

$$\xi\left(\bigcap_{i=1}^n W_i \cap \text{preim}_\pi(U)\right) = \bigcap_{i=1}^n \xi(W_i \cap \text{preim}_\pi(U)),$$

here, the equality holds because ξ is a bijection onto its image. This set is open in $\mathbb{R}^{2 \dim M}$ since each set $\xi(W_i \cap \text{preim}_\pi(U))$ is open in $\mathbb{R}^{2 \dim M}$ by the definition of open sets in $\mathcal{T}M$. Thus, the finite intersection $\bigcap_{i=1}^n W_i$ is open in $\mathcal{T}M$. Q.E.D.

Say $\mathcal{O}_{\mathcal{T}M}$ is the topology on the tangent bundle $\mathcal{T}M$ defined as above. Thus, we have made the tangent bundle $\mathcal{T}M$ a topological space $(\mathcal{T}M, \mathcal{O}_{\mathcal{T}M})$. Using the similar charts induced by the charts on the manifold M , we can also define a smooth structure on the tangent bundle $\mathcal{T}M$ as follows:

Proposition 4.75 (Smooth Atlas on Tangent Bundle)

Let $\mathcal{A}$ be a smooth atlas on the manifold M . Then the collection of charts

$$\mathcal{A}_{\mathcal{T}M} := \{(\text{preim}_\pi(U), \xi) \mid (U, x) \in \mathcal{A}\} \quad (4.160)$$

forms a smooth atlas on the tangent bundle $\mathcal{T}M$.

Proof:

We need to show that the charts in the collection $\mathcal{A}_{\mathcal{T}M}$ are smoothly compatible. Let (U, x) and (V, y) be two charts in the atlas $\mathcal{A}$ on the manifold M . Then the corresponding charts in the collection $\mathcal{A}_{\mathcal{T}M}$ on the tangent bundle $\mathcal{T}M$ are $(\text{preim}_\pi(U), \xi)$ and $(\text{preim}_\pi(V), \eta)$ respectively, where ξ and η are the coordinate maps defined as before. Now, we need to show that the transition map

$$\eta \circ \xi^{-1} : \xi(\text{preim}_\pi(U) \cap \text{preim}_\pi(V)) \rightarrow \eta(\text{preim}_\pi(U) \cap \text{preim}_\pi(V))$$

is a smooth map between open subsets of $\mathbb{R}^{2 \dim M}$.

Let $(a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M}) \in \xi(\text{preim}_\pi(U) \cap \text{preim}_\pi(V))$. by definition of the image, there exists a tangent vector $X \in \text{preim}_\pi(U) \cap \text{preim}_\pi(V)$ such that

$$\xi(X) = (a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M}).$$

Thus, we have

$$\begin{aligned} \eta \circ \xi^{-1}(a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M}) &= \eta(X) \\ &= (y^1(p), \dots, y^{\dim M}(p), \tilde{X}^1, \dots, \tilde{X}^{\dim M}), \end{aligned}$$

where $p = \pi(X) \in U \cap V$ and $\tilde{X}^i$ are the components of the tangent vector X in the basis induced by the chart (V, y) , that is

$$X = \tilde{X}^i \left(\frac{\partial}{\partial y^i} \right)_p.$$

Since the charts (U, x) and (V, y) are smoothly compatible, the map $y \circ x^{-1}$ is a smooth map between open subsets of $\mathbb{R}^{\dim M}$. Thus, the first $\dim M$ components of the map $\eta \circ \xi^{-1}$ are smooth functions of $(a^1, \dots, a^{\dim M})$.

For the last $\dim M$ components, we have the change of basis formula from the basis induced by the chart (U, x) to the basis induced by the chart (V, y) :

$$\tilde{X}^i = X^j \partial_j (y^i \circ x^{-1})(a^1, \dots, a^{\dim M}),$$

which are also smooth functions of $(a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M})$ since the map $y \circ x^{-1}$ is smooth. Thus, the transition map $\eta \circ \xi^{-1}$ is a smooth map between open subsets of $\mathbb{R}^{2 \dim M}$. Hence, the charts in the collection $\mathcal{A}_{\mathcal{T}M}$ are smoothly compatible. Therefore, the collection $\mathcal{A}_{\mathcal{T}M}$ forms a smooth atlas on the tangent bundle $\mathcal{T}M$. Q.E.D.

With this smooth atlas $\mathcal{A}_{\mathcal{T}M}$, we have made the tangent bundle $\mathcal{T}M$ a smooth manifold of dimension $2 \dim M$. Till now, we have not shown that the tangent bundle $\mathcal{T}M$ is indeed a bundle over the manifold M , for that we need to show that the bundle projection $\pi : \mathcal{T}M \rightarrow M$ is a continuous map.

Proposition 4.76 (Tangent Bundle is a Topological Bundle)

The bundle projection $\pi : \mathcal{T}M \rightarrow M$ is a continuous map.

Proof:

Let $V \subseteq M$ be an open set in the manifold M . We need to show that the preimage $\text{preim}_\pi(V) \subseteq \mathcal{T}M$ is an open set in the tangent bundle $\mathcal{T}M$. For that, let (U, x) be any chart on the manifold M . Then we have

$$\xi(\text{preim}_\pi(V) \cap \text{preim}_\pi(U)) = \xi(\text{preim}_\pi(V \cap U)) = x(V \cap U) \times \mathbb{R}^{\dim M}.$$

Since V and U are open sets in the manifold M , so $V \cap U$ is also an open set in the manifold M . Thus, the set $x(V \cap U)$ is an open set in $\mathbb{R}^{\dim M}$. Therefore, the product set $x(V \cap U) \times \mathbb{R}^{\dim M}$ is also an open set in $\mathbb{R}^{2 \dim M}$. Since this is true for any chart (U, x) on the manifold M , we have that the preimage $\text{preim}_\pi(V)$ is an open set in the tangent bundle $\mathcal{T}M$. Hence, the bundle projection $\pi : \mathcal{T}M \rightarrow M$ is a continuous map. Q.E.D.

With this, we have shown that the triple $(\mathcal{T}M, M, \pi)$ forms a topological bundle over the manifold M . Before concluding that the tangent bundle $\mathcal{T}M$ is a smooth bundle over the manifold M , let's take a look at the local trivializations of the tangent bundle $\mathcal{T}M$. Let's recall the definition of local trivializations of a (topological) bundle:

Definition 4.77 (Locally Trivial Bundle):

Let (E, M, π) be a (topological) bundle. The bundle is called locally trivial with typical fiber F if for every point $p \in M$, there exists an open neighborhood $U \subseteq M$ of the point p and a homeomorphism $\varphi : \text{preim}_\pi(U) \rightarrow U \times F$ such that the following diagram commutes:

$$\begin{array}{ccc} \text{preim}_\pi(U) & \xrightarrow{\varphi} & U \times F \\ & \searrow \pi & \swarrow \pi_1 \\ & U & \end{array}$$

Proposition 4.78 (Tangent Bundle is Locally Trivial)

The tangent bundle $(\mathcal{T}M, M, \pi)$ is a locally trivial bundle with typical fiber $\mathbb{R}^{\dim M}$.

Proof:

Let $p \in M$ be any point on the manifold M . Since M is a smooth manifold, there exists a chart (U, x) on the manifold M such that $p \in U$. Now, consider the map

$$\varphi : \text{preim}_\pi(U) \rightarrow U \times \mathbb{R}^{\dim M}$$

defined by

$$\varphi(X) := (\pi(X), X^1, \dots, X^{\dim M}),$$

where $X \in \mathcal{T}_q M$ is a tangent vector at the point $q = \pi(X) \in U$, and X^i are the components of the tangent vector X in the basis induced by the chart (U, x) . We need to show that the map φ is a homeomorphism such that the following diagram commutes:

$$\begin{array}{ccc} \text{preim}_\pi(U) & \xrightarrow{\varphi} & U \times \mathbb{R}^{\dim M} \\ & \searrow \pi & \swarrow \pi_1 \\ & U & \end{array}$$

To show that the map φ is a homeomorphism, we need to show that it is a bijection and both the map and its inverse are continuous. The map φ is injective since if $\varphi(X) = \varphi(Y)$ for some $X, Y \in \text{preim}_\pi(U)$, then we have

$$(\pi(X), X^1, \dots, X^{\dim M}) = (\pi(Y), Y^1, \dots, Y^{\dim M}),$$

from which we get $\pi(X) = \pi(Y)$ and $X^i = Y^i$ for all $i = 1, 2, \dots, \dim M$. Thus, we have

$$X = X^i \left(\frac{\partial}{\partial x^i} \right)_{\pi(X)} = Y^i \left(\frac{\partial}{\partial x^i} \right)_{\pi(Y)} = Y.$$

Hence, the map φ is injective. To show that the map φ is surjective, let $(q, v^1, \dots, v^{\dim M}) \in U \times \mathbb{R}^{\dim M}$. Then we can define a tangent vector $X \in \mathcal{T}_q M$ as

$$X := v^i \left(\frac{\partial}{\partial x^i} \right)_q.$$

Thus, we have

$$\varphi(X) = (\pi(X), X^1, \dots, X^{\dim M}) = (q, v^1, \dots, v^{\dim M}).$$

Hence, the map φ is surjective. Therefore, the map φ is a bijection. To show that the map φ is continuous, let $W \subseteq U \times \mathbb{R}^{\dim M}$ be an open set. We need to show that the preimage $\text{preim}_\varphi(W) \subseteq \text{preim}_\pi(U)$ is an open set. For that, let (U', x') be any chart on the manifold M . Then we have

$$\xi(\text{preim}_\varphi(W) \cap \text{preim}_\pi(U')) = \xi(\text{preim}_\pi(\pi(\text{preim}_\varphi(W))) \cap \text{preim}_\pi(U')).$$

Since $\pi(\text{preim}_\varphi(W))$ is an open set in U (as it is the image of an open set under the projection map), the set $\pi(\text{preim}_\varphi(W)) \cap U'$ is also an open set in the manifold M . Thus, the set

$$\xi(\text{preim}_\pi(\pi(\text{preim}_\varphi(W))) \cap \text{preim}_\pi(U')) = x'(\pi(\text{preim}_\varphi(W)) \cap U') \times \mathbb{R}^{\dim M}$$

is an open set in $\mathbb{R}^{2 \dim M}$. Since this is true for any chart (U', x') on the manifold M , we have that the preimage $\text{preim}_\varphi(W)$ is an open set in $\text{preim}_\pi(U)$. Hence, the map φ is continuous. To show that the inverse map φ^{-1} is continuous, let $W' \subseteq \text{preim}_\pi(U)$ be an open set. We need to show that the preimage $\text{preim}_{\varphi^{-1}}(W') \subseteq U \times \mathbb{R}^{\dim M}$ is an open set. For that, let (U', x') be any chart on the manifold M . Then we have

$$\pi_1(\text{preim}_{\varphi^{-1}}(W') \cap (U' \times \mathbb{R}^{\dim M})) = \pi(\text{preim}_\pi(U) \cap W').$$

Since W' is an open set in $\text{preim}_\pi(U)$, the set $\pi(\text{preim}_\pi(U) \cap W')$ is an open set in U (as it is the image of an open set under the projection map). Thus, the set

$$\pi_1(\text{preim}_{\varphi^{-1}}(W') \cap (U' \times \mathbb{R}^{\dim M})) = \pi(\text{preim}_\pi(U) \cap W')$$

is an open set in $\mathbb{R}^{\dim M}$. Since this is true for any chart (U', x') on the manifold M , we have that the preimage $\text{preim}_{\varphi^{-1}}(W')$ is an open set in $U \times \mathbb{R}^{\dim M}$. Hence, the inverse map φ^{-1} is continuous. Thus, we have shown that the map φ is a homeomorphism. Finally, to show that the diagram commutes, let $X \in \text{preim}_\pi(U)$. Then we have

$$\pi_1 \circ \varphi(X) = \pi_1(\pi(X), X^1, \dots, X^{\dim M}) = \pi(X),$$

which shows that the diagram commutes. Hence, the tangent bundle $(\mathcal{T}M, M, \pi)$ is a locally trivial bundle with typical fiber $\mathbb{R}^{\dim M}$. Q.E.D.

Now, we are ready to show that the tangent bundle $\mathcal{T}M$ is indeed a smooth bundle over the manifold M . For that, we need to show that the bundle projection $\pi : \mathcal{T}M \rightarrow M$ is a smooth map.

Proposition 4.79 (Tangent Bundle is a Smooth Bundle)

The bundle projection $\pi : \mathcal{T}M \rightarrow M$ is a smooth map.

Proof:

Let (U, x) be any chart on the manifold M and $(\text{preim}_\pi(U), \xi)$ be the corresponding chart on the tangent bundle $\mathcal{T}M$. We need to show that the map

$$x \circ \pi \circ \xi^{-1} : \xi(\text{preim}_\pi(U)) \rightarrow x(U)$$

is a smooth map between open subsets of $\mathbb{R}^{\dim M}$.

Let $(a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M}) \in \xi(\text{preim}_\pi(U))$. By definition of the image, there exists a tangent vector $X \in \text{preim}_\pi(U)$ such that

$$\xi(X) = (a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M}).$$

Thus, we have

$$x \circ \pi \circ \xi^{-1}(a^1, \dots, a^{\dim M}, X^1, \dots, X^{\dim M}) = x(\pi(X)) = x(p) = (a^1, \dots, a^{\dim M}),$$

where $p = \pi(X) \in U$. This map is clearly smooth as each component is just a projection onto the first $\dim M$ components. Hence, the bundle projection $\pi : \mathcal{T}M \rightarrow M$ is a smooth map. Q.E.D.

Thus, we have shown that the triple $(\mathcal{T}M, M, \pi)$ forms a smooth bundle over the manifold M , called the *tangent bundle* of the manifold M .

§4.8 Tensor Fields and Modules

We started by defining the tangent space at a point $p \in M$, and then with appropriate operations we have shown that $T_p M$ is a vector space over $\mathbb{R}$. Then we defined the tangent bundle TM , now we want study how different tangent spaces relate to each other, *i.e.*, how to relate $T_p M$ and $T_q M$ for $p, q \in M$.

Definition 4.80 (Vector Field):

Let M be a smooth manifold and TM be its tangent bundle, *i.e.*, $TM \xrightarrow{\pi} M$ (where π is smooth). A ‘smooth’ vector field on M is a smooth section of the tangent bundle, *i.e.*, a smooth map $\sigma : M \rightarrow TM$ such that $\pi \circ \sigma = \text{id}_M$.

$$\begin{array}{c} TM \\ \sigma \text{ smooth} \uparrow \downarrow \pi \text{ smooth} \\ M \\ \uparrow \\ \text{id}_M \end{array}$$

The way we have defined the tangent bundle and the bundle projection map π , it is clear that the section σ will assign to each point $p \in M$ a tangent vector $\sigma(p) \in T_p M$. Thus, a vector field can be thought of as an assignment of a tangent vector to each point of the manifold.

Recall the definition of [Ring](#), we have a few different kinds of rings based on the properties of the multiplication operation.

Definition (Ring):

A ring is a set R equipped with two binary operations $(+, \cdot)$ which satisfy set of properties. If a ring R satisfies the following additional properties, then we have special kinds of rings:

- Commutative Ring: If the multiplication operation is commutative, *i.e.*, $a \cdot b = b \cdot a$ for all $a, b \in R$.
- Ring with Unity: If there exists multiplicative identity $1_R \in R$ such that $a \cdot 1_R = 1_R \cdot a = a$ for all $a \in R$.
- Division Ring: If every non-zero element $a \in R$ has a multiplicative inverse $a^{-1} \in R$ such that $a \cdot a^{-1} = a^{-1} \cdot a = 1_R$.
- Field: If R is a commutative division ring.

Remark 4.81 (Smooth functions as a ring). Recall that the set of smooth functions on a manifold M , denoted by $C^\infty(M)$, forms a vector space over $\mathbb{R}$ with the pointwise addition and scalar multiplication. Moreover, define the multiplication of two smooth functions pointwise as $\bullet : C^\infty(M) \times C^\infty(M) \rightarrow C^\infty(M)$ given by

$$(f \bullet g)(p) := f(p) \cdot g(p), \quad \forall p \in M.$$

With this multiplication, $(C^\infty(M), +, \bullet)$ forms a *unital commutative ring*, where the multiplicative identity is the constant function $1_M : M \rightarrow \mathbb{R}$ defined by $1_M(p) := 1$ for all $p \in M$.

However, $C^\infty(M)$ is not a field, since not all non-zero smooth functions have multiplicative inverses, for example, a function that is zero at some point in M cannot have a smooth inverse defined everywhere on M .

Definition 4.82 (Module):

Let R be a ring. A module over R is an Abelian group $(M, +)$ together with a scalar multiplication $\cdot : R \times M \rightarrow M$ satisfying the following properties for all $r, s \in R$ and $m, n \in M$:

1. $r \cdot (m + n) = r \cdot m + r \cdot n$ (Distributivity over module addition)
2. $(r + s) \cdot m = r \cdot m + s \cdot m$ (Distributivity over ring addition)
3. $(rs) \cdot m = r \cdot (s \cdot m)$ (Associativity of scalar multiplication)
4. $1_R \cdot m = m$ if R has a multiplicative identity 1_R (Identity element of scalar multiplication)

The above definition is almost similar to the definition of a vector space, except that in a vector space the scalars come from a field, whereas in a module they come from a ring. One may think why we need modules when we have vector spaces. The reason is that not all rings are fields (in our case, $\mathcal{C}^\infty(M)$ is a ring but not a field), and the fact that the rings may not have multiplicative inverses for all non-zero elements give rise to ‘widely’ different structures than vector spaces.

Proposition 4.83 (Set of vector fields as a module)

Let M be a smooth manifold. Then the set of all vector fields on M , denoted by $\mathfrak{X}(M)$, forms a module over the ring of smooth functions $\mathcal{C}^\infty(M)$, with the following operations:

$$\begin{aligned} \oplus : \mathfrak{X}(M) \times \mathfrak{X}(M) &\rightarrow \mathfrak{X}(M) & \text{where } \forall p \in M, (\sigma_1 \oplus \sigma_2)(p) &:= \sigma_1(p) + \sigma_2(p), \\ (\sigma_1, \sigma_2) &\mapsto \sigma_1 \oplus \sigma_2 \end{aligned} \quad (4.161)$$

and

$$\begin{aligned} \odot : \mathcal{C}^\infty(M) \times \mathfrak{X}(M) &\rightarrow \mathfrak{X}(M) & \text{where } \forall p \in M, (f \odot \sigma)(p) &:= f(p) \cdot \sigma(p). \\ (f, \sigma) &\mapsto f \odot \sigma \end{aligned} \quad (4.162)$$

A non-trivial fact here is that unlike vector spaces, a module generically does not have a basis (unless the ring is a division ring).

Theorem 4.84 (Module Basis Theorem)

If D is a division ring, then every module over D has a basis.

Corollary 4.85

Every vector space has a basis.

Pf: Since a vector space is a module over a field, and a field is a division ring, the theorem applies directly. $\square$

Before we actually prove the theorem, let’s see some examples of modules on non-division rings which have bases and which do not have bases.

Example 4.86 (Vector fields on $\mathbb{R}^2$)

Let our smooth manifold be $M = \mathbb{R}^2$. The set of smooth functions on $\mathbb{R}^2$ is $\mathcal{C}^\infty(\mathbb{R}^2)$, and the set of vector fields on $\mathbb{R}^2$ is $\mathfrak{X}(\mathbb{R}^2)$. Coincidentally, $\mathfrak{X}(\mathbb{R}^2)$ has a basis over the ring $\mathcal{C}^\infty(\mathbb{R}^2)$. Consider the two vector fields defined by

$$\mathbf{e}_1(p) := \left(\frac{\partial}{\partial x} \right)_p, \quad \mathbf{e}_2(p) := \left(\frac{\partial}{\partial y} \right)_p, \quad \forall p = (x, y) \in \mathbb{R}^2.$$

Then, any vector field $\sigma \in \mathfrak{X}(\mathbb{R}^2)$ can be expressed as

$$\sigma = \sigma^1 \odot \mathbf{e}_1 + \sigma^2 \odot \mathbf{e}_2,$$

where $\sigma^1, \sigma^2 \in \mathcal{C}^\infty(\mathbb{R}^2)$ are smooth functions on $\mathbb{R}^2$. Thus, $\{\mathbf{e}_1, \mathbf{e}_2\}$ forms a basis of the module $\mathfrak{X}(\mathbb{R}^2)$ over the ring $\mathcal{C}^\infty(\mathbb{R}^2)$.

In the above example, we were lucky to find a basis for the module of vector fields on $\mathbb{R}^2$. However, in the next example we will see a module which does not have a basis.

Example 4.87 (Vector fields on S^2)

Let our smooth manifold be $M = S^2$, the 2-sphere. The set of smooth functions on S^2 is $\mathcal{C}^\infty(S^2)$, and the set of vector fields on S^2 is $\mathfrak{X}(S^2)$. It turns out that $\mathfrak{X}(S^2)$ does not have a basis over the ring $\mathcal{C}^\infty(S^2)$. This is a consequence of the Hairy Ball Theorem, which states that there is no non-vanishing continuous tangent field on even-dimensional spheres. Since a basis would require the ability to express any vector field as a linear combination of basis elements, the non-existence of a non-vanishing vector

field implies that no such basis can exist for $\mathfrak{X}(S^2)$.

One may be tempted to think then why don't we just work with the set of vector fields as a vector space over $\mathbb{R}$ instead of as a module over $C^\infty(M)$. The reason is that if we consider $\mathfrak{X}(M)$ as a vector space over $\mathbb{R}$, then we are only allowed to do scalar multiplication with constant real numbers, which in turn means we are only considering 'constant' vector fields, which is not very useful. By considering $\mathfrak{X}(M)$ as a module over $C^\infty(M)$, we can have scalar multiplication with smooth functions, allowing for a much richer structure of vector fields that can vary from point to point on the manifold.

The notions, methods used in the proof of this theorem is so ubiquitous in mathematics that we are not going to skip it. However, the proof requires some preparation since it uses the [Axiom of Choice](#), in the incarnation of Zorn's Lemma.

§4.8.1 Zorn's Lemma

Let's first state the lemma and then define the necessary terms.

Lemma 4.88 (Zorn's Lemma)

Let $(P, \leq)$ be a partially ordered set in which every chain has an upper bound in P . Then, P contains at least one maximal element.

Definition 4.89 (Partially Ordered Set):

A partially ordered set (or poset) is a set P equipped with a binary relation $\leq$ that satisfies the following properties for all $a, b, c \in P$:

- Reflexivity: $a \leq a$.
- Antisymmetry: If $a \leq b$ and $b \leq a$, then $a = b$.
- Transitivity: If $a \leq b$ and $b \leq c$, then $a \leq c$.

Definition 4.90 (Chain):

A chain in a partially ordered set $(P, \leq)$ is a subset $C \subseteq P$ such that for every pair of elements $a, b \in C$, either $a \leq b$ or $b \leq a$. In other words, every two elements in the chain are comparable.

This can be thought of as a totally ordered subset of the poset. Giving us following property:

- Totality: For all $a, b \in C$, either $a \leq b$ or $b \leq a$.

Definition 4.91 (Upper Bound):

Let $(P, \leq)$ be a partially ordered set and $S \subseteq P$ be a subset. An element $u \in P$ is called an upper bound of S if

$$\forall s \in S : s \leq u. \quad (4.163)$$

Definition 4.92 (Maximal Element):

Let $(P, \leq)$ be a partially ordered set. An element $m \in P$ is called a maximal element of P if

$$\nexists x \in P : m \leq x. \quad (4.164)$$

We will not prove Zorn's Lemma here, but it can be shown that in Zermelo-Fraenkel set theory (without the Axiom of Choice), Zorn's Lemma is equivalent to the Axiom of Choice. Thus, if we accept the Axiom of Choice as an axiom of our set theory, we can use Zorn's Lemma as a theorem.

§4.8.2 Proof of Module Basis Theorem

Proof (Theorem 4.84):

Let M be a module over a division ring D . We want to show that M has a basis. Consider a generating system $S \subseteq M$ of M , *i.e.*,

$$\forall \mathbf{m} \in M, \exists \mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n \in S, \exists m^1, m^2, \dots, m^n \in D : \mathbf{m} = m^i \cdot \mathbf{e}_i. \quad (4.165)$$

There exists such a generating system since we can always take $S = M$. Now, consider the set P of all linearly independent subsets of S , *i.e.*,

$$P := \{L \in \mathcal{P}(S) \mid L \text{ is linearly independent}\}. \quad (4.166)$$

We can partially order P by set inclusion $\subseteq$. Now, we want to show that every chain in P has an upper bound in P . Let $C \subseteq P$ be a chain, and define

$$U := \bigcup_{L \in C} L. \quad (4.167)$$

We claim that U is an upper bound of C in P . From the definition of U , it is clear that for every $L \in C$, we have $L \subseteq U$. Now, we need to show that U is linearly independent, thus $U \in P$. Suppose not, then there exist finitely many elements $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n \in U$ and scalars $d^1, d^2, \dots, d^n \in D$, not all zero, such that

$$d^i \cdot \mathbf{u}_i = 0. \quad (4.168)$$

Since each $\mathbf{u}_i$ belongs to U , there exists some $L_i \in C$ such that $\mathbf{u}_i \in L_i$. Because C is a chain, there exists some $L^* \in C$ such that $L_i \subseteq L^*$ for all i . Thus, all $\mathbf{u}_i$ belong to L^* . But this contradicts the linear independence of L^* , therefore, U must be linearly independent, and hence $U \in P$. Thus, every chain in P has an upper bound in P . By Zorn's Lemma, P contains at least one maximal element, say $\mathcal{B}$.

We claim that $\mathcal{B}$ is a basis of M . To see this, it is enough to show that $\mathcal{B}$ generates S , since S generates M . Let $\mathbf{m} \in S$. If $\mathbf{m} \in \mathcal{B}$, then we are done. Suppose $\mathbf{m} \notin \mathcal{B}$. Consider the set $\mathcal{B}' := \mathcal{B} \cup \{\mathbf{m}\}$. If $\mathcal{B}'$ is linearly independent, then it contradicts the maximality of $\mathcal{B}$. Therefore, $\mathcal{B}'$ must be linearly dependent. Thus, there exist finitely many elements $\mathbf{b}_1, \mathbf{b}_2, \dots, \mathbf{b}_n \in \mathcal{B}$ and scalars $d, d^1, d^2, \dots, d^n \in D$, not all zero, such that

$$d \cdot \mathbf{m} + d^i \cdot \mathbf{b}_i = 0. \quad (4.169)$$

If $d = 0$, then we have a linear dependence relation among the $\mathbf{b}_i$, contradicting the linear independence of $\mathcal{B}$. Therefore, $d \neq 0$. Since D is a division ring, d has a multiplicative inverse d^{-1} . Multiplying the above equation by d^{-1} , we get

$$\mathbf{m} = -(d^{-1}d^i) \cdot \mathbf{b}_i. \quad (4.170)$$

This shows that $\mathbf{m}$ can be expressed as a linear combination of elements in $\mathcal{B}$. Thus, $\mathcal{B}$ generates S , and hence M . Therefore, $\mathcal{B}$ is a basis of M . Q.E.D.

Till now in our theoretical physics journey, whenever we have dealt with vector fields, we always thought of them in terms of their components with respect to some basis. However, with this formalism, we see that in general vector fields may not have a basis, and thus we cannot always express them in terms of components. Locally, we can still express vector fields in terms of components using local charts, and then to relate vector fields in different charts we use the concept of transition maps.

§4.8.3 Module Construction and Important Terms

Definition 4.93 (Direct Sum of Modules):

Let M and N be modules over a ring R . The direct sum of M and N , denoted by $M \oplus N$, is defined as the set of ordered pairs

$$M \oplus N := M \times N = \{(m, n) \mid m \in M, n \in N\}, \quad (4.171)$$

equipped with the component-wise addition and scalar multiplication:

- Addition: For $(m_1, n_1), (m_2, n_2) \in M \oplus N$,

$$(m_1, n_1) + (m_2, n_2) := (m_1 + m_2, n_1 + n_2). \quad (4.172)$$

- Scalar Multiplication: For $r \in R$ and $(m, n) \in M \oplus N$,

$$r \cdot (m, n) := (r \cdot m, r \cdot n). \quad (4.173)$$

Let's define some important terms associated with the modules.

Definition (Modules):

Let M be a module over a ring R . We define the following:

- M is finitely generated if it has a finite generating set.
- M is free if it has a basis.
- M is projective if it is a direct summand of a free module, i.e., there exists an R -module N such that $M \oplus N$ is a free module.

As we have seen, $\mathfrak{X}(\mathbb{R}^2)$ is a free module, while $\mathfrak{X}(S^2)$ is not free. Also, it is clear that every free module is projective, but the converse is not true in general. Similar to vector spaces, we can define linear maps between modules.

Definition 4.94 (Module Homomorphism and Isomorphism):

Let M and N be modules over a ring R . A function $f : M \rightarrow N$ is called a module homomorphism if for all $m_1, m_2 \in M$ and $r \in R$, the following properties hold:

- Additivity: $f(m_1 + m_2) = f(m_1) + f(m_2)$.
- Scalar Compatibility: $f(r \cdot m_1) = r \cdot f(m_1)$.

If a module homomorphism $f : M \rightarrow N$ is bijective, then it is called a module isomorphism, and we say that M and N are isomorphic modules, denoted by $M \cong_{\text{mod}} N$.

Proposition 4.95

Let F be a finitely generated free module over a ring R , and let n be the cardinality of the generating set (or basis) of F . Then,

$$F \cong_{\text{mod}} R^n := \underbrace{R \oplus R \oplus \dots \oplus R}_{n \text{ times}}.$$

Proof:

Let $\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\}$ be a basis of F . Define a map $f : F \rightarrow R^n$ by

$$f\left(\sum_{i=1}^n r^i \cdot \mathbf{e}_i\right) := (r^1, r^2, \dots, r^n),$$

for all $r^i \in R$. It is straightforward to verify that f is a module homomorphism. Moreover, f is bijective since every element in R^n can be uniquely expressed as a linear combination of the basis elements of F . Therefore, f is a module isomorphism, and hence $F \cong_{\text{mod}} R^n$. Q.E.D.

With this, we can define the notion of dimension for finitely generated free modules, as the cardinality of their basis. And once we show that $R^m \cong_{\text{mod}} R^n$ implies $m = n$, we can conclude that the dimension is well-defined.

Coming back to our main discussion on vector fields, we define vector fiber bundles.

Definition 4.96 (Vector Bundle):

Let M be a smooth manifold. A vector bundle over M is a smooth fiber bundle (E, π, M) such that for each $p \in M$, the fiber $F_p := \pi^{-1}(p)$ is a vector space. Moreover, the local trivializations $\varphi_U : \pi^{-1}(U) \rightarrow U \times V$ are required to be vector space isomorphisms on each fiber, where V is a fixed vector space called the typical fiber of the bundle.

Similar to vector fields, we can define smooth sections of a vector bundle.

Definition 4.97 (Smooth Section of a Vector Bundle):

Let (E, π, M) be a vector bundle over a smooth manifold M . A smooth section of the vector bundle is a smooth map $\sigma : M \rightarrow E$ such that $\pi \circ \sigma = \text{id}_M$.

The set of all smooth sections of a vector bundle (E, π, M) is denoted by $\Gamma(E)$. Similar to vector fields, $\Gamma(E)$ forms a module over the ring $\mathcal{C}^\infty(M)$ with pointwise addition and scalar multiplication.

With this setup, we can now state the Serre-Swan theorem.

Theorem 4.98 (Serre, Swan, et al.)

Let (E, M, π) be a vector bundle over a smooth manifold M . Then, the module of smooth sections $\Gamma(E)$ of the vector bundle E is a finitely generated projective module over the ring $\mathcal{C}^\infty(M)$.

Remark 4.99. Using the Serre-Swan theorem, there exists a $\mathcal{C}^\infty(M)$ -module Q such that $\Gamma(E) \oplus Q$ is a free module. If for some vector bundle E , we can choose Q to be the zero module, then $\Gamma(E)$ itself is a free module. This happens, for example, when E is the tangent bundle of $\mathbb{R}^2$, as we have seen earlier.

Thus, in some sense, Q quantifies the obstruction to the existence of a basis for the module of sections $\Gamma(E)$.

Proposition 4.100

Let P and Q be finitely generated (projective) modules over a commutative ring R . Then

$$\text{Hom}_R(P, Q) := \{\phi : P \rightarrow Q \mid \phi \text{ is an } R\text{-module homomorphism}\} \quad (4.174)$$

is also a finitely generated (projective) module over R .

We will not prove this proposition here, but it can be shown using the properties of finitely generated and projective modules, along with the definition of module homomorphisms. This result is important because it allows us to construct new modules from existing ones, like the module of covector fields from the module of vector fields.

Let us define dual of a module, specifically the dual of the module of vector fields, which will give us the module of covector fields.

$$\mathfrak{X}^*(M) := \text{Hom}_{\mathcal{C}^\infty(M)}(\mathfrak{X}(M), \mathcal{C}^\infty(M)). \quad (4.175)$$

Elements of $\mathfrak{X}^*(M)$ are called *covector fields*. In the notation of set of smooth sections of a vector bundle, we can write

$$\mathfrak{X}(M) = \Gamma(\mathcal{T}M), \quad \mathfrak{X}^*(M) = \Gamma(\mathcal{T}M)^*,$$

where $\mathcal{T}M$ is the tangent bundle of M , and $\Gamma(\mathcal{T}M)^*$ denotes the dual module of $\Gamma(\mathcal{T}M)$.

Proposition 4.101

The dual of the module of vector fields $\Gamma(\mathcal{T}M)^*$ same as the module of smooth sections of the cotangent bundle $\mathcal{T}^*M$ over M , i.e.,

$$\Gamma(\mathcal{T}M)^* \cong_{\text{mod}} \Gamma(\mathcal{T}^*M). \quad (4.176)$$

With this, we have established a solid foundation for understanding vector fields, covector fields, and their associated modules on smooth manifolds. Now, let's move on to discuss tensor fields, due to this new understanding of dual modules, we can define tensor fields in two equivalent ways.

Definition 4.102 (Tensor Field):

Let M be a smooth manifold. A tensor field t of type (r, s) on M is a smooth section of the tensor bundle $\mathcal{T}_s^r M$, where

$$\mathcal{T}_s^r M := \bigsqcup_{p \in M} \mathcal{T}_s^r(\mathcal{T}_p M). \quad (4.177)$$

Alternatively, a tensor field of type (r, s) can be defined as a $\mathcal{C}^\infty(M)$ -multilinear map

$$t : \underbrace{\mathfrak{X}^*(M) \times \dots \times \mathfrak{X}^*(M)}_{r \text{ times}} \times \underbrace{\mathfrak{X}(M) \times \dots \times \mathfrak{X}(M)}_{s \text{ times}} \rightarrow \mathcal{C}^\infty(M). \quad (4.178)$$

In other words, a tensor field of type (r, s) can be viewed either as a smooth assignment of a tensor of type (r, s) to each point of the manifold or as a multilinear map that takes r covector fields and s vector fields and produces a smooth function on the manifold.

Similar to module of vector fields and covector fields, the set of all tensor fields of type (r, s) on M , denoted by $\Gamma(\mathcal{T}_s^r M)$, forms a module over the ring $\mathcal{C}^\infty(M)$ with pointwise addition and scalar multiplication. Also, we can define tensor product of tensor fields

$$\otimes : \Gamma(\mathcal{T}_s^r M) \times \Gamma(\mathcal{T}_{s'}^{r'} M) \rightarrow \Gamma(\mathcal{T}_{s+s'}^{r+r'} M), \quad (4.179)$$

given by

$$\begin{aligned} (t \otimes t')(\omega^1, \dots, \omega^r, \omega^{r+1}, \dots, \omega^{r+r'}, X_1, \dots, X_s, X_{s+1}, \dots, X_{s+s'}) \\ := t(\omega^1, \dots, \omega^r, X_1, \dots, X_s) \cdot t'(\omega^{r+1}, \dots, \omega^{r+r'}, X_{s+1}, \dots, X_{s+s'}), \end{aligned} \quad (4.180)$$

for all $t \in \Gamma(\mathcal{T}_s^r M)$, $t' \in \Gamma(\mathcal{T}_{s'}^{r'} M)$, $\omega^i \in \mathfrak{X}^*(M)$, and $X_j \in \mathfrak{X}(M)$.

LECTURE 12

§4.9 Differential Forms and de Rham Cohomology

Definition 4.103 (Differential n -form):

Let M be a smooth manifold, and $0 \leq n \leq \dim M$. A differential n -form on M is $(0, n)$ -tensor field ω that is totally antisymmetric, i.e.,

$$\omega(X_1, \dots, X_n) = \text{sgn}(\sigma) \omega(X_{\sigma(1)}, \dots, X_{\sigma(n)})$$

where $\sigma \in S_n = \text{Perm}(n)$ is a permutation of n elements, $\text{sgn}(\sigma)$ is the sign of the permutation, and $X_1, \dots, X_n \in \Gamma(\mathcal{T}M)$ are vector fields on M .

Let's look at some examples of differential forms.

Example 4.104

- a) A manifold M is *orientable* if it admits an oriented atlas, i.e., an atlas $\{(U_\alpha, x_\alpha)\}$ in which all the transition maps $x_\beta \circ x_\alpha^{-1} : x_\alpha(U_\alpha \cap U_\beta) \rightarrow x_\beta(U_\alpha \cap U_\beta)$ have positive determinant.

If M is orientable, then there exists a nowhere vanishing *top-form* ($n = \dim M$) ω on M ,

providing a volume.

- b) The electromagnetic field strength F in electrodynamics is a differential 2-form.
- c) In classical mechanics, define the configuration space Q of a system as the space of all possible configurations of the system, *i.e.*, the space of all possible values of the generalized coordinates. Then the phase space of the system is the cotangent bundle $\mathcal{T}^*Q$ of the configuration space.

On phase space, we have a canonical 2-form ω called the *symplectic form*. To define it, we need some more machinery, but the anti-symmetry of ω is the reason why we have the minus sign in Hamilton's equations of motion.

Define the set of all differential n -forms on M as $\Omega^n(M)$, which by definition is a $\mathcal{C}^\infty(M)$ -module. Generally, taking a tensor product of a form doesn't give us a form, but we want to be able to multiply forms together to get one, so we define the *wedge product*.

Definition 4.105 (Wedge Product):

Let $\omega \in \Omega^n(M)$ and $\eta \in \Omega^m(M)$ be two differential forms on M . The wedge product of ω and η is a differential $(n+m)$ -form defined as

$$\begin{aligned} (\omega \wedge \eta)(X_1, \dots, X_{n+m}) &= \frac{1}{n!} \frac{1}{m!} \sum_{\sigma \in S_{n+m}} \text{sgn}(\sigma) (\omega \otimes \eta)(X_{\sigma(1)}, \dots, X_{\sigma(n+m)}) \\ &= \frac{1}{n!} \frac{1}{m!} \sum_{\sigma \in S_{n+m}} \text{sgn}(\sigma) \omega(X_{\sigma(1)}, \dots, X_{\sigma(n)}) \eta(X_{\sigma(n+1)}, \dots, X_{\sigma(n+m)}). \end{aligned}$$

Simplest example of wedge product is of two 1-forms ω and η , then

$$(\omega \wedge \eta)(X_1, X_2) = \sum_{\sigma \in S_2} \text{sgn}(\sigma) \omega(X_{\sigma(1)}) \eta(X_{\sigma(2)}) = \omega(X_1) \eta(X_2) - \omega(X_2) \eta(X_1).$$

Thus, we can write $\omega \wedge \eta = \omega \otimes \eta - \eta \otimes \omega$.

Recall that given a smooth map $\phi : M \rightarrow N$ between two smooth manifolds, we induce pullbacks $\phi_p^* : \mathcal{T}_{\phi(p)}^* N \rightarrow \mathcal{T}_p^* M$ on the cotangent spaces for each $p \in M$. With this global pull-back map $\phi^* : \Gamma(\mathcal{T}^* N) \equiv \Omega^1(N) \rightarrow \Gamma(\mathcal{T}^* M) \equiv \Omega^1(M)$ is defined pointwise.

Definition 4.106 (Pull-back of Forms):

Let $\phi : M \rightarrow N$ be a smooth map between two smooth manifolds, and let $\omega \in \Omega^n(N)$ be a differential n -form on N . The pull-back of ω by ϕ is a differential n -form on M defined as

$$(\phi^* \omega)(X_1, \dots, X_n) = \omega(\phi_*(X_1), \dots, \phi_*(X_n))$$

where $\phi_* : \mathcal{T}M \rightarrow \mathcal{T}N$ is the (global) push-forward map induced by ϕ .

This can be equivalently defined pointwise as $(\phi^* \omega)_p = \phi_{\phi(p)}^* \omega_{\phi(p)}$ for each $p \in M$.