

$$0 = 1 - 1 = -1 + 1 = 0$$

From Elementary Arithmetic to Intrinsic Skein-Lasagna

Keyao Peng

Université de Bourgogne

Your favorite equations?

Your favorite equations?

- $a^2 + b^2 = c^2$
- $e^{i\pi} + 1 = 0$
- $\int_{\partial D} \omega = \int_D d\omega$
- ...

Your favorite equations?

- $a^2 + b^2 = c^2$
- $e^{i\pi} + 1 = 0$
- $\int_{\partial D} \omega = \int_D d\omega$
- ...

But, what is an **equation**?

Your favorite equations?

- $a^2 + b^2 = c^2$
- $e^{i\pi} + 1 = 0$
- $\int_{\partial D} \omega = \int_D d\omega$
- ...

But, what is an **equation**?

$A = B$ says A **is** equal to B , **how** they are equal depends on the *proof*.

Your favorite equations?

- $a^2 + b^2 = c^2$
- $e^{i\pi} + 1 = 0$
- $\int_{\partial D} \omega = \int_D d\omega$
- ...

But, what is an **equation**?

$A = B$ says A **is** equal to B , **how** they are equal depends on the *proof*.

Question

Can you prove $0 = 0$, non-trivially?

Test your math level

Test your math level

$$S^3 \sim \mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{CP}^1 \sim S^2$$



$$\pi_4(S^3) \cong \mathbb{Z}/2\mathbb{Z}$$

$$\pi_1^s(S) \cong \Omega_1^{\text{fr}}$$

$$K_1(\mathbb{F}_1) \cong S_\infty^{\text{ab}}$$

$$0=1-1=1+1=0$$

imgflip.com



Hopf fibration is a map defined by

$$H : \mathbb{S}^3 \subset \mathbb{C}^2 - \{0\} \rightarrow \mathbb{CP}^1 \cong \mathbb{S}^2, (z_1, z_2) \mapsto [z_1, z_2]$$


For any point $x \in \mathbb{S}^2$, preimage $H^{-1}(x)$ is a circle $\mathbb{S}^1$.

Hopf fibration

Hopf fibration is a map defined by

$$H : \mathbb{S}^3 \subset \mathbb{C}^2 - \{0\} \rightarrow \mathbb{CP}^1 \cong \mathbb{S}^2, (z_1, z_2) \mapsto [z_1, z_2]$$

For any point $x \in \mathbb{S}^2$, preimage $H^{-1}(x)$ is a circle $\mathbb{S}^1$. We can visualize this map by considering $H : \mathbb{R}^3 \cong \mathbb{S}^3 - \{\infty\} \rightarrow \mathbb{S}^2$:



Homotopy groups

To understand a space X , we use some homotopy invariants:


To understand a space X , we use some homotopy invariants:

- $\pi_0 X \in \text{Set}$, connected component, how many "pieces" X has.

Homotopy groups

To understand a space X , we use some homotopy invariants:

- $\pi_0 X \in \text{Set}$, connected component, how many "pieces" X has.
- Chose a base point $x \in X$, Let $\text{Map}_*(S^1, X) \in \text{Space}$ be the **space** of loops in X started from x . Here Map_* is the **space** of pointed continue maps, the continue structure of it comes from deform maps.



To understand a space X , we use some homotopy invariants:

- $\pi_0 X \in \text{Set}$, connected component, how many "pieces" X has.
- Chose a base point $x \in X$, Let $\text{Map}_*(\mathbb{S}^1, X) \in \text{Space}$ be the **space** of loops in X started from x . Here Map_* is the **space** of pointed continuous maps, the continuous structure of it comes from deformation maps.
- $\pi_1(X, x) := \pi_0 \text{Map}_*(\mathbb{S}^1, X) \in \text{Group}$, how many loops in X up to deformation. This is fundamental group.

To understand a space X , we use some homotopy invariants:

- $\pi_0 X \in \text{Set}$, connected component, how many "pieces" X has.
- Chose a base point $x \in X$, Let $\text{Map}_*(\mathbb{S}^1, X) \in \text{Space}$ be the **space** of loops in X started from x . Here Map_* is the **space** of pointed continuous maps, the continuous structure of it comes from deformation maps.
- $\pi_1(X, x) := \pi_0 \text{Map}_*(\mathbb{S}^1, X) \in \text{Group}$, how many loops in X up to deformation. This is fundamental group.
- $i > 1$, $\pi_i(X, x) := \pi_0 \text{Map}_*(\mathbb{S}^i, X) \cong \pi_0 \text{Map}_*(\mathbb{S}^1, \text{Map}_*(\mathbb{S}^{i-1}, X)) \in \text{Abel}$. This is homotopy group.

To understand a space X , we use some homotopy invariants:

- $\pi_0 X \in \text{Set}$, connected component, how many "pieces" X has.
- Chose a base point $x \in X$, Let $\text{Map}_*(\mathbb{S}^1, X) \in \text{Space}$ be the **space** of loops in X started from x . Here Map_* is the **space** of pointed continuous maps, the continuous structure of it comes from deformation maps.
- $\pi_1(X, x) := \pi_0 \text{Map}_*(\mathbb{S}^1, X) \in \text{Group}$, how many loops in X up to deformation. This is fundamental group.
- $i > 1, \pi_i(X, x) := \pi_0 \text{Map}_*(\mathbb{S}^i, X) \cong \pi_0 \text{Map}_*(\mathbb{S}^1, \text{Map}_*(\mathbb{S}^{i-1}, X)) \in \text{Abel}$. This is homotopy group.



Let us study the cases $X = \mathbb{S}^n$.



- For $i < n$, $\pi_i(\mathbb{S}^n) = \{0\}$, where $0 : \mathbb{S}^i \rightarrow * \xrightarrow{\text{base}} \mathbb{S}^n$.

- For $i < n$, $\pi_i(\mathbb{S}^n) = \{0\}$, where $0 : \mathbb{S}^i \rightarrow * \xrightarrow{\text{base}} \mathbb{S}^n$.
- For $\pi_n(\mathbb{S}^n)$, let's consider the space $\text{End}_*(\mathbb{S}^n) := \text{Map}_*(\mathbb{S}^n, \mathbb{S}^n)$.



- For $i < n$, $\pi_i(\mathbb{S}^n) = \{0\}$, where $0 : \mathbb{S}^i \rightarrow * \xrightarrow{\text{base}} \mathbb{S}^n$.
- For $\pi_n(\mathbb{S}^n)$, let's consider the space $\text{End}_*(\mathbb{S}^n) := \text{Map}_*(\mathbb{S}^n, \mathbb{S}^n)$.

Let $f \in \text{End}_*(\mathbb{S}^n)$, take a small disk $D_\epsilon^n \subset \mathbb{S}^n$ away from the base point. Then the preimage $f^{-1}(D_\epsilon^n) \subset \mathbb{S}^n - \{*\} \cong \mathbb{R}^n$ is a disjoint union of disks for ϵ small enough. Notice that the disks in the union have orientations $+$ and $-$.






deform



Homotopy groups of spheres

- For $i < n$, $\pi_i(\mathbb{S}^n) = \{0\}$, where $0 : \mathbb{S}^i \rightarrow * \xrightarrow{\text{base}} \mathbb{S}^n$.
- For $\pi_n(\mathbb{S}^n)$, let's consider the space $\text{End}_*(\mathbb{S}^n) := \text{Map}_*(\mathbb{S}^n, \mathbb{S}^n)$.

Let $f \in \text{End}_*(\mathbb{S}^n)$, take a small disk $D_\epsilon^n \subset \mathbb{S}^n$ away from the base point. Then the preimage $f^{-1}(D_\epsilon^n) \subset \mathbb{S}^n - \{*\} \cong \mathbb{R}^n$ is a disjoint union of disks for ϵ small enough. Notice that the disks in the union have orientations $+$ and $-$.



- For $i < n$, $\pi_i(\mathbb{S}^n) = \{0\}$, where $0 : \mathbb{S}^i \rightarrow * \xrightarrow{\text{base}} \mathbb{S}^n$.
- For $\pi_n(\mathbb{S}^n)$, let's consider the space $\text{End}_*(\mathbb{S}^n) := \text{Map}_*(\mathbb{S}^n, \mathbb{S}^n)$.

Let $f \in \text{End}_*(\mathbb{S}^n)$, take a small disk $D_\epsilon^n \subset \mathbb{S}^n$ away from the base point. Then the preimage $f^{-1}(D_\epsilon^n) \subset \mathbb{S}^n - \{*\} \cong \mathbb{R}^n$ is a disjoint union of disks for ϵ small enough. Notice that the disks in the union have orientations $+$ and $-$.

This gives an isomorphism to the configuration space of disks.

$$\text{End}_*(\mathbb{S}^n) \cong \text{Conf}^{fr}(\mathbb{R}^n) = \bigcup_{j,k} \text{Emb}(\bigsqcup_j D_+^n \sqcup \bigsqcup_k D_-^n, \mathbb{R}^n)$$

Therefore, $\pi_0 \text{End}_*(\mathbb{S}^n) \cong \mathbb{Z}$, $(j, k) \mapsto j - k$

Higher homotopy groups of spheres

- For $i > n$, $\pi_i(\mathbb{S}^n) = ?$

Higher homotopy groups of spheres

- For $i > n$, $\pi_i(\mathbb{S}^n) = ?$
- For $i > 1$, $\pi_i(\mathbb{S}^1) = \{0\}$.

Higher homotopy groups of spheres

- For $i > n$, $\pi_i(\mathbb{S}^n) = ?$
- For $i > 1$, $\pi_i(\mathbb{S}^1) = \{0\}$.
- But $\pi_3(\mathbb{S}^2) = \pi_0 \text{Map}_*(\mathbb{S}^3, \mathbb{S}^2) \ni H$, and $H \not\approx 0$, i.e. H can not deform to 0

How to see that?

- For $i > n$, $\pi_i(\mathbb{S}^n) = ?$
- For $i > 1$, $\pi_i(\mathbb{S}^1) = \{0\}$.
- But $\pi_3(\mathbb{S}^2) = \pi_0 \text{Map}_*(\mathbb{S}^3, \mathbb{S}^2) \ni H$, and $H \not\approx 0$, i.e. H can not deform to 0


How to see that?

$$\text{Map}_*(\mathbb{S}^{n+1}, \mathbb{S}^n) \cong \text{Map}_*(\mathbb{S}^1, \text{End}_*(\mathbb{S}^n)) \rightarrow \text{Map}_*([0, 1], \text{End}_*(\mathbb{S}^n))$$

Then H corresponds to a path (deformation or homotopy) from $0 \in \text{End}_*(\mathbb{S}^2)$ to 0 itself.


Visualize H as deformation

I made an animation to visualize this deformation.



Visualize H as deformation

I made an animation to visualize this deformation.



We can also visualize it as a movement in the configuration space.
We will show it in the end.

How to define $\mathbb{Z}$?

How to define $\mathbb{Z}$?

We can define natural number $\mathbb{N}$ as isomorphic classes of finite sets.

$$\text{FinSet}_{/\cong} \xrightarrow{|\cdot|} \mathbb{N}, |X \sqcup Y| = |X| + |Y|$$

How to define integral number $\mathbb{Z}$ with finite sets?

How to define $\mathbb{Z}$?

We can define natural number $\mathbb{N}$ as isomorphic classes of finite sets.

$$\text{FinSet}_{/\cong} \xrightarrow{|\cdot|} \mathbb{N}, |X \sqcup Y| = |X| + |Y|$$

How to define integral number $\mathbb{Z}$ with finite sets?

We use the **Grothendieck group** $K_0(\text{FinSet})$:

$$K_0(\text{FinSet}) = \{(X, Y) \in (\text{FinSet}_{/\cong})^2\} / \{(X \sqcup Z, Y \sqcup Z) \sim (X, Y)\}$$

We can think we formally define " $X - Y$ " in this way.

How to define $\mathbb{Z}$?

We can define natural number $\mathbb{N}$ as isomorphic classes of finite sets.

$$\text{FinSet}_{/\cong} \xrightarrow{|\cdot|} \mathbb{N}, |X \sqcup Y| = |X| + |Y|$$

How to define integral number $\mathbb{Z}$ with finite sets?

We use the **Grothendieck group** $K_0(\text{FinSet})$:

$$K_0(\text{FinSet}) = \{(X, Y) \in (\text{FinSet}_{/\cong})^2\} / \{(X \sqcup Z, Y \sqcup Z) \sim (X, Y)\}$$

We can think we formally define " $X - Y$ " in this way.


But what is $K_i(\text{FinSet})$ for $i > 0$?

Space of "finite sets"

Instead of thinking the set of "finite sets", we now consider the space of them $F = \text{FinSet}^{\cong}$: every point $x \in F$ corresponds to a finite set, and a path between two points corresponds to an isomorphism $f : x \xrightarrow{\cong} y$ of two finite sets.


Space of "finite sets"

Instead of thinking the set of "finite sets", we now consider the space of them $F = \text{FinSet}^{\cong}$: every point $x \in F$ corresponds to a finite set, and a path between two points corresponds to an isomorphism $f : x \xrightarrow{\cong} y$ of two finite sets.



Space of "finite sets"

Instead of thinking the set of "finite sets", we now consider the space of them $F = \text{FinSet}^{\cong}$: every point $x \in F$ corresponds to a finite set, and a path between two points corresponds to an isomorphism $f : x \xrightarrow{\cong} y$ of two finite sets.




- $\pi_0(F) = \text{FinSet}_{/\cong} \cong \mathbb{N}$.
- $\pi_1(F, x) = S_{|x|}$, the permutation group.
- $\pi_i(F, x) = \{0\}$ for $i > 1$ (one can never deform an isomorphism of set to another).

K-space (spectrum) of "finite sets"

We want to add "negative" to F . We can consider the space $K(F)$ of (ordered) finite sets, whose elements are marked by $+$ or $-$, and we also want to identify the sets with same "value" $\|X\| = |X_+| - |X_-|$, and these identifications will be the paths.

K-space (spectrum) of "finite sets"

We want to add "negative" to F . We can consider the space $K(F)$ of (ordered) finite sets, whose elements are marked by $+$ or $-$, and we also want to identify the sets with same "value" $\|X\| = |X_+| - |X_-|$, and these identifications will be the paths.



K-groups of "finite sets"

Let $K_i(\text{FinSet}) = \pi_i K(F)$ to be the higher K-groups, we have the following observations:

K-groups of "finite sets"

Let $K_i(\text{FinSet}) = \pi_i K(F)$ to be the higher K-groups, we have the following observations:

- $\pi_0 K(F) = \mathbb{Z}$, by $x \mapsto \|x\|$. This coincides with $K_0(\text{FinSet})$.

Let $K_i(\text{FinSet}) = \pi_i K(F)$ to be the higher K-groups, we have the following observations:

- $\pi_0 K(F) = \mathbb{Z}$, by $x \mapsto \|x\|$. This coincides with $K_0(\text{FinSet})$.
- We can construct a loop $h \in \pi_1 K(F)$ by using identifications

$$h : \emptyset \sim \{+, -\} \sim \{-, +\} \sim \emptyset$$

Let $K_i(\text{FinSet}) = \pi_i K(F)$ to be the higher K-groups, we have the following observations:

- $\pi_0 K(F) = \mathbb{Z}$, by $x \mapsto \|x\|$. This coincides with $K_0(\text{FinSet})$.
- We can construct a loop $h \in \pi_1 K(F)$ by using identifications

$$h : \emptyset \sim \{+, -\} \sim \{-, +\} \sim \emptyset$$

- In fact $\pi_1 K(F) = S_n^{ab} = \mathbb{Z}/2$ with h as the generator. That is to say, h is a non-trivial identification.

K-groups of "finite sets"

Let $K_i(\text{FinSet}) = \pi_i K(F)$ to be the higher K-groups, we have the following observations:

- $\pi_0 K(F) = \mathbb{Z}$, by $x \mapsto \|x\|$. This coincides with $K_0(\text{FinSet})$.
- We can construct a loop $h \in \pi_1 K(F)$ by using identifications

$$h : \emptyset \sim \{+, -\} \sim \{-, +\} \sim \emptyset$$

- In fact $\pi_1 K(F) = S_n^{ab} = \mathbb{Z}/2$ with h as the generator. That is to say, h is a non-trivial identification.

Answer

We can prove $0 = 0$ non-trivially with $h : 0 = 1 - 1 = -1 + 1 = 0$!

Algebraic K-groups of a field

Let k be a field, we can follow the same construction for finite dimensional k vector space. Let $V_k = \text{FinVect}_k^{\cong}$, we define **Algebraic K-groups** of k as $K_i(k) = \pi_i K(V_k)$.
(Notice that we can think finite sets as $\mathbb{F}_1$ vector spaces)

Let k be a field, we can follow the same construction for finite dimensional k vector space. Let $V_k = \text{FinVect}_k \cong$, we define

Algebraic K-groups of k as $K_i(k) = \pi_i K(V_k)$.

(Notice that we can think finite sets as $\mathbb{F}_1$ vector spaces)

Algebraic K-groups are important invariants with connections to geometry, topology, ring theory, and number theory, but are notoriously difficult to compute.

Let k be a field, we can follow the same construction for finite dimensional k vector space. Let $V_k = \text{FinVect}_k^{\cong}$, we define

Algebraic K-groups of k as $K_i(k) = \pi_i K(V_k)$.

(Notice that we can think finite sets as $\mathbb{F}_1$ vector spaces)

Algebraic K-groups are important invariants with connections to geometry, topology, ring theory, and number theory, but are notoriously difficult to compute.

- $K_0(k) = \mathbb{Z}$
- $K_1(k) = k^\times$
- $K_2(k) = k^\times \otimes_{\mathbb{Z}} k^\times / \langle x \otimes (1 - x) : x \in k \setminus \{0, 1\} \rangle$

Let k be a field, we can follow the same construction for finite dimensional k vector space. Let $V_k = \text{FinVect}_k^{\cong}$, we define

Algebraic K-groups of k as $K_i(k) = \pi_i K(V_k)$.

(Notice that we can think finite sets as $\mathbb{F}_1$ vector spaces)

Algebraic K-groups are important invariants with connections to geometry, topology, ring theory, and number theory, but are notoriously difficult to compute.

- $K_0(k) = \mathbb{Z}$
- $K_1(k) = k^\times$
- $K_2(k) = k^\times \otimes_{\mathbb{Z}} k^\times / \langle x \otimes (1 - x) : x \in k \setminus \{0, 1\} \rangle$
- $K_3(k) = ???$

What's more?

But what is the connection between these two topics?

What's more?

But what is the connection between these two topics?

- In fact, $K_i(\text{FinSet}) \cong \lim_{n \rightarrow \infty} \pi_{i+n}(\mathbb{S}^n) = \pi_i^s(\mathbb{S})$.

But what is the connection between these two topics?


- In fact, $K_i(\text{FinSet}) \cong \lim_{n \rightarrow \infty} \pi_{i+n}(\mathbb{S}^n) = \pi_i^s(\mathbb{S})$.
- So, $K(F)$ can be also seen as $\text{Conf}^{fr}(\mathbb{R}^\infty)$, and both the Hopf map H and h can be represented as movements within it.

What's more?

But what is the connection between these two topics?

- In fact, $K_i(\text{FinSet}) \cong \lim_{n \rightarrow \infty} \pi_{i+n}(\mathbb{S}^n) = \pi_i^s(\mathbb{S})$.
- So, $K(F)$ can be also seen as $\text{Conf}^{fr}(\mathbb{R}^\infty)$, and both the Hopf map H and h can be represented as movements within it.

And they are the same!




What's more?

But what is the connection between these two topics?

- In fact, $K_i(\text{FinSet}) \cong \lim_{n \rightarrow \infty} \pi_{i+n}(\mathbb{S}^n) = \pi_i^s(\mathbb{S})$.
- So, $K(F)$ can be also seen as $\text{Conf}^{fr}(\mathbb{R}^\infty)$, and both the Hopf map H and h can be represented as movements within it.

And they are the same!




New Question

Can we find geometric interpretation for general K-groups $K_i(k)$?


What about cobordism?

If we put the movement in 3D space, we get a circle:



What about cobordism?


If we put the movement in 3D space, we get a circle:



This provides a framed circle, meaning an embedding $S^1 \hookrightarrow \mathbb{R}^n$ with a **trivialization** τ of the normal bundle. We denote Ω_i^{fr} as the group of cobordism classes of i -dimensional framed manifolds.

What about cobordism?

If we put the movement in 3D space, we get a circle:



This provides a framed circle, meaning an embedding $S^1 \hookrightarrow \mathbb{R}^n$ with a **trivialization** τ of the normal bundle. We denote Ω_i^{fr} as the group of cobordism classes of i -dimensional framed manifolds. And we also have:

$$\Omega_1^{fr} \cong \pi_1^s(\mathbb{S})$$

Therefore, this framed circle is non-trivial and is the generator of Ω_1^{fr} .

A n -**treefold** is a space generally locally homeomorphic to $T \times \mathbb{R}^{n-1}$, where T is a tree.


A n -**treefold** is a space generally locally homeomorphic to $T \times \mathbb{R}^{n-1}$, where T is a tree. Give a symmetric monoidal category $(\mathcal{C}, \otimes)$, define an n - $(\mathcal{C}, \otimes)$ -**Lasagna** L as a framed n -treeifold $\mathcal{T}$ with a stratification $\mathcal{S}$ and the strata are marked compatibly with category $\mathcal{C}$:

Intrinsic Skein-Lasagna

A n -**treefold** is a space generally locally homeomorphic to $T \times \mathbb{R}^{n-1}$, where T is a tree. Give a symmetric monoidal category $(\mathcal{C}, \otimes)$, define an n - $(\mathcal{C}, \otimes)$ -**Lasagna** L as a framed n -treefold $\mathcal{T}$ with a stratification $\mathcal{S}$ and the strata are marked compatibly with category $\mathcal{C}$:

- 0-codimensional strata are marked with objects; For $i > 0$, i -codimensional strata are marked with i -morphism,
- At the branches of treefold marked with the monoidal product $\otimes$.

Example



Relation with "Skein-Lasagna module"

We do not ask for an ambient space X where lasagna embeds (or think $X = \mathbb{R}^\infty$). This means we study the intrinsic geometry of Lasagna itself.

Relation with "Skein-Lasagna module"

We do not ask for an ambient space X where lasagna embeds (or think $X = \mathbb{R}^\infty$). This means we study the intrinsic geometry of Lasagna itself.

We can think a Lasagna L as a string diagram of $\mathcal{C}$ drawing on $\mathcal{T}$, or another way to say $L \in \int_{\mathcal{T}}(\mathcal{C}, \otimes)$.

Cobordism of Lasagna

As one expects, we can define the cobordism relation between L_1, L_2 , if there is a Lasagna W with boundary such that $\partial W = L_1 \sqcup -L_2$.

Finally, we define the n -th $(\mathcal{C}, \otimes)$ -**cobordism group** $\Omega_n^{\mathcal{C}, \otimes}$ to be the abelian group generated by n -Lasagna modulo cobordism relations.

$(V_k, \oplus)$ -Cobodism Group

As an example $\Omega_n^{V_k, \oplus}$, where $V_k = \text{FinVect}_k^{\cong}$ is the subcategory of finite-dimensional vector space where the morphisms are all isomorphisms.

$(V_k, \oplus)$ -Cobodism Group


As an example $\Omega_n^{V_k, \oplus}$, where $V_k = \text{FinVect}_k^{\cong}$ is the subcategory of finite-dimensional vector space where the morphisms are all isomorphisms.

- It is clear $\Omega_0^{V_k, \oplus} \cong \mathbb{Z}$ is classified dimension.


$(V_k, \oplus)$ -Cobordism Group

As an example $\Omega_n^{V_k, \oplus}$, where $V_k = \text{FinVect}_k \cong$ is the subcategory of finite-dimensional vector space where the morphisms are all isomorphisms.

- It is clear $\Omega_0^{V_k, \oplus} \cong \mathbb{Z}$ is classified dimension.
- $\Omega_1^{V_k, \oplus}$ is generated by circle $L_V(f)$ marked with vector spaces V and matrix $f \in \text{GL}_V(k)$. And by adding identity matrices, we find we have a homomorphism $L : \text{GL}_\infty(k) \rightarrow \Omega_1^{V_k, \oplus}$.




We have one more relation given by surfaces:



$$L([f, g]) \sim 0$$

We have one more relation given by surfaces:



$$L([f, g]) \sim 0$$

Notice that $k^\times \cong \mathrm{GL}_\infty(k)/[\mathrm{GL}_\infty(k), \mathrm{GL}_\infty(k)]$. In fact, one can show the induced homomorphism $L : k^\times \xrightarrow{\cong} \Omega_1^{V_k, \oplus}$ is isomorphism. And the lasagna induced by Hopf map H is given by $L(-1)$.

Generalized Pontryagin–Thom Isomorphism

Recall that $K_0(k) = \mathbb{Z}$, $K_1(k) = k^\times$. This is not just a coincidence:


Generalized Pontryagin–Thom Isomorphism

We have the isomorphism

$$K_n(k) \xrightarrow{\cong} \Omega_n^{V_k, \oplus}$$

This gives us an abstract isomorphism, which indicates we can view Lasagna as elements in K -groups. So what about $n = 2$?


For matrices $f, g \in \mathrm{GL}(k)$ such that $[f, g] = 1$, one can define following 2-Lasagna $C(f, g)$:



Recall $K_2(k) = k^\times \otimes k^\times / \langle a \otimes 1 - a \rangle$, which means we can find a 3-Lasagna who bounds $C(a, 1 - a)$!

Thank you, hope you have got the brain upgrades!

$$S^3 \sim \mathbb{C}^2 \setminus \{0\} \rightarrow \mathbb{CP}^1 \sim S^2$$



$$\pi_4(S^3) \cong \mathbb{Z}/2\mathbb{Z}$$

$$\pi_1^s(S) \cong \Omega_1^{\text{fr}}$$

$$K_1(\mathbb{F}_1) \cong S_\infty^{\text{ab}}$$

$$0=1-1=1+1=0$$

imgfip.com

