

CAUSES OF DEATH.

CCXXXV

higher than that of the native whites (17). The death rates of males from these diseases (white, 26.9; colored, 29.9) exceeded those of the females (white, 18.6; colored, 14.1), particularly among the colored.

In comparison with 1890 there was a small decrease in the death rates from diseases of the liver in most of

the areas, the greatest being in the cities in the registration states.

The following table shows, for the registration states, in the aggregate, and for the cities and rural districts, the death rates from diseases of the liver in the census year, per 100,000 of population, in comparison with 1890:

DEATH RATES IN CITIES AND RURAL DISTRICTS.

REGISTRATION STATES.	AGGREGATE.			MALES.			FEMALES.		
	Total.	Cities.	Rural.	Total.	Cities.	Rural.	Total.	Cities.	Rural.
Total	21.7	22.6	20.4	25.0	27.2	22.3	18.3	18.1	18.4
1890..	23.2	26.5	18.1	24.9	29.3	18.3	21.5	23.7	17.9
Connecticut	20.6	21.6	18.8	25.5	27.0	23.0	15.6	16.2	14.6
1890..	25.2	26.8	24.1	26.0	32.2	21.6	24.4	21.5	26.5
District of Columbia.....	24.3	24.3		37.1	37.1		12.9	12.9	
1890..	24.3	24.3		32.9	32.9		16.6	16.6	
Maine ¹	19.8	21.2	19.4	22.8	28.3	21.7	16.5	14.5	17.0
Massachusetts	17.6	17.1	19.4	19.9	20.5	17.6	15.5	13.8	20.9
1890..	20.1	20.2	20.0	19.5	19.6	19.3	20.8	20.8	20.8
Michigan ¹	23.3	21.7	24.0	26.2	27.3	25.8	20.2	16.1	22.1
New Hampshire.....	15.9	18.9	13.9	20.5	24.9	17.8	11.2	13.4	9.7
1890..	24.4	29.0	22.6	28.9	30.7	28.3	20.0	27.4	16.7
New Jersey.....	20.4	23.0	16.9	23.9	27.7	19.0	16.9	18.6	14.6
1890..	25.3	29.2	20.3	27.3	33.6	19.3	23.3	24.9	21.3
New York.....	23.7	25.3	20.4	27.2	29.5	22.3	20.3	21.0	18.6
1890..	24.0	28.8	16.2	26.3	32.1	17.2	21.7	25.6	15.2
Rhode Island.....	22.8	22.5	23.4	27.5	27.0	28.7	18.4	18.5	17.9
1890..	25.5	32.0	16.5	27.4	39.6	11.1	23.7	25.0	21.8
Vermont	18.4	14.9	18.8	20.5	8.9	22.3	16.1	20.7	15.3
1890..	14.4	14.1	14.5	12.4	22.2	11.6	16.6	6.8	17.5

¹ Nonregistration in 1890.

The preceding table shows that the death rates from diseases of the liver in the registration states were highest in the District of Columbia (24.3) and New York (23.7), and lowest in New Hampshire (15.9) and Massachusetts (17.6). There was but little difference in the death rates from these diseases in the cities and rural districts. The death rate of males from diseases of the liver (25)

was considerably higher than that of females (18.3) in all of the registration states.

The following table shows, for the registration area and its subdivisions, the death rates from diseases of the liver among the whites during the census year, per 100,000 of white population, by birthplaces of mothers:

DEATH RATES BY BIRTHPLACES OF MOTHERS.

BIRTHPLACES OF MOTHERS.	REGISTRATION RECORD.					
	Total.	Cities.	States.			Cities in other states.
			Total.	Cities.	Rural.	
United States.....	15.6	13.8	16.0	13.9	17.7	13.5
Ireland	34.7	36.0	34.6	36.0	29.2	35.8
Germany	28.7	29.7	28.1	29.9	23.1	29.5
England and Wales	26.3	25.7	26.1	25.3	27.4	27.3
Canada.....	13.9	12.6	14.2	13.0	15.7	8.7
Scandinavia.....	12.4	12.3	15.6	17.4	12.6	6.7
Scotland	24.8	25.8	24.6	25.6	22.3	26.3
Italy	18.7	20.7	16.5	18.3	7.2	41.8
France	52.9	57.1	35.7	34.9	37.5	99.0
Hungary and Bohemia.....	21.6	23.3	17.5	19.5	5.3	29.3
Russia and Poland.....	9.8	9.8	9.6	9.4	11.1	10.8
Other foreign.....	17.2	16.7	16.5	15.7	18.7	20.6

The preceding table shows that the death rates due to diseases of the liver in the registration area were highest among those whose mothers were born in France (52.9), in Ireland (34.7), and in Germany (28.7),

and lowest among those whose mothers were born in Russia and Poland (9.8), in Scandinavia (12.4), and in Canada (13.9).