

## ESO 208A: Computational Methods in Engineering

### Problem Set 1

1. Let

$$F(x) = x + \frac{a^3}{x^2}, \text{ where } a = \text{constant parameter} \quad (1)$$

and let  $x_1$  and  $x_2$  be two positive values of  $x$  that satisfy the following relation:

$$a^3 \left( \frac{1}{x_1} - \frac{1}{x_2} \right) = \frac{1}{4} (x_2^2 - x_1^2), \quad x_2 > x_1 \quad (2)$$

It is desired to compute the difference  $\Delta F = F(x_1) - F(x_2)$ . with low precision arithmetic, by simulating a computer that performs floating point operations rounding *mantissa* to 6 decimals.

For the case in which  $a = 6.870429497$  and  $x_1 = 8.583454139$ , compute  $\Delta F$  directly from the definition of  $F$  in (1) and using (2) to compute  $x_2$ . True value of  $\Delta F$  computed using double precision algorithm is  $0.103622333 \times 10^{-4}$ . What is the true relative error in  $\Delta F$  using this algorithm?

Using (1) and (2), express  $\Delta F$  as:

$$\Delta F = \frac{x_1(r-1)^3}{4r}, \text{ where } r = x_2/x_1 \quad (3)$$

Estimate  $r$  using (2) and  $\Delta F$  using (3) with the same low precision arithmetic. What is the true relative error in  $\Delta F$  using this algorithm? Comment on your result.

2. Consider computation of the following expression:

$$G(\varepsilon) = \frac{\sqrt{1+8\varepsilon^2} - 1}{2}$$

Compute the value of  $G$  for  $\varepsilon = 0.001$  and estimate the corresponding relative error, performing operations by rounding all mantissas to six decimals. Improve your computation of  $G$  by employing a Taylor series expansion, and estimate the corresponding relative error. Use double precision calculation to obtain the true solution that is needed to calculate the relative errors.

3. One has measured the two sides and the included angle of a triangle as:  $a = 100.0 \pm 0.1$ ,  $b = 101.0 \pm 0.1$ , and the angle  $C = 1.00^\circ \pm 0.01^\circ$ . How accurately is it possible to give the third side  $c$ ?

4. The following set of equations is to be solved to get the value of  $x$  for a given  $\delta$ . For what values of  $\delta$  will this problem be well-conditioned?

$$\begin{aligned} x + y &= 2 \\ x + (1 - \delta)y &= 1 \end{aligned}$$