

MTL122 - Real and complex analysis

Assignment-3



Department of Mathematics
Indian Institute of Technology Delhi

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Proof

- Here $A \cap B = \emptyset$ and A, B are closed sets of (X, d) .

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Proof

- Here $A \cap B = \emptyset$ and A, B are closed sets of (X, d) .
- Then $X - A$ and $X - B$ are open sets in (X, d) and $B \subset X - A$, $A \subset X - B$.

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Proof

- Here $A \cap B = \emptyset$ and A, B are closed sets of (X, d) .
- Then $X - A$ and $X - B$ are open sets in (X, d) and $B \subset X - A$, $A \subset X - B$.
- If $A = \emptyset$, take $U = \emptyset$ and $V = X$.

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Proof

- Here $A \cap B = \emptyset$ and A, B are closed sets of (X, d) .
- Then $X - A$ and $X - B$ are open sets in (X, d) and $B \subset X - A$, $A \subset X - B$.
- If $A = \emptyset$, take $U = \emptyset$ and $V = X$.
- If $B = \emptyset$, take $V = \emptyset$ and $U = X$.

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Proof

- Here $A \cap B = \emptyset$ and A, B are closed sets of (X, d) .
- Then $X - A$ and $X - B$ are open sets in (X, d) and $B \subset X - A$, $A \subset X - B$.
- If $A = \emptyset$, take $U = \emptyset$ and $V = X$.
- If $B = \emptyset$, take $V = \emptyset$ and $U = X$.
- Now assume A and B both are non-empty disjoint closed subsets of X , consider $U = \{x \in X : d(x, A) < d(x, B)\}$ and $V = \{x \in X : d(x, B) < d(x, A)\}$.

Question 1

Question 1

Let A and B be disjoint closed subsets of a metric spaces (X, d) . Prove that there are disjoint open subsets U and V of X such that $A \subseteq U$ and $B \subseteq V$.

Proof

- Here $A \cap B = \emptyset$ and A, B are closed sets of (X, d) .
- Then $X - A$ and $X - B$ are open sets in (X, d) and $B \subset X - A$, $A \subset X - B$.
- If $A = \emptyset$, take $U = \emptyset$ and $V = X$.
- If $B = \emptyset$, take $V = \emptyset$ and $U = X$.
- Now assume A and B both are non-empty disjoint closed subsets of X , consider $U = \{x \in X : d(x, A) < d(x, B)\}$ and $V = \{x \in X : d(x, B) < d(x, A)\}$.

Claim: We need to show $A \subseteq U$, $B \subseteq V$ and $U \cap V = \emptyset$.

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$,

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$, then there exist $y \in U \cap V$. Thus

$$d(y, A) < d(y, B) \text{ and } d(y, B) < d(y, A),$$

which is impossible. Hence $U \cap V = \emptyset$.

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$, then there exist $y \in U \cap V$. Thus

$$d(y, A) < d(y, B) \text{ and } d(y, B) < d(y, A),$$

which is impossible. Hence $U \cap V = \emptyset$.

Claim: Now we have to show U and V are open in (X, d) .

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$, then there exist $y \in U \cap V$. Thus

$$d(y, A) < d(y, B) \text{ and } d(y, B) < d(y, A),$$

which is impossible. Hence $U \cap V = \emptyset$.

Claim: Now we have to show U and V are open in (X, d) .

We have

$$U = \{x \in X : d(x, A) < d(x, B)\}$$

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$, then there exist $y \in U \cap V$. Thus

$$d(y, A) < d(y, B) \text{ and } d(y, B) < d(y, A),$$

which is impossible. Hence $U \cap V = \emptyset$.

Claim: Now we have to show U and V are open in (X, d) .

We have

$$\begin{aligned} U &= \{x \in X : d(x, A) < d(x, B)\} \\ &= \bigcup_{r>0} \{x \in X : d(x, A) < r < d(x, B)\} \end{aligned}$$

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$, then there exist $y \in U \cap V$. Thus

$$d(y, A) < d(y, B) \text{ and } d(y, B) < d(y, A),$$

which is impossible. Hence $U \cap V = \emptyset$.

Claim: Now we have to show U and V are open in (X, d) .

We have

$$\begin{aligned} U &= \{x \in X : d(x, A) < d(x, B)\} \\ &= \bigcup_{r>0} \{x \in X : d(x, A) < r < d(x, B)\} \\ &= \bigcup_{r>0} \left(\{x \in X : d(x, A) < r\} \cap \{x \in X : d(x, B) > r\} \right). \end{aligned}$$

Question 1 Contd.

- Let $x \in A$. Then $d(x, A) = 0 < d(x, B)$, thus $x \in U$. Hence $A \subseteq U$.
- Let $x \in B$. Then $d(x, B) = 0 < d(x, A)$, thus $x \in V$. Hence $B \subseteq V$.
- Let us assume $U \cap V \neq \emptyset$, then there exist $y \in U \cap V$. Thus

$$d(y, A) < d(y, B) \text{ and } d(y, B) < d(y, A),$$

which is impossible. Hence $U \cap V = \emptyset$.

Claim: Now we have to show U and V are open in (X, d) .

We have

$$\begin{aligned} U &= \{x \in X : d(x, A) < d(x, B)\} \\ &= \bigcup_{r>0} \{x \in X : d(x, A) < r < d(x, B)\} \\ &= \bigcup_{r>0} \left(\{x \in X : d(x, A) < r\} \cap \{x \in X : d(x, B) > r\} \right). \end{aligned}$$

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.
- We know $d(p, A) \leq d(p, z) + d(z, A) < r - s + s = r$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.
- We know $d(p, A) \leq d(p, z) + d(z, A) < r - s + s = r$.
- Now we have $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.
- We know $d(p, A) \leq d(p, z) + d(z, A) < r - s + s = r$.
- Now we have $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Hence $\{x \in X : d(x, A) < r\}$ is open in X .

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.
- We know $d(p, A) \leq d(p, z) + d(z, A) < r - s + s = r$.
- Now we have $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Hence $\{x \in X : d(x, A) < r\}$ is open in X .
- Similarly, we can show $\{x \in X : d(x, B) > r\}$ is open in X .

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.
- We know $d(p, A) \leq d(p, z) + d(z, A) < r - s + s = r$.
- Now we have $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Hence $\{x \in X : d(x, A) < r\}$ is open in X .
- Similarly, we can show $\{x \in X : d(x, B) > r\}$ is open in X . Hence U is open in X .

Question 1 Contd.

To show U is open, we need show $\{x \in X : d(x, A) < r\}$ is open.

- Since A is closed, thus $\exists y \in A$ such that $d(x, A) = d(x, y)$. (Check!)
- Let $z \in \{x \in X : d(x, A) < r\}$. Then $d(z, A) < r$. Assume $d(z, A) = s < r$.
- Now the proof reduces to show, $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Let $p \in B(z, r - s)$, then $d(p, z) < r - s$.
- We know $d(p, A) \leq d(p, z) + d(z, A) < r - s + s = r$.
- Now we have $B(z, r - s) \subset \{x \in X : d(x, A) < r\}$.
- Hence $\{x \in X : d(x, A) < r\}$ is open in X .
- Similarly, we can show $\{x \in X : d(x, B) > r\}$ is open in X . Hence U is open in X .
- Similarly V can be open in X .

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c .

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$,

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.
- If $x \notin E^c$ and $x \in (E^\circ)^c$, we need to show $x \in (E^c)'$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.
- If $x \notin E^c$ and $x \in (E^\circ)^c$, we need to show $x \in (E^c)'$.
- To the contrary suppose that $x \notin (E^c)'$,

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.
- If $x \notin E^c$ and $x \in (E^\circ)^c$, we need to show $x \in (E^c)'$.
- To the contrary suppose that $x \notin (E^c)'$, then there exist $\epsilon > 0$ such that $B(x, \epsilon) \cap E^c = \emptyset$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.
- If $x \notin E^c$ and $x \in (E^\circ)^c$, we need to show $x \in (E^c)'$.
- To the contrary suppose that $x \notin (E^c)'$, then there exist $\epsilon > 0$ such that $B(x, \epsilon) \cap E^c = \emptyset$. So $B(x, \epsilon) \subset E$, hence $x \in E^\circ$. Which contradict the fact $x \in (E^\circ)^c$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.
- If $x \notin E^c$ and $x \in (E^\circ)^c$, we need to show $x \in (E^c)'$.
- To the contrary suppose that $x \notin (E^c)'$, then there exist $\epsilon > 0$ such that $B(x, \epsilon) \cap E^c = \emptyset$. So $B(x, \epsilon) \subset E$, hence $x \in E^\circ$. Which contradict the fact $x \in (E^\circ)^c$.
- So $x \in (E^c)' \subset \overline{(E^c)}$.

Question 2

Question 2

Let (X, d) be a metric space with $E \subset X$. Prove that $(E^\circ)^c = \overline{(E^c)}$.

Proof

- We know $E^\circ \subset E$, then $E^c \subset (E^\circ)^c$.
- Since $(E^\circ)^c$ is closed set and $\overline{(E^c)}$ is the smallest closed set containing E^c . Thus $\overline{(E^c)} \subset (E^\circ)^c$.
- Let $x \in (E^\circ)^c$. Since $E^c \subset (E^\circ)^c$, then $x \in E^c$ or $x \notin E^c$.
- If $x \in E^c$, then $x \in \overline{(E^c)}$. We are done.
- If $x \notin E^c$ and $x \in (E^\circ)^c$, we need to show $x \in (E^c)'$.
- To the contrary suppose that $x \notin (E^c)'$, then there exist $\epsilon > 0$ such that $B(x, \epsilon) \cap E^c = \emptyset$. So $B(x, \epsilon) \subset E$, hence $x \in E^\circ$. Which contradict the fact $x \in (E^\circ)^c$.
- So $x \in (E^c)' \subset \overline{(E^c)}$. Thus $(E^\circ)^c \subset \overline{(E^c)}$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.
- If $x \in A'$, then for each $n \in \mathbb{N}$, $B(x, \frac{1}{n}) \cap A \neq \emptyset$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.
- If $x \in A'$, then for each $n \in \mathbb{N}$, $B(x, \frac{1}{n}) \cap A \neq \emptyset$.
- Choose $y_n \in B(x, \frac{1}{n}) \cap A$, for each $n \in \mathbb{N}$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.
- If $x \in A'$, then for each $n \in \mathbb{N}$, $B(x, \frac{1}{n}) \cap A \neq \emptyset$.
- Choose $y_n \in B(x, \frac{1}{n}) \cap A$, for each $n \in \mathbb{N}$. Then $y_n \in A$ and $y_n \rightarrow x$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.
- If $x \in A'$, then for each $n \in \mathbb{N}$, $B(x, \frac{1}{n}) \cap A \neq \emptyset$.
- Choose $y_n \in B(x, \frac{1}{n}) \cap A$, for each $n \in \mathbb{N}$. Then $y_n \in A$ and $y_n \rightarrow x$.
- We know $d(x, A) = \inf\{d(x, y) : y \in A\}$, but $\inf\{d(x, y_n) : y_n \in A\} = 0$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.
- If $x \in A'$, then for each $n \in \mathbb{N}$, $B(x, \frac{1}{n}) \cap A \neq \emptyset$.
- Choose $y_n \in B(x, \frac{1}{n}) \cap A$, for each $n \in \mathbb{N}$. Then $y_n \in A$ and $y_n \rightarrow x$.
- We know $d(x, A) = \inf\{d(x, y) : y \in A\}$, but $\inf\{d(x, y_n) : y_n \in A\} = 0$. Hence $d(x, A) = 0$.

Question 3

Question 3

A point x not belonging to a closed set $M \subset (X, d)$ always has a nonzero distance from M . (Hint: To prove this, show that $x \in \overline{A}$ if and only if $D(x, A) = \text{dist}(x, A) = \inf_{y \in A} d(x, y) = 0$; here A is any nonempty subset of X).

Proof

- We need to show $x \in \overline{A} \Leftrightarrow d(x, A) = 0$.
- Let $x \in \overline{A}$. Then $x \in A \cup A'$. If $x \in A$, then $d(x, A) = 0$.
- If $x \in A'$, then for each $n \in \mathbb{N}$, $B(x, \frac{1}{n}) \cap A \neq \emptyset$.
- Choose $y_n \in B(x, \frac{1}{n}) \cap A$, for each $n \in \mathbb{N}$. Then $y_n \in A$ and $y_n \rightarrow x$.
- We know $d(x, A) = \inf\{d(x, y) : y \in A\}$, but $\inf\{d(x, y_n) : y_n \in A\} = 0$. Hence $d(x, A) = 0$.

Question 4

Question 4

Let A and B be non-empty subsets of a metric space $(X; d)$. Prove that:

- (i) $A \subset B \implies \text{diam}(A) \leq \text{diam}(B)$
- (ii) $\text{diam}(A) = 0$ if and only if for some $x \in X$, $A = \{x\}$.
- (iii) If $a \in A$ and $b \in B$, then
$$\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b).$$

Proof(i)

- Since $A \subseteq B$, this implies

Question 4

Question 4

Let A and B be non-empty subsets of a metric space $(X; d)$. Prove that:

- (i) $A \subset B \implies \text{diam}(A) \leq \text{diam}(B)$
- (ii) $\text{diam}(A) = 0$ if and only if for some $x \in X$, $A = \{x\}$.
- (iii) If $a \in A$ and $b \in B$, then
 $\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b)$.

Proof(i)

- Since $A \subseteq B$, this implies

$$\{d(x, y) : x, y \in A\} \subset \{d(x, y) : x, y \in B\}$$

Question 4

Question 4

Let A and B be non-empty subsets of a metric space $(X; d)$. Prove that:

- (i) $A \subset B \implies \text{diam}(A) \leq \text{diam}(B)$
- (ii) $\text{diam}(A) = 0$ if and only if for some $x \in X$, $A = \{x\}$.
- (iii) If $a \in A$ and $b \in B$, then
 $\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b)$.

Proof(i)

- Since $A \subseteq B$, this implies

$$\begin{aligned} \{d(x, y) : x, y \in A\} &\subset \{d(x, y) : x, y \in B\} \\ \sup\{d(x, y) : x, y \in A\} &\leq \sup\{d(x, y) : x, y \in B\} \end{aligned}$$

Question 4

Question 4

Let A and B be non-empty subsets of a metric space $(X; d)$. Prove that:

- (i) $A \subset B \implies \text{diam}(A) \leq \text{diam}(B)$
- (ii) $\text{diam}(A) = 0$ if and only if for some $x \in X$, $A = \{x\}$.
- (iii) If $a \in A$ and $b \in B$, then
 $\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b)$.

Proof(i)

- Since $A \subseteq B$, this implies

$$\begin{aligned}\{d(x, y) : x, y \in A\} &\subset \{d(x, y) : x, y \in B\} \\ \sup\{d(x, y) : x, y \in A\} &\leq \sup\{d(x, y) : x, y \in B\} \\ \text{diam}(A) &\leq \text{diam}(B)\end{aligned}$$

Question 4

Question 4

Let A and B be non-empty subsets of a metric space $(X; d)$. Prove that:

- (i) $A \subset B \implies \text{diam}(A) \leq \text{diam}(B)$
- (ii) $\text{diam}(A) = 0$ if and only if for some $x \in X$, $A = \{x\}$.
- (iii) If $a \in A$ and $b \in B$, then
 $\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b)$.

Proof(i)

- Since $A \subseteq B$, this implies

$$\begin{aligned}\{d(x, y) : x, y \in A\} &\subset \{d(x, y) : x, y \in B\} \\ \sup\{d(x, y) : x, y \in A\} &\leq \sup\{d(x, y) : x, y \in B\} \\ \text{diam}(A) &\leq \text{diam}(B)\end{aligned}$$

Proof(ii)

Question 4

Question 4

Let A and B be non-empty subsets of a metric space $(X; d)$. Prove that:

- (i) $A \subset B \implies \text{diam}(A) \leq \text{diam}(B)$
- (ii) $\text{diam}(A) = 0$ if and only if for some $x \in X$, $A = \{x\}$.
- (iii) If $a \in A$ and $b \in B$, then
 $\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b).$

Proof(i)

- Since $A \subseteq B$, this implies

$$\begin{aligned}\{d(x, y) : x, y \in A\} &\subset \{d(x, y) : x, y \in B\} \\ \sup\{d(x, y) : x, y \in A\} &\leq \sup\{d(x, y) : x, y \in B\} \\ \text{diam}(A) &\leq \text{diam}(B)\end{aligned}$$

Proof(ii) Its proof is obvious. Do yourself.

Question 4 Contd...

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.

Question 4 Contd...

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.
- Consider two points x, y in $A \cup B$.

Question 4 Contd...

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.
- Consider two points x, y in $A \cup B$.
- If they are both in A , then $d(x, y) \leq \text{diam}(A)$.

Question 4 Contd...

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.
- Consider two points x, y in $A \cup B$.
- If they are both in A , then $d(x, y) \leq \text{diam}(A)$.
- If they are both in B , then $d(x, y) \leq \text{diam}(B)$.

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.
- Consider two points x, y in $A \cup B$.
- If they are both in A , then $d(x, y) \leq \text{diam}(A)$.
- If they are both in B , then $d(x, y) \leq \text{diam}(B)$.
- Now suppose $x \in A$ and $y \in B$, then we have

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.
- Consider two points x, y in $A \cup B$.
- If they are both in A , then $d(x, y) \leq \text{diam}(A)$.
- If they are both in B , then $d(x, y) \leq \text{diam}(B)$.
- Now suppose $x \in A$ and $y \in B$, then we have

$$d(x, y) \leq d(x, a) + d(a, b) + d(b, y) \leq \text{diam}(A) + d(a, b) + \text{diam}(B)$$

Proof(iii)

- We know $\text{diam}(A \cup B) = \sup\{d(x, y) : x, y \in A \cup B\}$.
- Consider two points x, y in $A \cup B$.
- If they are both in A , then $d(x, y) \leq \text{diam}(A)$.
- If they are both in B , then $d(x, y) \leq \text{diam}(B)$.
- Now suppose $x \in A$ and $y \in B$, then we have

$$d(x, y) \leq d(x, a) + d(a, b) + d(b, y) \leq \text{diam}(A) + d(a, b) + \text{diam}(B)$$

- By taking supremum bothside over $x, y \in A \cup B$, we have

$$\text{diam}(A \cup B) \leq \text{diam}(A) + \text{diam}(B) + d(a, b).$$

Question 5

Question 5

Let (X, d) be a metric space with the property that every bounded sequence has a convergent subsequence. Prove that X is complete. Does Bolzano-Weierstrass theorem holds holds for any metric space? Give reasons/counterexamples.

Question 5

Question 5

Let (X, d) be a metric space with the property that every bounded sequence has a convergent subsequence. Prove that X is complete. Does Bolzano-Weierstrass theorem holds for any metric space? Give reasons/counterexamples.

Solution: To prove if Bolzano-Weierstrass theorem holds in a metric space then the metric space is complete .

Question 5

Question 5

Let (X, d) be a metric space with the property that every bounded sequence has a convergent subsequence. Prove that X is complete. Does Bolzano-Weierstrass theorem holds for any metric space? Give reasons/counterexamples.

Solution: To prove if Bolzano-Weierstrass theorem holds in a metric space then the metric space is complete .

- Let $\{x_n\}_{n>0}$ be a cauchy sequence in X .

Question 5

Question 5

Let (X, d) be a metric space with the property that every bounded sequence has a convergent subsequence. Prove that X is complete. Does Bolzano-Weierstrass theorem holds for any metric space? Give reasons/counterexamples.

Solution: To prove if Bolzano-Weierstrass theorem holds in a metric space then the metric space is complete .

- Let $\{x_n\}_{n>0}$ be a cauchy sequence in X .
- For $\epsilon > 0$, there exists $N_\epsilon > 0$ such that $d(x_n, x_m) < \epsilon$ for all $n, m > N_\epsilon$.
- Since cauchy sequences are bounded. Hence $\{x_n\}_{n>0}$ is a bounded sequence.

Question 5

Question 5

Let (X, d) be a metric space with the property that every bounded sequence has a convergent subsequence. Prove that X is complete. Does Bolzano-Weierstrass theorem holds for any metric space? Give reasons/counterexamples.

Solution: To prove if Bolzano-Weierstrass theorem holds in a metric space then the metric space is complete .

- Let $\{x_n\}_{n>0}$ be a cauchy sequence in X .
- For $\epsilon > 0$, there exists $N_\epsilon > 0$ such that $d(x_n, x_m) < \epsilon$ for all $n, m > N_\epsilon$.
- Since cauchy sequences are bounded. Hence $\{x_n\}_{n>0}$ is a bounded sequence.
- Let $\{x_{n_k}\}_{k>0} \rightarrow a$, be a convergent subsequence of x_n where $a \in X$.

Question 5

Question 5

Let (X, d) be a metric space with the property that every bounded sequence has a convergent subsequence. Prove that X is complete. Does Bolzano-Weierstrass theorem holds for any metric space? Give reasons/counterexamples.

Solution: To prove if Bolzano-Weierstrass theorem holds in a metric space then the metric space is complete .

- Let $\{x_n\}_{n>0}$ be a cauchy sequence in X .
- For $\epsilon > 0$, there exists $N_\epsilon > 0$ such that $d(x_n, x_m) < \epsilon$ for all $n, m > N_\epsilon$.
- Since cauchy sequences are bounded. Hence $\{x_n\}_{n>0}$ is a bounded sequence.
- Let $\{x_{n_k}\}_{k>0} \rightarrow a$, be a convergent subsequence of x_n where $a \in X$.

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.
- Choose $N = \max\{N_\epsilon, M_\epsilon\}$. (Why?)

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.
- Choose $N = \max\{N_\epsilon, M_\epsilon\}$. (Why?)
- Now, $d(x_n, a) \leq d(x_n, x_{n_k}) + d(x_{n_k}, a) < \epsilon$ for all $n > N$.

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.
- Choose $N = \max\{N_\epsilon, M_\epsilon\}$. (Why?)
- Now, $d(x_n, a) \leq d(x_n, x_{n_k}) + d(x_{n_k}, a) < \epsilon$ for all $n > N$.
- Hence every cauchy sequence converges and the metric space is complete.

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.
- Choose $N = \max\{N_\epsilon, M_\epsilon\}$. (Why?)
- Now, $d(x_n, a) \leq d(x_n, x_{n_k}) + d(x_{n_k}, a) < \epsilon$ for all $n > N$.
- Hence every cauchy sequence converges and the metric space is complete.
- Let $M = \mathbb{Q}$ and $d(x, y) = |x - y|$

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.
- Choose $N = \max\{N_\epsilon, M_\epsilon\}$. (Why?)
- Now, $d(x_n, a) \leq d(x_n, x_{n_k}) + d(x_{n_k}, a) < \epsilon$ for all $n > N$.
- Hence every Cauchy sequence converges and the metric space is complete.
- Let $M = \mathbb{Q}$ and $d(x, y) = |x - y|$
- Consider the sequence $x_{n+1} = 2^{\frac{1+x_n}{2+x_n}}$ with $x_1 = 1$

Question 5 Contd...

Claim: $x_n \rightarrow a$

- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $d(x_{n_k}, a) < \epsilon/2$ for all $n_k > M_\epsilon$.
- Choose $N = \max\{N_\epsilon, M_\epsilon\}$. (Why?)
- Now, $d(x_n, a) \leq d(x_n, x_{n_k}) + d(x_{n_k}, a) < \epsilon$ for all $n > N$.
- Hence every cauchy sequence converges and the metric space is complete.
- Let $M = \mathbb{Q}$ and $d(x, y) = |x - y|$
- Consider the sequence $x_{n+1} = 2^{\frac{1+x_n}{2+x_n}}$ with $x_1 = 1$
- This counterexample is enough. (Why?)

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,
 $|a_m - a_n| = |d(x_m, y_m) - d(x_n, y_n)|$

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,
$$|a_m - a_n| = |d(x_m, y_m) - d(x_n, y_n)|$$
$$= |\{d(x_m, y_m) - d(y_m, x_n)\} - \{d(x_n, y_n) - d(y_n, x_n)\}|$$

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,

$$\begin{aligned} |a_m - a_n| &= |d(x_m, y_m) - d(x_n, y_n)| \\ &= |\{d(x_m, y_m) - d(y_m, x_n)\} - \{d(x_n, y_n) - d(y_m, x_n)\}| \\ &\leq |d(x_m, y_m) - d(y_m, x_n)| + |d(x_n, y_n) - d(y_m, x_n)| \end{aligned}$$

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,

$$\begin{aligned} |a_m - a_n| &= |d(x_m, y_m) - d(x_n, y_n)| \\ &= |\{d(x_m, y_m) - d(y_m, x_n)\} - \{d(x_n, y_n) - d(y_m, x_n)\}| \\ &\leq |d(x_m, y_m) - d(y_m, x_n)| + |d(x_n, y_n) - d(y_m, x_n)| \\ &\leq d(x_m, x_n) + d(y_m, y_n) \end{aligned}$$

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,

$$\begin{aligned} |a_m - a_n| &= |d(x_m, y_m) - d(x_n, y_n)| \\ &= |\{d(x_m, y_m) - d(y_m, x_n)\} - \{d(x_n, y_n) - d(y_m, x_n)\}| \\ &\leq |d(x_m, y_m) - d(y_m, x_n)| + |d(x_n, y_n) - d(y_m, x_n)| \\ &\leq d(x_m, x_n) + d(y_m, y_n) < \epsilon \end{aligned}$$

Question 6

Question 6

If (x_n) and (y_n) are Cauchy sequences in a metric space (X, d) , show that (a_n) , where $a_n = d(x_n, y_n)$, converges.

Solution:

- First we try to prove that (a_n) is Cauchy in \mathbb{R} and that essentially proves what we want. (Why?)
- $\forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $d(x_m, x_n) < \epsilon/2$
- $\forall \epsilon > 0, \exists M_\epsilon \in \mathbb{N}$ such that $\forall m, n > M_\epsilon$ we have $d(y_m, y_n) < \epsilon/2$.
- Now, $\forall m, n > \max\{N_\epsilon, M_\epsilon\}$, we have,
$$\begin{aligned} |a_m - a_n| &= |d(x_m, y_m) - d(x_n, y_n)| \\ &= |\{d(x_m, y_m) - d(y_m, x_n)\} - \{d(x_n, y_n) - d(y_m, x_n)\}| \\ &\leq |d(x_m, y_m) - d(y_m, x_n)| + |d(x_n, y_n) - d(y_m, x_n)| \\ &\leq d(x_m, x_n) + d(y_m, y_n) < \epsilon \end{aligned}$$
- Hence proved

Question 7

Question 7

Let $X = (X, d)$ be a metric space and $CS(X)$ the collection of all Cauchy sequences in X . For (x_n) and (y_n) in $CS(X)$, define

$(x_n) \sim (y_n)$ if and only if $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$

Prove that \sim is an equivalence relation

Question 7

Question 7

Let $X = (X, d)$ be a metric space and $CS(X)$ the collection of all Cauchy sequences in X . For (x_n) and (y_n) in $CS(X)$, define $(x_n) \sim (y_n)$ if and only if $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$. Prove that \sim is an equivalence relation.

Solution: As usual we try to prove the three properties .

Question 7

Question 7

Let $X = (X, d)$ be a metric space and $CS(X)$ the collection of all Cauchy sequences in X . For (x_n) and (y_n) in $CS(X)$, define

$(x_n) \sim (y_n)$ if and only if $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$

Prove that \sim is an equivalence relation

Solution: As usual we try to prove the three properties .

- Reflexive: $d(x_n, x_n) = 0 \forall n > 0$. Hence $(x_n) \sim (y_n)$

Question 7

Question 7

Let $X = (X, d)$ be a metric space and $CS(X)$ the collection of all Cauchy sequences in X . For (x_n) and (y_n) in $CS(X)$, define $(x_n) \sim (y_n)$ if and only if $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$. Prove that \sim is an equivalence relation.

Solution: As usual we try to prove the three properties .

- Reflexive: $d(x_n, x_n) = 0 \ \forall n > 0$. Hence $(x_n) \sim (y_n)$
- Symmetric: Since d is symmetric we have $\lim_{n \rightarrow \infty} d(x_n, y_n) = \lim_{n \rightarrow \infty} d(y_n, x_n) = 0$.

Question 7

Question 7

Let $X = (X, d)$ be a metric space and $CS(X)$ the collection of all Cauchy sequences in X . For (x_n) and (y_n) in $CS(X)$, define

$(x_n) \sim (y_n)$ if and only if $\lim_{n \rightarrow \infty} d(x_n, y_n) = 0$

Prove that \sim is an equivalence relation

Solution: As usual we try to prove the three properties .

- Reflexive: $d(x_n, x_n) = 0 \forall n > 0$. Hence $(x_n) \sim (y_n)$
- Symmetric: Since d is symmetric we have $\lim_{n \rightarrow \infty} d(x_n, y_n) = \lim_{n \rightarrow \infty} d(y_n, x_n) = 0$.
- Transitive: Let $(x_n) \sim (y_n)$ and $(y_n) \sim (z_n)$. Now $0 \leq d(x_n, z_n) \leq d(x_n, y_n) + d(y_n, z_n)$ which using sandwich theorem gives that $(x_n) \sim (z_n)$

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .
- $\therefore \forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $|x_m - x_n| < \epsilon$.

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .
- $\therefore \forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $|x_m - x_n| < \epsilon$.
- For $\epsilon = 1$, we have N_1 such that $\forall n, m > N_1, |x_m - x_n| < 1$.

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .
- $\therefore \forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $|x_m - x_n| < \epsilon$.
- For $\epsilon = 1$, we have N_1 such that $\forall n, m > N_1, |x_m - x_n| < 1$.
- Since our space is set of integers and $|x_m - x_n| < 1$, this implies $|x_m - x_n| = 0$ for all $n, m > N_1$.

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .
- $\therefore \forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $|x_m - x_n| < \epsilon$.
- For $\epsilon = 1$, we have N_1 such that $\forall n, m > N_1, |x_m - x_n| < 1$.
- Since our space is set of integers and $|x_m - x_n| < 1$, this implies $|x_m - x_n| = 0$ for all $n, m > N_1$.
- That means (x_n) is eventually constant sequence, which is convergent.

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .
- $\therefore \forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $|x_m - x_n| < \epsilon$.
- For $\epsilon = 1$, we have N_1 such that $\forall n, m > N_1, |x_m - x_n| < 1$.
- Since our space is set of integers and $|x_m - x_n| < 1$, this implies $|x_m - x_n| = 0$ for all $n, m > N_1$.
- That means (x_n) is eventually constant sequence, which is convergent.
- Hence (X, d) is a complete metric space.

Question 8

Question 8

Show that the set X of all integers, with metric d defined by $d(m, n) = |m - n|$, is a complete metric space.

Solution:

- Let $(x_n)_{n \geq 0}$ be a Cauchy Sequence in X .
- $\therefore \forall \epsilon > 0, \exists N_\epsilon \in \mathbb{N}$ such that $\forall m, n > N_\epsilon$ we have $|x_m - x_n| < \epsilon$.
- For $\epsilon = 1$, we have N_1 such that $\forall n, m > N_1, |x_m - x_n| < 1$.
- Since our space is set of integers and $|x_m - x_n| < 1$, this implies $|x_m - x_n| = 0$ for all $n, m > N_1$.
- That means (x_n) is eventually constant sequence, which is convergent.
- Hence (X, d) is a complete metric space.

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Solution:

- Let $\{x_n\}$ be the cauchy sequence in ℓ^∞ , where $x_n = (x_1^{(n)}, x_2^{(n)}, \dots)$.

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Solution:

- Let $\{x_n\}$ be the cauchy sequence in ℓ^∞ , where $x_n = (x_1^{(n)}, x_2^{(n)}, \dots)$.
- Since $\{x_n\}$ is cauchy, this implies for $\epsilon > 0$, there exists an $N_0 \in \mathbb{N}$ such that

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Solution:

- Let $\{x_n\}$ be the cauchy sequence in ℓ^∞ , where $x_n = (x_1^{(n)}, x_2^{(n)}, \dots)$.
- Since $\{x_n\}$ is cauchy, this implies for $\epsilon > 0$, there exists an $N_0 \in \mathbb{N}$ such that

$$d_\infty(x_n, x_m) = \sup_{i \in \mathbb{N}} |x_i^{(n)} - x_i^{(m)}| < \epsilon \quad \text{for all } n, m > N_0.$$

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Solution:

- Let $\{x_n\}$ be the cauchy sequence in ℓ^∞ , where $x_n = (x_1^{(n)}, x_2^{(n)}, \dots)$.
- Since $\{x_n\}$ is cauchy, this implies for $\epsilon > 0$, there exists an $N_0 \in \mathbb{N}$ such that

$$d_\infty(x_n, x_m) = \sup_{i \in \mathbb{N}} |x_i^{(n)} - x_i^{(m)}| < \epsilon \text{ for all } n, m > N_0.$$

- Thus

$$|x_i^{(n)} - x_i^{(m)}| < \epsilon \text{ for all } n, m > N_0 \text{ and for all } i \in \mathbb{N} \quad (1)$$

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Solution:

- Let $\{x_n\}$ be the cauchy sequence in ℓ^∞ , where $x_n = (x_1^{(n)}, x_2^{(n)}, \dots)$.
- Since $\{x_n\}$ is cauchy, this implies for $\epsilon > 0$, there exists an $N_0 \in \mathbb{N}$ such that

$$d_\infty(x_n, x_m) = \sup_{i \in \mathbb{N}} |x_i^{(n)} - x_i^{(m)}| < \epsilon \text{ for all } n, m > N_0.$$

- Thus

$$|x_i^{(n)} - x_i^{(m)}| < \epsilon \text{ for all } n, m > N_0 \text{ and for all } i \in \mathbb{N} \quad (1)$$

- This implies for all $i \in \mathbb{N}$, $\{x_i^{(n)}\}$ is a cauchy sequence in \mathbb{R} .

Question 9

Question 9

Show that (ℓ^∞, d_∞) is a complete metric space.

Solution:

- Let $\{x_n\}$ be the cauchy sequence in ℓ^∞ , where $x_n = (x_1^{(n)}, x_2^{(n)}, \dots)$.
- Since $\{x_n\}$ is cauchy, this implies for $\epsilon > 0$, there exists an $N_0 \in \mathbb{N}$ such that

$$d_\infty(x_n, x_m) = \sup_{i \in \mathbb{N}} |x_i^{(n)} - x_i^{(m)}| < \epsilon \quad \text{for all } n, m > N_0.$$

- Thus

$$|x_i^{(n)} - x_i^{(m)}| < \epsilon \quad \text{for all } n, m > N_0 \quad \text{and for all } i \in \mathbb{N} \quad (1)$$

- This implies for all $i \in \mathbb{N}$, $\{x_i^{(n)}\}$ is a cauchy sequence in \mathbb{R} .

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.
- Hence for some $m > N_0$,

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.
- Hence for some $m > N_0$,

$$|l_i| = \left| x_i^{(m)} + (l_i - x_i^{(m)}) \right| \leq \left| x_i^{(m)} \right| + \epsilon \quad \text{for all } i \in \mathbb{N}$$

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.
- Hence for some $m > N_0$,

$$|l_i| = \left| x_i^{(m)} + (l_i - x_i^{(m)}) \right| \leq \left| x_i^{(m)} \right| + \epsilon \quad \text{for all } i \in \mathbb{N}$$

Since the sequence $x^{(m)}$ is bounded, so is the sequence L .

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.
- Hence for some $m > N_0$,

$$|l_i| = \left| x_i^{(m)} + (l_i - x_i^{(m)}) \right| \leq \left| x_i^{(m)} \right| + \epsilon \quad \text{for all } i \in \mathbb{N}$$

Since the sequence $x^{(m)}$ is bounded, so is the sequence L .

- Therefore, $L \in \ell_\infty$.

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.
- Hence for some $m > N_0$,

$$|l_i| = \left| x_i^{(m)} + (l_i - x_i^{(m)}) \right| \leq \left| x_i^{(m)} \right| + \epsilon \quad \text{for all } i \in \mathbb{N}$$

Since the sequence $x^{(m)}$ is bounded, so is the sequence L .

- Therefore, $L \in \ell_\infty$.
- Hence (ℓ_∞, d_∞) is complete.

Question 9 contd...

- Since \mathbb{R} with euclidean metric is complete. Thus for all $i \in \mathbb{N}$ the sequence $\{x_i^{(n)}\}$ converges in \mathbb{R} .
- Suppose that $x_i^{(n)} \rightarrow l_i$ for all $i \in \mathbb{N}$.
- Now we will prove that $x_n \rightarrow L$ where $L = (l_1, l_2, \dots)$ and $L \in \ell^\infty$.
- By taking $m \rightarrow \infty$ in (1), we obtain

$$\left| x_i^{(n)} - l_i \right| < \epsilon \quad \text{for all } n > N_0 \quad \text{and } i \in \mathbb{N}$$

- Thus $x_n \rightarrow L$ as $n \rightarrow \infty$.
- Hence for some $m > N_0$,

$$|l_i| = \left| x_i^{(m)} + (l_i - x_i^{(m)}) \right| \leq \left| x_i^{(m)} \right| + \epsilon \quad \text{for all } i \in \mathbb{N}$$

Since the sequence $x^{(m)}$ is bounded, so is the sequence L .

- Therefore, $L \in \ell_\infty$.
- Hence (ℓ_∞, d_∞) is complete.

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.
- We will prove that $\{x_n\}$ is cauchy.

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.
- We will prove that $\{x_n\}$ is cauchy.
- Since $\lim_{n \rightarrow \infty} \arctan(n) = \frac{\pi}{2}$.

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.
- We will prove that $\{x_n\}$ is cauchy.
- Since $\lim_{n \rightarrow \infty} \arctan(n) = \frac{\pi}{2}$. Thus for $\epsilon > 0$, there exists a natural number $n_0 \in \mathbb{N}$ such that $|\arctan(n) - \frac{\pi}{2}| < \frac{\epsilon}{2}$ for all $n > n_0$.

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.
- We will prove that $\{x_n\}$ is cauchy.
- Since $\lim_{n \rightarrow \infty} \arctan(n) = \frac{\pi}{2}$. Thus for $\epsilon > 0$, there exists a natural number $n_0 \in \mathbb{N}$ such that $|\arctan(n) - \frac{\pi}{2}| < \frac{\epsilon}{2}$ for all $n > n_0$.
- Now consider $m, n > n_0$

$$d(x_m, x_n) = |\arctan m - \arctan n|$$

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.
- We will prove that $\{x_n\}$ is cauchy.
- Since $\lim_{n \rightarrow \infty} \arctan(n) = \frac{\pi}{2}$. Thus for $\epsilon > 0$, there exists a natural number $n_0 \in \mathbb{N}$ such that $|\arctan(n) - \frac{\pi}{2}| < \frac{\epsilon}{2}$ for all $n > n_0$.
- Now consider $m, n > n_0$

$$\begin{aligned} d(x_m, x_n) &= |\arctan m - \arctan n| \\ &\leq \left| \arctan(m) - \frac{\pi}{2} \right| + \left| \arctan(n) - \frac{\pi}{2} \right| \end{aligned}$$

Question 10

Question 10

Show that the set of all real numbers constitutes an incomplete metric space if we choose $d(x, y) = |\arctan x - \arctan y|$.

Solution:

- Take $\{x_n\} = n$.
- We will prove that $\{x_n\}$ is cauchy.
- Since $\lim_{n \rightarrow \infty} \arctan(n) = \frac{\pi}{2}$. Thus for $\epsilon > 0$, there exists a natural number $n_0 \in \mathbb{N}$ such that $|\arctan(n) - \frac{\pi}{2}| < \frac{\epsilon}{2}$ for all $n > n_0$.
- Now consider $m, n > n_0$

$$\begin{aligned} d(x_m, x_n) &= |\arctan m - \arctan n| \\ &\leq \left| \arctan(m) - \frac{\pi}{2} \right| + \left| \arctan(n) - \frac{\pi}{2} \right| \\ &< \epsilon \end{aligned}$$

Question 10 Contd...

- Hence it is cauchy.

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\left| \frac{\pi}{2} - \arctan(x) \right|$$

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\left| \frac{\pi}{2} - \arctan(x) \right| \leq \left| \frac{\pi}{2} - \arctan(n) \right| + |\arctan(n) - \arctan(x)|$$

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\begin{aligned} \left| \frac{\pi}{2} - \arctan(x) \right| &\leq \left| \frac{\pi}{2} - \arctan(n) \right| + |\arctan(n) - \arctan(x)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\begin{aligned} \left| \frac{\pi}{2} - \arctan(x) \right| &\leq \left| \frac{\pi}{2} - \arctan(n) \right| + |\arctan(n) - \arctan(x)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

- Thus $\left| \frac{\pi}{2} - \arctan(x) \right| < \epsilon$ and $\epsilon > 0$ is arbitrary.

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\begin{aligned} \left| \frac{\pi}{2} - \arctan(x) \right| &\leq \left| \frac{\pi}{2} - \arctan(n) \right| + |\arctan(n) - \arctan(x)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

- Thus $\left| \frac{\pi}{2} - \arctan(x) \right| < \epsilon$ and $\epsilon > 0$ is arbitrary.
- This gives $\arctan(x) = \frac{\pi}{2}$,

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\begin{aligned} \left| \frac{\pi}{2} - \arctan(x) \right| &\leq \left| \frac{\pi}{2} - \arctan(n) \right| + |\arctan(n) - \arctan(x)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

- Thus $\left| \frac{\pi}{2} - \arctan(x) \right| < \epsilon$ and $\epsilon > 0$ is arbitrary.
- This gives $\arctan(x) = \frac{\pi}{2}$, but there doesn't exist any real number such that $\arctan(x) = \frac{\pi}{2}$.

Question 10 Contd...

- Hence it is cauchy.
- Suppose that $\{x_n\} \rightarrow x$ i.e $d(x_n, x) \rightarrow 0$ as $n \rightarrow \infty$.
- Let $\epsilon > 0$ and $N > n_0$,

$$d(x_n, x) < \frac{\epsilon}{2} \quad \text{for all } n > n_0,$$

- Consider

$$\begin{aligned} \left| \frac{\pi}{2} - \arctan(x) \right| &\leq \left| \frac{\pi}{2} - \arctan(n) \right| + |\arctan(n) - \arctan(x)| \\ &< \frac{\epsilon}{2} + \frac{\epsilon}{2} = \epsilon \end{aligned}$$

- Thus $\left| \frac{\pi}{2} - \arctan(x) \right| < \epsilon$ and $\epsilon > 0$ is arbitrary.
- This gives $\arctan(x) = \frac{\pi}{2}$, but there doesn't exist any real number such that $\arctan(x) = \frac{\pi}{2}$. contradiction.
- Hence it is not complete.