

MTL122 - Real and complex analysis

Assignment-2



Department of Mathematics
Indian Institute of Technology Delhi

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq (A \cap B)$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq (A \cap B)$.
- This implies $N_\epsilon(x) \subseteq A$ and $N_\epsilon(x) \subseteq B$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq (A \cap B)$.
- This implies $N_\epsilon(x) \subseteq A$ and $N_\epsilon(x) \subseteq B$.
- Thus we have $x \in \text{Int}A$ and $x \in \text{Int}B$

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq (A \cap B)$.
- This implies $N_\epsilon(x) \subseteq A$ and $N_\epsilon(x) \subseteq B$.
- Thus we have $x \in \text{Int}A$ and $x \in \text{Int}B$
- $\therefore x \in \text{Int}A \cap \text{Int}B$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq (A \cap B)$.
- This implies $N_\epsilon(x) \subseteq A$ and $N_\epsilon(x) \subseteq B$.
- Thus we have $x \in \text{Int}A$ and $x \in \text{Int}B$
- $\therefore x \in \text{Int}A \cap \text{Int}B$.
- Thus we finally have $\text{Int}(A \cap B) \subseteq \text{Int}A \cap \text{Int}B$.

Question 1

Question 1

Let $A \subset \mathbb{R}$ and $B \subset \mathbb{R}$ be sets.

- (i) Prove that $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.
- (ii) Prove that $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.
- (iii) Give an example of two sets A and B such that $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Solution: (i) To prove $\text{Int}(A \cap B) = \text{Int}A \cap \text{Int}B$.

- Let $x \in \text{Int}(A \cap B)$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq (A \cap B)$.
- This implies $N_\epsilon(x) \subseteq A$ and $N_\epsilon(x) \subseteq B$.
- Thus we have $x \in \text{Int}A$ and $x \in \text{Int}B$
- $\therefore x \in \text{Int}A \cap \text{Int}B$.
- Thus we finally have $\text{Int}(A \cap B) \subseteq \text{Int}A \cap \text{Int}B$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

- Let $x \in \text{Int}A \cup \text{Int}B$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

- Let $x \in \text{Int}A \cup \text{Int}B$.
- WLOG lets assume $x \in \text{Int}A$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

- Let $x \in \text{Int}A \cup \text{Int}B$.
- WLOG lets assume $x \in \text{Int}A$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq A$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

- Let $x \in \text{Int}A \cup \text{Int}B$.
- WLOG lets assume $x \in \text{Int}A$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq A$.
- Thus we have $N_\epsilon(x) \subseteq A \cup B \implies x \in \text{Int}(A \cup B)$.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

- Let $x \in \text{Int}A \cup \text{Int}B$.
- WLOG lets assume $x \in \text{Int}A$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq A$.
- Thus we have $N_\epsilon(x) \subseteq A \cup B \implies x \in \text{Int}(A \cup B)$.
- Hence Proved.

Question 1 Contd...

Now we prove the other side inclusion

- Let $x \in \text{Int}A \cap \text{Int}B$.
- $\therefore x \in \text{Int}A$ and $x \in \text{Int}B$.
- This implies $\exists \epsilon_1 > 0$ and $\epsilon_2 > 0$ such that $N_{\epsilon_1}(x) \subseteq A$ and $N_{\epsilon_2}(x) \subseteq B$.
- For $\epsilon = \min\{\epsilon_1, \epsilon_2\}$, we have $N_\epsilon(x) \subseteq (A \cap B)$.
- Thus proved. (Why?)

(ii) To prove $\text{Int}A \cup \text{Int}B \subseteq \text{Int}(A \cup B)$.

- Let $x \in \text{Int}A \cup \text{Int}B$.
- WLOG lets assume $x \in \text{Int}A$.
- $\therefore \exists \epsilon > 0$ such that $N_\epsilon(x) \subseteq A$.
- Thus we have $N_\epsilon(x) \subseteq A \cup B \implies x \in \text{Int}(A \cup B)$.
- Hence Proved.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.
- $\text{Int}(A \cup B) = (0, 2)$ and $\text{Int}A \cup \text{Int}B = (0, 1) \cup (1, 2)$.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.
- $\text{Int}(A \cup B) = (0, 2)$ and $\text{Int}A \cup \text{Int}B = (0, 1) \cup (1, 2)$.
- Hence $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.
- $\text{Int}(A \cup B) = (0, 2)$ and $\text{Int}A \cup \text{Int}B = (0, 1) \cup (1, 2)$.
- Hence $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Question 2

Prove that

- (i) If $A \subseteq \mathbb{R}$ is bounded above then $\sup A \in \text{Bd}(A)$.
- (ii) If $a < b < c$ and the two sets A and B has the property that $A \cap (a, c) = B \cap (a, c)$. Show that $b \in \text{Bd}(A)$ if and only if $b \in \text{Bd}(B)$.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.
- $\text{Int}(A \cup B) = (0, 2)$ and $\text{Int}A \cup \text{Int}B = (0, 1) \cup (1, 2)$.
- Hence $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Question 2

Prove that

- (i) If $A \subseteq \mathbb{R}$ is bounded above then $\sup A \in \text{Bd}(A)$.
- (ii) If $a < b < c$ and the two sets A and B has the property that $A \cap (a, c) = B \cap (a, c)$. Show that $b \in \text{Bd}(A)$ if and only if $b \in \text{Bd}(B)$.

Solution: (i)

- Let $M = \sup A$.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.
- $\text{Int}(A \cup B) = (0, 2)$ and $\text{Int}A \cup \text{Int}B = (0, 1) \cup (1, 2)$.
- Hence $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Question 2

Prove that

- (i) If $A \subseteq \mathbb{R}$ is bounded above then $\sup A \in \text{Bd}(A)$.
- (ii) If $a < b < c$ and the two sets A and B has the property that $A \cap (a, c) = B \cap (a, c)$. Show that $b \in \text{Bd}(A)$ if and only if $b \in \text{Bd}(B)$.

Solution: (i)

- Let $M = \sup A$.
- Thus $\forall \epsilon > 0, \exists a \in A \cap N_\epsilon(M)$ by the definition of supremum.

Question 2

(iii)

- Take $A = [0, 1]$ and $B = (1, 2)$.
- $\text{Int}(A \cup B) = (0, 2)$ and $\text{Int}A \cup \text{Int}B = (0, 1) \cup (1, 2)$.
- Hence $\text{Int}A \cup \text{Int}B \neq \text{Int}(A \cup B)$.

Question 2

Prove that

- (i) If $A \subseteq \mathbb{R}$ is bounded above then $\sup A \in \text{Bd}(A)$.
- (ii) If $a < b < c$ and the two sets A and B has the property that $A \cap (a, c) = B \cap (a, c)$. Show that $b \in \text{Bd}(A)$ if and only if $b \in \text{Bd}(B)$.

Solution: (i)

- Let $M = \sup A$.
- Thus $\forall \epsilon > 0, \exists a \in A \cap N_\epsilon(M)$ by the definition of supremum.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in Bd(A)$.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in Bd(A)$.

(ii)

- Let $\epsilon > 0$ be given,

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \emptyset$ and $N_\epsilon(M) \cap A^c \neq \emptyset$
- Hence $M = \sup A \in Bd(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in Bd(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in \text{Bd}(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

- Now $b \in \text{Bd}(A)$

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in Bd(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

- Now $b \in Bd(A)$ if and only if $A \cap (b - \epsilon, b + \epsilon) \neq \phi$
 $A^c \cap (b - \epsilon, b + \epsilon) \neq \phi$.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in \text{Bd}(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

- Now $b \in \text{Bd}(A)$ if and only if $A \cap (b - \epsilon, b + \epsilon) \neq \phi$
 $A^c \cap (b - \epsilon, b + \epsilon) \neq \phi$.
- We have to prove that $b \in \text{Bd}(B)$.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in \text{Bd}(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

- Now $b \in \text{Bd}(A)$ if and only if $A \cap (b - \epsilon, b + \epsilon) \neq \phi$
 $A^c \cap (b - \epsilon, b + \epsilon) \neq \phi$.
- We have to prove that $b \in \text{Bd}(B)$.
- To the contrary suppose that $B^c \cap (b - \epsilon, b + \epsilon) = \phi$,

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in Bd(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

- Now $b \in Bd(A)$ if and only if $A \cap (b - \epsilon, b + \epsilon) \neq \phi$
 $A^c \cap (b - \epsilon, b + \epsilon) \neq \phi$.
- We have to prove that $b \in Bd(B)$.
- To the contrary suppose that $B^c \cap (b - \epsilon, b + \epsilon) = \phi$, this implies
 $(b - \epsilon, b + \epsilon) \subseteq B$.

Question 2 Contd...

- We also note that $\forall \epsilon > 0, M + \frac{\epsilon}{2} \in N_\epsilon(M)$ but $M + \frac{\epsilon}{2} \notin A$.
- Thus $N_\epsilon(M) \cap A \neq \phi$ and $N_\epsilon(M) \cap A^c \neq \phi$
- Hence $M = \sup A \in \text{Bd}(A)$.

(ii)

- Let $\epsilon > 0$ be given, clearly $b - \epsilon < b < b + \epsilon$.
- Then by the given property, we have

$$A \cap (b - \epsilon, b + \epsilon) = B \cap (b - \epsilon, b + \epsilon).$$

- Now $b \in \text{Bd}(A)$ if and only if $A \cap (b - \epsilon, b + \epsilon) \neq \phi$
 $A^c \cap (b - \epsilon, b + \epsilon) \neq \phi$.
- We have to prove that $b \in \text{Bd}(B)$.
- To the contrary suppose that $B^c \cap (b - \epsilon, b + \epsilon) = \phi$, this implies
 $(b - \epsilon, b + \epsilon) \subseteq B$.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon)$

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon)$

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Solution:

a) Ans:

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Solution:

a) **Ans: False.** Counterexample.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Solution:

a) **Ans: False.** Counterexample. $\bigcup_{n=1}^{\infty} [\frac{1}{n}, 1] = (0, 1]$.

b) **Ans:**

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Solution:

- a) **Ans: False.** Counterexample. $\bigcup_{n=1}^{\infty} [\frac{1}{n}, 1] = (0, 1]$.
- b) **Ans: False.** Counterexample.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Solution:

a) **Ans: False.** Counterexample. $\bigcup_{n=1}^{\infty} [\frac{1}{n}, 1] = (0, 1]$.

b) **Ans: False.** Counterexample. $S = [-1, 0) \cup (0, 1]$. The sequence $\frac{1}{n} \in S$ but limit is not in S . Hence not closed not compact.

Question 2 Contd...

- This implies, $(b - \epsilon, b + \epsilon) \subseteq B \cap (b - \epsilon, b + \epsilon) = A \cap (b - \epsilon, b + \epsilon)$.
- This gives $(b - \epsilon, b + \epsilon) \subseteq A \implies A^c \cap (b - \epsilon, b + \epsilon) = \phi$, contradiction. (why?)
- Hence $b \in \text{Bd}(B)$.
- Similarly we can prove other direction as well.

Question 3

- The union of infinitely many compact sets is compact.
- A non-empty subset S of real numbers which has both a largest and a smallest element is compact.

Solution:

a) **Ans: False.** Counterexample. $\bigcup_{n=1}^{\infty} [\frac{1}{n}, 1] = (0, 1]$.

b) **Ans: False.** Counterexample. $S = [-1, 0) \cup (0, 1]$. The sequence $\frac{1}{n} \in S$ but limit is not in S . Hence not closed not compact.

Question 4

Question 4

For $A \subseteq \mathbb{R}$, $B \subseteq \mathbb{R}$, let

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let A be closed set, B be a compact set. Show that $A + B$ is closed.

Question 4

Question 4

For $A \subseteq \mathbb{R}$, $B \subseteq \mathbb{R}$, let

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let A be closed set, B be a compact set. Show that $A + B$ is closed.

Solution:

- Let $x + y$ be the limit point of $A + B$.

Question 4

Question 4

For $A \subseteq \mathbb{R}$, $B \subseteq \mathbb{R}$, let

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let A be closed set, B be a compact set. Show that $A + B$ is closed.

Solution:

- Let $x + y$ be the limit point of $A + B$.
- To show that $A + B$ is closed, it is sufficient to show that $x + y \in A + B$.

Question 4

Question 4

For $A \subseteq \mathbb{R}$, $B \subseteq \mathbb{R}$, let

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let A be closed set, B be a compact set. Show that $A + B$ is closed.

Solution:

- Let $x + y$ be the limit point of $A + B$.
- To show that $A + B$ is closed, it is sufficient to show that $x + y \in A + B$.
- Since $x + y$ is limit point of $A + B$. This implies there exists a sequence $\{x_n\} \in A$ and $\{y_n\} \in B$ such that $\{x_n + y_n\} \rightarrow x + y$ as $n \rightarrow \infty$.

Question 4

Question 4

For $A \subseteq \mathbb{R}$, $B \subseteq \mathbb{R}$, let

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let A be closed set, B be a compact set. Show that $A + B$ is closed.

Solution:

- Let $x + y$ be the limit point of $A + B$.
- To show that $A + B$ is closed, it is sufficient to show that $x + y \in A + B$.
- Since $x + y$ is limit point of $A + B$. This implies there exists a sequence $\{x_n\} \in A$ and $\{y_n\} \in B$ such that $\{x_n + y_n\} \rightarrow x + y$ as $n \rightarrow \infty$.
- Since B is compact, this implies B is closed and bounded.

Question 4

Question 4

For $A \subseteq \mathbb{R}$, $B \subseteq \mathbb{R}$, let

$$A + B = \{a + b : a \in A, b \in B\}.$$

Let A be closed set, B be a compact set. Show that $A + B$ is closed.

Solution:

- Let $x + y$ be the limit point of $A + B$.
- To show that $A + B$ is closed, it is sufficient to show that $x + y \in A + B$.
- Since $x + y$ is limit point of $A + B$. This implies there exists a sequence $\{x_n\} \in A$ and $\{y_n\} \in B$ such that $\{x_n + y_n\} \rightarrow x + y$ as $n \rightarrow \infty$.
- Since B is compact, this implies B is closed and bounded.

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)
- Note that $|x_{n_k}| \leq |x_{n_k} + y_{n_k}| + |y_{n_k}|$.

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)
- Note that $|x_{n_k}| \leq |x_{n_k} + y_{n_k}| + |y_{n_k}|$. Since $\{x_n + y_n\}$ and $\{y_n\}$ are bounded, this implies $\{x_{n_k}\}$ is also bounded.

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)
- Note that $|x_{n_k}| \leq |x_{n_k} + y_{n_k}| + |y_{n_k}|$. Since $\{x_n + y_n\}$ and $\{y_n\}$ are bounded, this implies $\{x_{n_k}\}$ is also bounded.
- By BWT, there exists a subsequence $x_{n_{k_j}}$ of x_{n_k} such that $x_{n_{k_j}} \rightarrow a \in A$. (why?)

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)
- Note that $|x_{n_k}| \leq |x_{n_k} + y_{n_k}| + |y_{n_k}|$. Since $\{x_n + y_n\}$ and $\{y_n\}$ are bounded, this implies $\{x_{n_k}\}$ is also bounded.
- By BWT, there exists a subsequence $x_{n_{k_j}}$ of x_{n_k} such that $x_{n_{k_j}} \rightarrow a \in A$. (why?)
- This implies $x_{n_{k_j}} + y_{n_{k_j}} \rightarrow a + b \in A + B$.

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)
- Note that $|x_{n_k}| \leq |x_{n_k} + y_{n_k}| + |y_{n_k}|$. Since $\{x_n + y_n\}$ and $\{y_n\}$ are bounded, this implies $\{x_{n_k}\}$ is also bounded.
- By BWT, there exists a subsequence $x_{n_{k_j}}$ of x_{n_k} such that $x_{n_{k_j}} \rightarrow a \in A$. (why?)
- This implies $x_{n_{k_j}} + y_{n_{k_j}} \rightarrow a + b \in A + B$.
- Since $\{x_{n_{k_j}} + y_{n_{k_j}}\}$ is a subsequence of $\{x_n + y_n\}$, but $x_n + y_n \rightarrow x + y$.

Question 4 Contd...

- By BWT, there exists a subsequence $\{y_{n_k}\}$ of $\{y_n\}$ such that $y_{n_k} \rightarrow b \in B$. (why?)
- Note that $|x_{n_k}| \leq |x_{n_k} + y_{n_k}| + |y_{n_k}|$. Since $\{x_n + y_n\}$ and $\{y_n\}$ are bounded, this implies $\{x_{n_k}\}$ is also bounded.
- By BWT, there exists a subsequence $x_{n_{k_j}}$ of x_{n_k} such that $x_{n_{k_j}} \rightarrow a \in A$. (why?)
- This implies $x_{n_{k_j}} + y_{n_{k_j}} \rightarrow a + b \in A + B$.
- Since $\{x_{n_{k_j}} + y_{n_{k_j}}\}$ is a subsequence of $\{x_n + y_n\}$, but $x_n + y_n \rightarrow x + y$.
- Hence $x + y = a + b \in A + B$.

Question 5

Question 5

Let (X, d) be a metric space. Define

$$\bar{d}(x, y) = \begin{cases} d(x, y) & \text{when } d(x, y) < 1 \\ 1 & \text{when } d(x, y) \geq 1. \end{cases}$$

Prove that \bar{d} is a metric on X .

Question 5

Question 5

Let (X, d) be a metric space. Define

$$\bar{d}(x, y) = \begin{cases} d(x, y) & \text{when } d(x, y) < 1 \\ 1 & \text{when } d(x, y) \geq 1. \end{cases}$$

Prove that \bar{d} is a metric on X .

Question 5

Question 5

Let (X, d) be a metric space. Define

$$\bar{d}(x, y) = \begin{cases} d(x, y) & \text{when } d(x, y) < 1 \\ 1 & \text{when } d(x, y) \geq 1. \end{cases}$$

Prove that \bar{d} is a metric on X .

Solution:

- Since d is metric, clearly \bar{d} satisfies positivity.

Question 5

Question 5

Let (X, d) be a metric space. Define

$$\bar{d}(x, y) = \begin{cases} d(x, y) & \text{when } d(x, y) < 1 \\ 1 & \text{when } d(x, y) \geq 1. \end{cases}$$

Prove that \bar{d} is a metric on X .

Solution:

- Since d is metric, clearly \bar{d} satisfies positivity.
- Since d satisfies symmetricity, so is \bar{d} .

Question 5

Question 5

Let (X, d) be a metric space. Define

$$\bar{d}(x, y) = \begin{cases} d(x, y) & \text{when } d(x, y) < 1 \\ 1 & \text{when } d(x, y) \geq 1. \end{cases}$$

Prove that \bar{d} is a metric on X .

Solution:

- Since d is metric, clearly \bar{d} satisfies positivity.
- Since d satisfies symmetricity, so is \bar{d} .
- Now we will check triangle inequality.

Question 5

Question 5

Let (X, d) be a metric space. Define

$$\bar{d}(x, y) = \begin{cases} d(x, y) & \text{when } d(x, y) < 1 \\ 1 & \text{when } d(x, y) \geq 1. \end{cases}$$

Prove that \bar{d} is a metric on X .

Solution:

- Since d is metric, clearly \bar{d} satisfies positivity.
- Since d satisfies symmetricity, so is \bar{d} .
- Now we will check triangle inequality.

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes



Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes
> 1	> 1	> 1	1	1	1	Yes

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes
> 1	> 1	> 1	1	1	1	Yes
> 1	> 1	< 1	1	1	$d(y, z)$	Yes

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes
> 1	> 1	> 1	1	1	1	Yes
> 1	> 1	< 1	1	1	$d(y, z)$	Yes
> 1	< 1	< 1	1	$d(x, y)$	$d(y, z)$	Yes

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes
> 1	> 1	> 1	1	1	1	Yes
> 1	> 1	< 1	1	1	$d(y, z)$	Yes
> 1	< 1	< 1	1	$d(x, y)$	$d(y, z)$	Yes
< 1	< 1	> 1	$d(x, z)$	$d(x, y)$	1	Yes

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes
> 1	> 1	> 1	1	1	1	Yes
> 1	> 1	< 1	1	1	$d(y, z)$	Yes
> 1	< 1	< 1	1	$d(x, y)$	$d(y, z)$	Yes
< 1	< 1	> 1	$d(x, z)$	$d(x, y)$	1	Yes

- Clearly from above table \bar{d} satisfies triangle inequality.

Question 5 Contd...

- We need to verify $\bar{d}(x, z) \leq \bar{d}(x, y) + \bar{d}(y, z)$.

$d(x, z)$	$d(x, y)$	$d(y, z)$	$\bar{d}(x, z)$	$\bar{d}(x, y)$	$\bar{d}(y, z)$	Yes/No
< 1	< 1	< 1	$d(x, z)$	$d(x, y)$	$d(y, z)$	Yes
> 1	> 1	> 1	1	1	1	Yes
> 1	> 1	< 1	1	1	$d(y, z)$	Yes
> 1	< 1	< 1	1	$d(x, y)$	$d(y, z)$	Yes
< 1	< 1	> 1	$d(x, z)$	$d(x, y)$	1	Yes

- Clearly from above table \bar{d} satisfies triangle inequality.
- Hence \bar{d} is a metric.

Question 6

Question 6

Suppose that $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies $\phi(0) = 0$, $\phi(r) > 0$ for all $r > 0$ and for all $a, b \in [0, \infty)$:

- 1) $\phi(a + b) \leq \phi(a) + \phi(b)$
- 2) If $a \leq b$ then $\phi(a) \leq \phi(b)$.

Let (X, d) be a metric space and let $D : X \times X \rightarrow \mathbb{R}$ be defined by $D(x, y) := \phi(d(x, y))$. Prove that D is a metric on X .

Question 6

Question 6

Suppose that $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies $\phi(0) = 0$, $\phi(r) > 0$ for all $r > 0$ and for all $a, b \in [0, \infty)$:

- 1) $\phi(a + b) \leq \phi(a) + \phi(b)$
- 2) If $a \leq b$ then $\phi(a) \leq \phi(b)$.

Let (X, d) be a metric space and let $D : X \times X \rightarrow \mathbb{R}$ be defined by $D(x, y) := \phi(d(x, y))$. Prove that D is a metric on X .

Solution:

- Since ϕ is non-negative function, this implies D satisfies positivity.

Question 6

Question 6

Suppose that $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies $\phi(0) = 0, \phi(r) > 0$ for all $r > 0$ and for all $a, b \in [0, \infty)$:

- 1) $\phi(a + b) \leq \phi(a) + \phi(b)$
- 2) If $a \leq b$ then $\phi(a) \leq \phi(b)$.

Let (X, d) be a metric space and let $D : X \times X \rightarrow \mathbb{R}$ be defined by $D(x, y) := \phi(d(x, y))$. Prove that D is a metric on X .

Solution:

- Since ϕ is non-negative function, this implies D satisfies positivity.
- $D(x, y) = 0 \iff \phi(d(x, y)) = 0 \iff d(x, y) = 0 \iff x = y$.

Question 6

Question 6

Suppose that $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies $\phi(0) = 0, \phi(r) > 0$ for all $r > 0$ and for all $a, b \in [0, \infty)$:

- 1) $\phi(a + b) \leq \phi(a) + \phi(b)$
- 2) If $a \leq b$ then $\phi(a) \leq \phi(b)$.

Let (X, d) be a metric space and let $D : X \times X \rightarrow \mathbb{R}$ be defined by $D(x, y) := \phi(d(x, y))$. Prove that D is a metric on X .

Solution:

- Since ϕ is non-negative function, this implies D satisfies positivity.
- $D(x, y) = 0 \iff \phi(d(x, y)) = 0 \iff d(x, y) = 0 \iff x = y$.
- Symmetric

$$D(x, y) = \phi(d(x, y)) = \phi(d(y, x)) = D(y, x).$$

Question 6

Question 6

Suppose that $\phi : [0, \infty) \rightarrow [0, \infty)$ satisfies $\phi(0) = 0, \phi(r) > 0$ for all $r > 0$ and for all $a, b \in [0, \infty)$:

- 1) $\phi(a + b) \leq \phi(a) + \phi(b)$
- 2) If $a \leq b$ then $\phi(a) \leq \phi(b)$.

Let (X, d) be a metric space and let $D : X \times X \rightarrow \mathbb{R}$ be defined by $D(x, y) := \phi(d(x, y))$. Prove that D is a metric on X .

Solution:

- Since ϕ is non-negative function, this implies D satisfies positivity.
- $D(x, y) = 0 \iff \phi(d(x, y)) = 0 \iff d(x, y) = 0 \iff x = y$.
- Symmetric

$$D(x, y) = \phi(d(x, y)) = \phi(d(y, x)) = D(y, x).$$

Question 6 Contd...

- Triangle inequality

$$D(x, z) = \phi(d(x, z))$$

Question 6 Contd...

- Triangle inequality

$$\begin{aligned} D(x, z) &= \phi(d(x, z)) \\ &\leq \phi(d(x, z) + d(z, y)) \text{ (why?)} \end{aligned}$$

- Triangle inequality

$$\begin{aligned} D(x, z) &= \phi(d(x, z)) \\ &\leq \phi(d(x, z) + d(z, y))(\text{why?}) \\ &\leq \phi(d(x, z)) + \phi(d(z, y))(\text{why?}) \end{aligned}$$

Question 6 Contd...

- Triangle inequality

$$\begin{aligned} D(x, z) &= \phi(d(x, z)) \\ &\leq \phi(d(x, z) + d(z, y)) \text{ (why?)} \\ &\leq \phi(d(x, z)) + \phi(d(z, y)) \text{ (why?)} \\ &= D(x, y) + D(y, z) \end{aligned}$$

- Hence D is a metric.

Question 7

Question 7

Let $(X_1, d_1), (X_2, d_2), \dots$ be a sequence of metric spaces. Let $X = \prod_{n \in \mathbb{N}} X_n$ i.e, X is the set of all sequences $x = (x_1, x_2, \dots)$ with $x_n \in X_n$ for all $n \in \mathbb{N}$. Prove that the function $d : X \times X \rightarrow \mathbb{R}$ defined by

$$d(x, y) = \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)}$$

is a metric on X .

Question 7

Question 7

Let $(X_1, d_1), (X_2, d_2), \dots$ be a sequence of metric spaces. Let $X = \prod_{n \in \mathbb{N}} X_n$ i.e, X is the set of all sequences $x = (x_1, x_2, \dots)$ with $x_n \in X_n$ for all $n \in \mathbb{N}$. Prove that the function $d : X \times X \rightarrow \mathbb{R}$ defined by

$$d(x, y) = \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)}$$

is a metric on X .

Solution:

- **Well-defined:** Since $\frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} < 1$. This implies

$$d(x, y) = \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)}$$

$$\begin{aligned} d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\ &< \sum_{n=1}^{\infty} 2^{-n} \end{aligned}$$

$$\begin{aligned} d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\ &< \sum_{n=1}^{\infty} 2^{-n} \\ &< \infty. \text{ (why?)} \end{aligned}$$

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.
- $d(x, y) = 0$

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.
- $d(x, y) = 0 \iff \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} = 0$

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.

- $d(x, y) = 0 \iff \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} = 0 \iff d_n(x_n, y_n) = 0$ for all $n \in \mathbb{N}$

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.

$$\begin{aligned}
 d(x, y) = 0 &\iff \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} = 0 \iff d_n(x_n, y_n) = 0 \text{ for all} \\
 n \in \mathbb{N} &\iff x_n = y_n \text{ for all } n \in \mathbb{N}
 \end{aligned}$$

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{ (why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.

$$\begin{aligned}
 d(x, y) = 0 &\iff \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} = 0 \iff d_n(x_n, y_n) = 0 \text{ for all} \\
 n \in \mathbb{N} &\iff x_n = y_n \text{ for all } n \in \mathbb{N} \iff x = y.
 \end{aligned}$$

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{(why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.
- $d(x, y) = 0 \iff \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} = 0 \iff d_n(x_n, y_n) = 0$ for all $n \in \mathbb{N} \iff x_n = y_n$ for all $n \in \mathbb{N} \iff x = y$.
- Clearly d is symmetric as well. (why?)

$$\begin{aligned}
 d(x, y) &= \sum_{n=1}^{\infty} 2^{-n} \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} \\
 &< \sum_{n=1}^{\infty} 2^{-n} \\
 &< \infty. \text{(why?)}
 \end{aligned}$$

Thus d is well-defined.

- Clear d is non-negative.
- $d(x, y) = 0 \iff \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} = 0 \iff d_n(x_n, y_n) = 0$ for all $n \in \mathbb{N} \iff x_n = y_n$ for all $n \in \mathbb{N} \iff x = y$.
- Clearly d is symmetric as well. (why?)

Question 7 Contd...

- **Triangle inequality**

Question 7 Contd...

- **Triangle inequality**

$$d_n(x_n, z_n) \leq d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

Question 7 Contd...

- **Triangle inequality**

$$d_n(x_n, z_n) \leq d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

$$1 + d_n(x_n, z_n) \leq 1 + d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

Question 7 Contd...

- **Triangle inequality**

$$d_n(x_n, z_n) \leq d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

$$1 + d_n(x_n, z_n) \leq 1 + d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

$$1 - \frac{1}{1 + d_n(x_n, z_n)} \leq 1 - \frac{1}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

Question 7 Contd...

- **Triangle inequality**

$$d_n(x_n, z_n) \leq d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

$$1 + d_n(x_n, z_n) \leq 1 + d_n(x_n, y_n) + d_n(y_n, z_n) (\text{why?})$$

$$1 - \frac{1}{1 + d_n(x_n, z_n)} \leq 1 - \frac{1}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$\frac{d_n(x_n, z_n)}{1 + d_n(x_n, z_n)} \leq \frac{d_n(x_n, y_n) + d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

Question 7 Contd...

- Triangle inequality

$$d_n(x_n, z_n) \leq d_n(x_n, y_n) + d_n(y_n, z_n) \text{ (why?)}$$

$$1 + d_n(x_n, z_n) \leq 1 + d_n(x_n, y_n) + d_n(y_n, z_n) \text{ (why?)}$$

$$1 - \frac{1}{1 + d_n(x_n, z_n)} \leq 1 - \frac{1}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$\frac{d_n(x_n, z_n)}{1 + d_n(x_n, z_n)} \leq \frac{d_n(x_n, y_n) + d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$\leq \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$+ \frac{d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

Question 7 Contd...

- Triangle inequality

$$d_n(x_n, z_n) \leq d_n(x_n, y_n) + d_n(y_n, z_n) \text{ (why?)}$$

$$1 + d_n(x_n, z_n) \leq 1 + d_n(x_n, y_n) + d_n(y_n, z_n) \text{ (why?)}$$

$$1 - \frac{1}{1 + d_n(x_n, z_n)} \leq 1 - \frac{1}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$\frac{d_n(x_n, z_n)}{1 + d_n(x_n, z_n)} \leq \frac{d_n(x_n, y_n) + d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$\leq \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$+ \frac{d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)}$$

$$\leq \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} + \frac{d_n(y_n, z_n)}{1 + d_n(y_n, z_n)}$$

Question 7 Contd...

- **Triangle inequality**

$$\begin{aligned}d_n(x_n, z_n) &\leq d_n(x_n, y_n) + d_n(y_n, z_n) \text{ (why?)} \\1 + d_n(x_n, z_n) &\leq 1 + d_n(x_n, y_n) + d_n(y_n, z_n) \text{ (why?)} \\1 - \frac{1}{1 + d_n(x_n, z_n)} &\leq 1 - \frac{1}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)} \\ \frac{d_n(x_n, z_n)}{1 + d_n(x_n, z_n)} &\leq \frac{d_n(x_n, y_n) + d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)} \\ &\leq \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)} \\ &\quad + \frac{d_n(y_n, z_n)}{1 + d_n(x_n, y_n) + d_n(y_n, z_n)} \\ &\leq \frac{d_n(x_n, y_n)}{1 + d_n(x_n, y_n)} + \frac{d_n(y_n, z_n)}{1 + d_n(y_n, z_n)}\end{aligned}$$

- Now by multiplying by 2^{-n} then taking sum over \mathbb{N} , we will get our required triangle inequality. Hence d is a metric.

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0$

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0$

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0 \iff n = m.$

Question 8

Question 8

Prove that the function $d(m, n) = \left| \frac{1}{m} - \frac{1}{n} \right|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff \left| \frac{1}{m} - \frac{1}{n} \right| = 0 \iff n = m.$
- $d(m, n) = \left| \frac{1}{m} - \frac{1}{n} \right| = \left| \frac{1}{n} - \frac{1}{m} \right| = d(n, m),$ hence symmetric.

Question 8

Question 8

Prove that the function $d(m, n) = \left| \frac{1}{m} - \frac{1}{n} \right|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff \left| \frac{1}{m} - \frac{1}{n} \right| = 0 \iff n = m$.
- $d(m, n) = \left| \frac{1}{m} - \frac{1}{n} \right| = \left| \frac{1}{n} - \frac{1}{m} \right| = d(n, m)$, hence symmetric.
- $d(m, n) = \left| \frac{1}{m} - \frac{1}{k} + \frac{1}{k} - \frac{1}{n} \right| \leq \left| \frac{1}{m} - \frac{1}{k} \right| + \left| \frac{1}{k} - \frac{1}{n} \right| \leq d(m, k) + d(k, n)$.

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0 \iff n = m$.
- $d(m, n) = |\frac{1}{m} - \frac{1}{n}| = |\frac{1}{n} - \frac{1}{m}| = d(n, m)$, hence symmetric.
- $d(m, n) = |\frac{1}{m} - \frac{1}{k} + \frac{1}{k} - \frac{1}{n}| \leq |\frac{1}{m} - \frac{1}{k}| + |\frac{1}{k} - \frac{1}{n}| \leq d(m, k) + d(k, n)$.
Thus d satisfies triangle inequality.

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0 \iff n = m$.
- $d(m, n) = |\frac{1}{m} - \frac{1}{n}| = |\frac{1}{n} - \frac{1}{m}| = d(n, m)$, hence symmetric.
- $d(m, n) = |\frac{1}{m} - \frac{1}{k} + \frac{1}{k} - \frac{1}{n}| \leq |\frac{1}{m} - \frac{1}{k}| + |\frac{1}{k} - \frac{1}{n}| \leq d(m, k) + d(k, n)$.
Thus d satisfies triangle inequality.
- Hence d is a metric.

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0 \iff n = m$.
- $d(m, n) = |\frac{1}{m} - \frac{1}{n}| = |\frac{1}{n} - \frac{1}{m}| = d(n, m)$, hence symmetric.
- $d(m, n) = |\frac{1}{m} - \frac{1}{k} + \frac{1}{k} - \frac{1}{n}| \leq |\frac{1}{m} - \frac{1}{k}| + |\frac{1}{k} - \frac{1}{n}| \leq d(m, k) + d(k, n)$.
Thus d satisfies triangle inequality.
- Hence d is a metric.

Does this metric extend to \mathbb{R}^+ .

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0 \iff n = m$.
- $d(m, n) = |\frac{1}{m} - \frac{1}{n}| = |\frac{1}{n} - \frac{1}{m}| = d(n, m)$, hence symmetric.
- $d(m, n) = |\frac{1}{m} - \frac{1}{k} + \frac{1}{k} - \frac{1}{n}| \leq |\frac{1}{m} - \frac{1}{k}| + |\frac{1}{k} - \frac{1}{n}| \leq d(m, k) + d(k, n)$.
Thus d satisfies triangle inequality.
- Hence d is a metric.

Does this metric extend to \mathbb{R}^+ .

Ans:

Question 8

Question 8

Prove that the function $d(m, n) = |\frac{1}{m} - \frac{1}{n}|$ for any $m, n \in \mathbb{N}$ defines a metric on the set of natural numbers. Does this metric extend to \mathbb{R}^+ .

Solution:

- Clearly d is non-negative.
- $d(n, m) = 0 \iff |\frac{1}{m} - \frac{1}{n}| = 0 \iff n = m$.
- $d(m, n) = |\frac{1}{m} - \frac{1}{n}| = |\frac{1}{n} - \frac{1}{m}| = d(n, m)$, hence symmetric.
- $d(m, n) = |\frac{1}{m} - \frac{1}{k} + \frac{1}{k} - \frac{1}{n}| \leq |\frac{1}{m} - \frac{1}{k}| + |\frac{1}{k} - \frac{1}{n}| \leq d(m, k) + d(k, n)$.
Thus d satisfies triangle inequality.
- Hence d is a metric.

Does this metric extend to \mathbb{R}^+ .

Ans: Yes.

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Solution: (i)

- Let $x \in \partial A$,

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Solution: (i)

- Let $x \in \partial A$, i.e. for $\epsilon > 0$, we have $N_\epsilon(x) \cap A \neq \emptyset$ and $N_\epsilon(x) \cap A^c \neq \emptyset$.

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Solution: (i)

- Let $x \in \partial A$, i.e. for $\epsilon > 0$, we have $N_\epsilon(x) \cap A \neq \emptyset$ and $N_\epsilon(x) \cap A^c \neq \emptyset$.
- In particular, for $\epsilon_n = \frac{1}{n} > 0$, we have $N_{\frac{1}{n}}(x) \cap A \neq \emptyset$.

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Solution: (i)

- Let $x \in \partial A$, i.e. for $\epsilon > 0$, we have $N_\epsilon(x) \cap A \neq \emptyset$ and $N_\epsilon(x) \cap A^c \neq \emptyset$.
- In particular, for $\epsilon_n = \frac{1}{n} > 0$, we have $N_{\frac{1}{n}}(x) \cap A \neq \emptyset$.
- This implies there exists $x_n \in N_{\frac{1}{n}}(x) \cap A$. This gives $x \in \bar{A}$.

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Solution: (i)

- Let $x \in \partial A$, i.e. for $\epsilon > 0$, we have $N_\epsilon(x) \cap A \neq \emptyset$ and $N_\epsilon(x) \cap A^c \neq \emptyset$.
- In particular, for $\epsilon_n = \frac{1}{n} > 0$, we have $N_{\frac{1}{n}}(x) \cap A \neq \emptyset$.
- This implies there exists $x_n \in N_{\frac{1}{n}}(x) \cap A$. This gives $x \in \bar{A}$.
- Since $N_\epsilon(x) \cap A^c \neq \emptyset$, i.e. $N_{\frac{1}{n}}(x)$ always contains element of A^c .

Question 9

Question 9

Let A be a subset of a metric space X with closure \bar{A} and boundary of A by ∂A . Show that

- (i) Show that $\partial A = \bar{A} \setminus A^\circ$ and ∂A is closed.
- (ii) Prove that $X \setminus \bar{A} = (X \setminus A)^\circ$.
- (iii) Prove that A is closed if and only if $\partial A \subset A$, and A is open if and only if $\partial A \subset A^c$.
- (iv) If A is open, does it follow that $(\bar{A})^\circ = A$?

Solution: (i)

- Let $x \in \partial A$, i.e. for $\epsilon > 0$, we have $N_\epsilon(x) \cap A \neq \emptyset$ and $N_\epsilon(x) \cap A^c \neq \emptyset$.
- In particular, for $\epsilon_n = \frac{1}{n} > 0$, we have $N_{\frac{1}{n}}(x) \cap A \neq \emptyset$.
- This implies there exists $x_n \in N_{\frac{1}{n}}(x) \cap A$. This gives $x \in \bar{A}$.
- Since $N_\epsilon(x) \cap A^c \neq \emptyset$, i.e. $N_{\frac{1}{n}}(x)$ always contains element of A^c .

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$.

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$.

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$,

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

(ii)

$$x \in X \setminus \bar{A} \iff x \in X \text{ but } x \notin \bar{A}$$

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

(ii)

$$\begin{aligned} x \in X \setminus \bar{A} &\iff x \in X \text{ but } x \notin \bar{A} \\ &\iff x \text{ is not a limit point of } A \end{aligned}$$

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

(ii)

$$\begin{aligned} x \in X \setminus \bar{A} &\iff x \in X \text{ but } x \notin \bar{A} \\ &\iff x \text{ is not a limit point of } A \\ &\iff \exists \epsilon > 0 \text{ such that } N_\epsilon(x) \cap A = \emptyset \end{aligned}$$

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

(ii)

$$\begin{aligned} x \in X \setminus \bar{A} &\iff x \in X \text{ but } x \notin \bar{A} \\ &\iff x \text{ is not a limit point of } A \\ &\iff \exists \epsilon > 0 \text{ such that } N_\epsilon(x) \cap A = \emptyset \\ &\iff N_\epsilon(x) \subset A^c \end{aligned}$$

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

(ii)

$$\begin{aligned} x \in X \setminus \bar{A} &\iff x \in X \text{ but } x \notin \bar{A} \\ &\iff x \text{ is not a limit point of } A \\ &\iff \exists \epsilon > 0 \text{ such that } N_\epsilon(x) \cap A = \emptyset \\ &\iff N_\epsilon(x) \subset A^c \\ &\iff x \text{ is interior point of } X \setminus A \text{ .i.e } x \in (X \setminus A)^\circ \end{aligned}$$

Question 9 Contd...

- Thus $N_{\frac{1}{n}}(x) \not\subseteq A$. Therefore $x \notin A^\circ$. Hence $x \in \bar{A} \setminus A^\circ$.
- Since every step was following if and only if, Thus we get $\partial A = \bar{A} \setminus A^\circ$.

Now we will prove that ∂A is closed.

- Since $\partial A = \bar{A} \setminus A^\circ = \bar{A} \cap (A^\circ)^c$, \bar{A} is always closed and A° is always open.
- Hence ∂A is closed

(ii)

$$\begin{aligned} x \in X \setminus \bar{A} &\iff x \in X \text{ but } x \notin \bar{A} \\ &\iff x \text{ is not a limit point of } A \\ &\iff \exists \epsilon > 0 \text{ such that } N_\epsilon(x) \cap A = \emptyset \\ &\iff N_\epsilon(x) \subset A^c \\ &\iff x \text{ is interior point of } X \setminus A \text{ .i.e } x \in (X \setminus A)^\circ \end{aligned}$$

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.
Assume that A is open

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open,

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that
$$N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset.$$

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A .

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

- Let $x \in A$

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

- Let $x \in A \implies x$ is not a boundary point.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

- Let $x \in A \implies x$ is not a boundary point.
- There exists $\epsilon > 0$ such that $N_\epsilon(x) \cap A = \emptyset$ or $N_\epsilon(x) \cap A^c = \emptyset$.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

- Let $x \in A \implies x$ is not a boundary point.
- There exists $\epsilon > 0$ such that $N_\epsilon(x) \cap A = \emptyset$ or $N_\epsilon(x) \cap A^c = \emptyset$.
- But $x \in N_\epsilon \cap A \neq \emptyset$. Thus $N_\epsilon(x) \cap A^c = \emptyset \implies N_\epsilon(x) \subset A$.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

- Let $x \in A \implies x$ is not a boundary point.
- There exists $\epsilon > 0$ such that $N_\epsilon(x) \cap A = \emptyset$ or $N_\epsilon(x) \cap A^c = \emptyset$.
- But $x \in N_\epsilon \cap A \neq \emptyset$. Thus $N_\epsilon(x) \cap A^c = \emptyset \implies N_\epsilon(x) \subset A$.
- Hence A is open.

Question 9 Contd...

(iii) To show that A is open if and only if $\partial A \subset A^c$.

Assume that A is open

- To the contrary suppose that $x \in \partial A$ but $x \notin A^c$. i.e. $x \in A$.
- Since A is open, there exists $\epsilon > 0$ such that $N_\epsilon(x) \subseteq A \implies N_\epsilon(x) \cap A^c = \emptyset$.
- Thus x cannot be boundary point of A . Contradiction.
- Hence $\partial A \subset A^c$

Assume that $\partial A \subset A^c$

- Let $x \in A \implies x$ is not a boundary point.
- There exists $\epsilon > 0$ such that $N_\epsilon(x) \cap A = \emptyset$ or $N_\epsilon(x) \cap A^c = \emptyset$.
- But $x \in N_\epsilon \cap A \neq \emptyset$. Thus $N_\epsilon(x) \cap A^c = \emptyset \implies N_\epsilon(x) \subset A$.
- Hence A is open.

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$A \text{ is closed} \iff A^c \text{ is open}$$

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \end{aligned}$$

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

(iv)

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

(iv) Ans:

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

(iv) **Ans:** No Counterexample

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

(iv) **Ans:** No Counterexample

- Take $A = (-\infty, 0) \cup (0, \infty)$.

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

(iv) **Ans:** No Counterexample

- Take $A = (-\infty, 0) \cup (0, \infty)$.
- Clearly $\bar{A} = \mathbb{R}$ and $(\bar{A})^\circ = \mathbb{R}$ thus $(\bar{A})^\circ \neq A$.

Question 9 Contd...

Now we will prove that A is closed if and only if $\partial A \subset A$.

$$\begin{aligned} A \text{ is closed} &\iff A^c \text{ is open} \\ &\iff \partial A^c \subset (A^c)^c \quad (\text{why?}) \\ &\iff \partial A \subset A \quad (\text{why?}) \end{aligned}$$

(iv) **Ans:** No Counterexample

- Take $A = (-\infty, 0) \cup (0, \infty)$.
- Clearly $\bar{A} = \mathbb{R}$ and $(\bar{A})^\circ = \mathbb{R}$ thus $(\bar{A})^\circ \neq A$.

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}$$

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

- Clearly E is bounded.

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

- Clearly E is bounded.
- Clearly there doesn't exist any rational number q such that $q^2 = 2$ or $q^2 = 3$.

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

- Clearly E is bounded.
- Clearly there doesn't exist any rational number q such that $q^2 = 2$ or $q^2 = 3$.
- Let $q \in \mathbb{Q}$ and $q^2 < 2$.

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

- Clearly E is bounded.
- Clearly there doesn't exist any rational number q such that $q^2 = 2$ or $q^2 = 3$.
- Let $q \in \mathbb{Q}$ and $q^2 < 2$.
- Take $r_1 = \frac{\sqrt{2}-|q|}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

- Clearly E is bounded.
- Clearly there doesn't exist any rational number q such that $q^2 = 2$ or $q^2 = 3$.
- Let $q \in \mathbb{Q}$ and $q^2 < 2$.
- Take $r_1 = \frac{\sqrt{2}-|q|}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.
- Thus $q \in \mathbb{Q}$ with $q^2 < 2$ can't be the limit point of E .

Question 10 Contd...

Question 10

Let \mathbb{Q} , the set of rational numbers, as a metric space with the Euclidean distance $d(p, q) = |p - q|$. Consider the set

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\}.$$

Show that E is closed and bounded in \mathbb{Q} .

Solution: We have

$$E = \{p \in \mathbb{Q} : 2 < p^2 < 3\} = \mathbb{Q} \cap (-\sqrt{3}, -\sqrt{2}) \cup (\sqrt{2}, \sqrt{3}).$$

- Clearly E is bounded.
- Clearly there doesn't exist any rational number q such that $q^2 = 2$ or $q^2 = 3$.
- Let $q \in \mathbb{Q}$ and $q^2 < 2$.
- Take $r_1 = \frac{\sqrt{2}-|q|}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.
- Thus $q \in \mathbb{Q}$ with $q^2 < 2$ can't be the limit point of E .

Question 10 Contd...

- Let $q \in \mathbb{Q}$ and $q^2 > 3$.

Question 10 Contd...

- Let $q \in \mathbb{Q}$ and $q^2 > 3$.
- Take $r_1 = \frac{|q| - \sqrt{3}}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.

Question 10 Contd...

- Let $q \in \mathbb{Q}$ and $q^2 > 3$.
- Take $r_1 = \frac{|q| - \sqrt{3}}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.
- Thus $q \in \mathbb{Q}$ with $q^2 > 3$ can't be the limit point of E .

Question 10 Contd...

- Let $q \in \mathbb{Q}$ and $q^2 > 3$.
- Take $r_1 = \frac{|q| - \sqrt{3}}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.
- Thus $q \in \mathbb{Q}$ with $q^2 > 3$ can't be the limit point of E .
- Therefore only possible limit point of E are $q \in \mathbb{Q}$ with $2 < q^2 < 3$.

Question 10 Contd...

- Let $q \in \mathbb{Q}$ and $q^2 > 3$.
- Take $r_1 = \frac{|q| - \sqrt{3}}{2} > 0$. then $N_{r_1}(q) \cap E = \emptyset$.
- Thus $q \in \mathbb{Q}$ with $q^2 > 3$ can't be the limit point of E .
- Therefore only possible limit point of E are $q \in \mathbb{Q}$ with $2 < q^2 < 3$.
- Hence E is closed.