

MTL122 - Real and complex analysis

Assignment-6



Department of Mathematics
Indian Institute of Technology Delhi

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Recall

For $z = x + iy$, we have $z = re^{i\theta}$ where $r = \sqrt{x^2 + y^2}$ and $\theta = \arctan(\frac{y}{x})$.

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Recall

For $z = x + iy$, we have $z = re^{i\theta}$ where $r = \sqrt{x^2 + y^2}$ and $\theta = \arctan(\frac{y}{x})$.

Proof(a)

- We need to find r and θ , such that $-3 + i3 = re^{i\theta}$.

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Recall

For $z = x + iy$, we have $z = re^{i\theta}$ where $r = \sqrt{x^2 + y^2}$ and $\theta = \arctan(\frac{y}{x})$.

Proof(a)

- We need to find r and θ , such that $-3 + i3 = re^{i\theta}$. We have $r = 3\sqrt{2}$

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Recall

For $z = x + iy$, we have $z = re^{i\theta}$ where $r = \sqrt{x^2 + y^2}$ and $\theta = \arctan(\frac{y}{x})$.

Proof(a)

- We need to find r and θ , such that $-3 + i3 = re^{i\theta}$. We have $r = 3\sqrt{2}$ and $\theta = \arctan(\frac{3}{-3}) = \arctan(-1) = \frac{3\pi}{4}$.

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Recall

For $z = x + iy$, we have $z = re^{i\theta}$ where $r = \sqrt{x^2 + y^2}$ and $\theta = \arctan(\frac{y}{x})$.

Proof(a)

- We need to find r and θ , such that $-3 + i3 = re^{i\theta}$. We have $r = 3\sqrt{2}$ and $\theta = \arctan(\frac{3}{-3}) = \arctan(-1) = \frac{3\pi}{4}$.
- This gives $z = -3 + i3 = 3\sqrt{2}e^{\frac{3\pi}{4}}$.

Question 1

Question 1

Find the argument for each of the following complex numbers.

(a) $-3 + i3$

(b) $(1 - i)(-\sqrt{3} + i)$

(c) $\frac{-1 + i\sqrt{3}}{2 + i2}$

Recall

For $z = x + iy$, we have $z = re^{i\theta}$ where $r = \sqrt{x^2 + y^2}$ and $\theta = \arctan(\frac{y}{x})$.

Proof(a)

- We need to find r and θ , such that $-3 + i3 = re^{i\theta}$. We have $r = 3\sqrt{2}$ and $\theta = \arctan(\frac{3}{-3}) = \arctan(-1) = \frac{3\pi}{4}$.
- This gives $z = -3 + i3 = 3\sqrt{2}e^{\frac{3\pi}{4}}$.

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$ and
 $-\sqrt{3} + i = 2(-\frac{\sqrt{3}}{2} + i\frac{1}{2}) = 2e^{i\frac{2\pi}{3}}.$

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$ and
 $-\sqrt{3} + i = 2(-\frac{\sqrt{3}}{2} + i\frac{1}{2}) = 2e^{i\frac{2\pi}{3}}$.
- Hence $(1 - i)(-\sqrt{3} + i) = 2\sqrt{2}e^{i\frac{5\pi}{12}}$.

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$ and
 $-\sqrt{3} + i = 2(-\frac{\sqrt{3}}{2} + i\frac{1}{2}) = 2e^{i\frac{2\pi}{3}}$.
- Hence $(1 - i)(-\sqrt{3} + i) = 2\sqrt{2}e^{i\frac{5\pi}{12}}$.

Proof(c)

- Write $\frac{-1 + i\sqrt{3}}{2 + i2} = \frac{r_1 e^{i\theta_1}}{r_2 e^{i\theta_2}}$

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$ and
 $-\sqrt{3} + i = 2(-\frac{\sqrt{3}}{2} + i\frac{1}{2}) = 2e^{i\frac{2\pi}{3}}$.
- Hence $(1 - i)(-\sqrt{3} + i) = 2\sqrt{2}e^{i\frac{5\pi}{12}}$.

Proof(c)

- Write $\frac{-1 + i\sqrt{3}}{2 + i2} = \frac{r_1 e^{i\theta_1}}{r_2 e^{i\theta_2}} = \frac{2e^{i\frac{5\pi}{6}}}{2\sqrt{2}e^{i\frac{\pi}{4}}}$

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$ and
 $-\sqrt{3} + i = 2(-\frac{\sqrt{3}}{2} + i\frac{1}{2}) = 2e^{i\frac{2\pi}{3}}$.
- Hence $(1 - i)(-\sqrt{3} + i) = 2\sqrt{2}e^{i\frac{5\pi}{12}}$.

Proof(c)

- Write $\frac{-1 + i\sqrt{3}}{2 + i2} = \frac{r_1 e^{i\theta_1}}{r_2 e^{i\theta_2}} = \frac{2e^{i\frac{5\pi}{6}}}{2\sqrt{2}e^{i\frac{\pi}{4}}} = \frac{1}{\sqrt{2}}e^{i\frac{7\pi}{12}}$.

Question 1 Contd...

Proof(b)

- Write $(1 - i)(-\sqrt{3} + i) = r_1 e^{i\theta_1} r_2 e^{i\theta_2} = r_1 r_2 e^{i(\theta_1 + \theta_2)}$.
- Now $1 - i = \sqrt{2}(\frac{1}{\sqrt{2}} - i\frac{1}{\sqrt{2}}) = \sqrt{2}e^{-i\frac{\pi}{4}}$ and
 $-\sqrt{3} + i = 2(-\frac{\sqrt{3}}{2} + i\frac{1}{2}) = 2e^{i\frac{2\pi}{3}}$.
- Hence $(1 - i)(-\sqrt{3} + i) = 2\sqrt{2}e^{i\frac{5\pi}{12}}$.

Proof(c)

- Write $\frac{-1 + i\sqrt{3}}{2 + i2} = \frac{r_1 e^{i\theta_1}}{r_2 e^{i\theta_2}} = \frac{2e^{i\frac{5\pi}{6}}}{2\sqrt{2}e^{i\frac{\pi}{4}}} = \frac{1}{\sqrt{2}}e^{i\frac{7\pi}{12}}$.

Question 2

Question 2

Solve the equation $z^5 = 1$ for all complex numbers z .

Question 2

Question 2

Solve the equation $z^5 = 1$ for all complex numbers z .

Recall: De Moivre's formula

If z is a complex number, written in polar form as $z = r(\cos x + i \sin x)$. then the n -th roots of z are given by $r^{\frac{1}{n}} \left(\cos \frac{x+2\pi k}{n} + i \sin \frac{x+2\pi k}{n} \right)$ where k varies over the integer values from 0 to $n - 1$.

Proof

- We know $z^5 = 1 = \cos(0) + i \sin(0)$.

Question 2

Question 2

Solve the equation $z^5 = 1$ for all complex numbers z .

Recall: De Moivre's formula

If z is a complex number, written in polar form as $z = r(\cos x + i \sin x)$. then the n -th roots of z are given by $r^{\frac{1}{n}} \left(\cos \frac{x+2\pi k}{n} + i \sin \frac{x+2\pi k}{n} \right)$ where k varies over the integer values from 0 to $n - 1$.

Proof

- We know $z^5 = 1 = \cos(0) + i \sin(0)$.
- By De Moivre's formula, we have
$$z = (1)^{\frac{1}{5}} = \cos\left(\frac{2\pi k}{5}\right) + i \sin\left(\frac{2\pi k}{5}\right) = e^{i\frac{2\pi k}{5}} \text{ where } k = 0, 1, 2, 3, 4.$$

Question 2

Question 2

Solve the equation $z^5 = 1$ for all complex numbers z .

Recall: De Moivre's formula

If z is a complex number, written in polar form as $z = r(\cos x + i \sin x)$. then the n -th roots of z are given by $r^{\frac{1}{n}} \left(\cos \frac{x+2\pi k}{n} + i \sin \frac{x+2\pi k}{n} \right)$ where k varies over the integer values from 0 to $n - 1$.

Proof

- We know $z^5 = 1 = \cos(0) + i \sin(0)$.
- By De Moivre's formula, we have
$$z = (1)^{\frac{1}{5}} = \cos\left(\frac{2\pi k}{5}\right) + i \sin\left(\frac{2\pi k}{5}\right) = e^{i\frac{2\pi k}{5}} \text{ where } k = 0, 1, 2, 3, 4.$$
- So $z = 1, e^{\frac{2\pi i}{5}}, e^{\frac{4\pi i}{5}}, e^{\frac{6\pi i}{5}}, e^{\frac{8\pi i}{5}}$.

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|},$

(ii) $f(z) = xy^2 + iyx^2.$

Proof 3(i)

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|},$

(ii) $f(z) = xy^2 + iyx^2.$

Proof 3(i)

- Here $u(x, y) = \sqrt{|xy|}, v(x, y) = 0.$

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|},$

(ii) $f(z) = xy^2 + iyx^2.$

Proof 3(i)

- Here $u(x, y) = \sqrt{|xy|}, v(x, y) = 0$. Clearly $v_x(0, 0) = 0, v_y(0, 0) = 0$.

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|}$,

(ii) $f(z) = xy^2 + iyx^2$.

Proof 3(i)

- Here $u(x, y) = \sqrt{|xy|}$, $v(x, y) = 0$. Clearly $v_x(0, 0) = 0$, $v_y(0, 0) = 0$.
- Now

$$u_x(0, 0) = \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} = 0$$

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|}$,

(ii) $f(z) = xy^2 + iyx^2$.

Proof 3(i)

- Here $u(x, y) = \sqrt{|xy|}$, $v(x, y) = 0$. Clearly $v_x(0, 0) = 0$, $v_y(0, 0) = 0$.
- Now

$$u_x(0, 0) = \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} = 0$$

and

$$u_y(0, 0) = \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} = 0.$$

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|}$,

(ii) $f(z) = xy^2 + iyx^2$.

Proof 3(i)

- Here $u(x, y) = \sqrt{|xy|}$, $v(x, y) = 0$. Clearly $v_x(0, 0) = 0$, $v_y(0, 0) = 0$.
- Now

$$u_x(0, 0) = \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} = 0$$

and

$$u_y(0, 0) = \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} = 0.$$

Hence Cauchy-Riemann equations are satisfied.

Question 3

Question 3

Show that for each of the following functions, Cauchy-Riemann equations are satisfied at the origin. Also determine whether these functions are differentiable at $z = 0$. Are these functions analytic at $z = 0$?

(i) $f(z) = \sqrt{|Re(z)Im(z)|}$,

(ii) $f(z) = xy^2 + iyx^2$.

Proof 3(i)

- Here $u(x, y) = \sqrt{|xy|}$, $v(x, y) = 0$. Clearly $v_x(0, 0) = 0$, $v_y(0, 0) = 0$.
- Now

$$u_x(0, 0) = \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} = 0$$

and

$$u_y(0, 0) = \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} = 0.$$

Hence Cauchy-Riemann equations are satisfied.

Question 3(i) Contd...

• Now

$$\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} = \lim_{h \rightarrow 0} \frac{\sqrt{|h_1 h_2|}}{h_1 + ih_2}$$

Question 3(i) Contd...

• Now

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} &= \lim_{h \rightarrow 0} \frac{\sqrt{|h_1 h_2|}}{h_1 + ih_2} \\ &= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m|h_1^2}}{(1+im)h_1} \quad (\text{Put } h_2 = mh_1),\end{aligned}$$

Question 3(i) Contd...

• Now

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} &= \lim_{h \rightarrow 0} \frac{\sqrt{|h_1 h_2|}}{h_1 + ih_2} \\ &= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m| h_1^2}}{(1 + im)h_1} \quad (\text{Put } h_2 = mh_1), \\ &= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m|}}{(1 + im)}\end{aligned}$$

Question 3(i) Contd...

• Now

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} &= \lim_{h \rightarrow 0} \frac{\sqrt{|h_1 h_2|}}{h_1 + ih_2} \\&= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m| h_1^2}}{(1 + im)h_1} \quad (\text{Put } h_2 = mh_1), \\&= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m|}}{(1 + im)} \\&= \frac{\sqrt{|m|}}{1 + im}.\end{aligned}$$

Question 3(i) Contd...

- Now

$$\begin{aligned}\lim_{h \rightarrow 0} \frac{f(0+h) - f(0)}{h} &= \lim_{h \rightarrow 0} \frac{\sqrt{|h_1 h_2|}}{h_1 + ih_2} \\&= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m| h_1^2}}{(1 + im)h_1} \quad (\text{Put } h_2 = mh_1), \\&= \lim_{h_1 \rightarrow 0} \frac{\sqrt{|m|}}{(1 + im)} \\&= \frac{\sqrt{|m|}}{1 + im}.\end{aligned}$$

- So f is not differentiable at $z = 0$, and hence f is not analytic at $z = 0$.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.
- Then $u_x = y^2$, $v_x = 2xy$, $u_y = 2yx$, $v_y = x^2$.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.
- Then $u_x = y^2$, $v_x = 2xy$, $u_y = 2yx$, $v_y = x^2$. Clearly partial derivative exist and continuous at origin and satisfy Cauchy-Riemann equations. Hence f is differentiable at $z = 0$.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.
- Then $u_x = y^2$, $v_x = 2xy$, $u_y = 2yx$, $v_y = x^2$. Clearly partial derivative exist and continuous at origin and satisfy Cauchy-Riemann equations. Hence f is differentiable at $z = 0$.
- $u_x(a, b) = v_y(a, b)$ if $a = \pm b$ and $u_y(a, b) = -v_x(a, b)$ if $a = 0$ or $b = 0$.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.
- Then $u_x = y^2$, $v_x = 2xy$, $u_y = 2yx$, $v_y = x^2$. Clearly partial derivative exist and continuous at origin and satisfy Cauchy-Riemann equations. Hence f is differentiable at $z = 0$.
- $u_x(a, b) = v_y(a, b)$ if $a = \pm b$ and $u_y(a, b) = -v_x(a, b)$ if $a = 0$ or $b = 0$. Now combine both the case, the Cauchy-Riemann equations are satisfied at origin only.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.
- Then $u_x = y^2$, $v_x = 2xy$, $u_y = 2yx$, $v_y = x^2$. Clearly partial derivative exist and continuous at origin and satisfy Cauchy-Riemann equations. Hence f is differentiable at $z = 0$.
- $u_x(a, b) = v_y(a, b)$ if $a = \pm b$ and $u_y(a, b) = -v_x(a, b)$ if $a = 0$ or $b = 0$. Now combine both the case, the Cauchy-Riemann equations are satisfied at origin only.
- So f can not be differentiable except $z = 0$, hence f is not analytic at $z = 0$.

Question 3(ii)

- $f(z) = xy^2 + iyx^2$, here $u(x, y) = xy^2$, $v(x, y) = yx^2$.
- Then $u_x = y^2$, $v_x = 2xy$, $u_y = 2yx$, $v_y = x^2$. Clearly partial derivative exist and continuous at origin and satisfy Cauchy-Riemann equations. Hence f is differentiable at $z = 0$.
- $u_x(a, b) = v_y(a, b)$ if $a = \pm b$ and $u_y(a, b) = -v_x(a, b)$ if $a = 0$ or $b = 0$. Now combine both the case, the Cauchy-Riemann equations are satisfied at origin only.
- So f can not be differentiable except $z = 0$, hence f is not analytic at $z = 0$.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.
- To prove the required result, it is sufficient to show that

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.
- To prove the required result, it is sufficient to show that whenever f is differentiable then its derivative must be zero.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.
- To prove the required result, it is sufficient to show that whenever f is differentiable then it's derivative must be zero.
- Since f is real valued, thus $f(x, y) = u(x, y) + iv(x, y)$ where $v(x, y) \equiv 0$.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.
- To prove the required result, it is sufficient to show that whenever f is differentiable then its derivative must be zero.
- Since f is real valued, thus $f(x, y) = u(x, y) + iv(x, y)$ where $v(x, y) \equiv 0$.
- Since f is differentiable, it must satisfy CR-equations.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.
- To prove the required result, it is sufficient to show that whenever f is differentiable then its derivative must be zero.
- Since f is real valued, thus $f(x, y) = u(x, y) + iv(x, y)$ where $v(x, y) \equiv 0$.
- Since f is differentiable, it must satisfy CR-equations.
- Then f must be constant, which gives $f'(z_0) = 0$.

Question 4

Question 4

Show that the derivative of a real valued function $f(z)$ of a complex variable z , at any point, is either zero or it does not exist.

Proof

- Let $f(z)$ be a real valued function of complex variable.
- To prove the required result, it is sufficient to show that whenever f is differentiable then its derivative must be zero.
- Since f is real valued, thus $f(x, y) = u(x, y) + iv(x, y)$ where $v(x, y) \equiv 0$.
- Since f is differentiable, it must satisfy CR-equations.
- Then f must be constant, which gives $f'(z_0) = 0$.

Question 5

Question 5

Show that the function

$$f(z) = \begin{cases} \frac{z^5}{|z|^4} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

is continuous at $z = 0$, first order partial derivatives of its real and imaginary part exist at $z = 0$, but $f(z)$ is not differentiable at $z = 0$.

Solution: Differentiability at $z = 0$

Question 5

Question 5

Show that the function

$$f(z) = \begin{cases} \frac{z^5}{|z|^4} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

is continuous at $z = 0$, first order partial derivatives of its real and imaginary part exist at $z = 0$, but $f(z)$ is not differentiable at $z = 0$.

Solution: Differentiability at $z = 0$

- Consider

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0}$$

Question 5

Question 5

Show that the function

$$f(z) = \begin{cases} \frac{z^5}{|z|^4} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

is continuous at $z = 0$, first order partial derivatives of its real and imaginary part exist at $z = 0$, but $f(z)$ is not differentiable at $z = 0$.

Solution: Differentiability at $z = 0$

- Consider

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{r \rightarrow 0} \frac{r^4 e^{i4\theta}}{r^4}$$

Question 5

Question 5

Show that the function

$$f(z) = \begin{cases} \frac{z^5}{|z|^4} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

is continuous at $z = 0$, first order partial derivatives of its real and imaginary part exist at $z = 0$, but $f(z)$ is not differentiable at $z = 0$.

Solution: Differentiability at $z = 0$

- Consider

$$\begin{aligned} \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} &= \lim_{r \rightarrow 0} \frac{r^4 e^{i4\theta}}{r^4} \\ &= \lim_{r \rightarrow 0} e^{i4\theta} \end{aligned}$$

Question 5

Question 5

Show that the function

$$f(z) = \begin{cases} \frac{z^5}{|z|^4} & \text{if } z \neq 0 \\ 0 & \text{if } z = 0 \end{cases}$$

is continuous at $z = 0$, first order partial derivatives of its real and imaginary part exist at $z = 0$, but $f(z)$ is not differentiable at $z = 0$.

Solution: Differentiability at $z = 0$

- Consider

$$\begin{aligned} \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} &= \lim_{r \rightarrow 0} \frac{r^4 e^{i4\theta}}{r^4} \\ &= \lim_{r \rightarrow 0} e^{i4\theta} \end{aligned}$$

Question 6 Contd...

- When $\theta = 0$ then limit will be 1,

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Continuity at $z = 0$

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Continuity at $z = 0$

- Consider

$$\lim_{z \rightarrow 0} f(z) - f(0)$$

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Continuity at $z = 0$

- Consider

$$\lim_{z \rightarrow 0} f(z) - f(0) = \lim_{z \rightarrow 0} \frac{z^5}{|z|^4}$$

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Continuity at $z = 0$

- Consider

$$\begin{aligned}\lim_{z \rightarrow 0} f(z) - f(0) &= \lim_{z \rightarrow 0} \frac{z^5}{|z|^4} \\ &= \lim_{r \rightarrow 0} r e^{i5\theta}\end{aligned}$$

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Continuity at $z = 0$

- Consider

$$\begin{aligned}\lim_{z \rightarrow 0} f(z) - f(0) &= \lim_{z \rightarrow 0} \frac{z^5}{|z|^4} \\ &= \lim_{r \rightarrow 0} r e^{i5\theta} \\ &= 0,\end{aligned}$$

for all values of θ .

Question 6 Contd...

- When $\theta = 0$ then limit will be 1, and when $\theta = \frac{\pi}{4}$ then limit will be -1 .
- Hence $f(z)$ is not differentiable at $z = 0$.

Continuity at $z = 0$

- Consider

$$\begin{aligned}\lim_{z \rightarrow 0} f(z) - f(0) &= \lim_{z \rightarrow 0} \frac{z^5}{|z|^4} \\ &= \lim_{r \rightarrow 0} r e^{i5\theta} \\ &= 0,\end{aligned}$$

for all values of θ .

- Hence f is continuous at $z = 0$.

Question 6

Partial derivatives of real and imaginary part



$$f(z) = \frac{z^5}{|z|^4}$$

Question 6

Partial derivatives of real and imaginary part



$$\begin{aligned} f(z) &= \frac{z^5}{|z|^4} \\ &= \frac{(x + iy)^5}{(x^2 + y^2)^2} \end{aligned}$$

Question 6

Partial derivatives of real and imaginary part



$$\begin{aligned} f(z) &= \frac{z^5}{|z|^4} \\ &= \frac{(x + iy)^5}{(x^2 + y^2)^2} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2 + i(y^5 + x^4y - 10x^2y^3)}{x^4 + 2x^2y^2 + y^4} \end{aligned}$$

Question 6

Partial derivatives of real and imaginary part



$$\begin{aligned} f(z) &= \frac{z^5}{|z|^4} \\ &= \frac{(x + iy)^5}{(x^2 + y^2)^2} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2 + i(y^5 + x^4y - 10x^2y^3)}{x^4 + 2x^2y^2 + y^4} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4} + i \frac{y^5 + x^4y - 10x^2y^3}{x^4 + 2x^2y^2 + y^4} \end{aligned}$$

Question 6

Partial derivatives of real and imaginary part



$$\begin{aligned}f(z) &= \frac{z^5}{|z|^4} \\&= \frac{(x + iy)^5}{(x^2 + y^2)^2} \\&= \frac{x^5 + 5xy^4 - 10x^3y^2 + i(y^5 + x^4y - 10x^2y^3)}{x^4 + 2x^2y^2 + y^4} \\&= \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4} + i \frac{y^5 + x^4y - 10x^2y^3}{x^4 + 2x^2y^2 + y^4} \\&= u(x, y) + iv(x, y)\end{aligned}$$

$$\text{where } u(x, y) = \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4}$$

Question 6

Partial derivatives of real and imaginary part



$$\begin{aligned} f(z) &= \frac{z^5}{|z|^4} \\ &= \frac{(x + iy)^5}{(x^2 + y^2)^2} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2 + i(y^5 + x^4y - 10x^2y^3)}{x^4 + 2x^2y^2 + y^4} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4} + i \frac{y^5 + x^4y - 10x^2y^3}{x^4 + 2x^2y^2 + y^4} \\ &= u(x, y) + iv(x, y) \end{aligned}$$

$$\text{where } u(x, y) = \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4} \text{ and } v(x, y) = \frac{y^5 + x^4y - 10x^2y^3}{x^4 + 2x^2y^2 + y^4}.$$

Question 6

Partial derivatives of real and imaginary part



$$\begin{aligned} f(z) &= \frac{z^5}{|z|^4} \\ &= \frac{(x + iy)^5}{(x^2 + y^2)^2} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2 + i(y^5 + x^4y - 10x^2y^3)}{x^4 + 2x^2y^2 + y^4} \\ &= \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4} + i \frac{y^5 + x^4y - 10x^2y^3}{x^4 + 2x^2y^2 + y^4} \\ &= u(x, y) + iv(x, y) \end{aligned}$$

$$\text{where } u(x, y) = \frac{x^5 + 5xy^4 - 10x^3y^2}{x^4 + 2x^2y^2 + y^4} \text{ and } v(x, y) = \frac{y^5 + x^4y - 10x^2y^3}{x^4 + 2x^2y^2 + y^4}.$$

Question 6 Contd...

- $u_x(0, 0)$

Question 6 Contd...

- $u_x(0, 0)$

$$\begin{aligned}u_x(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} \\&= \lim_{h \rightarrow 0} \frac{h^5}{h^5} \\&= 1\end{aligned}$$

Question 6 Contd...

- $u_x(0, 0)$

$$\begin{aligned}u_x(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} \\&= \lim_{h \rightarrow 0} \frac{h^5}{h^5} \\&= 1\end{aligned}$$

- $u_y(0, 0)$

Question 6 Contd...

- $u_x(0, 0)$

$$\begin{aligned}u_x(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} \\&= \lim_{h \rightarrow 0} \frac{h^5}{h^5} \\&= 1\end{aligned}$$

- $u_y(0, 0)$

$$\begin{aligned}u_y(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} \\&= 0\end{aligned}$$

Question 6 Contd...

- $u_x(0, 0)$

$$\begin{aligned}u_x(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0 + h, 0) - u(0, 0)}{h} \\&= \lim_{h \rightarrow 0} \frac{h^5}{h^5} \\&= 1\end{aligned}$$

- $u_y(0, 0)$

$$\begin{aligned}u_y(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} \\&= 0\end{aligned}$$

Question 6 Contd...

- $v_x(0,0)$

Question 6 Contd...

- $v_x(0, 0)$

$$\begin{aligned} v_x(0, 0) &= \lim_{h \rightarrow 0} \frac{v(0 + h, 0) - v(0, 0)}{h} \\ &= 0 \end{aligned}$$

Question 6 Contd...

- $v_x(0, 0)$

$$\begin{aligned} v_x(0, 0) &= \lim_{h \rightarrow 0} \frac{v(0 + h, 0) - v(0, 0)}{h} \\ &= 0 \end{aligned}$$

- $u_y(0, 0)$

Question 6 Contd...

- $v_x(0, 0)$

$$\begin{aligned}v_x(0, 0) &= \lim_{h \rightarrow 0} \frac{v(0 + h, 0) - v(0, 0)}{h} \\&= 0\end{aligned}$$

- $u_y(0, 0)$

$$\begin{aligned}v_y(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} \\&= \lim_{h \rightarrow 0} \frac{h^5}{h^5} \\&= 1\end{aligned}$$

Question 6 Contd...

- $v_x(0, 0)$

$$\begin{aligned}v_x(0, 0) &= \lim_{h \rightarrow 0} \frac{v(0 + h, 0) - v(0, 0)}{h} \\&= 0\end{aligned}$$

- $u_y(0, 0)$

$$\begin{aligned}v_y(0, 0) &= \lim_{h \rightarrow 0} \frac{u(0, 0 + h) - u(0, 0)}{h} \\&= \lim_{h \rightarrow 0} \frac{h^5}{h^5} \\&= 1\end{aligned}$$

Question 6

Question 6

Is there an analytic function $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$, defined on some open subset of \mathbb{C} with $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$? If so, find all such $f(z)$.

Solution

- We have $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$, this gives $u_{xx} = 6x - 4$ and $u_{yy} = -6x + 4$.

Question 6

Question 6

Is there an analytic function $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$, defined on some open subset of \mathbb{C} with $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$? If so, find all such $f(z)$.

Solution

- We have $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$, this gives $u_{xx} = 6x - 4$ and $u_{yy} = -6x + 4$.
- $u_{xx} + u_{yy} = 0$, this implies u is harmonic.

Question 6

Question 6

Is there an analytic function $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$, defined on some open subset of \mathbb{C} with $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$? If so, find all such $f(z)$.

Solution

- We have $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$, this gives $u_{xx} = 6x - 4$ and $u_{yy} = -6x + 4$.
- $u_{xx} + u_{yy} = 0$, this implies u is harmonic.
- Now we will find out $v(x, y)$ of an analytic function whose real part u .

Question 6

Question 6

Is there an analytic function $f(z) = u(x, y) + iv(x, y)$, where $z = x + iy$, defined on some open subset of \mathbb{C} with $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$? If so, find all such $f(z)$.

Solution

- We have $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$, this gives $u_{xx} = 6x - 4$ and $u_{yy} = -6x + 4$.
- $u_{xx} + u_{yy} = 0$, this implies u is harmonic.
- Now we will find out $v(x, y)$ of an analytic function whose real part u .
- Since f is analytic, it must satisfy CR-equations, i.e $u_x = v_y$ and $u_y = -v_x$.
- Consider $v_y = u_x = 3x^2 - 3y^2 - 4x$, on integrating both sides w.r.t y we get.

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x).$

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x)$.
- Now using $u_y = -v_x$, we have

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x)$.
- Now using $u_y = -v_x$, we have $-6xy + 4y = -6xy + 4x + g'(x)$, this implies $g' = 0 \implies$

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x)$.
- Now using $u_y = -v_x$, we have $-6xy + 4y = -6xy + 4x + g'(x)$, this implies $g' = 0 \implies g$ is constant.
- Hence we have $f(z) = u(x, y) + iv(x, y)$

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x)$.
- Now using $u_y = -v_x$, we have $-6xy + 4y = -6xy + 4x + g'(x)$, this implies $g' = 0 \implies g$ is constant.
- Hence we have $f(z) = u(x, y) + iv(x, y)$ where $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$ and

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x)$.
- Now using $u_y = -v_x$, we have $-6xy + 4y = -6xy + 4x + g'(x)$, this implies $g' = 0 \implies g$ is constant.
- Hence we have $f(z) = u(x, y) + iv(x, y)$ where $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$ and $v(x, y) = 3x^2y - y^3 - 4xy + C$.

Question 6 Contd...

- $v = 3x^2y - y^3 - 4xy + g(x)$.
- Now using $u_y = -v_x$, we have $-6xy + 4y = -6xy + 4x + g'(x)$, this implies $g' = 0 \implies g$ is constant.
- Hence we have $f(z) = u(x, y) + iv(x, y)$ where $u = x^3 - 3xy^2 - 2x^2 + 2y^2 + 1$ and $v(x, y) = 3x^2y - y^3 - 4xy + C$.

Question 7

Question 7

Show that the function $\log(z - i)$ is analytic everywhere except on the half line $y = 1, x \leq 0$.

Question 7

Question 7

Show that the function $\log(z - i)$ is analytic everywhere except on the half line $y = 1, x \leq 0$.

Solution: In this question we consider the standard branch. Let $z = x + iy$

- We know $\log(z) = \ln(|z|) + i\arg(z)$ is analytic everywhere except when $(x < 0 \text{ and } y = 0)$ or $(z = 0)$

Question 7

Question 7

Show that the function $\log(z - i)$ is analytic everywhere except on the half line $y = 1, x \leq 0$.

Solution: In this question we consider the standard branch. Let $z = x + iy$

- We know $\log(z) = \ln(|z|) + i\arg(z)$ is analytic everywhere except when $(x < 0 \text{ and } y = 0)$ or $(z = 0)$
- Thus $f(x) = \log(z - i)$ is analytic everywhere except when $(x < 0 \text{ and } y - 1 = 0)$ or $(z = i)$

Question 7

Question 7

Show that the function $\log(z - i)$ is analytic everywhere except on the half line $y = 1, x \leq 0$.

Solution: In this question we consider the standard branch. Let $z = x + iy$

- We know $\log(z) = \ln(|z|) + i\arg(z)$ is analytic everywhere except when $(x < 0 \text{ and } y = 0)$ or $(z = 0)$
- Thus $f(x) = \log(z - i)$ is analytic everywhere except when $(x < 0 \text{ and } y - 1 = 0)$ or $(z = i)$
- Thus $f(z)$ is analytic everywhere except when $(x < 0 \text{ and } y = 1)$ or $(x = 0 \text{ and } y = 1)$

Question 7

Question 7

Show that the function $\log(z - i)$ is analytic everywhere except on the half line $y = 1, x \leq 0$.

Solution: In this question we consider the standard branch. Let $z = x + iy$

- We know $\log(z) = \ln(|z|) + i\arg(z)$ is analytic everywhere except when $(x < 0 \text{ and } y = 0)$ or $(z = 0)$
- Thus $f(x) = \log(z - i)$ is analytic everywhere except when $(x < 0 \text{ and } y - 1 = 0)$ or $(z = i)$
- Thus $f(z)$ is analytic everywhere except when $(x < 0 \text{ and } y = 1)$ or $(x = 0 \text{ and } y = 1)$
- Thus $f(z)$ is analytic everywhere except the half line $y = 1, x \leq 0$

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Solution: Let $z = x + iy$

- Therefore $\sin(z) = \sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy)$

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Solution: Let $z = x + iy$

- Therefore $\sin(z) = \sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy)$
- Now $\cos(iy) = \frac{e^{i*iy} + e^{-i*iy}}{2} = \frac{e^{-y} + e^y}{2}$

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Solution: Let $z = x + iy$

- Therefore $\sin(z) = \sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy)$
- Now $\cos(iy) = \frac{e^{i*iy} + e^{-i*iy}}{2} = \frac{e^{-y} + e^y}{2}$
- Now $\sin(iy) = \frac{e^{i*iy} - e^{-i*iy}}{2i} = \frac{e^{-y} - e^y}{2i} = i \frac{e^y - e^{-y}}{2}$

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Solution: Let $z = x + iy$

- Therefore $\sin(z) = \sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy)$
- Now $\cos(iy) = \frac{e^{i*iy} + e^{-i*iy}}{2} = \frac{e^{-y} + e^y}{2}$
- Now $\sin(iy) = \frac{e^{i*iy} - e^{-i*iy}}{2i} = \frac{e^{-y} - e^y}{2i} = i \frac{e^y - e^{-y}}{2}$
- Now $2 = \sin(x) \left(\frac{e^{-y} + e^y}{2} \right) + i \cos(x) \left(\frac{e^y - e^{-y}}{2} \right)$

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Solution: Let $z = x + iy$

- Therefore $\sin(z) = \sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy)$
- Now $\cos(iy) = \frac{e^{i*iy} + e^{-i*iy}}{2} = \frac{e^{-y} + e^y}{2}$
- Now $\sin(iy) = \frac{e^{i*iy} - e^{-i*iy}}{2i} = \frac{e^{-y} - e^y}{2i} = i \frac{e^y - e^{-y}}{2}$
- Now $2 = \sin(x) \left(\frac{e^{-y} + e^y}{2} \right) + i \cos(x) \left(\frac{e^y - e^{-y}}{2} \right)$
- Now either $\cos(x) = 0$ or $e^y = e^{-y} = 1$. If the second case is true then $\sin(x) = 2$ which is impossible. Thus $\cos(x) = 0$. Thus $x = (2n + 1) \frac{\pi}{2}$

Question 8

Question 8

Solve the equation $\sin z = 2$ for z by equating real and imaginary parts in that equation

Solution: Let $z = x + iy$

- Therefore $\sin(z) = \sin(x + iy) = \sin(x) \cos(iy) + \cos(x) \sin(iy)$
- Now $\cos(iy) = \frac{e^{i*iy} + e^{-i*iy}}{2} = \frac{e^{-y} + e^y}{2}$
- Now $\sin(iy) = \frac{e^{i*iy} - e^{-i*iy}}{2i} = \frac{e^{-y} - e^y}{2i} = i \frac{e^y - e^{-y}}{2}$
- Now $2 = \sin(x) \left(\frac{e^{-y} + e^y}{2} \right) + i \cos(x) \left(\frac{e^y - e^{-y}}{2} \right)$
- Now either $\cos(x) = 0$ or $e^y = e^{-y} = 1$. If the second case is true then $\sin(x) = 2$ which is impossible. Thus $\cos(x) = 0$. Thus $x = (2n + 1)\frac{\pi}{2}$
- Since $\sin(x) \left(\frac{e^{-y} + e^y}{2} \right) = 2$, we have $\sin(x)$ cannot be -1. Thus $x = (4n + 1)\frac{\pi}{2}$

Question 8 Contd...

- Now since $e^y + e^{-y} = 4$, solving quadratic equation we have $y = \pm \ln(2 + \sqrt{3})$.
- Thus $z = (4n + 1)\frac{\pi}{2} \pm i \ln(2 + \sqrt{3})$

Question 9

Question 9

Show that neither $\sin \bar{z}$ nor $\cos \bar{z}$ is an analytic function of z anywhere.

Question 9

Question 9

Show that neither $\sin \bar{z}$ nor $\cos \bar{z}$ is an analytic function of z anywhere.

Solution: I will only prove this for $\sin(\bar{z})$ and the proof for $\cos(\bar{z})$ which is very similar is left as an exercise. Let $z = x + iy$

$$\bullet \sin(\bar{z}) = \frac{e^{x-iy} - e^{-(x-iy)}}{2i}$$

Question 9

Question 9

Show that neither $\sin \bar{z}$ nor $\cos \bar{z}$ is an analytic function of z anywhere.

Solution: I will only prove this for $\sin(\bar{z})$ and the proof for $\cos(\bar{z})$ which is very similar is left as an exercise. Let $z = x + iy$

- $\sin(\bar{z}) = \frac{e^{x-iy} - e^{-(x-iy)}}{2i}$
- Thus if $f(z) = \sin(\bar{z}) = u(x, y) + iv(x, y)$,

Question 9

Question 9

Show that neither $\sin \bar{z}$ nor $\cos \bar{z}$ is an analytic function of z anywhere.

Solution: I will only prove this for $\sin(\bar{z})$ and the proof for $\cos(\bar{z})$ which is very similar is left as an exercise. Let $z = x + iy$

- $\sin(\bar{z}) = \frac{e^{x-iy} - e^{-(x-iy)}}{2i}$
- Thus if $f(z) = \sin(\bar{z}) = u(x, y) + iv(x, y)$, then $u(x, y) = \sin(x) \cosh(y)$ and $v(x, y) = -\cos(x) \sinh(y)$ (Why?)

Question 9

Question 9

Show that neither $\sin \bar{z}$ nor $\cos \bar{z}$ is an analytic function of z anywhere.

Solution: I will only prove this for $\sin(\bar{z})$ and the proof for $\cos(\bar{z})$ which is very similar is left as an exercise. Let $z = x + iy$

- $\sin(\bar{z}) = \frac{e^{x-iy} - e^{-(x-iy)}}{2i}$
- Thus if $f(z) = \sin(\bar{z}) = u(x, y) + iv(x, y)$, then $u(x, y) = \sin(x) \cosh(y)$ and $v(x, y) = -\cos(x) \sinh(y)$ (Why?)
- Now if the Cauchy Riemann equations are to hold then we can see that it will hold only on isolated points namely $z = (2n + 1)\frac{\pi}{2}$

Question 9

Question 9

Show that neither $\sin \bar{z}$ nor $\cos \bar{z}$ is an analytic function of z anywhere.

Solution: I will only prove this for $\sin(\bar{z})$ and the proof for $\cos(\bar{z})$ which is very similar is left as an exercise. Let $z = x + iy$

- $\sin(\bar{z}) = \frac{e^{x-iy} - e^{-(x-iy)}}{2i}$
- Thus if $f(z) = \sin(\bar{z}) = u(x, y) + iv(x, y)$, then $u(x, y) = \sin(x) \cosh(y)$ and $v(x, y) = -\cos(x) \sinh(y)$ (Why?)
- Now if the Cauchy Riemann equations are to hold then we can see that it will hold only on isolated points namely $z = (2n+1)\frac{\pi}{2}$
- Thus $f(z)$ is not analytic anywhere since cauchy riemann equations do not hold in any open set.