

# MTL122 - Real and complex analysis

## Assignment-4



Department of Mathematics  
Indian Institute of Technology Delhi

# Question 1

## Question 1

Let  $(E, d)$  be a metric space, and let  $f, g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  be bounded and uniformly continuous functions. Show that the product  $f \cdot g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  is bounded and uniformly continuous.

# Question 1

## Question 1

Let  $(E, d)$  be a metric space, and let  $f, g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  be bounded and uniformly continuous functions. Show that the product  $f \cdot g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  is bounded and uniformly continuous.

### Proof

- Since  $f$  and  $g$  are bounded functions, then there exist  $M_1 > 0$  and  $M_2 > 0$  such that  $|f(s)| \leq M_1, |g(s)| \leq M_2$ , for all  $s \in E$ .

# Question 1

## Question 1

Let  $(E, d)$  be a metric space, and let  $f, g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  be bounded and uniformly continuous functions. Show that the product  $f \cdot g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  is bounded and uniformly continuous.

### Proof

- Since  $f$  and  $g$  are bounded functions, then there exist  $M_1 > 0$  and  $M_2 > 0$  such that  $|f(s)| \leq M_1, |g(s)| \leq M_2$ , for all  $s \in E$ .
- As  $f$  is uniformly continuous, for each  $\epsilon > 0$ ,  $\exists \delta_1 > 0$  such that

$$d(s, t) < \delta_1 \Rightarrow |f(s) - f(t)| < \frac{\epsilon}{2M_2},$$

for all  $s, t \in E$ .

# Question 1

## Question 1

Let  $(E, d)$  be a metric space, and let  $f, g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  be bounded and uniformly continuous functions. Show that the product  $f \cdot g : (E, d) \rightarrow (\mathbb{R}, \text{usual metric})$  is bounded and uniformly continuous.

### Proof

- Since  $f$  and  $g$  are bounded functions, then there exist  $M_1 > 0$  and  $M_2 > 0$  such that  $|f(s)| \leq M_1, |g(s)| \leq M_2$ , for all  $s \in E$ .
- As  $f$  is uniformly continuous, for each  $\epsilon > 0$ ,  $\exists \delta_1 > 0$  such that

$$d(s, t) < \delta_1 \Rightarrow |f(s) - f(t)| < \frac{\epsilon}{2M_2},$$

for all  $s, t \in E$ .

- As  $g$  is uniformly continuous, for each  $\epsilon > 0$ ,  $\exists \delta_2 > 0$  such that

$$d(s, t) < \delta_2 \Rightarrow |g(s) - g(t)| < \frac{\epsilon}{2M_1},$$

for all  $s, t \in E$ .

## Question 1 Contd.

- Now for all  $s, t \in E$  and  $d(s, t) < \delta = \min(\delta_1, \delta_2)$ , then

$$|f \cdot g(s) - f \cdot g(t)| = |f(s)g(s) - f(t)g(s) + f(t)g(s) - f(t)g(t)|$$

## Question 1 Contd.

- Now for all  $s, t \in E$  and  $d(s, t) < \delta = \min(\delta_1, \delta_2)$ , then

$$\begin{aligned} |f \cdot g(s) - f \cdot g(t)| &= |f(s)g(s) - f(t)g(s) + f(t)g(s) - f(t)g(t)| \\ &\leq |g(s)||f(s) - f(t)| + |f(t)||g(s) - g(t)| \end{aligned}$$

## Question 1 Contd.

- Now for all  $s, t \in E$  and  $d(s, t) < \delta = \min(\delta_1, \delta_2)$ , then

$$\begin{aligned} |f \cdot g(s) - f \cdot g(t)| &= |f(s)g(s) - f(t)g(s) + f(t)g(s) - f(t)g(t)| \\ &\leq |g(s)||f(s) - f(t)| + |f(t)||g(s) - g(t)| \\ &< M_2 \times \frac{\epsilon}{2M_2} + M_1 \times \frac{\epsilon}{2M_1} \end{aligned}$$



## Question 1 Contd.

- Now for all  $s, t \in E$  and  $d(s, t) < \delta = \min(\delta_1, \delta_2)$ , then

$$\begin{aligned} |f \cdot g(s) - f \cdot g(t)| &= |f(s)g(s) - f(t)g(s) + f(t)g(s) - f(t)g(t)| \\ &\leq |g(s)||f(s) - f(t)| + |f(t)||g(s) - g(t)| \\ &< M_2 \times \frac{\epsilon}{2M_2} + M_1 \times \frac{\epsilon}{2M_1} \\ &= \epsilon. \end{aligned}$$

## Question 1 Contd.

- Now for all  $s, t \in E$  and  $d(s, t) < \delta = \min(\delta_1, \delta_2)$ , then

$$\begin{aligned} |f \cdot g(s) - f \cdot g(t)| &= |f(s)g(s) - f(t)g(s) + f(t)g(s) - f(t)g(t)| \\ &\leq |g(s)||f(s) - f(t)| + |f(t)||g(s) - g(t)| \\ &< M_2 \times \frac{\epsilon}{2M_2} + M_1 \times \frac{\epsilon}{2M_1} \\ &= \epsilon. \end{aligned}$$

- Hence  $f \cdot g$  is uniformly continuous.

## Question 2

### Question 2

Equip the interval  $(0, 1) \subset \mathbb{R}$  with the usual metric.

- Show that if  $f : (0, 1) \rightarrow \mathbb{R}$  is uniformly continuous, then it is bounded.
- Give an example of a function  $f : (0, 1) \rightarrow \mathbb{R}$  that is continuous but unbounded.

## Question 2

### Question 2

Equip the interval  $(0, 1) \subset \mathbb{R}$  with the usual metric.

- Show that if  $f : (0, 1) \rightarrow \mathbb{R}$  is uniformly continuous, then it is bounded.
- Give an example of a function  $f : (0, 1) \rightarrow \mathbb{R}$  that is continuous but unbounded.

**Solution:** (a)

- $\forall \epsilon > 0, \exists \delta_\epsilon > 0$  such that  $|x - y| < \delta_\epsilon \implies |f(x) - f(y)| < \epsilon$ .

## Question 2

### Question 2

Equip the interval  $(0, 1) \subset \mathbb{R}$  with the usual metric.

- Show that if  $f : (0, 1) \rightarrow \mathbb{R}$  is uniformly continuous, then it is bounded.
- Give an example of a function  $f : (0, 1) \rightarrow \mathbb{R}$  that is continuous but unbounded.

**Solution:** (a)

- $\forall \epsilon > 0, \exists \delta_\epsilon > 0$  such that  $|x - y| < \delta_\epsilon \implies |f(x) - f(y)| < \epsilon$ .
- Corresponding to  $\epsilon = 1$ , we have  $\delta_1$  such that whenever  $|x - y| < \delta_1$  we have  $|f(x) - f(y)| < 1$ .

## Question 2

### Question 2

Equip the interval  $(0, 1) \subset \mathbb{R}$  with the usual metric.

- Show that if  $f : (0, 1) \rightarrow \mathbb{R}$  is uniformly continuous, then it is bounded.
- Give an example of a function  $f : (0, 1) \rightarrow \mathbb{R}$  that is continuous but unbounded.

**Solution:** (a)

- $\forall \epsilon > 0, \exists \delta_\epsilon > 0$  such that  $|x - y| < \delta_\epsilon \implies |f(x) - f(y)| < \epsilon$ .
- Corresponding to  $\epsilon = 1$ , we have  $\delta_1$  such that whenever  $|x - y| < \delta_1$  we have  $|f(x) - f(y)| < 1$ .
- $\exists x_1, x_2, \dots, x_n$  such that  $(0, 1) \subset \bigcup_{i=1}^n N_{\delta_\epsilon}(x_i)$  and  $x_i \in (0, 1)$  for all  $i = 1, 2, \dots, n$ . (Why?)

## Question 2

### Question 2

Equip the interval  $(0, 1) \subset \mathbb{R}$  with the usual metric.

- Show that if  $f : (0, 1) \rightarrow \mathbb{R}$  is uniformly continuous, then it is bounded.
- Give an example of a function  $f : (0, 1) \rightarrow \mathbb{R}$  that is continuous but unbounded.

**Solution:** (a)

- $\forall \epsilon > 0, \exists \delta_\epsilon > 0$  such that  $|x - y| < \delta_\epsilon \implies |f(x) - f(y)| < \epsilon$ .
- Corresponding to  $\epsilon = 1$ , we have  $\delta_1$  such that whenever  $|x - y| < \delta_1$  we have  $|f(x) - f(y)| < 1$ .
- $\exists x_1, x_2, \dots, x_n$  such that  $(0, 1) \subset \bigcup_{i=1}^n N_{\delta_\epsilon}(x_i)$  and  $x_i \in (0, 1)$  for all  $i = 1, 2, \dots, n$ . (Why?)

## Question 2 Contd...

- $\forall x \in (0, 1), \exists x_j$  such that  $x \in N_{\delta_\epsilon}(x_j)$ .



## Question 2 Contd...

- $\forall x \in (0, 1), \exists x_j$  such that  $x \in N_{\delta_\epsilon}(x_j)$ .
- Thus for  $M = \max\{|f(x_1)|, |f(x_2)|, \dots, |f(x_n)|\}$ , we have

$$|f(x)| = |f(x) - f(x_j) + f(x_j)| \leq 1 + |f(x_j)| \leq 1 + M$$

for all  $x \in (0, 1)$

## Question 2 Contd...

- $\forall x \in (0, 1), \exists x_j$  such that  $x \in N_{\delta_\epsilon}(x_j)$ .
- Thus for  $M = \max\{|f(x_1)|, |f(x_2)|, \dots, |f(x_n)|\}$ , we have

$$|f(x)| = |f(x) - f(x_j) + f(x_j)| \leq 1 + |f(x_j)| \leq 1 + M$$

for all  $x \in (0, 1)$

- Hence we have that  $f$  is bounded.

## Question 2 Contd...

- $\forall x \in (0, 1), \exists x_j$  such that  $x \in N_{\delta_\epsilon}(x_j)$ .
- Thus for  $M = \max\{|f(x_1)|, |f(x_2)|, \dots, |f(x_n)|\}$ , we have

$$|f(x)| = |f(x) - f(x_j) + f(x_j)| \leq 1 + |f(x_j)| \leq 1 + M$$

for all  $x \in (0, 1)$

- Hence we have that  $f$  is bounded.
- (b) The obvious answer to this question is  $f(x) = 1/x$  (Why?)

# Question3

## Question 3

Prove that a continuous function on a compact metric space  $X$  is bounded and uniformly continuous.

# Question3

## Question 3

Prove that a continuous function on a compact metric space  $X$  is bounded and uniformly continuous.

## Recall: Lebesgue Number Lemma

If the metric space  $(X, d)$  is compact and an open cover of  $X$  is given, then there exists a number  $\delta > 0$  such that every subset of  $X$  having diameter less than  $\delta$  is contained in some member of the cover.

## Proof

# Question3

## Question 3

Prove that a continuous function on a compact metric space  $X$  is bounded and uniformly continuous.

## Recall: Lebesgue Number Lemma

If the metric space  $(X, d)$  is compact and an open cover of  $X$  is given, then there exists a number  $\delta > 0$  such that every subset of  $X$  having diameter less than  $\delta$  is contained in some member of the cover.

## Proof

- Let  $f : (X, d_1) \rightarrow (Y, d_2)$  be a continuous function and  $(X, d_1)$  be a compact metric space.

## Question 3

Prove that a continuous function on a compact metric space  $X$  is bounded and uniformly continuous.

## Recall: Lebesgue Number Lemma

If the metric space  $(X, d)$  is compact and an open cover of  $X$  is given, then there exists a number  $\delta > 0$  such that every subset of  $X$  having diameter less than  $\delta$  is contained in some member of the cover.

## Proof

- Let  $f : (X, d_1) \rightarrow (Y, d_2)$  be a continuous function and  $(X, d_1)$  be a compact metric space.
- We know continuous image of a compact set is compact, hence  $f(X)$  is bounded.

## Question 3

Prove that a continuous function on a compact metric space  $X$  is bounded and uniformly continuous.

## Recall: Lebesgue Number Lemma

If the metric space  $(X, d)$  is compact and an open cover of  $X$  is given, then there exists a number  $\delta > 0$  such that every subset of  $X$  having diameter less than  $\delta$  is contained in some member of the cover.

## Proof

- Let  $f : (X, d_1) \rightarrow (Y, d_2)$  be a continuous function and  $(X, d_1)$  be a compact metric space.
- We know continuous image of a compact set is compact, hence  $f(X)$  is bounded.



## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ ,

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ .

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$  such that for every  $x \in X$  we can find a element  $B_{\delta_a}(a)$  in open cover

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$  such that for every  $x \in X$  we can find a element  $B_{\delta_a}(a)$  in open cover such that  $B_\eta(x) \subset B_{\delta_a}(a)$ .



## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$  such that for every  $x \in X$  we can find a element  $B_{\delta_a}(a)$  in open cover such that  $B_\eta(x) \subset B_{\delta_a}(a)$ .
- Thus

$$d_1(x, y) < \eta \implies$$

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$  such that for every  $x \in X$  we can find a element  $B_{\delta_a}(a)$  in open cover such that  $B_\eta(x) \subset B_{\delta_a}(a)$ .
- Thus

$$d_1(x, y) < \eta \implies d_1(x, y) < \delta_a \implies$$

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$  such that for every  $x \in X$  we can find a element  $B_{\delta_a}(a)$  in open cover such that  $B_\eta(x) \subset B_{\delta_a}(a)$ .
- Thus

$$d_1(x, y) < \eta \implies d_1(x, y) < \delta_a \implies d_2(f(x), f(y)) < \epsilon.$$

## Question 3 Contd...

- Let  $f$  be continuous at  $a \in X$ , then for each  $\epsilon > 0$ , there exist  $\delta(\epsilon, a) = \delta_a > 0$  such that

$$d_1(x, a) < \delta_a \implies d_2(f(x), f(a)) < \epsilon.$$

- Clearly  $X = \bigcup_{a \in X} B_{\delta_a}(a)$ . Thus  $\mathcal{U} := \{B_{\delta_a}(a) : a \in X\}$  is an open cover of  $X$ .
- Now by lebesgue number lemma, there exists a  $\eta > 0$  such that for every  $x \in X$  we can find a element  $B_{\delta_a}(a)$  in open cover such that  $B_\eta(x) \subset B_{\delta_a}(a)$ .
- Thus

$$d_1(x, y) < \eta \implies d_1(x, y) < \delta_a \implies d_2(f(x), f(y)) < \epsilon.$$

- Hence  $f$  is uniformly continuous.

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

**Proof**

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

**Proof**

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

- (i) False,



## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

- (i) **False, Counterexample:**

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

(i) **False, Counterexample:**  $\bigcup_{k=1}^{\infty} [-k, k] = \mathbb{R}.$

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

(i) **False, Counterexample:**  $\bigcup_{k=1}^{\infty} [-k, k] = \mathbb{R}.$

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

- (i) **False, Counterexample:**  $\bigcup_{k=1}^{\infty} [-k, k] = \mathbb{R}.$
- (ii) **False,**

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

- (i) **False, Counterexample:**  $\bigcup_{k=1}^{\infty} [-k, k] = \mathbb{R}.$
- (ii) **False, Counterexample:**

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

- (i) **False, Counterexample:**  $\bigcup_{k=1}^{\infty} [-k, k] = \mathbb{R}.$
- (ii) **False, Counterexample:**  $[-k, 0) \cup (0, k].$

## Question 4

### Question 4

Prove or give a counter example:

- (i) The union of infinitely many compact sets is compact.
- (ii) An non-empty subset of real numbers which has both largest and a smallest element is compact.

### Proof

- (i) **False, Counterexample:**  $\bigcup_{k=1}^{\infty} [-k, k] = \mathbb{R}.$
- (ii) **False, Counterexample:**  $[-k, 0) \cup (0, k].$

## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .



## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .

**Solution:** Given that  $A$  is a totally bounded set and  $B \subset A$ .

- $\forall \epsilon > 0, \exists a_1, a_2, \dots, a_N$  and  $N \in \mathbb{N}$  such that  $A \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .

## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .

**Solution:** Given that  $A$  is a totally bounded set and  $B \subset A$ .

- $\forall \epsilon > 0, \exists a_1, a_2, \dots, a_N$  and  $N \in \mathbb{N}$  such that  $A \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Since  $B \subset A$ , we have  $B \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .

## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .

**Solution:** Given that  $A$  is a totally bounded set and  $B \subset A$ .

- $\forall \epsilon > 0, \exists a_1, a_2, \dots, a_N$  and  $N \in \mathbb{N}$  such that  $A \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Since  $B \subset A$ , we have  $B \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Let  $b_i \in B_{\epsilon/2}(a_i) \cap B$  whenever  $B_{\epsilon/2}(a_i) \cap B$  is non empty.
- Let there be  $M \leq N$  many such  $b_i$ .

## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .

**Solution:** Given that  $A$  is a totally bounded set and  $B \subset A$ .

- $\forall \epsilon > 0, \exists a_1, a_2, \dots, a_N$  and  $N \in \mathbb{N}$  such that  $A \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Since  $B \subset A$ , we have  $B \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Let  $b_i \in B_{\epsilon/2}(a_i) \cap B$  whenever  $B_{\epsilon/2}(a_i) \cap B$  is non empty.
- Let there be  $M \leq N$  many such  $b_i$ .
- Suppose  $b \in B$ , and let  $b \in B_{\epsilon/2}(a_l)$  for some  $1 \leq l \leq N$ .

## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .

**Solution:** Given that  $A$  is a totally bounded set and  $B \subset A$ .

- $\forall \epsilon > 0, \exists a_1, a_2, \dots, a_N$  and  $N \in \mathbb{N}$  such that  $A \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Since  $B \subset A$ , we have  $B \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Let  $b_i \in B_{\epsilon/2}(a_i) \cap B$  whenever  $B_{\epsilon/2}(a_i) \cap B$  is non empty.
- Let there be  $M \leq N$  many such  $b_i$ .
- Suppose  $b \in B$ , and let  $b \in B_{\epsilon/2}(a_l)$  for some  $1 \leq l \leq N$ .
- Since  $B_{\epsilon/2}(a_l) \cap B$  is non-empty, we have  $d(b, b_l) \leq d(b, a_l) + d(a_l, b_l) \leq \epsilon$ .

## Question 5

### Question 5

Prove that every subset of a totally bounded subset  $A$  of a metric space  $X$  is totally bounded in  $X$ .

**Solution:** Given that  $A$  is a totally bounded set and  $B \subset A$ .

- $\forall \epsilon > 0, \exists a_1, a_2, \dots, a_N$  and  $N \in \mathbb{N}$  such that  $A \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Since  $B \subset A$ , we have  $B \subset \bigcup_{i=1}^N B_{\epsilon/2}(a_i)$ .
- Let  $b_i \in B_{\epsilon/2}(a_i) \cap B$  whenever  $B_{\epsilon/2}(a_i) \cap B$  is non empty.
- Let there be  $M \leq N$  many such  $b_i$ .
- Suppose  $b \in B$ , and let  $b \in B_{\epsilon/2}(a_l)$  for some  $1 \leq l \leq N$ .
- Since  $B_{\epsilon/2}(a_l) \cap B$  is non-empty, we have  $d(b, b_l) \leq d(b, a_l) + d(a_l, b_l) \leq \epsilon$ .
- Thus the needful is proved. (Why?)

## Question 6

### Question 6

Let  $(X, d)$  be a metric space, and let  $A, B \subset X$ . Show that, if  $A$  and  $B$  are sequentially compact, then so is  $A \cap B$ .

## Question 6

### Question 6

Let  $(X, d)$  be a metric space, and let  $A, B \subset X$ . Show that, if  $A$  and  $B$  are sequentially compact, then so is  $A \cap B$ .

**Solution:** Given that  $A$  and  $B$  are sequentially compact sets.

- Let  $(x_n)_{n>0}$  be a sequence in  $A \cap B$ . Thus  $(x_n)_{n>0} \subset A$  and  $(x_n)_{n>0} \subset B$ .



## Question 6

### Question 6

Let  $(X, d)$  be a metric space, and let  $A, B \subset X$ . Show that, if  $A$  and  $B$  are sequentially compact, then so is  $A \cap B$ .

**Solution:** Given that  $A$  and  $B$  are sequentially compact sets.

- Let  $(x_n)_{n>0}$  be a sequence in  $A \cap B$ . Thus  $(x_n)_{n>0} \subset A$  and  $(x_n)_{n>0} \subset B$ .
- Thus we have  $(x_{n_k})_{k>0} \rightarrow a \in A$ .

## Question 6

### Question 6

Let  $(X, d)$  be a metric space, and let  $A, B \subset X$ . Show that, if  $A$  and  $B$  are sequentially compact, then so is  $A \cap B$ .

**Solution:** Given that  $A$  and  $B$  are sequentially compact sets.

- Let  $(x_n)_{n>0}$  be a sequence in  $A \cap B$ . Thus  $(x_n)_{n>0} \subset A$  and  $(x_n)_{n>0} \subset B$ .
- Thus we have  $(x_{n_k})_{k>0} \rightarrow a \in A$ .
- Now since  $(x_{n_k})_{k>0} \subset B$  we have  $x_{n_{k_j}} \rightarrow b \in B$ .

## Question 6

### Question 6

Let  $(X, d)$  be a metric space, and let  $A, B \subset X$ . Show that, if  $A$  and  $B$  are sequentially compact, then so is  $A \cap B$ .

**Solution:** Given that  $A$  and  $B$  are sequentially compact sets.

- Let  $(x_n)_{n>0}$  be a sequence in  $A \cap B$ . Thus  $(x_n)_{n>0} \subset A$  and  $(x_n)_{n>0} \subset B$ .
- Thus we have  $(x_{n_k})_{k>0} \rightarrow a \in A$ .
- Now since  $(x_{n_k})_{k>0} \subset B$  we have  $x_{n_{k_j}} \rightarrow b \in B$ .
- Since  $x_{n_{k_j}} \subset A$  and sequentially compact implies closed we have  $b \in A$  and thus  $b \in A \cap B$ .

## Question 6

### Question 6

Let  $(X, d)$  be a metric space, and let  $A, B \subset X$ . Show that, if  $A$  and  $B$  are sequentially compact, then so is  $A \cap B$ .

**Solution:** Given that  $A$  and  $B$  are sequentially compact sets.

- Let  $(x_n)_{n>0}$  be a sequence in  $A \cap B$ . Thus  $(x_n)_{n>0} \subset A$  and  $(x_n)_{n>0} \subset B$ .
- Thus we have  $(x_{n_k})_{k>0} \rightarrow a \in A$ .
- Now since  $(x_{n_k})_{k>0} \subset B$  we have  $x_{n_{k_j}} \rightarrow b \in B$ .
- Since  $x_{n_{k_j}} \subset A$  and sequentially compact implies closed we have  $b \in A$  and thus  $b \in A \cap B$ .
- Thus we have proven every sequence of  $A \cap B$  has a convergent subsequence and hence proved.

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded,

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e. for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .



# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given.

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e. for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous,

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous, there exists  $\delta > 0$  such that

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e. for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous, there exists  $\delta > 0$  such that  $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$ .

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous, there exists  $\delta > 0$  such that  $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$ .
- Since  $X$  is totally bounded,

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous, there exists  $\delta > 0$  such that  $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$ .
- Since  $X$  is totally bounded, there exists a finite number of balls  $B_X(a_1, \delta), \dots, B_X(a_n, \delta)$

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous, there exists  $\delta > 0$  such that  $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$ .
- Since  $X$  is totally bounded, there exists a finite number of balls  $B_X(a_1, \delta), \dots, B_X(a_n, \delta)$  such that

$$X = \bigcup_{i=1}^n B_X(a_i, \delta).$$

# Question 7

## Question 7

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is totally bounded. Prove that  $Y$  is totally bounded as well.

## Solution

- To show that  $Y$  is totally bounded, i.e. for any  $\epsilon > 0$ ,  $Y$  is contained in the union of a finite number of balls of radius  $\epsilon$ .
- Let  $\epsilon > 0$  be given. Since  $f$  is uniformly continuous, there exists  $\delta > 0$  such that  $d_X(x, y) < \delta \implies d_Y(f(x), f(y)) < \epsilon$ .
- Since  $X$  is totally bounded, there exists a finite number of balls  $B_X(a_1, \delta), \dots, B_X(a_n, \delta)$  such that

$$X = \bigcup_{i=1}^n B_X(a_i, \delta).$$



## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution:**

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans-**

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

- Take  $X = (\mathbb{R}, d_{dis})$

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

- Take  $X = (\mathbb{R}, d_{dis})$  and  $Y = (\mathbb{R}, d_{Euc})$ .



## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

- Take  $X = (\mathbb{R}, d_{dis})$  and  $Y = (\mathbb{R}, d_{Euc})$ .
- Since  $d_{dis}$  is a bounded metric,

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

- Take  $X = (\mathbb{R}, d_{dis})$  and  $Y = (\mathbb{R}, d_{Euc})$ .
- Since  $d_{dis}$  is a bounded metric, thus  $\mathbb{R}$  is bounded.

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

- Take  $X = (\mathbb{R}, d_{dis})$  and  $Y = (\mathbb{R}, d_{Euc})$ .
- Since  $d_{dis}$  is a bounded metric, thus  $\mathbb{R}$  is bounded.
- Let  $f$  be a identity map on  $X$ .

## Question 7 Contd

- Since  $f$  is surjective, this implies

$$Y = f(X) = \bigcup_{i=1}^n B_Y(f(a_i), \epsilon).$$

- Hence  $Y$  is totally bounded.

## Question 8

Let  $f : X \rightarrow Y$  be a uniformly continuous surjective mapping between two metric spaces. Assume  $X$  is bounded. Does it necessarily mean that  $Y$  is bounded as well? If yes, give a proof. if no, create a counter-example.

**Solution: Ans- No.**

- Take  $X = (\mathbb{R}, d_{dis})$  and  $Y = (\mathbb{R}, d_{Euc})$ .
- Since  $d_{dis}$  is a bounded metric, thus  $\mathbb{R}$  is bounded.
- Let  $f$  be a identity map on  $X$ .

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ ,

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,



## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$  and so  $d_{Euc}(f(x), f(y))$

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$  and so  $d_{Euc}(f(x), f(y)) = d_{Euc}(f(x), f(x))$

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$  and so  $d_{Euc}(f(x), f(y)) = d_{Euc}(f(x), f(x)) = 0 < \epsilon$ .

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$  and so  $d_{Euc}(f(x), f(y)) = d_{Euc}(f(x), f(x)) = 0 < \epsilon$ .
- Hence  $f : (\mathbb{R}, d_{dis}) \rightarrow (\mathbb{R}, d_{Euc})$  defined by  $f(x) = x$  is uniformly continuous and surjective as well.

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$  and so  $d_{Euc}(f(x), f(y)) = d_{Euc}(f(x), f(x)) = 0 < \epsilon$ .
- Hence  $f : (\mathbb{R}, d_{dis}) \rightarrow (\mathbb{R}, d_{Euc})$  defined by  $f(x) = x$  is uniformly continuous and surjective as well.
- Clearly  $X$  is bounded but  $Y$  is not bounded.

## Question 8 Contd...

- Now we will prove that  $f$  is uniformly continuous.
- More generally we will prove that every function from discrete metric space  $X$  to an arbitrary metric space  $Y$  is uniformly continuous.
- Let  $\epsilon > 0$ , then set  $\delta = 1$ , and then if  $d_{dis}(x, y) < 1$ ,  $\implies x = y$  and so  $d_{Euc}(f(x), f(y)) = d_{Euc}(f(x), f(x)) = 0 < \epsilon$ .
- Hence  $f : (\mathbb{R}, d_{dis}) \rightarrow (\mathbb{R}, d_{Euc})$  defined by  $f(x) = x$  is uniformly continuous and surjective as well.
- Clearly  $X$  is bounded but  $Y$  is not bounded.

# Question 9

## Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.



## Question 9

### Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

**Solution:(a)**

# Question 9

## Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.

## Question 9

### Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.
- Since  $A$  is totally bounded,

# Question 9

## Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.
- Since  $A$  is totally bounded, there exists  $x_1, \dots, x_N$  in  $X$  such that

# Question 9

## Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.
- Since  $A$  is totally bounded, there exists  $x_1, \dots, x_N$  in  $X$  such that

$$A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N).$$

## Question 9

### Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.
- Since  $A$  is totally bounded, there exists  $x_1, \dots, x_N$  in  $X$  such that

$$A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N).$$

- We will prove that

## Question 9

### Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.
- Since  $A$  is totally bounded, there exists  $x_1, \dots, x_N$  in  $X$  such that

$$A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N).$$

- We will prove that

$$\bar{A} \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N).$$

## Question 9

### Question 9

Let  $(X, d)$  be a metric space.

- (a) Show that if  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is also totally bounded.
- (b) Use (a) to show that if  $(X, d)$  is complete and  $A$  is a totally bounded subset of  $(X, d)$ , then  $\bar{A}$  is compact.

### Solution:(a)

- Let  $\epsilon > 0$  be given.
- Since  $A$  is totally bounded, there exists  $x_1, \dots, x_N$  in  $X$  such that

$$A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N).$$

- We will prove that

$$\bar{A} \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N).$$



## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ .

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ ,

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

- Thus it shows that

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

- Thus it shows that  $x \in B_{\epsilon}(x_i) \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

- Thus it shows that  $x \in B_{\epsilon}(x_i) \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .
- Therefore



## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

- Thus it shows that  $x \in B_{\epsilon}(x_i) \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .
- Therefore  $\bar{A} \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

- Thus it shows that  $x \in B_{\epsilon}(x_i) \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .
- Therefore  $\bar{A} \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .
- Hence  $\bar{A}$  is totally bounded

## Question 9 Contd...

- Let  $x \in \bar{A}$ . Thus  $B_{\epsilon/2}(x) \cap A \neq \emptyset$ .
- Let  $y \in B_{\epsilon/2}(x) \cap A$ . Thus  $y \in B_{\epsilon/2}(x_i)$  for some  $1 \leq i \leq N$ , since  $A \subset B_{\epsilon/2}(x_1) \cup \dots \cup B_{\epsilon/2}(x_N)$ .
- Then

$$d(x, x_i) \leq d(x, y) + d(y, x_i) < \epsilon/2 + \epsilon/2 = \epsilon.$$

- Thus it shows that  $x \in B_{\epsilon}(x_i) \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .
- Therefore  $\bar{A} \subset B_{\epsilon}(x_1) \cup \dots \cup B_{\epsilon}(x_N)$ .
- Hence  $\bar{A}$  is totally bounded

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

**Solution:(b)**

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

### Solution:(b)

- From above theorem, to show that  $\bar{A}$  is compact, it is sufficient to prove that  $\bar{A}$  is complete and totally bounded.

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

### Solution:(b)

- From above theorem, to show that  $\bar{A}$  is compact, it is sufficient to prove that  $\bar{A}$  is complete and totally bounded.
- Since  $(X, d)$  is complete and closed subsets of a complete metric space are complete,

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

### Solution:(b)

- From above theorem, to show that  $\bar{A}$  is compact, it is sufficient to prove that  $\bar{A}$  is complete and totally bounded.
- Since  $(X, d)$  is complete and closed subsets of a complete metric space are complete, Consequently,  $\bar{A}$  is complete.



## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

### Solution:(b)

- From above theorem, to show that  $\bar{A}$  is compact, it is sufficient to prove that  $\bar{A}$  is complete and totally bounded.
- Since  $(X, d)$  is complete and closed subsets of a complete metric space are complete, Consequently,  $\bar{A}$  is complete.
- From part(a)  $\bar{A}$  is totally bounded subset of  $(X, d)$ .

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

### Solution:(b)

- From above theorem, to show that  $\bar{A}$  is compact, it is sufficient to prove that  $\bar{A}$  is complete and totally bounded.
- Since  $(X, d)$  is complete and closed subsets of a complete metric space are complete, Consequently,  $\bar{A}$  is complete.
- From part(a)  $\bar{A}$  is totally bounded subset of  $(X, d)$ .
- Hence  $\bar{A}$  is compact.

## Question 9(b)

### Theorem

*For a metric space  $(X, \rho)$ , the following are equivalent:*

- (1)  $X$  is compact.*
- (2)  $X$  is sequentially compact.*
- (3)  $X$  is complete and totally bounded.*

### Solution:(b)

- From above theorem, to show that  $\bar{A}$  is compact, it is sufficient to prove that  $\bar{A}$  is complete and totally bounded.
- Since  $(X, d)$  is complete and closed subsets of a complete metric space are complete, Consequently,  $\bar{A}$  is complete.
- From part(a)  $\bar{A}$  is totally bounded subset of  $(X, d)$ .
- Hence  $\bar{A}$  is compact.

## Question 10

### Question 10

Let  $A$  be a non-empty subset of a metric space  $(X, d)$ . Recall that the distance of a point  $x \in X$  to a set  $A$  is defined by

$$d(x, A) := \inf\{d(x, y) : y \in A\}$$

Show that if  $A$  is compact subset of  $X$ , then there is  $y \in A$  such that  $d(x, A) = d(x, y)$ .

### Solution

- Since  $d(x, A) := \inf\{d(x, y) : y \in A\}$ .

## Question 10

### Question 10

Let  $A$  be a non-empty subset of a metric space  $(X, d)$ . Recall that the distance of a point  $x \in X$  to a set  $A$  is defined by

$$d(x, A) := \inf\{d(x, y) : y \in A\}$$

Show that if  $A$  is compact subset of  $X$ , then there is  $y \in A$  such that  $d(x, A) = d(x, y)$ .

### Solution

- Since  $d(x, A) := \inf\{d(x, y) : y \in A\}$ .
- By definition of infimum,

## Question 10

### Question 10

Let  $A$  be a non-empty subset of a metric space  $(X, d)$ . Recall that the distance of a point  $x \in X$  to a set  $A$  is defined by

$$d(x, A) := \inf\{d(x, y) : y \in A\}$$

Show that if  $A$  is compact subset of  $X$ , then there is  $y \in A$  such that  $d(x, A) = d(x, y)$ .

### Solution

- Since  $d(x, A) := \inf\{d(x, y) : y \in A\}$ .
- By definition of infimum, for every  $n \geq 1$ ,

## Question 10

### Question 10

Let  $A$  be a non-empty subset of a metric space  $(X, d)$ . Recall that the distance of a point  $x \in X$  to a set  $A$  is defined by

$$d(x, A) := \inf\{d(x, y) : y \in A\}$$

Show that if  $A$  is compact subset of  $X$ , then there is  $y \in A$  such that  $d(x, A) = d(x, y)$ .

### Solution

- Since  $d(x, A) := \inf\{d(x, y) : y \in A\}$ .
- By definition of infimum, for every  $n \geq 1$ , there exists  $y_n \in A$  such that

## Question 10

### Question 10

Let  $A$  be a non-empty subset of a metric space  $(X, d)$ . Recall that the distance of a point  $x \in X$  to a set  $A$  is defined by

$$d(x, A) := \inf\{d(x, y) : y \in A\}$$

Show that if  $A$  is compact subset of  $X$ , then there is  $y \in A$  such that  $d(x, A) = d(x, y)$ .

### Solution

- Since  $d(x, A) := \inf\{d(x, y) : y \in A\}$ .
- By definition of infimum, for every  $n \geq 1$ , there exists  $y_n \in A$  such that

$$d(x, A) \leq d(x, y_n) \leq d(x, A) + \frac{1}{n}.$$



## Question 10

### Question 10

Let  $A$  be a non-empty subset of a metric space  $(X, d)$ . Recall that the distance of a point  $x \in X$  to a set  $A$  is defined by

$$d(x, A) := \inf\{d(x, y) : y \in A\}$$

Show that if  $A$  is compact subset of  $X$ , then there is  $y \in A$  such that  $d(x, A) = d(x, y)$ .

### Solution

- Since  $d(x, A) := \inf\{d(x, y) : y \in A\}$ .
- By definition of infimum, for every  $n \geq 1$ , there exists  $y_n \in A$  such that

$$d(x, A) \leq d(x, y_n) \leq d(x, A) + \frac{1}{n}. \quad (1)$$

## Question 10 Contd...

- Since  $A$  is compact,

## Question 10 Contd...

- Since  $A$  is compact, there exists a subsequence  $(y_{n_k})$  converging to some  $y \in A$ .

## Question 10 Contd...

- Since  $A$  is compact, there exists a subsequence  $(y_{n_k})$  converging to some  $y \in A$ . i.e  $d(x, y_{n_k}) \rightarrow d(x, y)$  as  $k \rightarrow \infty$ .

## Question 10 Contd...

- Since  $A$  is compact, there exists a subsequence  $(y_{n_k})$  converging to some  $y \in A$ . i.e  $d(x, y_{n_k}) \rightarrow d(x, y)$  as  $k \rightarrow \infty$ .
- From (1), we have

## Question 10 Contd...

- Since  $A$  is compact, there exists a subsequence  $(y_{n_k})$  converging to some  $y \in A$ . i.e  $d(x, y_{n_k}) \rightarrow d(x, y)$  as  $k \rightarrow \infty$ .
- From (1), we have

$$d(x, A) \leq d(x, y_{n_k}) \leq d(x, A) + \frac{1}{n_k},$$

## Question 10 Contd...

- Since  $A$  is compact, there exists a subsequence  $(y_{n_k})$  converging to some  $y \in A$ . i.e  $d(x, y_{n_k}) \rightarrow d(x, y)$  as  $k \rightarrow \infty$ .
- From (1), we have

$$d(x, A) \leq d(x, y_{n_k}) \leq d(x, A) + \frac{1}{n_k},$$

- Letting  $k \rightarrow \infty$  gives  $d(x, y) = d(x, A)$ .

## Question 10 Contd...

- Since  $A$  is compact, there exists a subsequence  $(y_{n_k})$  converging to some  $y \in A$ . i.e  $d(x, y_{n_k}) \rightarrow d(x, y)$  as  $k \rightarrow \infty$ .
- From (1), we have

$$d(x, A) \leq d(x, y_{n_k}) \leq d(x, A) + \frac{1}{n_k},$$

- Letting  $k \rightarrow \infty$  gives  $d(x, y) = d(x, A)$ .