Conversion of Cellular Kinetic Data for Chimeric Antigen Receptor T-cell Therapy (CAR-T) into Interpretable Units

Anwesha Chaudhury,¹ Andrew Stein,¹ Stephan Grupp,²,³ John Levine,⁴,⁵ Michael Pulsipher,⁶ G. Doug Myers,² Edward Waldron,⁶ Xu Zhu,¹ Fraser McBlane,⁶ Rakesh Awasthi,¹ Edmund K. Waller¹⁰

¹Novartis Institute of Biomedical Research, Cambridge, MA, USA; ²Division of Oncology, Cell Therapy and Transplant Section, Children's Hospital of Philadelphia, PA, USA; ³Department of Pediatrics, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA, USA; ⁴University of Michigan, Ann Arbor, MI, USA; ⁵School of Medicine at Mount Sinai, NY, USA; ⁵Children's Hospital Los Angeles Cancer and Blood Disease Institute, Los Angeles, CA, USA; ⁵Novartis Pharmaceuticals Corporation, East Hanover, NJ, USA; ⁵Novartis Pharma AG, Basel, Switzerland;

¹ºBone Marrow and Stem Cell Transplant Center, Emory University Winship Cancer Institute, Atlanta, GA, USA

Introduction

- The cellular kinetics of CAR-T therapy has been described using both quantitative polymerase chain reaction (qPCR) and flow cytometry assays¹.
- The cellular kinetic measurements for CAR-T therapy by qPCR has been widely reported in units of copies of CAR transgene per µg of genomic DNA, while the flow cytometry assay quantifies the percentage of CAR-T cells as a function of either T cells or white blood cells (WBC) based on their surface expression. Neither measurement accounts for the fact that the WBCs may increase significantly following CAR-T infusion as the patient's hematological counts recover from prior lymphodepletion.
- This warrants the need for converting the measurements into units that are physiologically interpretable, not only for fully appreciating the CAR-T expansion and correlating with outcomes, but also to use in future mechanistic models for CAR-T therapy.
- We propose a formula to convert the qPCR measurements (from copies/µg), and flow cytometry (from percentage of cells) to the more physiologically interpretable cellular concentration in the unit of cells/µl units.

Methods

 Data from 79 relapsed/refractory pediatric acute lymphoblastic leukemia (ALL) patients in the ELIANA (CCTL019B2202) trial were included in this analysis. For converting qPCR measurements from copies/µg to cells/µl, the below equation is proposed. Parameters M, N, F are unknown.

Equation 1
$$\frac{\text{CAR}^+ \text{ cells}}{\mu \text{l blood}} = \frac{\text{WBC}}{\mu \text{l blood}} \times \frac{\text{CAR DNA copies}}{\mu \text{g DNA}} \times \frac{M \ \mu \text{g DNA}}{\text{WBC}} \times \frac{1 \ \text{CAR cell}}{\text{N CAR DNA copies}} \times F$$

M is the number of µg of DNA per white blood cell; N is the number of copies of CAR DNA per CAR-T cell; F is the fraction of cells with CAR DNA that express the CAR.

• For converting flow cytometry measurements from percent of CAR+ cells to cells/µl, the below equation was proposed.

Equation 2
$$\frac{\text{CD3+ CAR+ cells}}{\mu l \text{ blood}} = \frac{\text{WBCs}}{\mu l \text{ blood}} \times \frac{\text{CD3+ CAR+ cells}}{\text{WBC}}$$

 Upon conversion of data to cellular concentration, the exposure-response relationships have been revisited to assess any benefit in response/safety prediction using the exposure in converted units.


Results

• Three different approaches with varying complexity were taken to estimate (M*F)/N in order to perform the conversion from native units to cellular concentration.

Approach 1

- Assume values for the three parameters, M, N, F in Equation (1). We assume $M = 6.6e-6 \mu g$ DNA per cell, $N = 1 \frac{CAR}{CAR}$ cell, $F = 1 \frac{100\%}{CAR}$ of T cells with CAR DNA express the CAR).
- Plotting values for CAR+ T cells measured by flow cytometry and CAR content based upon qPCR estimates using M=6.6e-6 µg, N=1, F=1 yielded DNA-based values for CAR-T consistently higher than flow cytometry-based values for CAR-T (**Figure 1**). We adjusted the assumptions for M, N, and F in Approach 2.


Figure 1. Comparing cells/ μ l derived from qPCR and flow cytometry assay assuming M=6.6e-6 μ g, N=1, F=1



Approach 2

- We perform a regression between WBC x CAR copies/µg from qPCR and cells/µl from the flow cytometry assay to estimate the lumped value of (M*F)/N = 2.68e-6 that gives the best agreement between the flow and PCR estimates.
- Assuming that M = 6.6e-6 µg DNA per cell is correct, there are different possible values for N and F. At one extreme, N = 1 copy of CAR DNA per CAR-T cell, and F = 0.4, or only 40% of the cells with CAR DNA are expressing the CAR.
- At the other extreme, N = 2.46 copies of CAR DNA per CAR-T cell and F = 1, or 100% of CAR-T cells express the CAR (Figure 2).

Figure 2. Comparing cells/µl derived from qPCR and flow cytometry assay by estimating a population average (M*F)/N




• Upon conversion from copies/ μ g to cells/ μ l, it can be noted that the native measurements and cellular concentration values are highly correlated (R² = 0.752). Thus, a difference in the exposure-response relationship between the two units is not expected.

Approach 3

The population cellular kinetic (popCK) model proposed by Stein et al² was fit to both the qPCR and flow cytometry data together with an estimate for the (M*F)/N ratio both for the population and for each individual, in addition to estimating the other parameters that capture the CAR-T kinetics (e.g., CAR-T expansion) (Figure 3).

Figure 3. Schematic showing the mathematical model describing Tisagenlecleucel expansion



• The population estimate for the (M*F)/N ratio is 1.58e-6. A comparison of the parameters using the various data sources (unconverted copies/µg vs cells/µl) are shown in **Table 1**.

Table 1. Comparison of crucial popCK model parameters using the native qPCR units vs cells/µl


	Copies/µg fit	Cells/µl pooled fit (Approach 3)
Fold expansion	3366	39044
T _{max}	8.87 days	9.43 days
α	0.17/day	0.31/day
β	0.0026/day	0.0024/day

A 12x higher expansion of CAR-T cells can be observed based on the parameter estimates using copies/µg as compared to cells/µl. In this trial, this could be attributed to the simultaneous increase in WBCs as well as CAR⁺ cells that lead to a higher expansion when considering cellular concentration. The units of copies/µg gives the CAR expression normalized by the total WBC DNA, thus the expansion quantified in copies/µg units does not capture the increase in the total number of CAR⁺ cells in vivo.

Exposure-Response

- For assessing the exposure-response relationship, cellular kinetic parameters (C_{max} , T_{max} , AUC_{0-28d}) that were derived from the popCK model were used to fit a regression model with a response variable.
- The time-to-event response variables explored were event free survival (EFS), duration of remission (DOR), and B-cell recovery time. For time-to-event response variables, a Cox proportionality regression model was fit.
- Logistic regression models were fit to two categorical variables: 28-day response, grade ≥3 cytokine release syndrome (CRS). The odds ratio provides the propensity of a categorical response upon doubling the exposure (**Figure 4**).

Figure 4. Forest plot showing the hazard/odds ratio describing the exposure-response relationship of time-to-event and categorical variables



Conclusions

- We have proposed equations to convert cellular kinetic measurements from qPCR and flow cytometry assays to physiologically meaningful units of cells/µl.
- The CAR-T fold-expansion in cells/µl is observed to be ~12x higher as compared to copies/µg. This can be attributed to the simultaneous expansion in WBCs along with the CAR+T cells.
- When assessing the exposure-response relationship, using units of cells/µl provides a similar result compared to the native copies/µg units, likely due to a high correlation between exposure measurements by either set of units.

References

1. Thudium et al, Blood (2017) 130 (21): 2317-2325. 2. Stein et al, CPT: Pharmacometrics and Systems Pharmacology (2019), 8 (5): 285-295.

Acknowledgments

The authors sincerely thank: Patients and their families; Principal investigators and support staff. The authors would also like to thank Karen Thudium Mueller for continued discussions and support, and Chris del Corral for clinical inputs. Medical writing support was provided by Mukul Rastogi, of Novartis Healthcare Pvt. Ltd.

Disclosures

SG: Personal fees from Jazz, Janssen, CBMG, TCR2, Humanigen, Roche, Vertex/CRISPR, Adaptimmune, Allogene, Kite/Gilead, Servier, Juno. **JL**: Personal fees from Incyte, Kamada, Mesoblast, Biogen, Novartis, Bluebird Bio, Ironwood, Talaris, X4 Pharmaceuticals; **MP**: Participated in Study steering committee and educational activities, Novartis; **GDM**: Novartis. **EKW**: Personal fees from Novartis, Kite/Gilead, Juno/BMS, Pharmacyclics, Verastem, Karyopharm, Cambium Medical Technologies; **AC, AS, EW, XZ, FM, RA**: Employment at Novartis.

Poster presented at the AACR Virtual Annual Meeting, Week 1: April 10-15 and Week 2: May 17-21, 2021.

This study was sponsored by Novartis Pharmaceuticals Corporation.

3 ways to instantly download an electronic copy of this poster to your mobile device or e-mail a copy to your computer or tablet

Text: Q0609a

To: 8NOVA (86682) US Only

+18324604729 North, Central, and SouthAmericas; Caribbean; China +447860024038 UK, Europe, and Russia

Visit the web at:

http://Novartis.medicalcongressposters.com/Default.aspx?doc=0609a

Copies of this poster obtained through Quick Response (QR) code are for personal use only and may not be reproduced without written permission from AACR and the author of this poster

Presenter email address: anwesha.chaudhury@novartis.com

Standard data or message rates may apply

