

## COMP3670/6670: Introduction to Machine Learning

These exercises will concentrate on vector calculus, and how to compute derivatives of functions that live in higher dimensions.

### Preliminaries

The formal definition of the derivative of a function  $f : \mathbb{R} \rightarrow \mathbb{R}$  is given by

$$\frac{df}{dx} := \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  be a function of a vector  $\mathbf{x}$ . The derivative of  $f(\mathbf{x})$  with respect to  $\mathbf{x}$  is defined as

$$\nabla_{\mathbf{x}} f = \text{grad} f = \frac{df}{d\mathbf{x}} := \begin{bmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} & \frac{\partial f(\mathbf{x})}{\partial x_2} & \cdots & \frac{\partial f(\mathbf{x})}{\partial x_n} \end{bmatrix} \in (\mathbb{R}^n \rightarrow \mathbb{R})^{1 \times n}$$

Note that  $\frac{df}{d\mathbf{x}}$  is a row vector, where each element is a function of the form  $\mathbb{R}^n \rightarrow \mathbb{R}$ . We write  $\nabla_{\mathbf{x}} f \in (\mathbb{R}^n \rightarrow \mathbb{R})^{1 \times n}$ . Some authors write  $\nabla_{\mathbf{x}} f \in \mathbb{R}^{1 \times n}$  as an abuse of notation for the sake of brevity, and ease of matching dimensions. Keep in mind that each element of the row vector isn't a real number, but itself a function.

Let  $\mathbf{g} : \mathbb{R} \rightarrow \mathbb{R}^n$  be a function of a scalar  $t$ . The derivative of  $\mathbf{g}(t)$  with respect to  $t$  is defined as

$$\frac{d\mathbf{g}}{dt} := \begin{bmatrix} \frac{dg_1(t)}{dt} \\ \frac{dg_2(t)}{dt} \\ \vdots \\ \frac{dg_n(t)}{dt} \end{bmatrix} \in (\mathbb{R} \rightarrow \mathbb{R})^{n \times 1}$$

Note that  $\frac{d\mathbf{g}}{dt}$  is a column vector, where each element is itself a function of the form  $\mathbb{R} \rightarrow \mathbb{R}$ . As before, we notate this using an abuse of notation as  $\frac{d\mathbf{g}}{dt} \in \mathbb{R}^{n \times 1}$ ,

The reason why the derivatives are defined this way, is so that the dimensions match when we define the chain rule.

Given  $f : \mathbb{R}^n \rightarrow \mathbb{R}$  and  $\mathbf{g} : \mathbb{R} \rightarrow \mathbb{R}^n$ , we can define two new functions

$$h : \mathbb{R} \rightarrow \mathbb{R}, \quad h(t) = f(\mathbf{g}(t))$$

$$\mathbf{k} : \mathbb{R}^n \rightarrow \mathbb{R}^n \quad \mathbf{k}(\mathbf{x}) = \mathbf{g}(f(\mathbf{x}))$$

and we can define their derivatives as

$$\frac{dh}{dt} = \frac{df}{d\mathbf{g}} \frac{d\mathbf{g}}{dt} = \begin{bmatrix} \frac{\partial f(\mathbf{g})}{\partial g_1} & \cdots & \frac{\partial f(\mathbf{g})}{\partial g_n} \end{bmatrix} \begin{bmatrix} \frac{\partial g_1}{\partial t} \\ \vdots \\ \frac{\partial g_n}{\partial t} \end{bmatrix} = \sum_{i=1}^n \frac{\partial f(\mathbf{g})}{\partial g_i} \frac{\partial g_i}{\partial t}$$

and

$$\frac{d\mathbf{k}}{d\mathbf{x}} = \frac{d\mathbf{g}}{df} \frac{df}{d\mathbf{x}} = \begin{bmatrix} \frac{\partial g_1}{\partial f} \\ \vdots \\ \frac{\partial g_n}{\partial f} \end{bmatrix} \begin{bmatrix} \frac{\partial f(\mathbf{x})}{\partial x_1} & \cdots & \frac{\partial f(\mathbf{x})}{\partial x_n} \end{bmatrix} = \begin{bmatrix} \frac{\partial g_1}{\partial f} \frac{\partial f(\mathbf{x})}{\partial x_1} & \cdots & \frac{\partial g_1}{\partial f} \frac{\partial f(\mathbf{x})}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial g_n}{\partial f} \frac{\partial f(\mathbf{x})}{\partial x_1} & \cdots & \frac{\partial g_n}{\partial f} \frac{\partial f(\mathbf{x})}{\partial x_n} \end{bmatrix} = \mathbf{A}$$

where  $\mathbf{A}_{ij} = \frac{\partial g_i}{\partial f} \frac{\partial f(\mathbf{x})}{\partial x_j}$ .

(Here, the term  $\frac{\partial f(\mathbf{g})}{\partial g_i}$  means to substitute each output component of  $\mathbf{g}$  into the inputs for  $f$ , and take the partial derivative with respect to the  $g_i$ , the  $i^{\text{th}}$  component of  $\mathbf{g}$ .)

For a vector valued function  $\mathbf{f} : \mathbb{R}^n \rightarrow \mathbb{R}^m$ , we define the matrix of all first order derivatives as the *Jacobian*, which is given by

$$\mathbf{J} = \nabla_{\mathbf{x}} \mathbf{f} = \frac{d\mathbf{f}(\mathbf{x})}{d\mathbf{x}} = \begin{bmatrix} \frac{\partial f_1(\mathbf{x})}{\partial x_1} & \cdots & \frac{\partial f_1(\mathbf{x})}{\partial x_n} \\ \vdots & \ddots & \vdots \\ \frac{\partial f_m(\mathbf{x})}{\partial x_1} & \cdots & \frac{\partial f_m(\mathbf{x})}{\partial x_n} \end{bmatrix}, \quad \mathbf{J}_{ij} = \frac{\partial f_i(\mathbf{x})}{\partial x_j}$$

.

You may also need the definition of matrix multiplication.

If  $\mathbf{A} \in \mathbb{R}^{n \times m}$  and  $\mathbf{B} \in \mathbb{R}^{m \times p}$ , the product  $\mathbf{C} = \mathbf{AB}$  is a matrix in  $\mathbb{R}^{n \times p}$  satisfying

$$C_{ij} = \sum_{k=1}^m A_{ik} B_{kj}$$

If  $\mathbf{A} \in \mathbb{R}^{n \times m}$ ,  $\mathbf{b} \in \mathbb{R}^{m \times 1}$  and  $\mathbf{c} \in \mathbb{R}^{n \times 1}$  then the matrix vector products  $\mathbf{Ab}$  and  $\mathbf{c}^T \mathbf{A}$  satisfy the properties

$$(\mathbf{Ab})_k = \sum_{j=1}^m A_{kj} b_j$$

and

$$(\mathbf{c}^T \mathbf{A})_k = \sum_{i=1}^n A_{ik} c_i$$

For  $\mathbf{x} \in \mathbb{R}^n$ , the Euclidean norm  $\|\cdot\|_2$  is given by

$$\|\mathbf{x}\|_2 := \sqrt{\mathbf{x}^T \mathbf{x}}$$

For all problems below, state the dimension of the answer where appropriate.

**Question 1** **Formal definition of derivative**

Compute the derivative of  $f : \mathbb{R} \rightarrow \mathbb{R}, f(x) = x^2$  from the formal limit definition of the derivative.

**Question 2** **Vector Derivative of Scalar Function**

Given  $f : \mathbb{R}^2 \rightarrow \mathbb{R}, f(\mathbf{x}) = 2x_1x_2 + x_1 + 3x_2 + 5$ , compute  $\frac{df}{d\mathbf{x}}$ .

**Question 3** **Scalar Derivative of Vector Function**

Given  $\mathbf{g}(t) : \mathbb{R} \rightarrow \mathbb{R}^2, \mathbf{g}(t) = \begin{bmatrix} t^2 \\ e^t \end{bmatrix}$  compute  $\frac{d\mathbf{g}}{dt}$ .

**Question 4** **Derivative of the L2 Norm**

Let  $\mathbf{x} \in \mathbb{R}^n$ , and define  $k : \mathbb{R}^n \rightarrow \mathbb{R}, k(\mathbf{x}) = \|\mathbf{x}\|_2^2 := \mathbf{x}^T \mathbf{x}$ . Compute  $\frac{dk}{d\mathbf{x}}$ .

**Question 5** **Chain Rule, Scalar Derivative**

Let  $h : \mathbb{R} \rightarrow \mathbb{R}, h(t) = f(\mathbf{g}(t))$ , where  $f$  and  $\mathbf{g}$  are defined in Question 2 and Question 3 respectively.

1. Compute  $\frac{dh}{dt}$  by using the chain rule.
2. Compute  $\frac{dh}{dt}$  by evaluating  $f(\mathbf{g}(t))$  first, and then differentiating the entire expression by  $t$ . Compare your answer to the above and check that they match.

**Question 6** **Chain Rule, Vector Derivative**

Let  $\mathbf{k} : \mathbb{R}^n \rightarrow \mathbb{R}^n, \mathbf{k}(\mathbf{x}) = \mathbf{g}(f(\mathbf{x}))$ , where  $f$  and  $\mathbf{g}$  are defined in Question 2 and Question 3 respectively.

1. Compute  $\frac{d\mathbf{k}}{d\mathbf{x}}$  using the chain rule.
2. Compute  $\frac{d\mathbf{k}}{d\mathbf{x}}$  directly by using the Jacobian to differentiate  $\mathbf{g}(f(\mathbf{x}))$ . Check your answer matches the above using chain rule.

**Question 7** **More Derivatives**

1. Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}, f(\mathbf{x}) = (\mathbf{x}^T \mathbf{x} + 1)^2$ . Compute  $\frac{d}{d\mathbf{x}} f(\mathbf{x})$  using the chain rule. (You can use the previous questions to help you.)
2. Directly compute  $\frac{d}{d\mathbf{x}} f(\mathbf{x})$  by expanding out  $(\mathbf{x}^T \mathbf{x} + 1)^2$  first. Your result should match the above.

**Question 8** **Derivative of a Matrix-Vector product**

Let  $\mathbf{A} \in \mathbb{R}^{m \times n}$  and  $\mathbf{x} \in \mathbb{R}^{n \times 1}$ . Show that  $\frac{d}{d\mathbf{x}}(\mathbf{A}\mathbf{x}) = \mathbf{A}$ .

**Question 9** **Linear Regression**

Let  $\Phi \in \mathbb{R}^{n \times m}, \mathbf{w} \in \mathbb{R}^{n \times 1}, \mathbf{t} \in \mathbb{R}^{m \times 1}$ .

Let  $f : \mathbb{R}^n \rightarrow \mathbb{R}, f(\mathbf{w}) = \|((\mathbf{w}^T \Phi)^T - \mathbf{t})\|_2^2$

1. Verify that  $f$  is well defined (the dimensions of all the components match up).
2. Compute  $\frac{d}{d\mathbf{w}} f(\mathbf{w})$ .

**Question 10** **Matrix Gradient**

Given  $\mathbf{X} \in \mathbb{R}^{n \times m}$  and some vectors  $\mathbf{a} \in \mathbb{R}^{? \times ?}, \mathbf{b} \in \mathbb{R}^{? \times ?}$ .

1. What are the dimensions of  $\mathbf{a}$  and  $\mathbf{b}$  such that  $\mathbf{a}^T \mathbf{X} \mathbf{b}$  is well defined?<sup>1</sup> What is the dimension of the result?
2. Compute the matrix gradient  $\frac{d}{d\mathbf{X}} \mathbf{a}^T \mathbf{X} \mathbf{b}$ .

---

<sup>1</sup>Note that if  $\mathbf{X}$  is square, symmetric and positive definite, then defining  $\langle \mathbf{a}, \mathbf{b} \rangle := \mathbf{a}^T \mathbf{X} \mathbf{b}$  gives an inner product.