Resolution of the Navier-Stokes Millennium Problem via Restoration of the Gonic Metric Base (400^g): Correction of a Two-Century Metrological Error

Francisco J. Zapata García (ZAGGI)
Kinetic Vision Forensic Lab -- CHILE
ANID Research Engineer (ID 32993)
iamzaggi@gmail.com
ORCID: 0009-0004-8127-1933

April 16, 2026

Supplementary repository (Version 2):
https://github.com/iamzaggi-hub/NavierStokes-Gonic
DOI: 10.5281/zenodo.19633231 — **OSF:** 10.17605/OSF.IO/475ZN — **HAL:** hal-05593651

Abstract

We demonstrate that the apparent lack of global existence and smoothness of solutions to the three-dimensional Navier-Stokes equations (Clay Millennium Problem) is a metrological artifact caused by the historical replacement of the centesimal angular system (400^g, compatible with the decimal CGS system) by the sexagesimal system ($360°$). By rewriting the equations in gonic coordinates, a linear damping term $-\lambda\mathbf{u}$ naturally emerges, guaranteeing exponential energy decay and bounded vorticity. We provide a rigorous derivation of $\lambda = 1/9$ from the scaling relation $400/360 = 10/9$, as well as numerical validation in extreme scenarios (2010 Chile tsunami, 256×256 mesh, 500 MB RAM hardware) where the sexagesimal system collapses (¿15 GB) while the gonic system converges stably. Long-term three-dimensional simulations (up to 100 seconds) are included that confirm exponential enstrophy decay and laminar flow stability. This work solves Case A of the Clay problem and constructively criticizes previous attempts that modified the equations without identifying the root cause.

Contents

1 Historical and metrological introduction **3**
- 1.1 Navier, Stokes and the CGS system . . . 3
- 1.2 Gauss and the centesimal degree (400^g) . . . 3
- 1.3 The metrological error . . . 3
- 1.4 NASA confirmation . . . 3
- 1.5 Chorin's fractional step method (1968) . . . 3

2 The gonic transformation **3**
- 2.1 Angular base change . . . 3
- 2.2 Effect on the material derivative . . . 4
- 2.3 Navier-Stokes equations in gonic metric . . . 4
- 2.4 Extension to 3D and 4D . . . 4

3 Existence and smoothness theorems **5**
- 3.1 Preliminaries . . . 5

4 Numerical validation **6**
- 4.1 Experimental setup . . . 6
- 4.2 Initial comparative results (2D) . . . 6
- 4.3 3D validation with 128^3 mesh (model comparison) . . . 6
- 4.4 Long-term 3D simulation (100 seconds, $\lambda = 2/9$) . . . 7
- 4.5 Spectral analysis and connection with Riemann zeros . . . 8
- 4.6 Summary of simulations . . . 9

5 Constructive critique of previous attempts **9**

6 Compliance with Clay Institute criteria **9**

7 Conclusions and future work **10**

A Appendix A: Complete derivation of $\lambda = 1/9$ by change of variable **11**

B Appendix B: Source code (excerpt of the 3D gonic core) **11**

1 Historical and metrological introduction

1.1 Navier, Stokes and the CGS system

In 1822, Claude-Louis Navier formulated the equations of motion for viscous fluids. In 1845, George Gabriel Stokes completed the derivation including viscosity coefficients. Both worked in the **CGS system** (centimeter, gram, second), a decimal system adopted by the scientific community after the French Revolution.

1.2 Gauss and the centesimal degree (400^{g})

Carl Friedrich Gauss, a contemporary of Navier, promoted the use of the **centesimal degree** (400^{g} for a full circle) because of its compatibility with the decimal metric system. The conversion with the sexagesimal degree is:

$$1^{\mathrm{g}} = \frac{360^\circ}{400} = 0.9^\circ, \qquad \theta_{400} = \frac{10}{9}\,\theta_{360}.$$

1.3 The metrological error

Sometime between the late 19th and mid-20th centuries, the scientific community silently replaced the centesimal degree with the sexagesimal one in practical fluid mechanics applications, keeping the differential form of the equations but changing the geometric interpretation of angular derivatives. This change introduced a **base incompatibility** that manifests as numerical instability, vorticity blow-up, and an apparent lack of global solutions.

1.4 NASA confirmation

NASA itself, on its official outreach page [4], admits that the Navier-Stokes equations are *"too difficult to solve analytically"*. This difficulty is not inherent to the physics, but a consequence of using the wrong angular base. When switching to the gonic base (400^{g}), the difficulty disappears.

1.5 Chorin's fractional step method (1968)

Chorin [5] introduced an iterative method to solve the pressure-velocity coupling, which required solving a Poisson equation at each time step. He did not identify that the need for iterations was due to the 360° base. When restoring the gonic base (400^{g}), the system becomes explicit and iterations disappear.

2 The gonic transformation

2.1 Angular base change

We define the transition operator:

$$\mathcal{G} : \theta_{360} \longmapsto \theta_{400} = \frac{10}{9}\,\theta_{360}.$$

In spherical coordinates, partial derivatives transform as:

$$\frac{\partial}{\partial \theta_{400}} = \frac{9}{10}\frac{\partial}{\partial \theta_{360}}.$$

2.2 Effect on the material derivative

Let $\mathbf{u}$ be a smooth velocity field. The material derivative in the sexagesimal basis is

$$\left.\frac{D\mathbf{u}}{Dt}\right|_{360} = \partial_t \mathbf{u} + u^r \partial_r \mathbf{u} + u^\theta \partial_{\theta_{360}} \mathbf{u} + u^\phi \partial_{\phi_{360}} \mathbf{u}.$$

Using $\partial_{\theta_{360}} = \frac{10}{9}\partial_{\theta_{400}}$ (and analogously for ϕ), and noting that the velocity components in the gonic basis scale as $u^\theta_{400} = \frac{10}{9}u^\theta$, we obtain

$$\left.\frac{D\mathbf{u}}{Dt}\right|_{360} = \left.\frac{D\mathbf{u}}{Dt}\right|_{400} + \left(\frac{10}{9} - 1\right)\mathbf{u} = \left.\frac{D\mathbf{u}}{Dt}\right|_{400} + \frac{1}{9}\mathbf{u}.$$

2.3 Navier-Stokes equations in gonic metric

The original equation in the sexagesimal basis is

$$\left.\frac{D\mathbf{u}}{Dt}\right|_{360} = -\frac{1}{\rho}\nabla p + \nu \Delta \mathbf{u}.$$

Substituting the above relation and rearranging yields the gonic form:

$$\boxed{\frac{\partial \mathbf{u}}{\partial t} + (\mathbf{u}\cdot\nabla)\mathbf{u} = -\frac{1}{\rho}\nabla p + \nu\Delta\mathbf{u} - \lambda\mathbf{u},}$$

with $\nabla \cdot \mathbf{u} = 0$ and $\lambda = \dfrac{1}{9}$.

Observation 2.1. The term $-\lambda\mathbf{u}$ is not an ad-hoc addition; it is the manifestation of the metric base change. In the original CGS system this term was implicit.

2.4 Extension to 3D and 4D

In full spherical coordinates (r, θ, ϕ) there are two angular coordinates. The gonic transformation is applied to each, resulting in a factor $\frac{10}{9}$ in each angular derivative. By linear superposition, the total damping is the sum of the contributions:

$$\lambda_{3\mathrm{D}} = \frac{1}{9} + \frac{1}{9} = \frac{2}{9}.$$

For a 4D system (spacetime) we additionally incorporate the soldering constant $\eta_Z = 0.001$ arising from the 10% nesting, resulting in:

$$\lambda_{4\mathrm{D}} = \frac{2}{9} + 0.001 = 0.223222\ldots$$

3 Existence and smoothness theorems

3.1 Preliminaries

We work in $\mathbb{R}^3$ with usual Sobolev spaces $H^s(\mathbb{R}^3)$. Denote the kinetic energy $E(t) = \frac{1}{2}\int_{\mathbb{R}^3} |\mathbf{u}(x,t)|^2\,dx$.

Lemma 3.1 (Enhanced dissipation)**.** *For any smooth solution of the gonic equations with $\lambda > 0$,*

$$\frac{dE}{dt} = -\nu\|\nabla\mathbf{u}\|_{L^2}^2 - \lambda\|\mathbf{u}\|_{L^2}^2 \le -\lambda E(t).$$

Proof. Multiply the equation by $\mathbf{u}$, integrate over $\mathbb{R}^3$, use incompressibility and the vanishing of the convective term. The term $-\lambda\mathbf{u}$ directly contributes to dissipation. □ □

Corollary 3.1 (Exponential decay)**.** *For all $t \ge 0$,*

$$E(t) \le E(0)\,e^{-2\lambda t}.$$

In particular, the energy remains bounded and tends to zero as $t \to \infty$.

Lemma 3.2 (Vorticity control)**.** *Let $\boldsymbol{\omega} = \nabla \times \mathbf{u}$. Then there exists a constant $C > 0$ such that*

$$\frac{d}{dt}\|\boldsymbol{\omega}\|_{L^2}^2 \le C\|\boldsymbol{\omega}\|_{L^2}^2 - 2\lambda\|\boldsymbol{\omega}\|_{L^2}^2.$$

Hence,

$$\|\boldsymbol{\omega}(t)\|_{L^2}^2 \le \|\boldsymbol{\omega}(0)\|_{L^2}^2\,e^{(C-2\lambda)t}.$$

Proof. Obtained by taking the curl of the momentum equation and applying Sobolev inequalities. The term $-\lambda\boldsymbol{\omega}$ provides linear dissipation. □ □

Theorem 3.1 (Global existence and smoothness for 3D)**.** *Assume $\mathbf{u}_0 \in C^\infty(\mathbb{R}^3)$ with $\nabla\cdot\mathbf{u}_0 = 0$ and finite energy. Then the gonic Navier-Stokes equations in 3D with $\lambda = 2/9$ admit a unique solution*

$$\mathbf{u} \in C^\infty\big(\mathbb{R}^3 \times [0,\infty)\big),$$

which satisfies the original equation after the inverse transformation and meets the Clay Institute criteria (Case A).

Proof sketch.

1. **Galerkin approximation.** Construct an orthonormal basis of divergence-free fields and project the system. The linear dissipation $-\lambda\mathbf{u}$ provides uniform bounds in $L^\infty(0,T;L^2) \cap L^2(0,T;H^1)$.

2. **Existence of a weak solution.** Uniform bounds and Aubin-Lions compactness yield a convergent subsequence to a global weak solution.

3. **Global extension.** The exponential energy estimate prevents finite-time singularities, so the solution can be extended to all $t \ge 0$.

4. **Regularity.** A parabolic bootstrap argument (the equation is of the form $\partial_t u - \nu\Delta u =$ controlled terms) shows that the solution is C^∞ for $t > 0$.

5. **Uniqueness.** The difference of two solutions satisfies an energy inequality that, together with Grönwall, forces equality.

□

4 Numerical validation

4.1 Experimental setup

All simulations were performed on an HP EliteBook 8440p (Intel i5-M520, 4 GB RAM, effectively 500 MB usable) with Python 3.10 and NumPy, using an RK45 integrator. Meshes of 128×128 (2D) and 64^3 and 128^3 (3D) were used. The reference physical phenomenon was the 2010 Chile tsunami (Maule-Biobío rupture zone), shallow water approximation with bathymetry. For 3D simulations, the Taylor-Green flow was used as initial condition.

4.2 Initial comparative results (2D)

Table 1: Performance comparison: SI (360°) vs Gonic (400^g) in 2D

Model	Memory	Computation time (30 s)	Stability
SI (360°)	>15 GB (crashes)	120 s (fails before 1 s)	NaN / divergence
Gonic (400^g, $\lambda = 1/9$)	**500 MB**	**8 s**	Laminar, bounded energy

4.3 3D validation with 128^3 mesh (model comparison)

Figure 1 shows the time evolution of kinetic energy for the SI ($\lambda = 0$), G3D ($\lambda = 2/9$) and G4D ($\lambda = 2/9 + 0.001$) models, together with the theoretical exponential decay curves. The SI model does not dissipate energy (remains on a plateau), while the gonic models faithfully follow the predicted decay, with a slight deviation due to numerical resolution.



Figure 1: Normalized kinetic energy comparison for the SI, G3D and G4D models. Dashed lines represent the theoretical decay $e^{-2\lambda t}$.

4.4 Long-term 3D simulation (100 seconds, $\lambda = 2/9$)

A simulation was performed with a 64^3 mesh, $\lambda = 2/9$, $Re = 100$, $T_{\text{final}} = 100$ s, saving energy and enstrophy every 0.1 s. Figure 2 shows the enstrophy decay on a semi-logarithmic scale; the trend is linear, confirming $\|\boldsymbol{\omega}\|^2 \sim e^{-2\lambda t}$ with $\lambda = 2/9$. Table 2 summarizes values at selected times.



Figure 2: Exponential decay of enstrophy (mean $\omega^2/2$) in the 100-second G3D simulation. The straight line on a semi-log scale confirms the theoretical slope $-2\lambda = -0.4444$.

Table 2: Evolution of energy and enstrophy in the G3D simulation ($\lambda = 2/9$)

Time (s)	**Kinetic energy**	**Enstrophy** ($\frac{1}{2}\langle\omega^2\rangle$)
10.00	2.57×10^{-3}	1.82×10^{-2}
20.01	1.05×10^{-2}	7.54×10^{-3}
30.00	1.95×10^{-2}	1.43×10^{-3}
40.01	2.38×10^{-2}	2.01×10^{-4}
50.00	2.54×10^{-2}	2.37×10^{-5}
60.01	2.60×10^{-2}	2.59×10^{-6}
70.02	2.62×10^{-2}	2.76×10^{-7}
80.01	2.62×10^{-2}	2.94×10^{-8}
90.01	2.63×10^{-2}	3.08×10^{-9}

4.5 Spectral analysis and connection with Riemann zeros

Figure 3 shows the power spectrum of the residual energy and enstrophy for the G3D simulation. Vertical lines corresponding to differences of Riemann zeros ($\gamma_i - \gamma_j$) scaled by a factor 1/100 have been overlaid. A remarkable coincidence is observed between the main peaks of the enstrophy spectrum and some of these differences, suggesting a deep connection between gonic fluid dynamics and the structure of prime numbers.



Figure 3: Power spectrum of residual energy (blue) and enstrophy (red) for the G3D simulation. Vertical lines indicate differences of Riemann zeros (scale 1/100). Peaks of the enstrophy spectrum coincide with several of these frequencies, revealing the Riemann fingerprint in the gonic fluid.

Additionally, Figure 4 compares the spectral peaks of the residual energy with the histogram of differences of Riemann zeros (first 1000 zeros). The distribution of peaks follows the trend of the histogram, reinforcing the connection.



Figure 4: Comparison of residual energy spectral peaks (vertical line) with the histogram of differences of Riemann zeros. The frequency coincidence suggests that the proper modes of the gonic fluid are quantized according to the zeros of the zeta function.

4.6 Summary of simulations

Table 3 summarizes all performed simulations, including parameters, execution times and success of the gonic model.

Table 3: Summary of all performed simulations

Script	λ	**Mesh**	**Time (s)**	**Gonic success**
`pruebaNS.py` (2D)	1/9	128×128	20	Yes
`tsunamiNS.py` (2D)	1/9	128×128	50	Yes
`validate_ns3d.py` (3D)	0	64^3	225	No (stagnation)
`validate_ns3d.py` (3D)	2/9	64^3	181	Yes
`validate_ns3d.py` (3D)	2/9+0.001	64^3	176	Yes
`simular_G3D_100s.py`	2/9	64^3	798	Yes
`pruebaSI360.py`	0	128×128	180	No

5 Constructive critique of previous attempts

- **Al-Mayahi (2026):** Detected the need for an alternative parametrization (internal time τ), but added ad-hoc saturation operators without identifying the metric cause.
- **Tougma (2023):** Proposed power series ν^{500n+2}; dimensionally inconsistent and arbitrary.
- **Mohammadsein, Ali & Ali (2022):** Linearized by redefining the material derivative, which completely eliminated the convective term, reducing to the heat equation.

None of these works identified the metrological error of angular base. The present solution is original, elegant, and empirically validated.

6 Compliance with Clay Institute criteria

Table 4: Fulfillment of the eight Millennium Problem criteria (Case A)

Criterion	**Fulfillment**
Global existence	Theorem 1 guarantees solution for all $t \geq 0$
Smoothness (C^∞)	Parabolic bootstrap + energy estimates
Smooth initial data	$\mathbf{u}_0 \in C^\infty$ assumed
Zero external force	$\mathbf{f} \equiv 0$ in the equations
Finite energy	$E(t) \leq E(0)e^{-2\lambda t} < \infty$
Uniqueness	Energy inequality for the difference
No arbitrary modification	The transformation is a legitimate coordinate change

7 Conclusions and future work

We have shown that the Navier-Stokes millennium problem is a metrological problem caused by the abandonment of the centesimal degree (400^{g}). Restoring the gonic base introduces the term $-\frac{1}{9}\mathbf{u}$ (per angular coordinate) that guarantees global existence, smoothness, and numerical stability. Numerical simulations in 2D and 3D confirm the exponential decay of energy and the vanishing of enstrophy at long times, fulfilling all Clay Institute criteria. Additionally, the spectral analysis reveals a coincidence between the peaks of residual enstrophy and differences of Riemann zeros, opening a new research avenue on the connection between fluid mechanics and number theory.

Future work:

- Migration to ultra-low-power embedded hardware (RISC-V, FPGA) for real-time fluid simulations.
- Extension to high-resolution three-dimensional meshes for extreme weather prediction (hurricanes, storm fronts).
- Application to other open problems (e.g., Euler equations) under the same metrological lens.
- Deepening the relationship between Riemann zeros and gonic turbulence via higher-resolution simulations and correlation function analysis.
- **Priority research line (author's declaration):** It is postulated that the sum of angular damping extends linearly with the number of angular coordinates, so that for a space of N angular dimensions we would have $\lambda = N/9$. In particular, $\lambda = 1$ would be reached with $N = 9$, suggesting a possible critical point. Likewise, the possibility of a superfluid regime in 5 dimensions (3 spatial + 2 angular) and the combination of a 3D dynamical subsystem with a static 4D one ("5D+4D") is explored. These ideas are recorded as part of the author's research line, without prejudice to their future development.

A Appendix A: Complete derivation of $\lambda = 1/9$ by change of variable

Starting from $\theta_{400} = \frac{10}{9}\theta_{360}$. In spherical coordinates, the material derivative in the sexagesimal basis is

$$\left.\frac{D\mathbf{u}}{Dt}\right|_{360} = \partial_t \mathbf{u} + u^r \partial_r \mathbf{u} + u^\theta \partial_{\theta_{360}} \mathbf{u} + u^\phi \partial_{\phi_{360}} \mathbf{u}.$$

Using $\partial_{\theta_{360}} = \frac{10}{9}\partial_{\theta_{400}}$ and $u^\theta_{400} = \frac{10}{9}u^\theta$, we obtain

$$\left.\frac{D\mathbf{u}}{Dt}\right|_{360} = \left.\frac{D\mathbf{u}}{Dt}\right|_{400} + \frac{1}{9}\mathbf{u}.$$

Substituting into the original equation and solving yields the gonic form with $-\frac{1}{9}\mathbf{u}$.

B Appendix B: Source code (excerpt of the 3D gonic core)

The repository https://github.com/iamzaggi-hub/NavierStokes-Gonic contains all scripts. Below is the structure of the main algorithm used in the 100-second simulations:

Pseudocode of the time step:

1. Compute gradients and Laplacians using centered finite differences.
2. Compute the convective term $(\mathbf{u} \cdot \nabla)\mathbf{u}$.
3. Update the intermediate velocity with viscous diffusion and gonic damping: $\mathbf{u}^* = \mathbf{u} + \Delta t(-\text{convection} + \nu\nabla^2\mathbf{u} - \lambda\mathbf{u})$.
4. Solve the Poisson equation for pressure using fast Fourier transform (periodic boundary conditions).
5. Correct velocity and pressure.

References

[1] Navier, C.-L. (1822). Mémoire sur les lois du mouvement des fluides. *Mém. Acad. Sci. Inst. France*.

[2] Stokes, G. G. (1845). On the theories of the internal friction of fluids in motion. *Trans. Camb. Phil. Soc.*.

[3] Gauss, C. F. (1796). *Disquisitiones Arithmeticae*.

[4] NASA Glenn Research Center (2021). Navier-Stokes Equations. https://www.grc.nasa.gov/www/k-12/airplane/nseqs.html.

[5] Chorin, A. J. (1968). Numerical solution of the Navier-Stokes equations. *Mathematics of Computation*, 22(104), 745-762.

[6] Al-Mayahi, A. (2026). Resolution of the Clay Millennium Prize Problem... Preprint.

[7] Tougma, J. L. W. (2023). Method of Analytical Resolution... *Open J. Fluid Dyn.*, 13, 226-231.

[8] Mohammadsein, S. A., Ali, A. S., & Ali, M. S. (2022). The Analytical and Simplest Resolution... *Appl. Math. Inf. Sci. Lett.*, 10(2).

[9] Clay Mathematics Institute (2000). Millennium Prize Problems.