

# Laboratory 8 - Linear Algebra

Ian Hanus (ih52)  
Lab Section 1B, Tuesday 8:30-11:20  
5 November 2017

I understand and have adhered to all the tenets of the Duke Community Standard in completing every part of this assignment. I understand that a violation of any part of the Standard on any part of this assignment can result in failure of this assignment, failure of this course, and/or suspension from Duke University.

---

## Contents

<b>1 Palm Problem 8.1</b>	<b>2</b>
<b>2 Based on Chapra Problem 8.3</b>	<b>2</b>
<b>3 Based on Chapra Problem 8.10</b>	<b>3</b>
<b>4 Palm 8.5(b)</b>	<b>3</b>
<b>5 Based on Palm 8.9</b>	<b>3</b>
<b>6 Based on Palm 8.16(a)</b>	<b>3</b>
<b>A Codes and Output</b>	<b>4</b>
A.1 Chapra83.m . . . . .	4
A.2 Chapra810.m . . . . .	4
A.3 Palm85 . . . . .	5
A.4 Palm89 . . . . .	5
A.5 Palm816 . . . . .	6
<b>B Figures</b>	<b>7</b>

## List of Figures

1 Plot of Palm 8.5(b) . . . . .	7
2 Plot of Polynomial for Points (1,4), (4,73), (5,120) . . . . .	7
3 Plot of Polynomial for Points (1,4), (4,-73), (5,120) . . . . .	8
4 Plot of Polynomial for Points (1,4), (4,73), (5,-120) . . . . .	8

## 1 Palm Problem 8.1

Part	x	y	z
a	2.4762e+00	4.7619e-02	N/A
b	-1.1818e+00	1.0909e+00	N/A
c	3.4329e+00	-6.0390e+00	4.4935e+00
d	2.0035e+00	-2.6848e+00	5.2312e+00

## 2 Based on Chapra Problem 8.3

Equations in matrix form:

$$\begin{bmatrix} 0 & -7 & 5 \\ 0 & 4 & 7 \\ -4 & 3 & -7 \end{bmatrix} \begin{Bmatrix} x_1 \\ x_2 \\ x_3 \end{Bmatrix} = \begin{Bmatrix} 50 \\ -30 \\ 40 \end{Bmatrix}$$

Solutions for this system:

$$\begin{Bmatrix} x_1 \\ x_2 \\ x_3 \end{Bmatrix} = \begin{Bmatrix} -1.5181e + 01 \\ -7.2464e + 00 \\ -1.4493e - 01 \end{Bmatrix}$$

Transpose of A:

$$\begin{bmatrix} 0 & 0 & -4 \\ -7 & 4 & 3 \\ 5 & 7 & -7 \end{bmatrix}$$

Inverse of A:

$$\begin{bmatrix} -1.7754e - 01 & -1.2319e - 01 & -2.5000e - 01 \\ -1.0145e - 01 & 7.2464e - 02 & 0 \\ 5.7971e - 02 & 1.0145e - 01 & 0 \end{bmatrix}$$

Condition numbers:

$$\begin{aligned} \text{1-norm condition} &= 6.4022e + 00 \\ \text{2-norm condition} &= 4.0569e + 00 \\ \text{Frobenius condition} &= 5.4293e + 00 \\ \infty\text{-condition} &= 7.7101e + 00 \end{aligned}$$

The condition numbers signify how inaccurate the solution can be. Because the condition for each of the norms is greater than one, the system is ill-conditioned. This suggests that the solutions of the system could exhibit rounding errors. Because the magnitude of each of the condition numbers is below ten, the last digit of the solutions could exhibit roundoff errors as the logarithm of the condition numbers is between 0 and 1.

### 3 Based on Chapra Problem 8.10

$$\begin{bmatrix} -8.6603e-01 & 0 & 5.0000e-01 & 0 & 0 & 0 \\ -5.0000e-01 & 0 & -8.6603e-01 & 0 & 0 & 0 \\ 8.6603e-01 & 1.0000e+00 & 0 & 1.0000e+00 & 0 & 0 \\ 5.0000e-01 & 0 & 0 & 0 & 1.0000e+00 & 0 \\ 0 & -1.0000e+00 & -5.0000e-01 & 0 & 0 & 0 \\ 0 & 0 & 8.6603e-01 & 0 & 0 & 1.0000e+00 \end{bmatrix} \begin{Bmatrix} F_1 \\ F_2 \\ F_3 \\ H_2 \\ V_2 \\ V_3 \end{Bmatrix} = \begin{Bmatrix} 0 \\ 1000 \\ 0 \\ 0 \\ 0 \\ 0 \end{Bmatrix}$$

1 F1: -5.000e+02  
2 F2: +4.330e+02  
3 F3: -8.660e+02  
4 H2: +0.000e+00  
5 V2: +2.500e+02  
6 V3: +7.500e+02

### 4 Palm 8.5(b)

There is a c value with easily predicted solutions. For a c value of 0, x,y, and z are expected to be zero. This seems appears to be true on the graph. There are 201 points instead of 200 because using 200 creates an even spacing of 0.1 between all points. This would include the c value zero. Using 201 points, the spacing is different and does not include a calculated c value of zero.

### 5 Based on Palm 8.9

$$\begin{bmatrix} -1 & 1/3 & 1/3 & 0 \\ 1/2 & -1 & 0 & 1/2 \\ 1/3 & 0 & -1 & 1/3 \\ 0 & 1/2 & 1/2 & -1 \end{bmatrix} \begin{Bmatrix} T_1 \\ T_2 \\ T_3 \\ T_4 \end{Bmatrix} = \begin{Bmatrix} -50 \\ 0 \\ -20/3 \\ 0 \end{Bmatrix}$$

The following are the temperatures at each square.

1 T1: +1.027e+02 C  
2 T2: +9.091e+01 C  
3 T3: +6.727e+01 C  
4 T4: +7.909e+01 C

### 6 Based on Palm 8.16(a)

$$\begin{bmatrix} x_1^2 & x_1 & 1 \\ x_2^2 & x_2 & 1 \\ x_3^2 & x_3 & 1 \end{bmatrix} \begin{Bmatrix} a \\ b \\ c \end{Bmatrix} = \begin{Bmatrix} y_1 \\ y_2 \\ y_3 \end{Bmatrix}$$

The following are the coefficients of the quadratics containing the given points.

Points	a	b	c
(1,4), (4,73), (5,120)	6.00e+00	-7.00e+00	5.00e+00
(1,4), (4,-73), (5,120)	5.47e+01	-2.99e+02	2.48e+02
(1,4), (4,73), (4,120)	N/A	N/A	N/A
(1,4), (4,73), (5,-120)	-5.40e+01	2.93e+02	-2.35e+02

## A Codes and Output

### A.1 Chapra83.m

```
1 % I have adhered to all the tenets of the
2 % Duke Community Standard in creating this code.
3 % Signed: [ih52]
4 %% Initialize workspace
5 clear; format short e
6
7 %% Setup matrices
8 A = [0 -7 5 ;0 4 7; -4 3 -7]
9 b = [50; -30; 40]
10
11 %% Solutions for the system
12 Solutions = A\b
13
14 %% Transpose and inverse of A
15 ATrans = A'
16 AInv = inv(A)
17
18 %% Calculate the condition numbers
19 Norm1Cond = cond(A,1)
20 Norm2Cond = cond(A,2)
21 NormFroCond = cond(A,'fro')
22 NormInfCond = cond(A,inf)
```

### A.2 Chapra810.m

```
1 % I have adhered to all the tenets of the
2 % Duke Community Standard in creating this code.
3 % Signed: [ih52]
4 %% Initialize workspace
5 clear; format short e
6
7 %% Rewrite equations in matrix form
8 A = [-cosd(30) 0 cosd(60) 0 0 0
9      -sind(30) 0 -sind(60) 0 0 0
10      cosd(30) 1 0 1 0 0
11      sind(30) 0 0 0 1 0
12      0 -1 -cosd(60) 0 0 0
13      0 0 sind(60) 0 0 1]
14 b = [0; 1000; 0; 0; 0; 0]
15
16 %% Solve for unknowns
17 Solutions = A\b;
18
19 %% Save in text file
20 FID = fopen('TrussData.txt', 'w');
21
22 fprintf(FID, 'F1: %+1.3e \n', Solutions(1))
23 fprintf(FID, 'F2: %+1.3e \n', Solutions(2))
24 fprintf(FID, 'F3: %+1.3e \n', Solutions(3))
25 fprintf(FID, 'H2: %+1.3e \n', Solutions(4))
26 fprintf(FID, 'V2: %+1.3e \n', Solutions(5))
27 fprintf(FID, 'V3: %+1.3e \n', Solutions(6))
28
29 fclose(FID)
```

### A.3 Palm85

```
1 % I have adhered to all the tenets of the
2 % Duke Community Standard in creating this code.
3 % Signed: [ih52]
4 %% Initialize workspace
5 clear; format short e
6
7 %% Create coefficient matrix A
8 A = [1 -5 -2; 6 3 1; 7 3 -5];
9
10 %% Create c values for plotting
11 c = linspace(-10,10,201);
12
13 %% Loop through different c values
14 for k = 1:length(c)
15     b = [11*c(k); 13*c(k); 10*c(k)];
16     MyVals = A\b
17     x(k) = MyVals(1);
18     y(k) = MyVals(2);
19     z(k) = MyVals(3);
20 end
21
22 %% Make and save plot
23 plot(x,c,'k-',y,c,'c-.',z,c,'m--')
24 legend('x-values','y-values','z-values')
25 title('Plot of Palm 8.5(b)')
26 xlabel('c value')
27 ylabel('Solutions value')
28 print -depsc Palm85bPlot
```

### A.4 Palm89

```
1 % I have adhered to all the tenets of the
2 % Duke Community Standard in creating this code.
3 % Signed: [ih52]
4 %% Initialize workspace
5 clear; format short e
6
7 %% Create matrices
8 A = [-1 1/3 1/3 0; 1/2 -1 0 1/2; 1/3 0 -1 1/3; 0 1/2 1/2 -1]
9 b = [-1/3*150; 0; -1/3*20; 0]
10
11 %% Find temperatures
12 Temps = A\b
13
14 %% Print and save results
15 FID = fopen('TempData.txt','w')
16
17 fprintf(FID,'T1: %+1.3e C\n', Temps(1))
18 fprintf(FID,'T2: %+1.3e C\n', Temps(2))
19 fprintf(FID,'T3: %+1.3e C\n', Temps(3))
20 fprintf(FID,'T4: %+1.3e C\n', Temps(4))
21
22 fclose(FID)
```

## A.5 Palm816

```
1 % I have adhered to all the tenets of the
2 % Duke Community Standard in creating this code.
3 % Signed: [ih52]
4 %% Create function
5 function [a, b, c] = findquad(x,y,flag)
6
7 %% Check user inputs
8 if nargin < 2
9     error('Not enough arguments.')
```

10 end

11 if nargin == 2

12 flag = 0;

13 end

14

15 %% Create matrices

16 Amat = [x(1)^2, x(1), 1;x(2)^2, x(2), 1;x(3)^2, x(3), 1];

17 bmat = [y(1); y(2); y(3)];

18

19 %% Check condition number

20 if cond(Amat,2) > 10^5

21 error('Matrix is ill-conditioned')

22 end

23

24 %% Solve for a, b, & c

25 Solutions = Amat\bmat;

26 a = Solutions(1)

27 b = Solutions(2)

28 c = Solutions(3)

29

30 %% Plot of polynomial

31 if flag == 1

32 MyPoly = @(x) a\*x.^2+b\*x+c;

33 Xmin = min(x) - 0.1\*(max(x)-min(x));

34 Xmax = max(x) + 0.1\*(max(x)-min(x));

35 Xvals = linspace(Xmin,Xmax,100);

36 plot(Xvals,MyPoly(Xvals),'k-')

37 hold on

38 for k = 1:3

39 plot(x(k),y(k),'bs','LineWidth',3,'MarkerSize',12,'MarkerFaceColor','g')

40 end

41 hold off

42 grid on

43 xlabel('x')

44 ylabel('y')

45 title(sprintf('Graph of y =%0.2ex^2%+0.2ex%+0.2e',a,b,c))

46 end

47 print -depsc Palm816fig3

## B Figures




Figure 1: Plot of Palm 8.5(b)




Figure 2: Plot of Polynomial for Points (1,4), (4,73), (5,120)




Figure 3: Plot of Polynomial for Points (1,4), (4,-73), (5,120)




Figure 4: Plot of Polynomial for Points (1,4), (4,73), (5,-120)