

# INVESTIGATING THE STRUCTURE OF $M_2(\mathbb{F}_q)$

IAN GALLAGHER  
DR. ERIC BRUSSEL

ABSTRACT. We present and prove a count of the maximal commutative subalgebras of  $M_2(\mathbb{F}_q)$ , as well as counts for the individual isomorphism classes.

## 1. INTRODUCTION

For a finite field of  $q = p^m$  elements,  $p \neq 2$ , it is possible to count the number of vector subspaces of  $\mathbb{F}_q^n$  of a given dimension. These counts arise in problems involving the number of points of  $\mathbb{P}_q^n$ , the Grassmannian, and further generalizations.

This paper is meant to address a similar problem. Namely, the structure and count of the *maximal commutative* subalgebras of  $M_2(\mathbb{F}_q)$ , the vector space  $\mathbb{F}_q^4$  with matrix multiplication. Such commutative subalgebras are themselves two dimensional vector subspaces with the inherited multiplication of  $M_2(\mathbb{F}_q)$ . We find that there are three isomorphism classes of maximal commutative subalgebras and give explicit formulas for the size of the isomorphism classes based on the size of  $\mathbf{q}$ .

For the duration of the paper,  $q = p^n$  for some prime  $p \neq 2$ .

## 2. GENERALIZED QUATERNIONS AND $M_2(\mathbb{F}_q)$

Here, we introduce an alternative formulation of  $M_2(\mathbb{F}_q)$  based on the generalized quaternions, an abstraction of the well-known quaternion algebra. This formulation lends itself nicely to a more general result detailing the structure of commutative subalgebras of  $M_2(\mathbb{F}_q)$ .

**Definition 1** (Generalized Quaternions). Let  $\mathbb{F}$  be a field,  $\text{char } \mathbb{F} \neq 2$ , and fix  $a, b \in \mathbb{F} - \{0\}$ . Then define the *generalized quaternions* as

$$A_{a,b}(\mathbb{F}) = \{t + xi + yj + zk : t, x, y, z \in \mathbb{F}\}$$

where  $i^2 = a, j^2 = b$ , and  $ij = -ji = k$

The generalized quaternions form what is known as a central simple algebra over  $\mathbb{F}$ . That is, it is a finite dimensional  $\mathbb{F}$ -algebra with center  $\mathbb{F}$  and no nontrivial two-sided ideals. [possibly insert proof that the generalized quaternions are a csa here](#)

**Definition 2** (Generalized Quaternion Norm).

$$N_{a,b} : A_{a,b}(\mathbb{F}) \rightarrow \mathbb{F}$$
$$q \mapsto q\bar{q} = t^2 - ax^2 - by^2 + abz^2$$

It turns out that  $M_2(\mathbb{F}_q)$  is the only isomorphism class of generalized quaternions over  $\mathbb{F}_q$  but this fact is not immediately obvious. We will use the following two results to prove this.

**Theorem 3** (Wedderburn [3, pg. 854]). *Let  $k$  be a field and  $A$  a central simple algebra. Then  $A \simeq M_n(D)$  for some  $k$ -division algebra  $D$ .*

**Theorem 4** (Chevalley-Warning [2, pg. 5]). *Let  $f_\alpha \in \mathbb{F}_q[X_1, \dots, X_n]$  be polynomials in  $n$  variables such that  $\sum_\alpha \deg f_\alpha < n$  and let  $V$  be the set of their common zeros in  $\mathbb{F}_q^n$ . One has*

$$|V| \equiv 0 \pmod{p}$$

With these theorems, we now give a proof of the structure of the generalized quaternions over  $\mathbb{F}_q$ .

**Theorem 5.**  $A_{a,b}(\mathbb{F}_q) \cong M_2(\mathbb{F}_q)$

*Proof.* We first make use of Theorem 4. Let

$$\begin{aligned} f &= N_{a,b}(X_1 + iX_2 + jX_3 + kX_4) \\ &= X_1^2 - aX_2^2 - bX_3^2 + abX_4^2 \\ &\in \mathbb{F}_q[X_1, X_2, X_3, X_4] \end{aligned}$$

Here,  $\deg f = 2 < 4$ , so Chevalley-Warning holds and  $|V| \equiv 0 \pmod{p}$ , where  $V$  is the set of zeros of  $f$ . Note that  $f(0, 0, 0, 0) = 0$ , so we must have  $p$  divides  $|V|$ . Therefore,  $f$  has a nontrivial zero.

It follows that there exists  $q \in A_{a,b}(\mathbb{F}_q)$ ,  $q \neq 0$ , such that  $N_{a,b}(q) = q\bar{q} = 0$ . Therefore,  $q$  is a nontrivial zero divisor and  $A_{a,b}(\mathbb{F}_q)$  is not a  $\mathbb{F}_q$ -division algebra. By Theorem 3 we must have  $A_{a,b}(\mathbb{F}_q) \cong M_n(D)$  for some  $\mathbb{F}_q$ -division algebra  $D$  and natural number  $n$ . Since  $A_{a,b}(\mathbb{F}_q)$  is not a division algebra, we know  $n \neq 1$ .  $\dim_{\mathbb{F}_q} A_{a,b}(\mathbb{F}_q) = 4$ , so  $n$  must be 2 and  $D$  must have  $\mathbb{F}_q$ -dimension 1, so is isomorphic to  $\mathbb{F}_q$ . Therefore,  $A_{a,b}(\mathbb{F}_q) \cong M_2(\mathbb{F}_q)$ .  $\square$

### 3. PLANES IN $M_2(\mathbb{F}_q)$

**Definition 6.** For a matrix  $A \in M_2(\mathbb{F}_q)$ , define

- (1)  $A$ 's characteristic polynomial  $p_A(X) = X^2 - \text{tr}(A)X + \det(A)$ .
- (2)  $A$ 's minimal polynomial  $m_A(X)$  is the unique monic polynomial of smallest degree such that  $m_A(A) = 0$ .

By the Cayley-Hamilton Theorem [3, pg. 478], we know that  $m_A(X)$  divides  $p_A(X)$ . So for nontrivial  $A \in M_2(\mathbb{F}_q)$ , we observe that if  $A \in \mathbb{F}_q$ , then  $m_A(X) = X - A$ . Otherwise, no minimal polynomial of degree 1 exists, and we have  $m_A(X) = p_A(X)$ .

Define the evaluation map  $\varepsilon_A : \mathbb{F}_q[X] \rightarrow M_2(\mathbb{F}_q)$  taking  $f(X)$  to  $f(A)$ , which is a  $\mathbb{F}_q$ -algebra homomorphism with image  $\mathbb{F}_q[A] \subseteq M_2(\mathbb{F}_q)$ . The kernel is nontrivial, since  $M_2(\mathbb{F}_q)$  has finite  $\mathbb{F}_q$ -dimension and  $\mathbb{F}_q[X]$  does not. Since  $\mathbb{F}_q[X]$  is a principal ideal domain, it follows that  $\ker(\varepsilon_A) = (m_A(X))$ . [Include more about kernel being an ideal here?](#) By the First Isomorphism Theorem,

$$\frac{\mathbb{F}_q[X]}{(m_A(X))} \cong \mathbb{F}_q[A]$$

If  $A \in M_2(\mathbb{F}_q) - \mathbb{F}_q$ , we know  $m_A(X) = p_A(X)$  is degree 2 and  $\mathbb{F}_q[A]$  has  $\mathbb{F}_q$ -dimension 2.

$A$  then generates the two dimensional commutative subalgebra of  $M_2(\mathbb{F}_q)$  containing  $A$ . All possible 2 dimensional commutative subalgebras are therefore of the form

$$\frac{\mathbb{F}_q[X]}{(m_A(X))}$$

for some  $A \in M_2(\mathbb{F}_q) - \mathbb{F}_q$ . The set of two dimensional commutative subalgebras of  $M_2(\mathbb{F}_q)$  must then be composed of exactly three isomorphism classes based on  $m_A(X)$

- (1) If  $m_A(X)$  is irreducible, define  $\mathbb{F}_q[A] \cong \mathbb{F}_{q^2}$ .
- (2) If  $m_A(X)$  is reducible but not separable, define  $\mathbb{F}_q[A] \cong \mathbb{F}_q \times \mathbb{F}_q$ .
- (3) If  $m_A(X)$  is separable, define  $\mathbb{F}_q[A] \cong \mathbb{F}_{q_{nil}}^2$ .

The commutative subalgebra generated when  $m_A(X)$  is separable can be considered a degenerate case due to the presence of nontrivial nilpotent elements which square to 0. Since  $m_A(X)$  is separable and reducible, we have  $m_A(X) = (X - x)^2$  for some  $x \in \mathbb{F}_q$ . By the definition of the minimal polynomial, we must have that  $(A - x)^2 = 0$ .  $A$  is not an element of  $\mathbb{F}_q$ , so  $A - x$  is nonzero and is therefore a nontrivial nilpotent element. It is worth noting that the case that  $m_A(X)$  is both separable and irreducible does not require consideration since finite fields are perfect.

#### 4. IDENTIFYING MAXIMAL COMMUTATIVE SUBALGEBRAS

we have shown that each matrix  $A \in M_2(\mathbb{F}_q) - \mathbb{F}_q$  generates a commutative subalgebra containing it,  $\mathbb{F}_q[A]$ , and that this plane belongs to one of three given isomorphism classes. Since there exists subalgebras of dimension three, such as the set of upper triangular matrices, it is reasonable to ask whether this is the maximal commutative subalgebra of  $M_2(\mathbb{F}_q)$  containing  $A$ .

To prove that there are no commutative subalgebras of dimension 3, we return to the generalized quaternions formulation. We first introduce a useful vector notation for the generalized quaternions.

**Definition 7** (Generalized Quaternions, Vector Notation). Let  $p = (s, \mathbf{v})$ , and  $q = (t, \mathbf{w})$ , using the vector notation  $\mathbf{v} = \langle v_1i, v_2j, v_3l \rangle$  and  $\mathbf{w} = \langle w_1i, w_2j, w_3l \rangle$  with coefficients in  $A_{a,b}(k)$ . Modified dot and cross product formulas compute

$$\begin{aligned} \mathbf{v} \cdot_{a,b} \mathbf{w} &= v_1w_1a + v_2w_2b - av_3w_3 \\ \mathbf{v} \times_{a,b} \mathbf{w} &= \left\langle \begin{vmatrix} v_2j & v_3l \\ w_2j & w_3l \end{vmatrix}, \begin{vmatrix} v_1i & v_3l \\ w_1i & w_3l \end{vmatrix}, \begin{vmatrix} v_1i & v_2j \\ w_1i & w_2j \end{vmatrix} \right\rangle \\ &= \langle (-v_2w_3 + v_3w_2)bi, (v_1w_3 - v_3w_1)aj, (v_1w_2 - v_2w_1)l \rangle \end{aligned}$$

Compare this to  $\langle v_1, v_2, v_3 \rangle \times \langle w_1, w_2, w_3 \rangle = \langle v_2w_3 - v_3w_2, -v_1w_3 + v_3w_1, v_1w_2 - v_2w_1 \rangle$ . Thus

$$\begin{aligned} pq &= (st + v_1w_1a + v_2w_2b - av_3w_3) + (sw_1 + tv_1 - v_2w_3b + v_3w_2b)i \\ &\quad + (sw_2 + tv_2 + v_1w_3a - v_3w_1a)j + (sw_3 + tv_3 + v_1w_2 - v_2w_1)l \\ &= (st + \mathbf{v} \cdot \mathbf{w}, s\mathbf{w} + t\mathbf{v} + \mathbf{v} \times \mathbf{w}) \end{aligned}$$

**Lemma 8.** Let  $p = (0, \mathbf{v})$ , and  $q = (0, \mathbf{w})$ . Then  $pq = qp$  if and only if  $\mathbf{v} \times_{a,b} \mathbf{w} = 0$  if and only if  $\langle v_1, v_2, v_3 \rangle \times \langle w_1, w_2, w_3 \rangle = 0$ .

*Proof.* We have  $pq - qp = 2\mathbf{v} \times \mathbf{w}$ , and since  $i, j, k$  are linearly independent and  $\text{char } k \neq 2$ ,  $pq = qp$  if and only if  $\mathbf{v} \times \mathbf{w} = 0$

if and only if  $\langle v_1, v_2, v_3 \rangle \times \langle w_1, w_2, w_3 \rangle = 0$ .  $\square$

**Theorem 9.** If  $K \subseteq A_{a,b}(k)$  is a commutative subalgebra, then  $K = k[\mathbf{w}]$  for some pure imaginary  $\mathbf{w} \in A_{a,b}(k)$ , and  $\dim_k K = 1$  or 2.

*Proof.* If  $q = (t, \mathbf{w}) \in K$ , then  $q - t = \mathbf{w}$  is in  $k[q]$ , hence  $k[\mathbf{w}] = k[q] \subseteq K$ . If  $p = (s, \mathbf{v}) \in K$  then since  $pq = qp$ ,  $\mathbf{v}$  is a scalar multiple of  $\mathbf{w}$  by Lemma 8. Therefore  $K = k[\mathbf{w}]$ . Since  $\deg(m_{\mathbf{w}}(X))$  is either 1 or 2, this completes the proof.  $\square$

Now,  $A_{a,b}(\mathbb{F}_q) \cong M_2(\mathbb{F}_q)$ , so Theorem 9 suffices to show that no commutative subalgebra of dimension 3 exists and we may conclude that the three isomorphism classes of commutative planes are the maximal commutative subalgebras of  $M_2(\mathbb{F}_q)$

## 5. PLANE COUNTS

**Theorem 10.**  $M_2(\mathbb{F}_q)$  has  $q^2 + q + 1$  unique nontrivial commutative subalgebras.

*Proof.* Let  $E_x$  for  $x \in M_2(\mathbb{F}_q) - \mathbb{F}_q$  be the commutative subalgebra created by the span of 1 and  $x$ . Label  $\Sigma = \{E_x \mid x \in M_2(\mathbb{F}_q) - \mathbb{F}_q\}$  and set  $N = |\Sigma|$ , the number of unique 2D commutative subalgebras. Let  $x, y \in M_2(\mathbb{F}_q) - \mathbb{F}_q$ .  $E_x, E_y$  are both two dimensional  $\mathbb{F}_q$  algebras so we know that  $E_x \cap E_y$  is also a  $\mathbb{F}_q$ -algebra. By dimension arguments, either  $E_x = E_y$  or  $E_x \cap E_y = \mathbb{F}_q$ . Since  $\bigcup_{E \in \Sigma} E = M_2(\mathbb{F}_q)$ ,  $\{E_x - \mathbb{F}_q \mid x \in M_2(\mathbb{F}_q) - \mathbb{F}_q\}$  is a partition of  $M_2(\mathbb{F}_q) - \mathbb{F}_q$ . The following then holds,

$$\begin{aligned} N(|E_x| - |\mathbb{F}_q|) &= |M_2(\mathbb{F}_q)| - |\mathbb{F}_q| \\ N(q^2 - q) &= q^4 - q \\ N(q - 1) &= q^3 - 1 \\ N &= q^2 + q + 1 \end{aligned}$$

$\square$

**Lemma 11.** Consider the following elements of  $M_2(\mathbb{F}_q)$ .

$$\begin{aligned} x &= \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} \\ y &= \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \end{aligned}$$

Under the conjugation action of  $GL_2(\mathbb{F}_q)$  on  $M_2(\mathbb{F}_q)$ ,

$$\begin{aligned} |\mathcal{O}_x| &= q(q + 1) \\ |\mathcal{O}_y| &= (q + 1)(q - 1) \end{aligned}$$

*Proof.* We may compute the size of the orbits of  $x, y$  by first computing the size of their stabilizers,  $G_x, G_y$ , and then applying the orbit-coset correspondence theorem. Now,

$$\begin{aligned} G_x &= \{A \in GL_2(\mathbb{F}_q) \mid A \cdot x = x\} \\ &= \{A \in GL_2(\mathbb{F}_q) \mid Ax A^{-1} = x\} \\ &= \{A \in GL_2(\mathbb{F}_q) \mid Ax = xA\} \end{aligned}$$

Inevitably, matrices of the form  $sx + tI$  commute with  $x$  for  $s, t \in \mathbb{F}_q$ , so if  $sx + tI \in GL_2(\mathbb{F}_q)$ , then  $sx + tI \in G_x$ . These are also the only possible matrices in  $G_x$  since they are the elements of the maximal commutative subalgebra containing  $x$  (Theorem 9). It now suffices to determine when  $\det(sx + tI) = 0$ .

$$\det(sx + tI) = \begin{vmatrix} t & s \\ s & t \end{vmatrix} = t^2 - s^2$$

Therefore,  $\det(sx + tI) = 0$  when  $t^2 = s^2$ , so when  $t = \pm s$ . We have thus found precisely  $q^2 - 2q + 1 = (q - 1)^2$  elements of  $G_x$ . It follows that,

$$[GL_2(\mathbb{F}_q) : G_x] = \frac{(q^2 - 1)(q^2 - q)}{(q - 1)^2} = q(q + 1)$$

By the orbit coset correspondence theorem [3] we have that  $|\mathcal{O}_x| = q(q + 1)$

Similarly we have  $G_y = \{A \in GL_2(\mathbb{F}_q) | Ay = yA\}$  and the only possible elements of  $G_y$  are those of the form  $sy + tI$  for  $s, t \in \mathbb{F}_q$  where  $sy + tI \in GL_2(\mathbb{F}_q)$ .

$$\det(sy + tI) = \begin{vmatrix} t & 0 \\ s & t \end{vmatrix} = t^2$$

So  $\det(sy + tI) = 0$  if and only if  $t = 0$ . There are then  $q^2 - q$  elements of  $G_y$ . Therefore,

$$[GL_2(\mathbb{F}_q) : G_y] = \frac{(q^2 - 1)(q^2 - q)}{q^2 - q} = q^2 - 1$$

and we have  $|\mathcal{O}_y| = q(q - 1)$  □

**Theorem 12.**

- (1)  $||[\mathbb{F}_{q^2}]|| = \binom{q}{2}$
- (2)  $||[\mathbb{F}_q \times \mathbb{F}_q]|| = \binom{q+1}{2}$
- (3)  $||[\mathbb{F}_{q^n}^2 iI]|| = q + 1$

*Proof.* The elements  $x, y$  from Lemma 11 are the rational canonical forms for all matrices with minimum polynomials  $X^2 - 1$  and  $X^2$  respectively. Note that two matrices in  $M_2(k)$  are conjugate if and only if they have the same rational canonical form [3].

Since  $X^2 - 1 = (X + 1)(X - 1)$  is reducible and not separable over  $\mathbb{F}_q$ , and  $\text{char}(\mathbb{F}_q) \neq 2$ , all of the elements of  $\mathcal{O}_x$  generate an embedding  $K_1$ .

Any embedding  $K_1$  must also contain an element of  $\mathcal{O}_x$ . To see this, pick a nontrivial element  $A \in K_1$  such that  $A$  is orthogonal to the identity. Then  $\text{tr}(A) = 0$ , and  $m_A(X) = p_A(X) = X^2 + \det(A)$ . We picked  $A \in K_1$ , which means that  $m_A(X)$  must also be reducible and not separable. It follows that  $\det(A) = -\lambda^2$  for some  $\lambda \in \mathbb{F}_q$ . Let  $B = A/\lambda \in K_1$  and observe  $\text{tr}(B) = \text{tr}(A)/\lambda = 0$ ,  $\det(B) = \det(A)/\lambda^2 = -\lambda^2/\lambda^2 = -1$ . Therefore,  $m_B(X) = p_B(X) = X^2 - \text{tr}(B)X + \det(B) = X^2 - 1$  as desired.

Furthermore, suppose  $B' \in K_1$  has minimum polynomial  $X^2 - 1$ . Then  $B'$  is orthogonal to the identity as well and  $B' = \alpha B$  for some  $\alpha \in \mathbb{F}_q^\times$  due to  $K_1$  being dimension 2. We have  $-1 = \det(B') = \alpha^2 \det(B) = -\alpha^2$ , so  $\alpha = \pm 1$  and  $B' = \pm B$ . We have therefore shown that each embedding  $K_1$  contains exactly 2 elements of  $\mathcal{O}_x$ . Applying Lemma 11,

$$|[K_1]| = \frac{1}{2}|\mathcal{O}_x| = \frac{q(q+1)}{2} = \binom{q+1}{2}$$

Similarly, since  $X^2$  is separable over  $\mathbb{F}_q$ , all of the elements of  $\mathcal{O}_y$  generate an embedding of  $K_0$ . Any embedding  $K_0$  will always contain exactly  $q - 1$  elements of  $\mathcal{O}_y$ . These are precisely the nontrivial elements of  $K_0$  that are orthogonal to the identity. Applying Lemma 11,

$$|[K_0]| = \frac{1}{q-1}|\mathcal{O}_y| = \frac{q^2-1}{q-1} = q+1$$

It remains to show  $||[K_{-1}]|| = \binom{q}{2}$ . We may use the previous counts and Theorem 10 to do so.

$$\begin{aligned}
 ||[K_{-1}]|| &= q^2 + q + 1 - ||[K_1]|| - ||[K_0]|| \\
 &= q^2 + q + 1 - \binom{q+1}{2} - (q+1) \\
 &= q^2 - \frac{q(q+1)}{2} \\
 &= \frac{q(q-1)}{2} \\
 &= \binom{q}{2}
 \end{aligned}$$

This completes the proof.

**Corollary 13.** *As  $q \rightarrow \infty$ , the ratio  $||[K_{-1}]|| : ||[K_1]|| : ||[K_0]||$  approaches  $1 : 1 : 0$ .*

□

#### REFERENCES

- [1] D. W. Henderson, *A short proof of Wedderburn's theorem*, Amer. Math. Monthly **72** (1965), 385–386.
- [2] Jean-Pierre Serre, *A Course in Arithmetic*, Springer-Verlag, New York, 1973.
- [3] D. S. Dummit and R. M. Foote, *Abstract Algebra*, third edition, John Wiley & Sons, Inc., Hoboken, New Jersey, 2004.