

foieGras an R package for animal movement data: rapid quality control, behavioural estimation and simulation

Ian D. Jonsen^{1,*}, W. James Grecian², and Toby A. Patterson³

¹Department of Biological Sciences, Macquarie University, Sydney, NSW, Australia

²Scottish Oceans Institute, University of St Andrews, KY16 8LB, United Kingdom

³CSIRO Ocean and Atmosphere Research, Hobart, TAS, Australia

*corresponding author, ian.jonsen@mq.edu.au

Abstract

- 1.
- 2.
- 3.
- 4.

Keywords:

1 | Introduction

The R package `foieGras`, pronounced “*fwah grah*”, ...

2 | foieGras overview

The workflow for `foieGras` is deliberately simple, with much of the usual track data processing checks and formatting handled automatically. The main functions are listed in Table 1. When fitting a model, `foieGras` automatically detects the type of tracking data location quality classes designations that are typical of Argos data and that can be added to the data by the researcher for other types of track data. Based on the location quality classes and other, optional information on observation errors contained in the data, `foieGras` chooses an appropriate measurement error model for each observation. This capability allows for combinations of different tracking data types, e.g., Argos and GPS, in a single input data frame and to be fit in a single state-space model.

2.1 | Data preparation

Animal tracking data, consisting of a time-series of location coordinates, can be read into R as a data frame using standard functions such as `read.csv`. The canonical data format for Argos tracks consists of a data frame with 5 columns corresponding to the following named variables: `id` (individual id), `date` (date and time), `lc` (location class), `lon` (longitude), `lat` (latitude). Optionally, an additional 3 columns, `smaj` (semi-major axis), `smin` (semi-minor axis), `eor` (ellipse orientation), providing Argos error ellipse information may be included.

Table 1: Main functions for the R package `foieGras`

Function	Description
<code>fit_mpm</code>	Fit a Move Persistence Model to location data
<code>fit_ssm</code>	Fit a State-Space Model to location data
<code>fmap</code>	Plot fitted/predicted locations on a map with or without a defined projection
<code>grab</code>	Extract fitted/predicted/observed locations from a <code>foieGras</code> model, with or without projection information
<code>osar</code>	Estimate One-Step-Ahead Residuals from a <code>foieGras</code> SSM
<code>sim</code>	Simulate individual animal tracks with Argos LS or KF errors
<code>simfit</code>	Simulate animal tracks from 'fG_ssm' fit objects
<code>plot.fG_mpm</code>	Plot move persistence estimates as 1-D or 2-D (along track) time-series
<code>plot.fG_ssm</code>	Plot the fit of a <code>foieGras</code> SSM to data
<code>plot.fG_osar</code>	Plot One-Step-Ahead Residuals from a <code>foieGras</code> SSM

Other types of track data can be accommodated, for example, by including the `lc` column where all `lc = "G"` for GPS data. In this case, measurement error in the GPS locations is assumed to have a standard deviation of $0.1 \times \text{Argos class 3 locations}$ (approximately 30 m). Other types of track data can be considered in a similar manner (see the package vignette for further details).

2.2 | State-space model fitting - `fit_ssm`

State-space models are fit using `fit_ssm`. There are a large number of options that can be set in `fit_ssm` (see Suppl for details). We focus only the essential options here:

- `data` the input data structured as described in 2.1
- `vmax` a maximum threshold speed (ms^{-1}) to help identify potential outlier locations
- `model` the process model to be used
- `time.step` the prediction time interval (h)

The function first invokes an automated data processing stage where the following occurs: 1) data type (Argos Least-Squares, Argos Kalman Filter/Smoothing, GPS, or General (e.g., processed light-level geolocations, acoustic telemetry, coded VHF telemetry) is determined; 2) datetimes are converted to POSIXt format, chronological order is ensured, and duplicate datetime records are removed; 3) observations occurring less than `min.dt` seconds after a prior observation are removed; 4) a speed filter [`sda` from the `trip` R package; Sumner et al. (2009)] is used to identify potential outlier locations; 5) locations are projected from spherical lon-lat coordinates to planar x,y coordinates in km.

The function then fits a state-space model to the processed data, where the process model (currently, either a continuous-time `rw` or a continuous-time `crw`) is specified by the user and the measurement model(s) are selected automatically (see I. D. Jonsen et al., 2020 for model details). The model is fit by numerical optimization of the likelihood using either the `optim` or `nlm` R function. The R package `TMB`, Template Model Builder (Kristensen et al., 2016), is used to compute the gradient function in C++ via reverse-mode auto-differentiation and the Laplace Approximation is used to integrate out the latent states (random effects). Fits to a single versus multiple individuals are handled automatically, with sequential SSM fits occurring in the latter case. No hierarchical or pooled estimation among individuals is currently available.

61 `fit_ssm` returns a `foieGras` fit object (a nested data frame with class `fG_ssm`). The outer data
62 frame lists the individual id(s), basic convergence information and a list with class `ssm`. This list con-
63 tains dense information on the model parameter and state estimates, predictions, processed data,
64 optimizer results, and other diagnostic and contextual information. Users can extract a simple data
65 frame of SSM fitted (location estimates corresponding to the, typically irregular, observation times)
66 or predicted values (locations predicted at regular `time.step` intervals) using the `grab` function.

67 2.3 | Model checking and visualisation - `osar`, `plot`, `fmap`

68 Before using fitted or predicted locations, a model fit should be checked and visualised to confirm
69 that the model adequately describes the data. In linear regression and a variety of analogous
70 methods, goodness-of-fit can be assessed by calculating standard residuals such as Pearson or
71 deviance residuals. There is no simple way to calculate residuals for latent variable models that have
72 non-finite state-spaces and that may be nonlinear, but they can be computed based on iterative
73 forecasts of the model (Thygesen et al., 2017). The `osar` function computes one-step-ahead
74 (prediction) residuals and uses the `oneStepPredict` function from the TMB R package to make this
75 as efficient as possible. A set of residuals are calculated for the `x` and `y` values corresponding to
76 the fitted values from the SSM and returned as an `fG_osar` object.

77 A generic `plot` method provides an easy way to visualise the `fG_osar` residuals. Time-series plots
78 of the prediction residuals can be used to detect temporal changes in goodness-of-fit. Quantile-
79 quantile plots of residuals against standard normal quantiles can be used to detect departures
80 from normality. Sample autocorrelation function plots of the residuals are useful for detecting auto-
81 correlation not accounted for by the model. Assessing residual autocorrelation can be particularly
82 important as Argos locations, for example, are themselves derived from a time-series model (Lopez
83 et al., 2015) which can introduce additional autocorrelation in the location errors.

84 State-space model fits to data can also be visualised by using the generic `plot` function on an
85 `fG_ssm` data frame. Options exist to plot fitted or predicted values along with observations as
86 either paired, 1-D time-series or as 2-D tracks with confidence intervals or ellipses, respectively.
87 These plots provide a more intuitive and rapid method for assessing SSM fits to data, however,
88 they do not replace the residual diagnostics. Fitted `fG_ssm` data frames can be mapped using the
89 `fmap` function for single or multiple individuals. Estimated tracks can be displayed with or without
90 confidence ellipses, observations, and/or a projection and maps of single tracks can be coloured
91 by date.

92 2.4 | Behavioural estimation - `fit_mpm`

93 The `fit_mpm` function fits a simple move persistence model to estimate a continuous-valued, time-
94 varying latent variable that indexes changes in movement behaviour (I. Jonsen et al., 2019). This
95 variable measures the autocorrelation in speed and direction between consecutive pairs of move-
96 ments such that high values correspond to fast, directed movements at one end of the continuum
97 and low values correspond to slow, tortuous movements at the other end. It's important to note
98 that this approach is unlike hidden Markov models (McClintock & Michelot, 2018; Michelot et al.,
99 2016) and some state-space models (I. D. Jonsen, 2016) as there is no notion of discrete be-
100 havioural states that animals periodically switch between. Nonetheless, move persistence can be
101 used to identify objectively places where animals spend disproportionately more or less time, and
102 with extensions be correlated with environment or other covariates (See Examples 3.x).

103 The move persistence model assumes that locations are absent of measurement error and can
104 occur either irregularly or regularly in time. `fit_mpm` takes either a `fG_ssm` data frame as input or a

data frame with the follow variables: `id`, `date`, `x`, `y`, where `x` and `y` coordinates can be planar `x,y` or spherical `long,lat`. This latter input format allows the model to be fit easily to GPS or other tracking data with negligible measurement error. When the data contain multiple individuals, the default model is fit jointly by assuming all individuals share the same move persistence variance parameter. There is an option to fit the model separately to each individual. The time-series of estimated move persistence with confidence intervals can be visualized by using the generic `plot` function with the resulting `fG_mpm` data frame. Visualization of move persistence along the 2-D tracks can be plotted or mapped by using the `plot` or `fmap` functions, respectively, and supplying both the `fG_mpm` and `fG_ssm` nested data frames. When using `fit_mpm` on, for example, GPS tracking data that do not require state-space filtering, the movement persistence estimates can be extracted from the `fG_mpm` data frame using the `grab` function and subsequently merged with the observed track data for visualization.

2.5 | Simulation - `sim`, `simfit`

Track simulation can be a helpful, yet informal, way of evaluating the degree to which statistical movement models capture essential features of animal movement data (Michelot et al., 2017). Michelot et al. (2016) advocate comparison of simulated tracks from fitted hidden Markov models to the observed tracks as a means of identifying potential weakness in the hidden Markov model formulation. Here, we suggest that the `rw` and `crw` state-space models and the `mpm` model can be fit to track data simulated from different movement processes to evaluate robustness of location and movement persistence estimates to model mis-specification. We illustrate this idea in section 3.x by drawing on flexibility in the `sim` function that allows a variety of movement processes to be simulated.

Simulation is also used frequently to infer habitat availability, e.g., a null model of the distribution of foraging animals in the absence of external drivers, in habitat utilization studies (Hindell et al., 2020; Raymond et al., 2015). The `simfit` function extracts movement parameters from a `fG_ssm` fit object and simulates an arbitrary number of random tracks of the same duration from these parameters. The argument `cpf = TRUE` ensures that the simulated tracks start and end at approximately the same location, thereby simulating a central place forager.

3 | Examples

We illustrate the main capabilities of `foieGras` through a series of examples using real and simulated tracking data. These examples are purely for demonstration purposes and not intended as a comprehensive guide for conducting animal tracking data quality control or analysis with `foieGras`. Complete code for reproducing the examples and in-depth understanding of `foieGras` functions are provided as supplements.

3.1 | Southern Elephant seal - SSM validation with prediction residuals

We use a subadult male southern elephant seal track included in `foieGras` (`sese1`), sourced from the Australian Integrated Marine Observing System (IMOS; data publicly available via imos.aodn.org.au) deployments at Iles Kerguelen in collaboration with the French IPEV and SNO-MEMO programmes. We fit both the `rw` and `crw` models using `fit_ssm` with a speed filter threshold (`vmax`) of 4 ms^{-1} and a 12-h time step. We calculate prediction residuals using `osar`, and then use the generic `plot` method for `osar` residuals to assess and compare the model fits (Fig. 1). The plots of predicted states on top of the observations suggests both models yield similar fits (Fig.

1a,b), however, there are marked trends in the time-series of residuals for the `rw` model fit (Fig. 1c) and the `rw` ACF's reveal consistent positive autocorrelation in the prediction residuals (Fig. 1e). The corresponding `crw` prediction residuals show no apparent trends through time and have relatively little autocorrelation (Fig. 1d,f), implying that the `crw` provides a better fit to the data.

3.2 | Assessing SSM robustness with simulated data

Using the `sim` function, we simulate animal movement tracks with a variety of plausible movement patterns. We use these simulated data to examine the SSM's goodness-of-fit to tracks generated by processes that differ from the `rw` and `crw` process models within the SSM.

3.3 | Southern elephants seals - behavioural inference

Drawing on an expanded version of the data used in 3.1, we quality control and infer movement persistence along five southern elephant seal tracks. These data can be accessed in `foieGras` via the call: `data(sese, package = 'foieGras')`.

3.4 | Simulating tracks from `foieGras` model fits



Figure 1: Selected diagnostic plots for assessing `rw` (a,c,e) and `crw` (b,d,f) state-space model fits to a southern elephant seal track. Top panels (a,b) are plots of predicted states (red; regular 12-h time intervals) and observations (blue) with pre-filtered observations (orange; ignored by the SSM), using the `plot.fG_ssm` function. Panels c,d are time-series plots of the prediction residuals for the x and y coordinates of each fitted state. Panels e,f are autocorrelation functions of the prediction residuals. All residual plots generated using the `plot.fG_osar` function.




Figure 2: Animal tracks simulated from the movement persistence model `mpm` and a 2-state `crw` model.




Figure 3: Inferred move persistence, γ_t , along southern elephant seal tracks. Locations associated with low move persistence (blue) are indicative of slow, undirected movements, whereas high move persistence (yellow) is indicative of faster, directed movements. Plot generated using `fmap`.

4 | Discussion

Acknowledgements

We thank xx, xx for providing data used in the examples, and xx, xx for helpful comments on an earlier draft of this manuscript. We thank Marie Auger-Méthé for contributing code to the movement persistence models. IDJ acknowledges support from a Macquarie University co-Funded Fellowship and from partners: the US Office of Naval Research, Marine Mammal Program (grant N00014-18-1-2405); the Integrated Marine Observing System (IMOS); Taronga Conservation Society; the Ocean Tracking Network; Birds Canada; and Innovasea/VEMCO. TAP was supported by CSIRO Oceans & Atmosphere internal research funding scheme.

Author's Contributions

IDJ developed the R package; IDJ and TAP developed the state-space models and wrote the manuscript.

Data Accessibility

All code mentioned here is provided in the `foieGras` package for R available on CRAN at <https://CRAN.R-project.org/package=foieGras>. The development version of the package is available on GitHub at <https://github.com/ianjonsen/foieGras>. Data used in the examples are available at...

ORCID

Ian D Jonsen <https://orcid.org/0000-0001-5423-6076>
Toby A Patterson <https://orcid.org/0000-0002-7150-9205>

References

- Hindell, M. A., Reisinger, R. R., Ropert-Coudert, Y., Hückstädt, L. A., Trathan, P. N., Bornemann, H., Charrassin, J.-B., Chown, S. L., Costa, D. P., Danis, B. others. (2020). Tracking of marine predators to protect southern ocean ecosystems. *Nature*, 580(7801), 87–92.
- Jonsen, I. D. (2016). Joint estimation over multiple individuals improves behavioural state inference from animal movement data. *Scientific Reports*, 6, 20625.
- Jonsen, I. D., Patterson, T. A., Costa, D. P., Doherty, P. D., Godley, B. J., Grecian, W. J., Guinet, C., Hoenner, X., Kienle, S. S., Robinson, P. W., Votier, S. C., Whiting, S., Witt, M. J., Hindell, M. A., Harcourt, R. G., & McMahon, C. R. (2020). A continuous-time state-space model for rapid quality control of Argos locations from animal-borne tags. *Movement Ecology*, 8, 31.
- Jonsen, I., McMahon, C. R., Patterson, T. A., Auger-Méthé, M., Harcourt, R., Hindell, M. A., & Bestley, S. (2019). Movement responses to environment: Fast inference of variation among southern elephant seals with a mixed effects model. *Ecology*, 100, e02566.
- Kristensen, K., Nielsen, A., Berg, C. W., Skaug, H., & Bell, B. M. (2016). TMB: Automatic differentiation and Laplace approximation. *Journal of Statistical Software*, 70, 1–21.

- Lopez, R., Malardé, J.-P., Danès, P., & Gaspar, P. (2015). Improving Argos Doppler location using multiple-model smoothing. *Animal Biotelemetry*, 3, 32.
- McClintock, B. T., & Michelot, T. (2018). MomentuHMM: R package for generalized hidden Markov models of animal movement. *Methods in Ecology and Evolution*, 9, 1518–1530.
- Michelot, T., Langrock, R., Bestley, S., Jonsen, I. D., Photopoulou, T., & Patterson, T. A. (2017). Estimation and simulation of foraging trips in land-based marine predators. *Ecology*, 98(7), 1932–1944.
- Michelot, T., Langrock, R., & Patterson, T. A. (2016). MoveHMM: An R package for the statistical modelling of animal movement data using hidden Markov models. *Methods in Ecology and Evolution*, 7, 1308–1315.
- Raymond, B., Lea, M.-A., Patterson, T., Andrews-Goff, V., Sharples, R., Charrassin, J.-B., Cottin, M., Emmerson, L., Gales, N., Gales, R., Goldsworthy, S. D., Harcourt, R., Kato, A., Kirkwood, R., Lawton, K., Ropert-Coudert, Y., Southwell, C., Hoff, J. van den, Wienecke, B., ... Hindell, M. A. (2015). Important marine habitat off east Antarctica revealed by two decades of multi-species predator tracking. *Ecography*, 38, 121–129.
- Sumner, M. D., Wotherspoon, S. J., & Hindell, M. A. (2009). Bayesian estimation of animal movement from archival and satellite tags. *PLoS ONE*, 4(10). <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0007324>
- Thygesen, U. H., Albertsen, C. M., Berg, C. W., Kristensen, K., & Neilsen, A. (2017). Validation of ecological state space models using the Laplace approximation. *Environmental and Ecological Statistics*. <https://doi.org/DOI: 10.1007/s10651-017-0372-4>