

Surgical evacuation of the uterus for RPOC is not without morbidity and can result in uterine perforation (1.5%)^{180,191} and Asherman's syndrome.¹⁹² It is, therefore, recommended that surgical evacuation of retained placental tissue should be undertaken or supervised by an experienced clinician. An appropriately trained clinician may consider performing uterine evacuation under direct ultrasound guidance.

Evidence
level 3

A 2002 Cochrane review (assessed as up-to-date in January 2008) addressed treatments for secondary PPH.⁴ No trials were identified which met the review group's inclusion criteria and no recommendations were made regarding effective treatments. Uterotonics, such as misoprostol and ergometrine, have been recommended in the management of secondary PPH, although evidence to support their use is limited.¹⁷⁸ Transcatheter arterial embolisation¹⁹³ and balloon tamponade¹⁹⁴ have been employed in cases of secondary PPH with ongoing bleeding.

7. Risk management

7.1 *Training and preparation: what measures can be taken to ensure optimal management of PPH?*

Every maternity unit should have a multidisciplinary protocol for the management of PPH.



All staff involved in maternity care should receive training in the management of obstetric emergencies, including the management of PPH.



Training for PPH should be multiprofessional and include team rehearsals.



All cases of PPH involving a blood loss of greater than 1500 ml should be the subject of a formal clinical incident review.



To ensure optimal management of PPH, every unit should have a multidisciplinary protocol with which staff should be familiar (see section 5). Updates on the management of obstetric emergencies (including the management of PPH) are a proactive approach to risk management. Skills drills should ensure that all members of staff, including those working in the transfusion laboratory, are aware of their role in the management of PPH. A systematic review¹⁹⁵ of the effectiveness of multidisciplinary simulation training in obstetric emergencies (including PPH) showed that teamwork training in a simulation setting resulted in improvement of knowledge, practical skills, communication and team performance. Training in a simulation centre did not further improve outcome compared with training at a local unit.

Evidence
level 2++

The RCOG recommends that all cases of PPH with an estimated blood loss of more than 1500 ml should be the subject of a formal clinical incident review.¹⁹⁶

Evidence
level 4

7.2 Documentation

Accurate documentation of a delivery with PPH is essential.



Accurate documentation is important for further clinical management, continuity of care and team work. In addition, inadequate documentation can contribute to the likelihood of there being medicolegal consequences.¹⁹⁷ The team member recording events on the structured proforma, the scribe, is crucial in the management of PPH (see Appendix V); the proforma is effectively a checklist of available interventions, and team leaders should communicate with the scribe during the PPH to ensure that no steps have been omitted. PPH should be notified through a clinical incident reporting or risk management system.

Evidence
level 4

It is important to record:

- the staff in attendance and the time they arrived
- the sequence of events
- the administration of different pharmacological agents, their timing and sequence
- the time of surgical intervention, where relevant
- the condition of the mother throughout the different steps
- the timing of the fluid and blood products given.

7.3 Debriefing

An opportunity to discuss the events surrounding the obstetric haemorrhage should be offered to the woman (possibly with her birthing partner/s) at a mutually convenient time.



After obstetric emergencies, women can be psychologically affected by postnatal depression or fear of further childbirth. Major PPH can be traumatic to women and their families and has been associated with the subsequent development of post-traumatic stress disorder.¹⁹⁸ Women who have experienced a major PPH should be offered an opportunity to discuss the events surrounding their delivery. A discussion of future pregnancy, including the likelihood of a repeat PPH and any fears regarding pregnancy and childbirth that the woman may have should be addressed. This should include arrangements for appropriate investigations as necessary, such as testing for coagulopathies if there are other indicators and screening for the rare complication of postpartum hypopituitarism (Sheehan syndrome) secondary to hypotension.¹⁹⁹

Evidence
level 4

8. Recommendations for future research

- RCTs are required to identify the best drug combinations, route and dose of uterotonics for the treatment of primary PPH.
- The role of viscoelastometric point of care tests using TEG[®] and ROTEM[®] in the management of PPH requires evaluation.
- Studies are required to determine the optimal ratio of packed red cells to FFP in the management of obstetric haemorrhage.

- Studies are required to determine the role of fibrinogen concentrate in the management of PPH.
- The role of prothrombin complex concentrate in the management of PPH requires evaluation.
- RCTs are required to investigate the role of uterotonic agents (misoprostol and ergometrine) in the management of secondary PPH.

9. Auditable topics

- The proportion of women who are screened for antenatal anaemia (100%).
- The proportion of women who are offered uterotonics for the third stage of labour (100%).
- The proportion of women undergoing an assessment of risk factors for PPH when they present in labour (100%).
- Appropriate documentation of management, especially with the timing of events for women who have had PPH (100%).
- Notification to the risk management team of women with PPH involving a blood loss greater than 1500 ml (100%).
- Proportion of the multidisciplinary team who have undergone skills drills training in PPH (100%).

10. Useful links and support groups

- Royal College of Obstetricians and Gynaecologists. *Heavy bleeding after birth (postpartum haemorrhage). Information for you*. London: RCOG; 2016 [<https://www.rcog.org.uk/en/patients/patient-leaflets/heavy-bleeding-after-birth-postpartum-haemorrhage/>].
- Royal College of Obstetricians and Gynaecologists. *Blood transfusion, pregnancy and birth. Information for you*. London: RCOG; 2015 [<https://www.rcog.org.uk/en/patients/patient-leaflets/blood-transfusion-pregnancy-and-birth/>].
- Patient. *Postpartum Haemorrhage* [www.patient.co.uk/showdoc/40000261].
- Netdoctor.co.uk. *I suffered with postpartum haemorrhage* [www.netdoctor.co.uk/ate/womenshealth/207160.html].

References

1. Scottish Obstetric Guidelines and Audit Project. *The Management of Postpartum Haemorrhage: A Clinical Practice Guideline for Professionals involved in Maternity Care in Scotland*. Pilot edition. Glasgow: SOGAP, 1998. [<http://healthcareimprovementscotland.org/his/idoc.ashx?docid=84ee51e6-d441-4dba-8ebf-4fa6a2857e0d&version=-1>]. Accessed 2015 Jul 2.
2. Mousa HA, Blum J, Abou El Senoun G, Shakur H, Alfirevic Z. Treatment for primary postpartum haemorrhage. *Cochrane Database Syst Rev* 2014;(2):CD003249.
3. Knight M, Tuffnell D, Kenyon S, Shakespeare J, Gray R, Kurinczuk JJ, editors, on behalf of MBRRACE-UK. *Saving Lives, Improving Mothers' Care - Surveillance of maternal deaths in the UK 2011-13 and lessons learned to inform maternity care from the UK and Ireland Confidential Enquiries into Maternal Deaths and Morbidity 2009-13*. Oxford: National Perinatal Epidemiology Unit, University of Oxford; 2015.
4. Alexander J, Thomas PW, Sanghera J. Treatments for secondary postpartum haemorrhage. *Cochrane Database Syst Rev* 2002;(1):CD002867.
5. Demers C, Derzko C, David M, Douglas J; Society of Obstetricians and Gynaecologists of Canada. Gynaecological and obstetric management of women with inherited bleeding disorders. SOGC Clinical Practice Guidelines No 163. *J Obstet Gynaecol Can* 2005;27:707-18.
6. Baudo F, de Cataldo F; Italian Association of Haemophilia Centres (AICE); Register of acquired factor VIII inhibitors (RIIA). Acquired factor VIII inhibitors in pregnancy: data from the Italian Haemophilia Register relevant to clinical practice. *BJOG* 2003;110:311-4.
7. The Royal College of Surgeons of England. *Code of Practice for the Surgical Management of Jehovah's Witnesses*. London: The Royal College of Surgeons of England; 2002.
8. The Association of Anaesthetists of Great Britain and Ireland. *Management of Anaesthesia for Jehovah's Witnesses*, 2nd ed. London: AAGBI; 2005.
9. National Institute for Health and Care Excellence. *Intrapartum care: care of healthy women and their babies during childbirth*. NICE clinical guideline 190. Manchester: NICE; 2014.
10. Centre for Maternal and Child Enquiries (CMACE). *Saving Mothers' Lives: reviewing maternal deaths to make motherhood safer: 2006-08. The Eighth Report on Confidential Enquiries into Maternal Deaths in the United Kingdom*. *BJOG* 2011;118 Suppl 1:1-203.

11. Royal College of Obstetricians and Gynaecologists. *Antepartum Haemorrhage*. Green-top Guideline No. 63. London: RCOG; 2011.
12. Royal College of Obstetricians and Gynaecologists. *Placenta Praevia, Placenta Praevia Accreta and Vasa Praevia: Diagnosis and Management*. Green-top Guideline No. 27. London: RCOG; 2011.
13. Royal College of Obstetricians and Gynaecologists. *Blood Transfusion in Obstetrics*. Green-top Guideline No. 47. London: RCOG; 2015.
14. Calvert C, Thomas SL, Ronsmans C, Wagner KS, Adler AJ, Filippi V. Identifying regional variation in the prevalence of postpartum haemorrhage: a systematic review and meta-analysis. *PLoS ONE* 2012;7:e41114.
15. Bryant A, Mhyre JM, Leffert LR, Hoban RA, Yakoob MY, Bateman BT. The association of maternal race and ethnicity and the risk of postpartum hemorrhage. *Anesth Analg* 2012;115:1127–36.
16. Combs CA, Murphy EL, Laros RK Jr. Factors associated with postpartum hemorrhage with vaginal birth. *Obstet Gynecol* 1991;77:69–76.
17. Stones RW, Paterson CM, Saunders NJ. Risk factors for major obstetric haemorrhage. *Eur J Obstet Gynecol Reprod Biol* 1993;48:15–8.
18. Al-Zirqi I, Vangen S, Forsen L, Stray-Pedersen B. Prevalence and risk factors of severe obstetric haemorrhage. *BJOG* 2008;115:1265–72.
19. Tsu VD. Postpartum haemorrhage in Zimbabwe: a risk factor analysis. *Br J Obstet Gynaecol* 1993;100:327–33.
20. Bais JM, Eskes M, Pel M, Bonsel GJ, Bleker OP. Postpartum haemorrhage in nulliparous women: incidence and risk factors in low and high risk women. A Dutch population-based cohort study on standard (≥ 500 ml) and severe (≥ 1000 ml) postpartum haemorrhage. *Eur J Obstet Gynecol Reprod Biol* 2004;115:166–72.
21. Magann EF, Evans S, Hutchinson M, Collins R, Howard BC, Morrison JC. Postpartum hemorrhage after vaginal birth: an analysis of risk factors. *South Med J* 2005;98:419–22.
22. Magann EF, Evans S, Chauhan SP, Lanneau G, Fisk AD, Morrison JC. The length of the third stage of labor and the risk of postpartum hemorrhage. *Obstet Gynecol* 2005;105:290–3.
23. Sheiner E, Sarid L, Levy A, Seidman DS, Hallak M. Obstetric risk factors and outcome of pregnancies complicated with early postpartum hemorrhage: a population-based study. *J Matern Fetal Neonatal Med* 2005;18:149–54.
24. Sosa CG, Althabe F, Belizán JM, Buekens P. Risk factors for postpartum hemorrhage in vaginal deliveries in a Latin-American population. *Obstet Gynecol* 2009;113:1313–9.
25. Brinsden PR, Clark AD. Postpartum haemorrhage after induced and spontaneous labour. *Br Med J* 1978;ii:855–6.
26. Selo-Ojeme DO, Okonofua FE. Risk factors for primary postpartum haemorrhage. A case control study. *Arch Gynecol Obstet* 1997;259:179–87.
27. World Health Organization. *WHO recommendations for the prevention and treatment of postpartum haemorrhage*. Geneva: WHO; 2012.
28. Oyelese Y, Ananth CV. Postpartum hemorrhage: epidemiology, risk factors, and causes. *Clin Obstet Gynecol* 2010;53:147–56.
29. Confidential Enquiry into Maternal and Child Health. *Why Mothers Die 2000–2002. The Sixth Report of the Confidential Enquiries into Maternal Deaths in the United Kingdom*. London: RCOG Press; 2004.
30. Leduc D, Senikas V, Lalonde AB, Ballerman C, Biringner A, Delaney M, et al.; Clinical Practice Obstetrics Committee; Society of Obstetricians and Gynaecologists of Canada. Active management of the third stage of labour: prevention and treatment of postpartum hemorrhage. SOGC Clinical Practice Guideline No. 235. *J Obstet Gynaecol Can* 2009;31:980–93.
31. Combs CA, Murphy EL, Laros RK Jr. Factors associated with hemorrhage in cesarean deliveries. *Obstet Gynecol* 1991;77:77–82.
32. National Institute for Health and Care Excellence. *Antenatal Care*. NICE clinical guideline 62. Manchester: NICE; 2008.
33. Pavord S, Myers B, Robinson S, Allard S, Strong J, Oppenheimer C; British Committee for Standards in Haematology. UK guidelines on the management of iron deficiency in pregnancy. *Br J Haematol* 2012;156:588–600.
34. Kavle JA, Stoltzfus RJ, Witter F, Tielsch JM, Khalfan SS, Caulfield LE. Association between anaemia during pregnancy and blood loss at and after delivery among women with vaginal births in Pemba Island, Zanzibar, Tanzania. *J Health Popul Nutr* 2008;26:232–40.
35. Hofmeyr GJ, Abdel-Aleem H, Abdel-Aleem MA. Uterine massage for preventing postpartum haemorrhage. *Cochrane Database Syst Rev* 2013;(7):CD006431.
36. McDonald SJ, Abbott JM, Higgins SP. Prophylactic ergometrine-oxytocin versus oxytocin for the third stage of labour. *Cochrane Database Syst Rev* 2004;(1):CD000201.
37. Tunçalp Ö, Hofmeyr GJ, Gülmezoglu AM. Prostaglandins for preventing postpartum haemorrhage. *Cochrane Database Syst Rev* 2012;(8):CD000494.
38. Oladapo OT, Fawole B, Blum J, Abalos E. Advance misoprostol distribution for preventing and treating postpartum haemorrhage. *Cochrane Database Syst Rev* 2012;(2):CD009336.
39. Begley CM, Gyte GM, Devane D, McGuire W, Weeks A. Active versus expectant management for women in the third stage of labour. *Cochrane Database Syst Rev* 2011;(11):CD007412.
40. Hutton EK, Hassan ES. Late vs early clamping of the umbilical cord in full-term neonates: systematic review and meta-analysis of controlled trials. *JAMA* 2007;297:1241–52.
41. Royal College of Obstetricians and Gynaecologists. *Clamping of the Umbilical Cord and Placental Transfusion*. Scientific Impact Paper No. 14. London: RCOG; 2015.
42. Tita AT, Szychowski JM, Rouse DJ, Bean CM, Chapman V, Nothern A, et al. Higher-dose oxytocin and hemorrhage after vaginal delivery: a randomized controlled trial. *Obstet Gynecol* 2012;119:293–300.
43. Gizzo S, Patrelli TS, Di Gangi S, Carrozzini M, Saccardi C, Zambon A, et al. Which uterotonic is better to prevent the postpartum hemorrhage? Latest news in terms of clinical efficacy, side effects, and contraindications: a systematic review. *Reprod Sci* 2013;20:1011–9.
44. Su LL, Chong YS, Samuel M. Carbetocin for preventing postpartum haemorrhage. *Cochrane Database Syst Rev* 2012;(4):CD005457.
45. Dennehy KC, Rosaeg OP, Cicutti NJ, Krepski B, Sylvain JP. Oxytocin injection after caesarean delivery: intravenous or intramyometrial? *Can J Anaesth* 1998;45:635–9.
46. Munn MB, Owen J, Vincent R, Wakefield M, Chestnut DH, Hauth JC. Comparison of two oxytocin regimens to prevent uterine atony at caesarean delivery: a randomized controlled trial. *Obstet Gynecol* 2001;98:386–90.
47. Lokugamage AU, Paine M, Bassaw-Balroop K, Sullivan KR, El Refaey H, Rodeck CH. Active management of the third stage at caesarean section: a randomised controlled trial of misoprostol versus Syntocinon. *Aust N Z J Obstet Gynaecol* 2001;41:411–4.
48. Chou MM, MacKenzie IZ. A prospective, double-blind, randomized comparison of prophylactic intramyometrial 15-methyl prostaglandin F_{2α}, 125 micrograms, and intravenous oxytocin, 20 units, for the control of blood loss at elective cesarean section. *Am J Obstet Gynecol* 1994;171:1356–60.

49. Boucher M, Horbay GL, Griffin P, Deschamps Y, Desjardins C, Schulz M, et al. Double-blind, randomized comparison of the effect of carbetocin and oxytocin on intraoperative blood loss and uterine tone of patients undergoing cesarean section. *J Perinatol* 1998;18:202–7.
50. Dansereau J, Joshi AK, Helewa ME, Doran TA, Lange IR, Luther ER, et al. Double-blind comparison of carbetocin versus oxytocin in prevention of uterine atony after cesarean section. *Am J Obstet Gynecol* 1999;180:670–6.
51. National Collaborating Centre for Women's and Children's Health. *Caesarean Section*. London: RCOG; 2011.
52. Novikova N, Hofmeyr GJ, Cluver C. Tranexamic acid for preventing postpartum haemorrhage. *Cochrane Database Syst Rev* 2015;(6):CD007872.
53. Glover P. Blood loss at delivery: how accurate is your estimation? *Aust J Midwifery* 2003;16:21–4.
54. Toledo P, McCarthy RJ, Hewlett BJ, Fitzgerald PC, Wong CA. The accuracy of blood loss estimation after simulated vaginal delivery. *Anesth Analg* 2007;105:1736–40.
55. Patel A, Goudar SS, Geller SE, Kodkany BS, Edlavitch SA, Wagh K, et al. Drape estimation vs. visual assessment for estimating postpartum hemorrhage. *Int J Gynaecol Obstet* 2006;93:220–4.
56. Schorn MN. Measurement of blood loss: review of the literature. *J Midwifery Womens Health* 2010;55:20–7.
57. Zhang WH, Deneux-Tharaux C, Brocklehurst P, Juszczak E, Joslin M, Alexander S; EUPHRATES Group. Effect of a collector bag for measurement of postpartum blood loss after vaginal delivery: cluster randomised trial in 13 European countries. *BMJ* 2010;340:c293.
58. Buckland SS, Homer CS. Estimating blood loss after birth: using simulated clinical examples. *Women Birth* 2007;20:85–8.
59. Bose P, Regan F, Paterson-Brown S. Improving the accuracy of estimated blood loss at obstetric haemorrhage using clinical reconstructions. *BJOG* 2006;113:919–24.
60. Rath WH. Postpartum hemorrhage – update on problems of definitions and diagnosis. *Acta Obstet Gynecol Scand* 2011;90:421–8.
61. Allgöwer M, Burri C. ["Shock index"]. *Dtsch Med Wochenschr* 1967;92:1947–50. German.
62. Nathan HL, El Ayadi A, Hezelgrave NL, Seed P, Butrick E, Miller S, et al. Shock index: an effective predictor of outcome in postpartum haemorrhage? *BJOG* 2015;122:268–75.
63. Royal College of Obstetricians and Gynaecologists. *Responsibility of Consultant On-Call*. Good Practice No. 8. London: RCOG; 2009.
64. Duthie SJ, Ven D, Yung GL, Guang DZ, Chan SY, Ma HK. Discrepancy between laboratory determination and visual estimation of blood loss during normal delivery. *Eur J Obstet Gynecol Reprod Biol* 1991;38:119–24.
65. Ho AM, Karmakar MK, Dion PW. Are we giving enough coagulation factors during major trauma resuscitation? *Am J Surg* 2005;190:479–84.
66. Hirshberg A, Dugas M, Banez EI, Scott BG, Wall MJ Jr, Mattox KL. Minimizing dilutional coagulopathy in exsanguinating hemorrhage: a computer simulation. *J Trauma* 2003;54:454–63.
67. Hunt BJ, Allard S, Keeling D, Norfolk D, Stanworth SJ, Pendry K; British Committee for Standards in Haematology. A practical guideline for the haematological management of major haemorrhage. *Br J Haematol* 2015;170:788–803.
68. British Committee for Standards in Haematology, Stainsby D, MacLennan S, Thomas D, Isaac J, Hamilton PJ. Guidelines on the management of massive blood loss. *Br J Haematol* 2006;135:634–41.
69. Cochrane Injuries Group Albumin Reviewers. Human albumin administration in critically ill patients: systematic review of randomised controlled trials. *BMJ* 1998;317:235–40.
70. Schierhout G, Roberts I. Fluid resuscitation with colloid or crystalloid solutions in critically ill patients: a systematic review of randomised trials. *BMJ* 1998;316:961–4.
71. Association of Anaesthetists of Great Britain and Ireland, Thomas D, Wee M, Clyburn P, Walker I, Brohi K, et al. Blood transfusion and the anaesthetist: management of massive haemorrhage. *Anaesthesia* 2010;65:1153–61.
72. Luna GK, Maier RV, Pavlin EG, Anardi D, Copass MK, Oreskovich MR. Incidence and effect of hypothermia in seriously injured patients. *J Trauma* 1987;27:1014–8.
73. World Health Organization. *WHO guidelines for the management of postpartum haemorrhage and retained placenta*. Geneva: WHO; 2009. [http://apps.who.int/iris/bitstream/10665/44171/1/978924159-8514_eng.pdf]. Accessed 2016 Feb 4.
74. Perel P, Roberts I, Ker K. Colloids versus crystalloids for fluid resuscitation in critically ill patients. *Cochrane Database Syst Rev* 2013;(2):CD000567.
75. Murphy MF, Wallington TB, Kelsey P, Boulton F, Bruce M, Cohen H, et al.; British Committee for Standards in Haematology, Blood Transfusion Task Force. Guidelines for the clinical use of red cell transfusions. *Br J Haematol* 2001;113:24–31.
76. United Kingdom Blood Services; Norfolk D, editor. *Handbook of Transfusion Medicine*. 5th ed. Norwich: TSO; 2013.
77. Serious Hazards of Transfusion (SHOT). [www.shotuk.org/]. Accessed 2013 Aug 6.
78. Kozek-Langenecker SA, Afshari A, Albaladejo P, Santullano CA, De Robertis E, Filipescu DC, et al. Management of severe perioperative bleeding: guidelines from the European Society of Anaesthesiology. *Eur J Anaesthesiol* 2013;30:270–382.
79. British Committee for Standards in Haematology, Milkins C, Berryman J, Cantwell C, Elliott C, Haggas R, et al. Guidelines for pre-transfusion compatibility procedures in blood transfusion laboratories. *Transfus Med* 2013;23:3–35.
80. Royal College of Obstetricians and Gynaecologists. *The Management of Women with Red Cell Antibodies during Pregnancy*. Green-top Guideline No. 65. London: RCOG; 2014.
81. Laupacis A, Brown J, Costello B, Delage G, Freedman J, Hume H, et al. Prevention of posttransfusion CMV in the era of universal WBC reduction: a consensus statement. *Transfusion* 2001;41:560–9.
82. Advisory Committee on the Safety of Blood, Tissues and Organs. *Cytomegalovirus tested blood components - position statement*. London: Department of Health; 2012.
83. Carless PA, Henry DA, Moxey AJ, O'Connell D, Brown T, Fergusson DA. Cell salvage for minimising perioperative allogeneic blood transfusion. *Cochrane Database Syst Rev* 2010;(4):CD001888.
84. Catling SJ, Williams S, Fielding AM. Cell salvage in obstetrics: an evaluation of the ability of cell salvage combined with leucocyte depletion filtration to remove amniotic fluid from operative blood loss at caesarean section. *Int J Obstet Anesth* 1999;8:79–84.
85. National Institute for Health and Clinical Excellence. *Intraoperative blood cell salvage in obstetrics*. NICE interventional procedure guidance 144. Manchester: NICE; 2005.
86. Wee MY, Thomas D, Verma R, Jones J, Catling S, Isaac J, et al.; The Association of Anaesthetists of Great Britain and Ireland. *Blood Transfusion and the Anaesthetist: Intra-operative Cell Salvage*. AAGBI Safety Guideline. London: AAGBI; 2009.
87. Liumbruno GM, Liumbruno C, Rafanelli D. Intraoperative cell salvage in obstetrics: is it a real therapeutic option? *Transfusion* 2011;51:2244–56.
88. Geoghegan J, Middleton L, Moore P, Subeson G, Khan K, Daniels J. Routine cell salvage during elective caesarean section: a pilot randomised trial. *Int J Obstet Anesth* 2015;24:86–7.