


Appendix 4: Maternal collapse algorithm



Appendix 5: Recommended airway equipment

Routine airway equipment


- Face masks
- Oropharyngeal airways size 2, 3 and 4
- Endotracheal tubes in a range of sizes
- Laryngoscopes – Macintosh blades (sizes 3 and 4)
 - two working short handles
 - McCoy laryngoscopes (sizes 3 and 4 blades)
 - videolaryngoscopes (at least one type)
- Tracheal tube introducer – such as a bougie
- Malleable stylet
- Magill forceps
- Nasal cannula and oxygen tubing
- Equipment for ramping/pillows
- Monitoring equipment including capnography (see AAGBI guidelines- Recommendations for standards of monitoring during anaesthesia and recovery. 4th edition, 2007)

Recommended equipment for the management of unanticipated difficult intubation

- Endotracheal tubes – range of reinforced tubes, microlaryngeal tubes sizes 5.0 and 6.0 mm, LMA-Fastrach™ tracheal tubes
- Supraglottic airway devices (SAD) to include cLMA, and a second generation SAD (e.g. LMA Proseal™, LMA Supreme™ or l-gel®) - sizes 3, 4 and 5
- LMA cuff pressure manometer
- Fibreoptic scope, camera and monitor
- Aintree® intubating catheter
- Surgical cricothyroidotomy equipment for the ‘can’t intubate can’t oxygenate’ situation:
 - Scalpel with No. 10 blade
 - Bougie
 - Size 6.0 endotracheal tube
 - Tracheal hook
 - Forceps or tracheal dilator
- Equipment for awake fibreoptic intubation:
 - Equipment to deliver topical atomised local anaesthetic to the upper airway such as the Mucosal Atomization Device (MAD®) or Mackenzie technique set
 - Berman airway
 - Epidural catheter
 - Local anesthetic for topical anaesthesia (4% lidocaine, Instillagel®)
 - Vasoconstrictors for the nose – phenylephrine/lidocaine (Co-phenylcaine®) or Xylometazoline

Taken from the Obstetric Anaesthetists’ Association and Difficult Airway Society [http://www.oaa-anaes.ac.uk/assets/_managed/cms/files/03102015_Equipment_List%20final.docx].

Appendix 6: Post collapse management



This guideline was produced on behalf of the Royal College of Obstetricians and Gynaecologists by: **Dr J Chu PhD MRCOG, Birmingham; Dr TA Johnston MD FRCOG, Birmingham Women's and Children's NHS Foundation Trust; Dr J Geoghegan FRCA, Department of Anaesthetics, Queen Elizabeth Hospital, Birmingham**

and peer reviewed by: Royal College of Anaesthetists, London; RCOG Women's Network, London; Ms N Lucas, London; Ms C Bagness, London; Ms A Mansfield, London; Dr DA Gabbott, Gloucester; Mr P Steer FRCOG, London; Ms K Fitzpatrick, London; Ms L Vincent, Oxford; Mr D Fraser FRCOG, Norwich; Ms M Jokinen, London; Mr D Tuffnell FRCOG, Bradford; Mrs K Dent FRCOG, Derby; Dr K Calvert, Perth and Dr VA Beckett FRCOG, Bradford.

Committee lead reviewers were: Dr B Kumar FRCOG, Chester and Dr A Macleod FRCOG, Edinburgh

The chairs of the Guidelines Committee were: Dr MA Ledingham MRCOG, Glasgow¹; Dr B Magowan FRCOG, Melrose¹; and Dr AJ Thomson MRCOG, Paisley².

¹co-chairs from June 2018 ²until May 2018.

All RCOG guidance developers are asked to declare any conflicts of interest. A statement summarising any conflicts of interest for this guideline is available from: <https://www.rcog.org.uk/en/guidelines-research-services/guidelines/gtg56>.

The final version is the responsibility of the Guidelines Committee of the RCOG.

The guideline will be considered for update 3 years after publication, with an intermediate assessment of the need to update 2 years after publication.

DISCLAIMER

The Royal College of Obstetricians and Gynaecologists produces guidelines as an educational aid to good clinical practice. They present recognised methods and techniques of clinical practice, based on published evidence, for consideration by obstetricians and gynaecologists and other relevant health professionals. The ultimate judgement regarding a particular clinical procedure or treatment plan must be made by the doctor or other attendant in the light of clinical data presented by the patient and the diagnostic and treatment options available.

This means that RCOG Guidelines are unlike protocols or guidelines issued by employers, as they are not intended to be prescriptive directions defining a single course of management. Departure from the local prescriptive protocols or guidelines should be fully documented in the patient's case notes at the time the relevant decision is taken.