

4.2 Worsening the prognosis of an existing sarcoma

Morcellation (by any means of disruption or injury at open, vaginal or laparoscopic surgery) of an unexpected uterine sarcoma can potentially disseminate sarcoma into the pelvis and peritoneal cavity. A diagnosis of metastatic sarcoma carries a poor prognosis with a median survival of 18 months. Data from national cancer registries have reported that sarcoma mortality was higher in the morcellated group than in the non-morcellated group (age-adjusted HR 1.90, CI 1.05–3.44; multivariate HR, 2.50, 95% CI 0.57–10.9). Age-adjusted 10-year uterine sarcoma survival was 32.2% for women treated with morcellation compared with 57.2% for the non-morcellated group (difference 25.5%; CI -55.7 to 18.1).^{16,17,22}

4.3 Disseminated fibroids (presence of benign fibroids within the abdominal and pelvic cavity)

The range of risk is considered to be 1 in 120 (uncommon) to 1 in 1200 (rare).¹⁸

4.4 Damage to bowel, bladder, ureters and blood vessels

There is a risk of damage to the bladder, bowel, ureters and blood vessels with laparoscopic hysterectomy.²³ These data were published in 2007. Since then, there have been advances in techniques and equipment, and the possibility that these rates may now be lower has been highlighted.²⁴

Laparoscopic myomectomy, with and without morcellation also carries a risk of these injuries. However, these risks are unknown because the reported literature is based on case reports rather than large trials. The total number of cases of morcellation of fibroids is also unknown. Surgeons should declare the rate of such injuries from their personal or institutional data where available.

5. Any extra procedures which may become necessary

General additional procedures associated with myomectomy or hysterectomy should be detailed in local procedure specific consent forms where available. A laparotomy may be required if there is a major complication, or, if the surgeon considers that conversion is necessary for safety or access.

6. The benefits and risks of any available alternative treatments, including no treatment

6.1 Open myomectomy or hysterectomy

An abdominal (open) myomectomy or hysterectomy to remove a uterus or fibroid may be performed as an alternative. The benefit is that an unsuspected sarcoma would not be morcellated. However, if this was performed in everyone more harm may be caused overall due to the increased risks of open surgery over laparoscopy. These are, an increased risk of thromboembolism, wound infections, blood transfusion and incisional hernias, with a longer hospital stay and recovery.^{4,20,21,25}

It should be noted that when an open myomectomy or vaginal morcellation are performed, simply cutting into the fibroid will cause spillage of cells and if there is an undiagnosed uterine sarcoma, upstaging can still occur.

6.2 Contained retrieval during morcellation

The use of tissue retrieval bags for the contained removal of fibroids or the uterus, once they have been detached has been proposed.²⁶ However, there is no current evidence that bags reduce the incidence of disseminated fibroids or upstaging of an undiagnosed uterine sarcoma. There is also a theoretical risk that the bags obscure the laparoscopic view, with the potential of causing more intra-abdominal injuries.⁴ The decision regarding contained retrieval during morcellation should be based on local practice and individualised patient care.

6.3 *Conservative measures and uterine artery embolisation*

Any management option, including uterine artery embolisation, medical management and no treatment that results in uterine preservation, or preservation of some element of fibroid tissue, runs the risk of leaving an undiagnosed uterine sarcoma in situ;¹ see Section 4 above.

7. Patient statement

The woman should be given the opportunity to state in writing any procedures that should not be performed without further discussion. If other procedures are anticipated to become necessary during the planned procedure these should be discussed preoperatively, and a record of the woman's wishes made.

8. Preoperative information

A record should be made of any sources of information given to the woman prior to surgery (such as RCOG or locally produced information leaflets; <https://www.rcog.org.uk/en/patients/patient-leaflets/>).

Existing information sheets for procedures need to be reviewed to ensure they reflect the available current evidence.

9. Information and support for women and their families

All women should be provided with relevant and up-to-date information sources. Translation services should be sourced if a language-specific leaflet is not available to ensure that a woman has a full understanding of the procedures planned.

10. This consent advice

Please note that the content herein is based on the latest evidence available at the time and will be reviewed again approximately one year from publication.

References

1. Sizzi O, Manganaro L, Rossetti A, Saldari M, Florio G, Loddo A, Zurawin R, van Herendael B, Djokovic D. Assessing the risk of laparoscopic morcellation of occult uterine sarcomas during hysterectomy and myomectomy: Literature review and the ISGE recommendations. *Eur J Obstet Gynecol Reprod Biol* 2018;220:30–38.
2. Brohl AS, Li L, Andikyan V, Obi an SG, Cioffi A, Hao K, Dudley JT, Ascher-Walsh C, Kasarskis A, Maki RG. Age-stratified risk of unexpected uterine sarcoma following surgery for presumed benign leiomyoma. *Oncologist* 2015;20:433–439.
3. Halaska MJ, Haidopoulos D, Guyon F, Morice P, Zapardiel I, Kesic V; ESGO Council. European Society of Gynecological Oncology Statement on Fibroid and Uterine Morcellation. *Int J Gynecol Cancer* 2017;27:189–192.
4. ACOG Committee Opinion No. 770: Uterine Morcellation for Presumed Leiomyomas. *Obstet Gynecol* 2019;133:e238–e248.
5. Munro MG, Critchley HOD, Fraser IS; FIGO Menstrual Disorders Committee. The two FIGO systems for normal and abnormal uterine bleeding symptoms and classification of causes of abnormal uterine bleeding in the reproductive years: 2018 revisions. *Int J Gynaecol Obstet* 2018;143:393–408.
6. LUDOVISI M, Moro F, Pasciuto T, Di Noi S, et al. Imaging of gynecological disease: clinical and ultrasound characteristics of uterine sarcomas. *Ultrasound Obstet Gynecol* 2019. doi: 10.1002/uog.20270.
7. On behalf of AAGL. Morcellation during uterine tissue extraction: an update. *JMIG* 2019;25:543–550.
8. Wickerman DL, Fisher B, Wolmark N et al. Association of tamoxifen and uterine sarcoma. *J Clin Oncol* 2002;20:2758–2760.
9. Toro JR, Travis LB, Wu HJ, Zhu K, Fletcher CD, Devesa SS. Incidence patterns of soft tissue sarcomas, regardless of primary site, in the surveillance, epidemiology and end results program, 1978–2001: An analysis of 26,758 cases. *Int J Cancer* 2006;119:2922–2930.
10. Royal College of Obstetricians and Gynaecologists. *Obtaining Valid Consent. Clinical Governance Advice No. 6*. London: RCOG; 2015.
11. Desai VB, Wright JD, Schwartz PE, Jorgensen EM, Fan L, Litkouhi B, Lin H, Gross CP, Xu X. Occult Gynecologic Cancer in Women Undergoing Hysterectomy or Myomectomy for Benign Indications. *Obstet Gynecol* 2018;131:642–651.
12. Peters A, Sadecky AM, Winger DG, Guido RS, Lee TTM, Mansuria SM, Donnellan NM. Characterization and Preoperative Risk Analysis of Leiomyosarcomas at a High-Volume Tertiary Care Center. *Int J Gynecol Cancer* 2017;27:1183–1190.
13. Sarcoma UK webpage: <https://sarcoma.org.uk/>.
14. Mao J, Pfeifer S, Zheng XE, Schlegel P, Sedrakyan A. Population-based estimates of the prevalence of uterine sarcoma among patients with leiomyomata undergoing surgical treatment. *JAMA Surg* 2015;150:368–70.
15. Multinu F, Casarin J, Tortorella L, Huang Y, Weaver A, Angioni S, Melis GB, Mariani A, Stewart EA, Laughlin-Tommaso SK. Incidence of sarcoma in patients undergoing hysterectomy for benign indications: a population-based study. *Am J Obstet Gynecol* 2019;220:179.e1–10.
16. George S, Barysaukas C, Serrano C, et al. Retrospective cohort study evaluating the impact of intraperitoneal morcellation on outcomes of localized uterine leiomyosarcoma. *Cancer* 2014;120:3154–3158.
17. Park JY, Park SK, Kim DY, et al. The impact of tumor morcellation during surgery on the prognosis of patients with apparently early uterine leiomyosarcoma. *Gynecol Oncol* 2011;122:255–259.
18. Perri T, Korach J, Sadetzki S, et al. Uterine leiomyosarcoma: Does the primary surgical procedure matter?. *Int J Gynecol Cancer* 2009;19:257–260.
19. Montgomery v Lanarkshire Health Board [2015] SC 11 [2015] 1 AC 1430.
20. Aarts JWM, Nieboer TE, Johnson N, Tavender E, Garry R, Mol BJ, Kluivers KB. Surgical approach to hysterectomy for benign gynaecological disease. *Cochrane Database Syst Rev* 2015;8:CD003677.
21. Bhav Chittawar P, Franik S, Pouwer AW, Farquhar C. Minimally invasive surgical techniques versus open myomectomy for uterine fibroids. *Cochrane Database Syst Rev* 2014;10:CD004638.
22. Bretthauer M, Goderstad JM, Løberg M, Emilsson L, Ye W, Adami HO, Kalager M. Uterine morcellation and survival in uterine sarcomas. *Eur J Cancer* 2018;101:62–68.
23. NICE Interventional procedures guidance [IPG239]. *Laparoscopic techniques for hysterectomy*; 2007. Available from: www.nice.org.uk/ipg239.
24. Garry R. Re-evaluating the eVALuate study and the NICE guidelines: a personal review. *BJOG* 2016;123: 1796.
25. Siedhoff MT, Wheeler SB, Rutstein SE, Geller EJ, Doll KM, Wu JM, Clarke-Pearson DL. Laparoscopic hysterectomy with morcellation vs abdominal hysterectomy for presumed fibroid tumors in premenopausal women: a decision analysis. *Am J Obstet Gynecol* 2015;212:591.e1–8.
26. Rousseau M, Morel A, Dechoux S, Bouet PE, Catala L, Lefebvre Lacoëuille C, Descamps P, Legendre G. Can the risks associated with uterine sarcoma morcellation really be prevented? Overview of the role of uterine morcellation in 2018. *J Gynecol Obstet Hum Reprod* 2018;47:341–349.
27. Reference 27: Agency for Healthcare Research and Quality (AHRQ). *Comparative Effectiveness Review*. No.195. p.70. Rockville, MD: AHRQ; 2017 [<https://effectivehealthcare.ahrq.gov/sites/default/files/pdf/cer-195-uterine-fibroids-final-revision.pdf>].

Appendix 1

Definition of perimenopause referred to in this consent advice is as per https://www.imsociety.org/menopause_terminology.php.

Menopause terminology

The International Menopause Society (IMS) and the World Health Organization (WHO), with the aim of standardising the terminology of menopause have put forward the following definition:

Perimenopause: the period immediately prior to the menopause (when the endocrinological, biological, and clinical features of approaching menopause commence) and the first year after menopause

Appendix 2

The full extracted Table 4 from reference 2: Brohl AS, Li L, Andikyan V, Obian SG, Cioffi A, Hao K, Dudley JT, Ascher-Walsh C, Kasarskis A, Maki RG. Age-stratified risk of unexpected uterine sarcoma following surgery for presumed benign leiomyoma. *Oncologist*, 2015;20:433–439 [<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4391766/table/T4/?report=objectonly>].

Appendix 3

Reference 27: Agency for Healthcare Research and Quality (AHRQ). *Comparative Effectiveness Review*. No. 195. p. 70. Rockville, MD: AHRQ; 2017 [<https://effectivehealthcare.ahrq.gov/sites/default/files/pdf/cer-195-uterine-fibroids-final-revision.pdf>]. This reference has been used as a resource during preparation of the consent advice for morcellation for myomectomy or hysterectomy.

Appendix 4: Consent to treatment form

Patient identifier/label

Name of proposed procedure or course of treatment

Supplementary consent – morcellation for myomectomy or hysterectomy

Statement of health professional (to be filled in by health professional with appropriate knowledge of proposed procedure, as specified in consent policy).

I have explained the procedure to the patient, in particular, I have explained:

The intended benefits

The main benefit of the use of morcellation is the completion of the entire procedure laparoscopically or vaginally, which is associated with smaller incisions, less pain, reduced risk of infection, reduced risk of thromboembolism, shorter hospital stay and a quicker recovery.

Serious risks (as detailed in Section 4 above)

- Unintended morcellation of a uterine sarcoma.
- Worsening the prognosis of an existing sarcoma.
- Disseminated fibroids (presence of benign fibroids within the abdominal and pelvic cavity).
- Damage to bowel, bladder, ureters and blood vessels.

Any extra procedures which may become necessary

General additional procedures associated with myomectomy or hysterectomy should be detailed in local procedure specific consent forms where available. A laparotomy may be required if there is a major complication, or, if the surgeon considers that conversion is necessary for safety or access.

I have also discussed what the procedure is likely to involve, the benefits and risks of any available alternative treatments (including no treatment) and any particular concerns of this patient.

The following leaflet has been provided: *Information for you: Morcellation for Myomectomy or Hysterectomy*

This procedure will involve

☐ General and/or regional anaesthesia ☐ Local anaesthesia ☐ Sedation

Signed Date

Name (print)

Position

Contact details (if patient wishes to discuss options later)

.....

Statement of interpreter (where appropriate)

I have interpreted the information above to the patient to the best of my ability and in a way in which I believe they can understand.

Signed Date

Name (print)

Top copy accepted by patient? Yes / No (please ring)