

101.

147

$$\psi(x) = \frac{1}{x^2 + a^2}$$

$$\int_{-\infty}^{+\infty} \left| \frac{1}{x^2 + a^2} \right|^2 dx = 1$$

$$A^2 \int_{-\infty}^{+\infty} \left( \frac{1}{x^2 + a^2} \right)^2 dx = \left| \int_{-\infty}^{+\infty} \frac{1}{(ax^2 + b)^n} dx = \frac{2n-3}{2b(n-1)} \int_{-\infty}^{+\infty} \frac{1}{(ax^2 + b)^{n-1}} \right.$$

$$\left. + \left( \frac{x}{2b(n-1)(ax^2 + b)^{n-1}} \right) \right|_{-\infty}^{+\infty} =$$

$$= \frac{A^2}{2a^2 \cdot 1} \int_{-\infty}^{+\infty} \frac{1}{x^2 + a^2} dx + \frac{A^2 x}{2a^2 \cdot (x^2 + a^2)} \Big|_{-\infty}^{+\infty} =$$

$$= \frac{A^2}{2a^2} \cdot \frac{1}{a} \cdot \left( \arctg \frac{x}{a} \right) \Big|_{-\infty}^{+\infty} + (0 - 0) = \left| \arctg \frac{x}{a} = \frac{\pi}{2} \right|_{x \rightarrow \infty} =$$

$$= \frac{A^2 \pi}{2a^3} \cdot \int_{-\infty}^{+\infty} |\psi(x)|^2 dx = 1$$

$$\frac{A^2 \pi}{2a^3} = 1 \Rightarrow A = \sqrt{\frac{2a^3}{\pi}} \quad \text{Answer: } \sqrt{\frac{2a^3}{\pi}}$$

102.  $\varphi(x) = \frac{B}{x + ib}$

$$\int_{-\infty}^{+\infty} \frac{B B}{(x + ib)(x - ib)} dx = \int_{-\infty}^{+\infty} \frac{B^2}{x^2 - (ib)^2} dx =$$

$$= \int_{-\infty}^{+\infty} \frac{B^2}{x^2 + b^2} dx = \left| \text{modulus imaginary} \right| = \frac{B^2}{b} \arctg \frac{x}{b} \Big|_{-\infty}^{+\infty} =$$

$$= \frac{B^2}{b} \left( \frac{\pi}{2} - \left( -\frac{\pi}{2} \right) \right) = \frac{B^2}{b} \pi = 1 \Rightarrow B = \sqrt{\frac{b}{\pi}}$$

Answer:  $\sqrt{\frac{b}{\pi}}$

$$\langle \varphi | \varphi \rangle \quad \text{№ 3.}$$

$$\langle \varphi | \varphi \rangle = \int_{-\infty}^{+\infty} \varphi^*(x) \varphi(x) dx = 1$$

$$\int_{-\infty}^{+\infty} \frac{AB}{(x-ib)(x^2+a^2)} dx =$$

Далее я приведу вырезки из черновика, где раскладывал дробь на простейшие. Простите, если нужно, позже перепису в чистовик, но сегодня не успеваю.

$$\frac{A}{x-ib} + \frac{B}{x+ia} + \frac{C}{x+ia}$$

$$A(x^2+a^2) + B(x-ib)(x+ia) + C(x-ib)(x-ia)$$

$$A(x^2+a^2) + B(x^2+xia-xib+ab) + C(x^2-xia-xib-ab)$$

$$A+B+C=0$$

$$iB(a-b) + iC(1+a+ib) = 0$$

$$Aa^2 + abB + abC = 1$$

$$\begin{aligned} & \frac{a^2}{a} \quad -a^2 - \frac{a^3 - a^2b}{a+b} + ab - a^2b + ab^2 \\ & \frac{a}{a} \quad -a^3 + \cancel{a^2b} - a^3 + \cancel{a^2b} + \cancel{a^2b} + ab^2 - \cancel{a^2b} + ab^2 \\ & \frac{a}{a} \quad -2a^3 + 2ab^2 \\ & \frac{-a}{-a} \quad \frac{a+b}{-2a^3+2ab^2} = \frac{1}{3} \end{aligned}$$

$$\frac{a+b}{-2a^3+2ab^2} = B$$

$$C = \frac{a-b}{-2a^3+2ab^2}$$

$$A+B+C=0$$

$$A + a + b + a - b = 0$$

$$A = -2a$$

$$\frac{-2a}{-2a^3+2ab^2} = \frac{1}{b^2-a^2}$$



Возврат к заданию 3

$$\begin{aligned}
 & \frac{1}{c} \int_{-\infty}^{+\infty} \frac{2a}{x-ib} dx + \frac{1}{c} \int_{-\infty}^{+\infty} \frac{a+b}{x-ia} dx + \frac{1}{c} \int_{-\infty}^{+\infty} \frac{a-b}{x+ia} dx = \\
 & = \left| \text{где } c = -2a^2 + 2ab^2 \right| = \\
 & = \frac{1}{c} \left( 2a \ln|x-ib| \right) \Big|_{-\infty}^{+\infty} + \frac{a+b}{c} \ln|x-ia| \Big|_{-\infty}^{+\infty} + \\
 & + \frac{a-b}{c} \ln|x+ia| \Big|_{-\infty}^{+\infty} = \\
 & = \frac{1}{c} (2a \ln|x-ib| + (a+a+b-b) \ln|x-ia| + (a-a+b-b) \ln|x+ia|) \Big|_{-\infty}^{+\infty} = \\
 & = \frac{1}{c} (2a \ln|x-ib| + a \ln|x-ia| + a \ln|x+ia| + b \ln|x-ia| - b \ln|x+ia|) \Big|_{-\infty}^{+\infty} = \\
 & = \frac{1}{c} (2a \ln|x-ib| + a \ln(x^2+a^2) + b \ln \frac{x-ia}{x+ia}) \Big|_{-\infty}^{+\infty}
 \end{aligned}$$

$$\begin{aligned}
 & = \frac{1}{c} \left( \frac{2a}{c} \ln(x-ib) + \frac{a}{c} \cdot 0 + \frac{b}{c} \ln(x-ia) - \frac{b}{c} \ln \frac{(x+ia)}{(x-ia)} \right) = \\
 & = \frac{1}{c} \left( \frac{2a}{c} i\pi + \frac{b}{c} i\pi + \frac{b}{c} i\pi \right) =
 \end{aligned}$$

$$\begin{aligned}
 & = \frac{1}{c} \left( \frac{2a - 2b}{-2a^2 + 2ab^2} \right) i\pi = \\
 & = \frac{1}{c} \left( \frac{a-b}{-a(a^2-b^2)} \right) i\pi = \frac{-\sqrt{\frac{2a^3}{\pi}} \cdot \sqrt{\frac{b}{\pi}} i\pi}{(a+b) \cdot a} = \\
 & = \frac{-\sqrt{2} a^{\frac{3}{2}} \sqrt{b} i}{(a+b) a} = \frac{-\sqrt{2} a b^{\frac{1}{2}} i}{a+b}
 \end{aligned}$$

$$\text{пр. 4. } \delta(f(x)) = \delta\left(\sum f'(x_i) \cdot (x - x_i)\right) \quad \textcircled{1}$$

$$\text{и.к. } \delta(ax) = \frac{1}{|a|} \delta(x) \Rightarrow \textcircled{2} \sum \frac{\delta(x - x_i)}{|f'(x_i)|}$$

7.7.2

$$\text{пр. 5. } f_1(x) = \alpha_1 e^{i\pi x/a} \quad f_2(x) = \alpha_2 e^{-i\pi x/a}$$

$$\begin{aligned} 1. \langle f_1 | f_2 \rangle &= \int_0^a f_1^*(x) f_2(x) dx = \int_0^a \alpha_1 e^{-i\pi x/a} \alpha_2 e^{-i\pi x/a} dx = \\ &= \alpha_1 \alpha_2 \int_0^a e^{\frac{-i\pi x - i\pi x}{a}} dx = \alpha_1 \alpha_2 \int_0^a e^{\frac{-2i\pi x}{a}} dx = \\ &= \alpha_1 \alpha_2 \int_0^a \left( \cos \frac{2\pi x}{a} - i \sin \frac{2\pi x}{a} \right) dx = \alpha_1 \alpha_2 \frac{a}{2\pi} \cdot \\ &\cdot \left( \sin \frac{2\pi x}{a} + i \cos \frac{2\pi x}{a} \right) \Big|_0^a = \\ &= \frac{\alpha_1 \alpha_2 a}{2\pi} (\sin 2\pi - \sin 0 + i \cos 2\pi - i \cos 0) = \frac{\alpha_1 \alpha_2 a}{2\pi} \cdot 0 = 0 \end{aligned}$$

$$2. \int_0^a |\alpha_1 e^{i\pi x/a}|^2 dx = \alpha_1^2 \int_0^a e^{\frac{-i\pi x}{a}} \cdot e^{\frac{i\pi x}{a}} dx \stackrel{\text{бери}}{=} \alpha_1^2 \int_0^a 1 dx = \alpha_1^2 a$$

$$\textcircled{3} \alpha_1^2 \int_0^a e^{\frac{i\pi x}{a} - \frac{i\pi x}{a}} dx = \alpha_1^2 \int_0^a 1 dx = \alpha_1^2 (a - 0) = 1$$

$$\alpha_1^2 a = 1 \Rightarrow \alpha_1 = \sqrt{\frac{1}{a}}$$

$$\text{аналогично } \alpha_2 = \sqrt{\frac{1}{a}}$$

$$3. \psi(x) = \sqrt{\frac{2}{a}} \sin(\pi x/a) \quad |\psi\rangle = C_1 |f_1\rangle + C_2 |f_2\rangle$$

$$\sqrt{\frac{2}{a}} \sin\left(\frac{\pi x}{a}\right) = \frac{C_1}{\sqrt{a}} \left( \cos \frac{\pi x}{a} + i \sin \frac{\pi x}{a} \right) + \frac{C_2}{\sqrt{a}} \left( \cos \frac{\pi x}{a} - i \sin \frac{\pi x}{a} \right)$$

$$\sqrt{\frac{2}{a}} \sin\left(\frac{\pi x}{a}\right) = \frac{C_1 + C_2}{\sqrt{a}} \cos \frac{\pi x}{a} + i \frac{C_1 - C_2}{\sqrt{a}} \sin \frac{\pi x}{a}$$



$$\begin{cases} \frac{C_1 + C_2}{\sqrt{a}} = 0 \\ \frac{i(C_1 - C_2)}{\sqrt{a}} = \sqrt{\frac{2}{a}} \end{cases} \quad \begin{cases} C_2 = -C_1 \\ i(C_1 + C_2) = \sqrt{2} \end{cases} \quad \Rightarrow \begin{aligned} C_1 &= -\frac{\sqrt{2}}{2}i \\ C_2 &= +\frac{\sqrt{2}}{2}i \end{aligned}$$

Problem:  $|\psi\rangle = -i\frac{\sqrt{2}}{2}|f_1\rangle + i\frac{\sqrt{2}}{2}|f_2\rangle$