

POLS201 Spring 2019

Basic Hypothesis Testing

March 13

Agenda

POLS201
Spring 2019

- We're going to consider hypothesis testing and try it with R
- Log into your RStudio Cloud account and move into the Lewis & Clark workspace
- You can see these slides from the POLS201_Spring_2019_190313.pdf
- You will see a live example of data loaded raw from the Pew survey.

We can use our mastery of Normal Distributions to Test Hypotheses

POLS201
Spring 2019

- Suppose we compare two different values even though they were produced by the same data generating process.
- Or, they came from the same distribution
- Or, they are really different and weren't generated from the same distribution.
 - Which means: they represent two things that are truly different.
- How do we know?

Hypothesis Testing

POLS201
Spring 2019

- Null Hypothesis: A statement about what we would expect to observe if our values came from the same distribution.
- Alternative Hypothesis: A statement about what we would expect to observe if our values did not come from the same distribution.

Null Hypotheses:

POLS201
Spring 2019

- There is no difference in population means
- $H_o : \mu = \mu$ or $H_o : \mu - \mu = 0$
- H_o : There is no relationship between our independent and dependent variable.
- H_o : $\text{corr}(x, y) = 0$
- We may see some correlation in our samples, or not, but we conclude the populations have none.

Alternative Hypotheses:

POLS201
Spring 2019

- There is meaningful difference in population means
- $H_a : \mu \neq \mu_0$ or $H_o : \mu - \mu_0 \neq 0$
- H_a : There is a relationship between our independent and dependent variable.
- H_a : $\text{corr}(x, y) \neq 0$

Conventions

POLS201
Spring 2019

- By convention, when we engage in hypothesis testing, we seek to “reject the null hypothesis”
- We never accept the alternative hypothesis until demonstrated
- The burden of proof lies with the alternative hypothesis
- Note that we are generally interested in two-tailed tests, which as a practical matter, are harder to pass.

Different data require different tests

POLS201
Spring 2019

Table 8.1. Variable types and appropriate bivariate hypothesis tests

		Independent variable type	
		Categorical	Continuous
Dependent variable type	Categorical	<i>Tabular analysis</i>	Probit/logit
	Continuous	<i>Difference of means</i>	<i>Correlation coefficient;</i> bivariate regression model

Notes: Tests in italics are discussed in this chapter.

Different data require different tests

POLS201
Spring 2019

- All tests have a common logic: You are comparing the identified relationship of X and Y to what we would expect to see if NO relationship existed.
- Express Null and Alternative Hypotheses
- Central Limit Theorem gives us the parameters of what we would expect to see “by chance”

Another Convention

POLS201
Spring 2019

- An old fashioned convention says: Would we expect to see the findings we got less than 5% of the time?
- A P-value tell us this result.
- But beware. . . .there is no magic threshold that “proves” we have a relationship.

P-Values

POLS201
Spring 2019

- Tells us the probability we would expect to see a given relationship because of random chance
- Limitations:
- Does not tell us if the relationship is causal!
- Does not tell us the STRENGTH of the relationship if it exists.
- It merely says: “we have some basis for rejecting the null hypothesis”

How could we use this idea in a simple research project?

POLS201
Spring 2019

- Draw a sample of Facebook users
- Randomly assign users to control and treatment group
- Give treatment group a political ad. No ad for control group.
- Measure voter turnout of control and treatment groups.
- Could the mean turnout of the treatment group have been pulled from the same distribution, or is it different?

How could we use this idea in a simple research project?

POLS201
Spring 2019

- REMEMBER: The central limit theorem is the backbone for all hypothesis testing!
- We know how much random error will affect estimates. It's simply a function of standard deviation and sample size!
- We ask: "If they were pulled from the same distribution, what size of differences could be explained by mere randomness?"

Difference of Means Testing: Two Varieties

POLS201
Spring 2019

- Often, we can analyze sample means for an answer to this question. We might ask:
- Is a sample mean different than a population mean?
- Are two sample means different from each other?

Sample VS Population Means

POLS201
Spring 2019

- Say you are told told that $\mu = 12$

Sample VS Population Means

POLS201
Spring 2019

- Say you are told told that $\mu = 12$
- You are given a sample of 100 with a sample mean of 10 and a sample standard deviation of 4.

Sample VS Population Means

POLS201
Spring 2019

- Say you are told told that $\mu = 12$
- You are given a sample of 100 with a sample mean of 10 and a sample standard deviation of 4.
- You could build a confidence interval with this information OR calculate Z!

Let's ask:

POLS201
Spring 2019

- “How often would our sample distribution emerge from a population of with a mean of 12?”
- Our sample has a mean (\bar{x}) of 10 and an s of 4.
- Is it plausible that a distribution with $\mu = 12$ produced our sample?
- What are the ranges of plausible population means? How often would we see the sample mean of 10 produced from a population with mean of 10? Or 5? Or 20?

We compute the *confidence interval*


POLS201
Spring 2019

- The confidence interval defines _____ % of the possible population means that could have produced this result.
 - Often, we define this range as 95%.
- What is the range of means that defines these boundaries?

Observe the standard deviation of 2

POLS201
Spring 2019

- Notice that slightly more than 95% of the curve covers the area ± 2 standard deviations.
- With 1.96 standard deviations, we have covered 95% exactly.



How many standard deviations away is our population mean?

POLS201
Spring 2019

- A Z score of ± 1.96 will define those boundaries.
- The formula is this:

$$Z = \frac{(x - \mu)}{\sigma \div \sqrt{n}}$$

- For our example, the 95% interval is 11.216 to 12.784.
- Does 10 fall inside or outside that interval?
- Notice that we're figuring out a standard error, so we include \sqrt{n} in the denominator.

Remember: We apply the standard deviation of the sampling distribution

POLS201
Spring 2019

- Bigger samples will narrow the range of a confidence interval. Do you see why?

$$Z = \frac{(x - \mu)}{\sigma \div \sqrt{n}}$$

- In our example:

$$Z = (10 - 12) / (4 / 10) = -5$$

- A Z score of -5 is completely out of bounds. Which means: it's extremely unlikely that a population with a mean of 12 would have produced our sample!

Usually we compare two samples, instead of a sample v. population


POLS201
Spring 2019

- So we make a slight adjustment from our normal distribution
- We use the *Student-t Distribution* and compute a t-score instead of a Z-score.
- As the sample size increases, the t-distribution approaches the shape of a normal distribution.

The look of a t-distribution

POLS201
Spring 2019

- But when the sample size is small, the t-distribution is a bit flatter than normal, with fatter tails.



Interpretation of t

POLS201
Spring 2019

- We use t when we compare two *samples*
- The interpretation of t is identical to the interpretation of Z.
- High (or low) t values represent a difference that cannot be explained by chance alone.
- The “critical” t-value will vary on sample size, but with very large sample sizes it is precisely ± 1.96
- Considering sample size complicates the formula, however

Difference of Means and the T-Test

POLS201
Spring 2019

- With two samples, we ask a different question: what is the likelihood both samples came from the same population
- We can analyze this likelihood by comparing means, and using standard errors and sample sizes in the formula
- That formula is:

$$t = \frac{\overline{x_1} - \overline{x_2}}{\sqrt{\frac{s_1^2}{N_1} + \frac{s_2^2}{N_2}}}$$

But Use the Computer to Calculate a Difference of Means Test

POLS201
Spring 2019

- We want to suppose a distribution of means differences.
- In other words, a distribution of $\bar{x}_1 - \bar{x}_2$
- If the samples came from the same source, what is the confidence interval of this difference?
- Or: is the true difference of means $= 0$?
- At some point, the difference of means is too great, the same source couldn't have produced both samples.

Example

POLS201
Spring 2019

- A random sample of 300 men has an income of \$32,800 (sd of 4k)
- A random sample of 200 women has an income of \$32,000 (sd of 3k)
- Is this mean difference real or an artifact of random error?

R's t.test function can handle it

POLS201
Spring 2019

- Generate our samples

Suppose our means are identical. This really supports the null hypothesis

POLS201
Spring 2019

```
rnorm2 <- function(n,mean,sd) {mean+sd*scale(rnorm(n))

men <- rnorm2(300, 32000, 4000)
women <- rnorm2(200, 32000, 3000)
(d_mean <- t.test(men, women))

##
## Welch Two Sample t-test
##
## data:  men and women
## t = 0, df = 491.11, p-value = 1
## alternative hypothesis: true difference in means is
## 95 percent confidence interval:
## -616.1268  616.1268
## sample estimates:
## mean of x mean of y
```

Now let's say mean for men is 32,800 and 32,000 for women

POLS201
Spring 2019

```
rnorm2 <- function(n,mean,sd) {mean+sd*scale(rnorm(n))

men <- rnorm2(300, 32800, 4000)
women <- rnorm2(200, 32000, 3000)
(d_mean <- t.test(men, women))

##
## Welch Two Sample t-test
##
## data: men and women
## t = 2.5512, df = 491.11, p-value = 0.01104
## alternative hypothesis: true difference in means is
## 95 percent confidence interval:
## 183.8732 1416.1268
## sample estimates:
## mean of x mean of y
```

Let's bump the confidence level from .95 to .99

POLS201
Spring 2019

```
men <- rnorm2(300, 32800, 4000)
women <- rnorm2(200, 32000, 3000)
(d_mean <- t.test(men, women, conf.level = 0.99))

##
##  Welch Two Sample t-test
##
## data:  men and women
## t = 2.5512, df = 491.11, p-value = 0.01104
## alternative hypothesis: true difference in means is not equal to 0
## 99 percent confidence interval:
##    -10.88315 1610.88315
## sample estimates:
## mean of x mean of y
##      32800      32000
```


How to interpret?

POLS201
Spring 2019

- We see that the null hypothesis (difference = 0) is extremely unlikely.
- We would expect the differences to range between -10.88 and 1610.88 95% of the time.
- So (we argue) the difference is not just random error. Something else is producing the difference.

Do Catholics like Pope Francis better than non-Catholics?

POLS201
Spring 2019

- After some data wrangling from the Pew Survey, we see the following (1: Very favourable ... 4: Very unfavourable)
NB: Excuse the presentation

```
## # A tibble: 4 x 3
##               q84 Catholic `Not Catholic`
##               <dbl+lbl>      <int>      <int>
## 1 1 1 [Very favorable]          96        137
## 2 2 2 [Mostly favorable]       141        549
## 3 3 3 [Mostly unfavorable]    49         227
## 4 4 4 [Very unfavorable]     20         153
```

```
## # A tibble: 2 x 2
##   CATHOLIC      Mean
##   <chr>      <dbl>
## 1 Catholic    1.98
## 2 Not Catholic 2.37
```

```
t.test(q84 ~ CATHOLIC, sep_18)
```

```
##
```

```
## Welch Two Sample t-test
```

```
##
```

```
## data: q84 by CATHOLIC
```

```
## t = -7, df = 500, p-value = 6e-12
```

```
## alternative hypothesis: true difference in means is
```

```
## 95 percent confidence interval:
```

```
## -0.5044 -0.2843
```

```
## sample estimates:
```

```
## mean in group Catholic mean in group Not Catho
```

```
## 1.977
```

```
2.
```

This crude boxplot suggests a similar story

POLS201
Spring 2019

