

240 mm

↓ Front
210 mm

Size: 210 x 240 mm



LITHIUM CARBONATE

MICROLIT 400 mg Sustained-Release Tablet ANTIPSYCHOTIC

PRODUCT NAME: Microlit

NAME AND STRENGTH:

Lithium Carbonate Sustained Release tablets 400 mg

PHARMACOLOGIC CATEGORY: Psycholeptics, antipsychotics

PRODUCT DESCRIPTION:

White, flat, circular, bevel edged tablets with MICRO engraved on one side and plain on the other.

FORMULATION/COMPOSITION:

Each sustained-release tablet contains:

Lithium Carbonate BP400 mg

PHARMACODYNAMICS/PHARMACOKINETICS:

Pharmacodynamics:

Pharmacotherapeutic group: Psycholeptics, antipsychotics, ATC-code: N05AN01

Mechanism of action

The precise mechanism of action of lithium as a mood-stabilizing agent remains unknown, although many cellular actions of lithium have been characterised.

Pharmacokinetics:

Distribution

It crosses the placenta and is excreted in breast milk.

The pharmacokinetics of lithium are extremely well documented. A single oral dose of LITHIUM CARBONATE 400 gives a peak plasma level approximately 3-4 hours later, with the level at 24 hours being approximately 40% of peak levels.

Elimination

Lithium is excreted almost exclusively in the urine by the kidneys.

The half-life of lithium varies considerably between formulations, but generally is considered to be about 12 to 24 hours following a single dose.

Special populations

Elderly

Half-lives of up to 36 hours have been reported for elderly patients and 40 to 50 hours for patients with renal impairment. Steady-state concentrations may not be attained until 4 to 7 days after starting treatment.

INDICATIONS:

The treatment and prophylaxis of mania, manic-depressive illness and recurrent depression, and the treatment of aggressive or self mutilating behaviour.

DOSAGE AND MODEL ROUTE OF ADMINISTRATION:

Lithium Carbonate 400 mg tablets are usually administered according to a twice daily regimen.

When lithium levels have stabilized, a once daily regimen may be preferred

Lithium carbonate has a narrow therapeutic window. The dose required for treatment must be titrated and adjusted on the basis of regular monitoring of the serum concentration.

Lithium therapy should not be initiated unless adequate facilities for routine monitoring of plasma concentrations are available. On initiation of treatment, plasma therapy concentrations should be measured weekly until stabilization is achieved, then weekly for one month and at monthly intervals thereafter.

Additional measurements should be made if signs of lithium toxicity occur, on dosage alteration, development of significant intercurrent disease, signs of manic depressions or depressive relapse and if significant change in sodium or fluid intake occurs. More frequent monitoring is required if patients are receiving any drug treatment that affects renal clearance of lithium e.g. diuretics and NSAID. As bioavailability may vary between formulations, should a change of preparations be made, blood levels should be monitored weekly until destabilization is achieved.

Toxic symptoms are usually associated with concentrations exceeding 1.5 mmol/l and levels above 1.5mmol/l should be avoided. In the event of toxicity, lithium should be withdrawn immediately.

Withdrawal

If lithium is to be discontinued for other reasons, particularly in cases of high doses the dose should be reduced gradually over a suitable period of time, e.g. 2 weeks, to prevent the risk of relapse.

Posology

Acute mania:

Adults: Treatment should be initiated in hospital where regular monitoring of plasma lithium levels can be conducted. The dosage of Lithium Carbonate should be adjusted to produce a plasma lithium level between 0.6 and 1.0 mmol/l 12 hours after the last dose. The required plasma lithium level may be achieved in one of two ways but, whichever is adopted, regular estimations must be carried out to ensure maintenance of levels within the therapeutic range.

For consistent results it is essential that the blood samples for plasma lithium estimations are taken 12 hours after the last dose of lithium.

1. 1,000-1,500 mg of lithium carbonate are administered daily for the first five days. A blood sample for plasma lithium estimation is taken 12 hours after the last dose on the fifth day, and the dosage of Lithium Carbonate is adjusted to keep the plasma lithium level within the therapeutic range. Subsequently, regular plasma lithium estimations must be carried out and, where necessary, the dosage of Lithium Carbonate adjusted accordingly. The precise initial dose of lithium should be decided in the light of the age and weight of the patient; young patients often require a dose higher than average and older patients a lower dose.

2. A lithium clearance test is carried out and the initial dosage calculated from the results. Even when the initial dosage is calculated in this way, it is still desirable that plasma lithium levels should be determined at weekly intervals during the first three weeks of treatment, and any necessary adjustments to dosage made as a result of the levels actually obtained.

Most of the above applies in the treatment of hypomania as well as mania, but the patient (if not too ill) can be started on treatment as an outpatient provided that facilities for regular plasma lithium monitoring are available, and assays are initiated within one week.

Prophylaxis of recurrent affective disorders:

Adults: (Including unipolar mania & unipolar depressions and bipolar manic-depressive illness): A low dose of 300-400 mg of lithium carbonate can be administered daily for the first seven days. A blood sample for plasma lithium estimation is then taken 12 hours after the last dose, and the dosage of Lithium Carbonate is adjusted to keep the plasma lithium level within the range of 0.4-0.8 mmol/l.

Aggressive and self-mutilating behaviour:

Adults: Dosage is at the lower end of the range for the treatment for manic depressive illness.

Special Populations

Elderly

Start treatment with a low dose.

Elderly patients often require lower lithium dosage to achieve therapeutic serum levels. As for prophylaxis above, but 12-hour lithium levels should be kept in the range of 0.4-0.7 mmol/l as toxic symptoms are likely with plasma concentrations above 1.0 mmol/l. Toxic symptoms are more likely at lower concentrations than in the general population.

Paediatric population

Lithium Carbonate should not be used in children.

Method of administration

For oral administration

CONTRAINDICATIONS & PRECAUTION(S), WARNING(S):

Hypersensitivity to the active substance or to any of the excipients.

- Severely impaired renal function
- Untreated or untreatable hypothyroidism.
- Cardiac disease associated with rhythm disorder.
- Brugada syndrome or family history of Brugada syndrome
- Low body sodium levels for example dehydrated patients, those on low sodium diets, or those with Addison's disease.
- Breast-feeding.

PRECAUTIONS & WARNINGS:

Lithium carbonate has a narrow therapeutic window. The dose required for treatment must be titrated and adjusted on the basis of regular monitoring of serum concentration of lithium. Lithium therapy should not be initiated unless adequate facilities for routine monitoring of plasma concentrations are available.

Elderly patients are particularly liable to lithium toxicity. Use with care as lithium excretion may also be reduced. They may also exhibit adverse reactions at serum levels ordinarily tolerated by younger patients.

Before beginning a lithium treatment

-It is important to ensure that renal function is evaluated

-Thyroid function should be evaluated. Patients should be euthyroid before initiation of lithium therapy.

-Cardiac function should be assessed especially in patients with cardiovascular disease.

Renal, cardiac and thyroid functions should be re-assessed periodically.

Risk of convulsions

The risk of convulsions may be increased when lithium is co-administered with drugs that lower the epileptic threshold, or in epileptic patients.

Benign intracranial hypertension

There have been case reports of benign intracranial hypertension). Patients should be warned to report persistent headache and/or visual disturbances.

QT prolongation

As a precautionary measure, lithium should be avoided in patients with congenital long QT syndrome, and in patients concomitantly treated with drugs that are known to prolong the QT interval. Caution should be exercised in patients with risk factors for QT interval prolongation (which include cardiac disease, bradycardia, thyroid disease, hypokalaemia, hypomagnesaemia, hypocalcaemia, female sex and advanced age.

Brugada syndrome

Lithium may unmask or aggravate Brugada syndrome, a hereditary disease of the cardiac sodium channel with characteristic ECG changes (right bundle branch block and ST segment elevation in right precordial leads), which may lead to cardiac arrest or sudden death. Lithium should not be administered to patients with Brugada syndrome or a family history of Brugada syndrome Caution is advised in patients with a family history of cardiac arrest or sudden death.

Concomitant administration of antipsychotics

Concomitant administration of antipsychotics should be avoided.

Monitoring of blood lithium levels

Serum concentration of lithium should be measured on a sample taken just prior to the time when a dose of lithium is due to be taken (i.e. at trough level 12 hours following the last dose).

Toxic effects may be expected at serum-lithium concentrations of about 1.5

mmol/litre, although they can appear at lower concentrations. They call for immediate withdrawal of treatment and should always be considered very seriously

Serum concentration of lithium should be measured every 5 to 7 days from initiation until stabilization is achieved and at regular intervals for the duration of treatment.

Serum lithium concentrations should be monitored more frequently (revert to weekly monitoring) in the following circumstances:

- Dosage alteration or change of lithium formulation (bioavailability may differ)
- Significant intercurrent disease
- Intercurrent infection
- Significant change in sodium intake
- Significant change in fluid intake
- Treatment with drugs altering renal clearance of lithium
- Treatment with drugs likely to upset electrolyte balance.

Patients should also be warned to report if polyuria or polydipsia develops. Episodes of nausea and vomiting or other conditions leading to salt/water depletion (including severe dieting) should also be reported. Patients should be advised to maintain their usual salt and fluid intake.

Lithium should be stopped 24 hours before major surgery, but the normal dose can be continued for minor surgery if fluids and electrolytes are carefully monitored

Renal impairment

Lithium excretion is reduced in the presence of renal impairment. This increases the risk of toxicity. Lithium is contra-indicated in patients with severe renal impairment. If patients with mild or moderate renal impairment are being treated with lithium, serum levels should be closely monitored.

Renal function should be monitored in patients with renal impairment, and in patients with polyuria and polydipsia.

Warnings to be given to patients about signs and symptoms of toxicity

Clear instructions regarding the symptoms of impending toxicity should be given by the doctor to all patients receiving long-term lithium therapy and advice given for the need for urgency in seeking medical assistance if these symptoms appear.

Renal tumours: cases of microcysts, oncocytomas and collecting duct renal carcinoma have been reported in patients with severe renal impairment who received lithium for more than 10 years

PREGNANCY & LACTATION:

Pregnancy

Lithium therapy should not be used during pregnancy, especially during the first trimester, unless considered essential. There is epidemiological evidence that it may be harmful to the foetus in human pregnancy.

Lithium crosses the placental barrier. An increase in cardiac and other abnormalities, especially Ebstein anomaly, are reported. Therefore, a pre-natal diagnosis such as ultrasound and electrocardiogram examination is strongly recommended. In certain cases where a severe risk to the patient could exist if treatment were stopped, lithium has been continued during pregnancy. Studies in animals have shown reproductive toxicity.

If it is considered essential to maintain lithium treatment during pregnancy, serum lithium levels should be closely monitored and measured frequently since renal function changes gradually during pregnancy and suddenly at parturition. Dosage adjustments are required. It is recommended that lithium be discontinued shortly before delivery and reinitiated a few days *post-partum*.

Neonates may show signs of lithium toxicity necessitating fluid therapy in the neonatal period. Neonates born with low serum lithium concentrations may have a flaccid appearance that returns to normal without any treatment.

Women of child-bearing potential

It is advisable that women treated with lithium should adopt adequate

↓ Front
210 mm

Size: 210 x 240 mm

240 mm

contraceptive methods. In case of a planned pregnancy, it is strongly recommended to discontinue lithium therapy.

Lactation

Since adequate human data on use during lactation and adequate human reproduction studies are not available and as lithium is secreted in breast milk, bottle-feeding is recommended.

Fertility

Studies in animals have shown adverse effects on male fertility

INTERACTIONS:

Interactions may occur as a result of increased or decreased lithium levels, or may act through other mechanisms, the most important being neurotoxicity which may occur at therapeutic levels when other drugs which act centrally on the CNS are taken concurrently.

Interactions which increase lithium concentrations

Co-administration of the following drugs with lithium may lead to increased lithium concentrations and a risk of toxicity:

- Any drug which may cause renal impairment has the potential to cause lithium levels to rise, thereby causing toxicity. If the use of the drug is unavoidable, carefully monitor lithium blood level and adapt dosage as necessary.
- Antibiotics (metronidazole, tetracyclines, co-trimoxazole, trimethoprim), N.B. Toxic symptoms may also occur at low or normal levels when used in conjunction with co-trimoxazole or trimethoprim. Lithium toxicity has been reported on isolated occasions in patients receiving spectinomycin.
- Non-steroidal anti-inflammatory drugs (including selective cyclooxygenase (COX) II inhibitors); monitor serum lithium concentrations more frequently if NSAID therapy is initiated or discontinued.
- Drugs affecting the renin angiotensin system (ACE inhibitors, Angiotensin II receptor antagonists).
- Diuretics (including herbal preparations). In addition to the effects noted above, thiazide diuretics show a paradoxical antidiuretic effect resulting in possible water retention and lithium intoxication. Loop diuretics (furosemide and bumetanide, and ethacrynic acid) seem less likely to cause lithium retention, although caution is warranted.
- Other drugs affecting electrolyte balance, e.g. steroids, may alter lithium excretion and should therefore be avoided.

Interactions which decrease serum lithium concentrations:

Co-administration of the following drugs with lithium may lead to decreased lithium concentrations and a risk of loss of efficacy:

- xanthine derivatives (e.g. theophylline, caffeine).
- products containing large quantities of sodium e.g. sodium bicarbonate.
- carbonic anhydrase inhibitors.
- Urea.

Interactions which may not be associated with increased or reduced lithium levels:

Concomitant use of the following drugs may precipitate symptoms of toxicity when the lithium level is within the normal range:

- antipsychotics, including the atypical antipsychotics olanzapine, clozapine and haloperidol at high doses
- carbamazepine
- phenytoin
- methylodopa
- clonazepam
- tricyclic and tetracyclic antidepressants
- calcium channel blockers. These drugs may cause neurotoxic reactions at therapeutic levels
- neuromuscular blocking agents. Lithium may cause neurotoxic reactions at therapeutic lithium levels.

Selective serotonin re-uptake inhibitors (SSRIs): Concurrent use with lithium may precipitate a serotonergic syndrome.

Non-steroidal anti-inflammatory drugs including COX II inhibitors: monitor serum lithium concentrations more frequently if NSAID therapy is initiated or discontinued

Triptans: lithium toxicity reported suggestive of serotonin syndrome.

Neuromuscular blockers: Lithium may prolong the effects of neuromuscular blocking agents.

Drugs which lower seizure threshold

Caution is advised if lithium is co-administered with drugs that lower the epileptic threshold. e.g. antidepressants, antipsychotics, anaesthetics and theophylline.

Drugs which prolong the QT interval

Lithium can cause an increase in the QTc interval, particularly at higher blood levels. Therefore, concurrent use of drugs which have a risk of prolonging the QTc interval should be avoided, and consideration be made of other potential risk factors such as increasing age, female sex, congenital long QT syndrome, cardiac and thyroid disease and the following metabolic disturbances: hypocalcaemia, hypokalaemia, hypomagnesaemia.

The following products have a high risk of causing QT prolongation and torsade de pointes:

- Class Ia antiarrhythmics, (Ajmaline, cibenzoline, disopyramide, hydroquinidine, procainamide, quinidine),
- Class III antiarrhythmics (amiodarone, azimilide, cibenzoline, dofetilide, ibutilide, sotalol),
- Antipsychotics (Amisulpride, haloperidol, Droperidol, mesoridazine, pimozide, sertindole, thioridazine and Clozaril),

- Antibiotics (intravenous erythromycin, sparflloxacin),
- Serotonin antagonists (ketanserine, dolasetron mesylate),
- Antihistamines (astemizole, terfenadine),
- Antimalarials (artemisinin derivatives, mefloquine, halofantrine),
- Other: arsenic trioxide, cisapride and ranolazine.
- ECG should be performed after initiation of treatment and at any point where the patient becomes symptomatic or when there are changes in disease or treatment which may increase the risk of interaction or arrhythmia.

Non-Drug Interactions:

- Low sodium diet. Rapid reduction of sodium intake may cause raised lithium levels.
- Intercurrent illness may cause lithium toxicity.

ADVERSE EFFECTS:

Side effects are usually related to serum lithium concentrations and are less common in patients with plasma lithium concentrations below 1.0 mmol/l.

Initial Therapy: fine tremor of the hands, polyuria and thirst may occur.

Blood and lymphatic system disorders: leukocytosis.

Immune system disorders: increase in antinuclear antibodies.

Endocrine disorders: disturbances of thyroid function including (euthyroid) goiter, hypothyroidism and hyperthyroidism, hyperparathyroidism, parathyroid adenoma.

Metabolism and nutrition disorders: hypercalcaemia, hypomagnesaemia, Hyperglycaemia, anorexia, weight gain.

Nervous system disorders: coma, benign intracranial hypertension, syndrome of irreversible lithium effectuated neurotoxicity (SILENT), encephalopathy, stupor, seizures, neuroleptic malignant syndrome, myasthenia gravis, serotonin syndrome, parkinsonism, extrapyramidal symptoms, ataxia, dizziness, memory impairment, mild cognitive impairment may occur during long term use, giddiness, nystagmus, slurred speech, vertigo, hyperactive deep tendon reflexes, dazed feeling, fine hand tremors.

Eye Disorders: scotomata and blurred vision

Cardiac disorders: cardiac arrest, ventricular fibrillation, ventricular tachycardia, ventricular arrhythmias, Torsade de pointes, QT interval prolongation, cardiomyopathy, arrhythmia, bradycardia, sinus node dysfunction, ECG changes.

Vascular disorders: peripheral circulatory collapse, hypotension.

Gastrointestinal disorders: gastritis, nausea, diarrhoea, vomiting, dry mouth, excessive salivation. Lithium salts have been implicated in dysgeusia.

Skin and subcutaneous tissue disorders: Allergic rash, exacerbation of psoriasis, acneiform eruptions, alopecia, acne, papular skin disorder, folliculitis, pruritus, rash.

Musculoskeletal and connective tissue disorders: muscle weakness.

Renal and urinary disorders: symptoms of nephrogenic diabetes insipidus, impairment of renal function, permanent changes in the kidney, nephrotic syndrome, histological renal changes with interstitial fibrosis after long term treatment, polyuria, polydipsia. Frequency unknown: Microcysts, oncocytomas and collecting duct renal carcinoma (in long-term therapy).

Reproductive system and breast disorders: sexual dysfunction.

General disorders and administration site conditions: sudden unexplained death, oedema, asthenia, lethargy, thirst, fatigue, and malaise can occur due to lithium toxicity.

OVERDOSEAGE AND TREATMENT:

In patients with a raised lithium concentration, the risk of toxicity is greater in those with the following underlying medical conditions: hypertension, diabetes, congestive heart failure, chronic renal failure, schizophrenia, Addison's disease.

Acute

A single acute overdose usually carries low risk and patients tend to show mild symptoms only, irrespective of their serum lithium concentration. However more severe symptoms may occur after a delay if lithium elimination is reduced because of renal impairment, particularly if a slow-release preparation has been taken. The fatal dose, in a single overdose, is probably over 5g.

If an acute overdose has been taken by a patient on chronic lithium therapy, this can lead to serious toxicity occurring even after a modest overdose as the extravascular tissues are already saturated with lithium.

Chronic

Lithium toxicity can also occur in chronic accumulation for the following reasons: Acute or chronic overdose; dehydration e.g. due to intercurrent illness, deteriorating renal function, drug interactions, most commonly involving a thiazide diuretic or a non-steroidal anti-inflammatory drug (NSAID).

Symptoms

The onset of symptoms may be delayed, with peak effects not occurring for as long as 24 hours, especially in patients who are not receiving chronic lithium therapy or following the use of a sustained release preparation.

Symptoms of lithium intoxication include:

Mild: Nausea, diarrhoea, blurred vision, polyuria, light headedness, fine resting tremor, muscular weakness and drowsiness.

Moderate: Increasing confusion, blackouts, fasciculation and increased deep tendon reflexes, myoclonic twitches and jerks, choreoathetosis movements, urinary or faecal incontinence,

increasing restlessness followed by stupor. Hyponatremia

Severe: Coma, convulsions, cerebellar signs, cardiac dysrhythmias including sinoatrial block, sinus and junctional bradycardia and first degree heart block. Hypotension or rarely hypertension, circulatory collapse and renal failure

Others:

Gastrointestinal disorders: increasing anorexia and vomiting.

Nervous system disorders: Encephalopathy, cerebellar syndrome with symptoms such as muscle weakness, lack of coordination, drowsiness or lethargy, giddiness, ataxia, nystagmus, coarse tremor. Tinnitus, dysarthria, twitching, myoclonus, extrapyramidal disorders ECG changes (flat or inverted T waves, QT prolongation), AV block, dehydration and electrolyte disturbances

At blood levels above 2-3 mmol/l, there may be a large output of dilute urine and renal insufficiency, with increasing confusion, convulsions, coma and death.

Management

There is no specific antidote to lithium. In the event of lithium overdose, lithium should be discontinued and lithium serum levels monitored closely.

Supportive treatment should be initiated, which includes correction of fluid and electrolyte balance, if necessary.

Diuretics should not be used. All patients should be observed for a minimum of 24 hours. ECG should be monitored in symptomatic patients. Steps should be taken to correct hypotension.

Consider gastric lavage for non-sustained-release preparations if more than 4 g has been ingested by an adult within 1 hour or definite ingestion of a significant amount by a child. Slow-release tablets do not disintegrate in the stomach and most are too large to pass up a lavage tube. Gut decontamination is not useful for chronic accumulation. Activated charcoal does not adsorb lithium.

Hemodialysis is the treatment of choice for severe lithium intoxication (especially in patients manifesting with severe nervous system disorders), or in cases of overdose accompanied by renal impairment. Hemodialysis should be continued until there is no lithium in the serum or dialysis fluid. Serum lithium levels should be monitored for at least another week to take account of any possible rebound in serum lithium levels as a result of delayed diffusion from the body tissues.

In cases of acute on chronic overdose or in cases of chronic lithium toxicity if the lithium concentration is >4.0 mmol/l, discuss with your local poisons service.

Clinical improvement generally takes longer than reduction of serum lithium concentrations regardless of the method used.

STORAGE CONDITION:

Store at temperatures not exceeding 30°C.

DOSAGE FORMS AND PACKAGING AVAILABLE:

Foil Strip of 10's (Box of 30 and 100 Tablets)

INSTRUCTIONS AND SPECIAL PRECAUTIONS FOR HANDLING AND DISPOSAL (IF APPLICABLE):

Not Applicable

NAME AND ADDRESS OF MARKETING AUTHORIZATION HOLDER:

Marketing Authorization Holder

Brown & Burk Philippines Inc
U-501, 5/F., SEDCCO 1 Bldg., 120 Rada cor.,
Legaspi Sts., Legaspi Village, Makati City, Philippines

NAME AND ADDRESS OF MANUFACTURER:

MICRO LABS LIMITED

92, Sipcot Industrial Complex,
Hosur – 635 126, India.

CAUTION STATEMENT:

FOODS, DRUGS, DEVICES, AND COSMETICS ACT PROHIBITS DISPENSING WITHOUT PRESCRIPTION.

ADR REPORTING STATEMENT:

"FOR SUSPECTED ADVERSE DRUG REACTION, REPORT TO THE

FDA: www.fda.gov/ph

Seek medical attention immediately at the first sign of Adverse Drug Reaction.

REGISTRATION NUMBER:

DR No.: XY41516

DATE OF FIRST AUTHORIZATION:

27 December 2012

DATE OF REVISION OF PACKAGE INSERT:

Feb. 2017.

EXG-ML011-1102/A