

# STAT40810 — Stochastic Models


Brendan Murphy

Week 10

## Multistate Markov Models


# Multistate Markov Model

- Consider the following process being proposed to model a health/illness scenario.



# Multistate Markov Model


- Consider the following process being proposed to model a health/illness scenario.



- The process has four states:
  - healthy (h)
  - ill (i)
  - seriously ill (s)
  - dead (d)

# Multistate Markov Model


- Consider the following process being proposed to model a health/illness scenario.



- The process has four states:
  - healthy (h)
  - ill (i)
  - seriously ill (s)
  - dead (d)
- And people can move between states in *continuous time*.

# Multistate Markov Model

- Consider the following process being proposed to model a health/illness scenario.



- The process has four states:
  - healthy (h)
  - ill (i)
  - seriously ill (s)
  - dead (d)
- And people can move between states in *continuous time*.
- The flow between states is controlled by unknown parameters.

# Assumptions

- Consider the process at time  $t$  and what happens before time  $t + dt$ .
- We make the following assumptions:
  - ① The probabilities of being in any state at any subsequent time depends only on the time involved and on the state currently occupied.

# Assumptions

- Consider the process at time  $t$  and what happens before time  $t + dt$ .
- We make the following assumptions:
  - ① The probabilities of being in any state at any subsequent time depends only on the time involved and on the state currently occupied.
  - ② We assume a constant transition intensity  $\lambda_{gh}$ .

# Assumptions

- Consider the process at time  $t$  and what happens before time  $t + dt$ .
- We make the following assumptions:
  - ① The probabilities of being in any state at any subsequent time depends only on the time involved and on the state currently occupied.
  - ② We assume a constant transition intensity  $\lambda_{gh}$ .
  - ③ The probability of transitioning from state  $g$  to state  $h$  between time  $t$  and  $t + dt$  is  $\lambda_{gh}dt + o(dt)$  as  $dt \rightarrow 0$ .



# Assumptions

- Consider the process at time  $t$  and what happens before time  $t + dt$ .
- We make the following assumptions:
  - ① The probabilities of being in any state at any subsequent time depends only on the time involved and on the state currently occupied.
  - ② We assume a constant transition intensity  $\lambda_{gh}$ .
  - ③ The probability of transitioning from state  $g$  to state  $h$  between time  $t$  and  $t + dt$  is  $\lambda_{gh}dt + o(dt)$  as  $dt \rightarrow 0$ .
  - ④ The probability of transitioning between two or more states is  $o(dt)$  as  $dt \rightarrow 0$ .

# Data

- Suppose for a cohort of people we observed the following data:

- Suppose for a cohort of people we observed the following data:

Transitions	Count
$h$ to $i$	$n_{hi}$
$i$ to $h$	$n_{ih}$
$i$ to $s$	$n_{is}$
$s$ to $i$	$n_{si}$
$i$ to $d$	$n_{id}$
$s$ to $d$	$n_{sd}$

# Data

- Suppose for a cohort of people we observed the following data:

Transitions	Count
$h$ to $i$	$n_{hi}$
$i$ to $h$	$n_{ih}$
$i$ to $s$	$n_{is}$
$s$ to $i$	$n_{si}$
$i$ to $d$	$n_{id}$
$s$ to $d$	$n_{sd}$

- And...

State	Time
$h$	$t_h$
$i$	$t_i$
$s$	$t_s$
$d$	$t_d$

# Data

- Suppose for a cohort of people we observed the following data:

Transitions	Count
$h$ to $i$	$n_{hi}$
$i$ to $h$	$n_{ih}$
$i$ to $s$	$n_{is}$
$s$ to $i$	$n_{si}$
$i$ to $d$	$n_{id}$
$s$ to $d$	$n_{sd}$

- And...

State	Time
$h$	$t_h$
$i$	$t_i$
$s$	$t_s$
$d$	$t_d$

- We have enough information to form the likelihood function.


# Example

- The following data were collected from 100 individuals who were monitored:
  - total waiting time in state h: 4,100 months
  - total waiting time in state i: 1,000 months
  - total waiting time in state s: 500 months
  - total number of transfers from state h to state i: 110
  - total number of transfers from state i to state h: 90
  - total number of transfers from state i to state s: 40
  - total number of transfers from state s to state i: 20
  - total number of transfers from state i to state d: 7
  - total number of transfers from state s to state d: 10

# Likelihood

- The likelihood is of the form

$$L = \exp(-\lambda_{hi}t_h) \exp(-(\lambda_{ih} + \lambda_{is} + \lambda_{id})t_i) \exp(-(\lambda_{si} + \lambda_{sd})t_s) \\ \lambda_{hi}^{n_{hi}} \lambda_{ih}^{n_{ih}} \lambda_{is}^{n_{is}} \lambda_{id}^{n_{id}} \lambda_{si}^{n_{si}} \lambda_{sd}^{n_{sd}}$$



- Note that  $t_d$  doesn't enter the expression.

# Parameter Estimates

- The model parameters can be easily found using maximum likelihood.
- It turns out (check) that

$$\hat{\lambda}_{gh} = \frac{n_{gh}}{t_g}$$

for any pair of states  $g$  and  $h$ .



# Parameter Estimates

- The model parameters can be easily found using maximum likelihood.
- It turns out (check) that

$$\hat{\lambda}_{gh} = \frac{n_{gh}}{t_g}$$

for any pair of states  $g$  and  $h$ .

- Furthermore,

$$SE(\hat{\lambda}_{gh}) = \frac{\sqrt{n_{gh}}}{t_g}.$$

- Hence, the MLEs are

$$\hat{\lambda}_{hi} = \frac{n_{hi}}{t_h} = \frac{110}{4100}$$

$$\hat{\lambda}_{ih} = \frac{n_{ih}}{t_i} = \frac{90}{1000}$$

$$\hat{\lambda}_{is} = \frac{n_{is}}{t_i} = \frac{40}{1000}$$

$$\hat{\lambda}_{id} = \frac{n_{id}}{t_i} = \frac{7}{1000}$$

$$\hat{\lambda}_{si} = \frac{n_{si}}{t_s} = \frac{20}{500}$$

$$\hat{\lambda}_{sd} = \frac{n_{sd}}{t_s} = \frac{10}{500}$$

- Hence, approximate 95% confidence intervals for the intensities are

$$\lambda_{hi} \in (0.022, 0.032)$$

$$\lambda_{ih} \in (0.071, 0.109)$$

$$\lambda_{is} \in (0.028, 0.052)$$

$$\lambda_{id} \in (0.002, 0.012)$$

$$\lambda_{si} \in (0.022, 0.058)$$

$$\lambda_{sd} \in (0.008, 0.032)$$