

¹ Status of Big Skate (*Beringraja binoculata*)
² Off the U.S. Pacific Coast in 2019



⁴ Ian G. Taylor¹
⁵ Vladlena Gertseva¹
⁶ Andi Stephens²
⁷ Joseph Bizzarro³

⁸ ¹Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and
⁹ Atmospheric Administration, National Marine Fisheries Service, 2725 Montlake Boulevard East,
¹⁰ Seattle, Washington 98112

¹¹ ²Northwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and
¹² Atmospheric Administration, National Marine Fisheries Service, 2032 S.E. OSU Drive Newport,
¹³ Oregon 97365

¹⁴ ³Southwest Fisheries Science Center, U.S. Department of Commerce, National Oceanic and
¹⁵ Atmospheric Administration, National Marine Fisheries Service, 110 Shaffer Road, Santa Cruz,
¹⁶ California 95060

¹⁷ DRAFT SAFE

¹⁸ Disclaimer: This information is distributed solely for the purpose of pre-dissemination peer review
¹⁹ under applicable information quality guidelines. It has not been formally disseminated by NOAA
²⁰ Fisheries. It does not represent and should not be construed to represent any agency
²¹ determination or policy.

²² This report may be cited as:

²³ Taylor, I.G., Gertseva, V., Stephens, A. and Bizzarro, J. Status of Big Skate (*Beringraja*
²⁴ *binoculata*) Off the U.S. West Coast, 2019. Pacific Fishery Management Council, Portland, OR.
²⁵ Available from <http://www.p council.org/groundfish/stock-assessments/>

²⁶ **Acronyms used in this Document**

ABC	Allowable Biological Catch
ACL	Annual Catch Limit
ADFG	Alaska Department of Fish and Game
AFSC	Alaska Fisheries Science Center
A-SHOP	At-Sea Hake Observer Program
CalCOM	California Cooperative Groundfish Survey
CDFW	California Department of Fish and Wildlife
CPFV	Commercial Passenger Fishing Vessel
CRFS	California Recreational Fisheries Survey
DFO	Canada's Department of Fisheries and Oceans
IFQ	Individual Fishing Quota
IPHC	International Pacific Halibut Commission
MRFSS	Marine Recreational Fisheries Statistics Survey
NMFS	National Marine Fisheries Service
NORPAC	the North Pacific Database Program
NWFSC	Northwest Fisheries Science Center
ODFW	Oregon Department of Fish and Wildlife
OFL	Overfishing Limit
ORBS	Oregon Recreational Boat Survey
OY	Optimum Yield
PacFIN	Pacific Fisheries Information Network
PFMC	Pacific Fishery Management Council
PSMFC	Pacific States Marine Fisheries Commission
RecFIN	Recreational Fisheries Information Network
SPR	Spawning Potential Ratio
SSC	Scientific and Statistical Committee
SWFSC	Southwest Fisheries Science Center
WCGOP	West Coast Groundfish Observer Program
WDFW	Washington Department of Fish and Wildlife

27 Status of Big Skate (*Beringraja binoculata*)
28 Off the U.S. Pacific Coast in 2019

29 **Contents**

30 Executive Summary	1
31 Stock	1
32 Catches	2
33 Data and Assessment	4
34 Stock Biomass	5
35 Recruitment	7
36 Exploitation Status	9
37 Reference Points	11
38 Ecosystem Considerations	12
39 Management Performance	12
40 Unresolved Problems and Major Uncertainties	12
41 Decision Table	12
42 Projected Landings, OFLs and Time-varying ACLs	13
43 Research and Data Needs	17
44 1 Introduction	18
45 1.1 Biology	18
46 1.2 Distribution and Life History	19
47 1.3 Ecosystem Considerations	20
48 1.4 Fishery Information	21
49 1.5 Stock Status and Management History	21
50 1.6 Fisheries Off Alaska, Canada and Mexico	22
51 1.6.1 Alaska	22
52 1.6.2 Canada	23
53 1.6.3 Mexico	23

54	2 Fishery Data	24
55	2.1 Data	24
56	2.2 Fishery Landings and Discards	24
57	2.2.1 Washington Commercial Skate Landings Reconstruction	24
58	2.2.2 Oregon Commercial Skate Landings Reconstruction	25
59	2.2.3 California Catch Reconstruction	26
60	2.2.4 Tribal Catch in Washington	26
61	2.2.5 Fishery Discards	26
62	3 Fishery-Independent Data Sources	28
63	3.1 Indices of abundance	28
64	3.1.1 Alaska Fisheries Science Center (AFSC) Triennial Shelf Survey	28
65	3.1.2 Northwest Fisheries Science Center West Coast Groundfish Bottom	
66	Trawl Survey	28
67	3.1.3 Index Standardization	29
68	3.1.4 Internation Pacific Halibut Commission Longline Survey	29
69	4 Biological Parameters and Data	31
70	4.1 Measurement Details and Conversion Factors	31
71	4.2 Fishery dependent length and age composition data	31
72	4.3 Survey length and age composition data	31
73	4.4 Environmental or Ecosystem Data Included in the Assessment	32
74	5 Assessment	33
75	5.1 Previous Assessments	33
76	5.2 Model Description	33
77	5.2.1 Modeling Software	33
78	5.2.2 Summary of Data for Fleets and Areas	33
79	5.2.3 Other Specifications	34
80	5.2.4 Data Weighting	34
81	5.2.5 Priors	34
82	5.2.6 Estimated Parameters	35
83	5.2.7 Fixed Parameters	36
84	5.3 Model Selection and Evaluation	37

85	5.3.1 Key Assumptions and Structural Choices	37
86	5.3.2 Alternate Models Considered	37
87	5.3.3 Convergence	37
88	5.4 Response to the Current STAR Panel Requests	37
89	5.5 Base Case Model Results	38
90	5.5.1 Parameter Estimates	38
91	5.5.2 Fits to the Data	38
92	5.5.3 Uncertainty and Sensitivity Analyses	39
93	5.5.4 Retrospective Analysis	39
94	5.5.5 Likelihood Profiles	40
95	5.5.6 Reference Points	40
96	6 Harvest Projections and Decision Tables	41
97	7 Regional Management Considerations	42
98	8 Research Needs	43
99	9 Acknowledgments	43
100	10 Tables	44
101	10.1 Data Tables	44
102	10.2 Model Results Tables	50
103	11 Figures	66
104	11.1 Data Figures	66
105	11.2 Biology Figures	74
106	11.3 Model Results Figures	82
107	11.3.1 Growth and Selectivity	82
108	11.3.2 Fits to the Data	85
109	11.3.3 Time Series Figures	96
110	11.3.4 Sensitivity Analyses and Retrospectives	102
111	11.3.5 Likelihood Profiles	106
112	11.3.6 Reference Points and Forecasts	113

¹¹³ Appendix A. Detailed fits to length composition data

A-1

¹¹⁴ References

¹¹⁵ **Executive Summary**

¹¹⁶ **Stock**

¹¹⁷ This assessment reports the status of the Big Skate (*Beringraja binoculata*) resource in U.S.
¹¹⁸ waters off the West Coast using data through 2018. A map showing the area of the U.S.
¹¹⁹ West Coast Exclusive Economic Zone covered by this stock assessment is provided in Figure
¹²⁰ a.



Figure a: U.S. West Coast Exclusive Economic zone covering the area in which this stock assessment is focused.

¹²¹ **Catches**

¹²² Landings and estimated discards of Big Skate were reconstructed for this assessment from
¹²³ historical records of other species and from species composition data collected in the recent
¹²⁴ fishery. These reflect the fishery from 1916-1994. The current fishery started in 1995. For
¹²⁵ records from 1995-2017, Big Skate landings were estimated from species-composition samples
¹²⁶ and the landings of “Unspecified Skates”. Beginning in 2017, Big Skate have been recorded
¹²⁷ in species-specific landings.

¹²⁸ In the current fishery (since 1995), annual total landings of Big Skate have ranged between
¹²⁹ 135-528 mt, with landings in 2018 totaling 173 mt.

Table a: Recent Big Skate landings (mt)

Year	Landings
2008	366.00
2009	205.70
2010	196.20
2011	268.40
2012	269.60
2013	135.00
2014	372.40
2015	331.50
2016	411.50
2017	277.60
2018	172.60




Figure b: Catch history of Big Skate in the model.

130 **Data and Assessment**

131 This the first full assessment for Big Skate. It is currently managed using an OFL which was
132 based on a proxy for F_{MSY} and a 3-year recent average of survey biomass. This assessment
133 uses the newest version of Stock Synthesis (3.30.13). The model begins in 1916, and assumes
134 the stock was at an unfished equilibrium that year.

135 The assessment relies on two bottom trawl survey indices of abundance, the Triennial Survey
136 from an index covering the period 1980–2004 was used here and the West Coast Groundfish
137 Bottom Trawl (WCGBT) Survey, which began in 2003 and for which data is available through
138 2018. The triennial survey shows an increasing trend over the 25 year period it covers, which
139 the model is not able to fit as this includes the peak period of the fishery when the stock
140 would have been expected to be declining. The WCGBT Survey also shows an increasing
141 trend, with the 5 most recent observations (2014–2018) all falling in the top 6 ever observed
142 (2004 was the 5th highest observation). The most estimates an increasing trend during this
143 period but the slope is more gradual than the trend in the survey. The misfit to these
144 survey indices could be due to some combination of incorrect estimation of the catch history,
145 variability in recruitment which is not modeled here, or biological or ecological changes for
146 which data are not available.

147 Length composition data from the fishery is available starting in 1995 but is sparse until the
148 past decade. Most of the ages are also from 2008 onward. This limits the ability of the model
149 to estimate any changes composition of the population during the majority of the history of
150 the fishery. Estimates of discard rates and mean body weight of discards are available for
151 the years 2002 onward and discard length compositions are available starting in 2010.

152 The age and length data provide evidence for growth patterns and sex-specific differences
153 in selectivity that are unusual among groundfish stocks that have been assessed within the
154 U.S. West Coast and are not found in Longnose Skate where the data show little difference
155 between the sexes. Growth appears to be almost linear and similar between females and
156 males up to about age 7 or over 100 cm at which point male growth appears to stabilize
157 while females continue to grow. However, in spite of the similar growth pattern for ages prior
158 to 7, males are observed more frequently, with the 70–100 cm length bins often showing 60%
159 males. Sex-specific differences in selectivity were included in the model in order to better
160 match patterns in the sex ratios in the length composition data. The length and age data
161 do not cover enough years or show enough evidence of distinct cohorts to reliably estimate
162 deviations in recruitment around the stock-recruit curve.

163 The scale of the population is not reliably informed by the data due to the combination of
164 surveys that show trends which can't be matched by the structure of the model and length
165 and age data which inform growth and selectivity but provide relatively little information
166 about changes in stock structure over time. Therefore, a prior on catchability of the WCGBT
167 Survey (centered at 0.83) was applied in order to provide more stable results.

168 Although the assessment model requires numerous simplifying assumptions, it represents an

¹⁶⁹ improvement over the simplistic status-quo method of setting management limits, which re-
¹⁷⁰ lies on average survey biomass and an assumption about F_{MSY} . The use of an age-structured
¹⁷¹ model with estimated growth, selectivity, and natural mortality likely provide a better esti-
¹⁷² mate of past dynamics and the impacts of fishing in the future.

¹⁷³ Stock Biomass

¹⁷⁴ The 2018 estimated spawning biomass relative to unfished equilibrium spawning biomass is
¹⁷⁵ above the target of 40% of unfished spawning biomass at 75.0% (95% asymptotic interval:
¹⁷⁶ $\pm 63.9\%-86.0\%$) (Figure ??). Approximate confidence intervals based on the asymptotic
¹⁷⁷ variance estimates show that the uncertainty in the estimated spawning biomass is high,
¹⁷⁸ although even the lower range of the 95% interval for %unfished is above the 40% reference
¹⁷⁹ point, and all sensitivity analyses explore also show the stock to be at a high level.

Table b: Recent trend in beginning of the year spawning output and (spawning biomass relative to unfished equilibrium spawning biomass)

Year	Spawning Output (mt)	~ 95% confidence interval	Estimated %unfished	~ 95% confidence interval
2010	1603.690	(726.67-2480.71)	0.721	(0.6-0.842)
2011	1617.560	(738.18-2496.94)	0.727	(0.608-0.847)
2012	1625.610	(745.16-2506.06)	0.731	(0.613-0.849)
2013	1634.790	(753.17-2516.41)	0.735	(0.618-0.852)
2014	1657.340	(772.22-2542.46)	0.745	(0.631-0.859)
2015	1657.020	(772.35-2541.69)	0.745	(0.632-0.859)
2016	1659.820	(774.79-2544.85)	0.746	(0.634-0.859)
2017	1652.180	(768.4-2535.96)	0.743	(0.63-0.856)
2018	1655.400	(770.86-2539.94)	0.744	(0.632-0.857)
2019	1667.190	(780.42-2553.96)	0.750	(0.639-0.86)

Spawning biomass (mt) with ~95% asymptotic intervals




Figure c: Time series of spawning biomass trajectory (circles and line: median; light broken lines: 95% credibility intervals) for the base case assessment model.

¹⁸⁰ **Recruitment**

¹⁸¹ Recruitment was assumed to follow the Beverton-Holt stock recruit curve, so uncertainty in
¹⁸² estimated recruitment is due to uncertainty in spawning biomass and the unfished equilibrium
¹⁸³ recruitment R_0 (Figure d and Table c).

Table c: Recent recruitment for the model.

Year	Estimated Recruitment (1,000s)	~ 95% confidence interval
2010	5393.54	(2966.19 - 9807.3)
2011	5414.62	(2982.16 - 9831.17)
2012	5426.77	(2991.46 - 9844.65)
2013	5440.55	(3002 - 9859.94)
2014	5474.01	(3027.33 - 9898.09)
2015	5473.54	(3027.82 - 9894.77)
2016	5477.66	(3031.98 - 9896.09)
2017	5466.40	(3024.88 - 9878.58)
2018	5471.15	(3029.62 - 9880.26)
2019	5488.48	(3043.55 - 9897.46)

Age-0 recruits (1,000s) with ~95% asymptotic intervals




Figure d: Time series of estimated Big Skate recruitments for the base-case model with 95% confidence or credibility intervals.

¹⁸⁴ **Exploitation Status**

¹⁸⁵ Harvest rates estimated by the base model indicate catch levels have been below the limits
¹⁸⁶ that would be associated with the SPR = 50% target (Table d and Figure e).

Table d: Recent trend in spawning potential ratio and exploitation for Big Skate in the model. Relative fishing intensity is (1-SPR) divided by 50% (the SPR target) and exploitation is catch divided by age 2+ biomass.

Year	Relative fishing intensity	~ 95% confidence interval	Exploitation rate	~ 95% confidence interval
2009	0.21	(0.12-0.3)	0.01	(0.01-0.02)
2010	0.20	(0.11-0.29)	0.01	(0.01-0.02)
2011	0.26	(0.15-0.38)	0.01	(0.01-0.02)
2012	0.26	(0.15-0.38)	0.01	(0.01-0.02)
2013	0.14	(0.08-0.2)	0.01	(0-0.01)
2014	0.36	(0.21-0.5)	0.02	(0.01-0.03)
2015	0.32	(0.19-0.45)	0.02	(0.01-0.03)
2016	0.39	(0.23-0.55)	0.02	(0.01-0.03)
2017	0.28	(0.16-0.39)	0.02	(0.01-0.02)
2018	0.18	(0.1-0.25)	0.01	(0.01-0.01)




Figure e: Estimated spawning potential ratio (SPR) for the base-case model. One minus SPR is plotted so that higher exploitation rates occur on the upper portion of the y-axis. The management target is plotted as a red horizontal line and values above this reflect harvests in excess of the overfishing proxy based on the SPR_{50%} harvest rate. The last year in the time series is 2018.

187 Reference Points

188 This stock assessment estimates that Big Skate in the model is above the biomass target
 189 ($SB_{40\%}$), and well above the minimum stock size threshold ($SB_{25\%}$). The estimated %un-
 190 fished level for the base model in 2019 is 75.0% (95% asymptotic interval: $\pm 63.9\%-86.0\%$,
 191 corresponding to an unfished spawning biomass of 1667.19 mt (95% asymptotic interval:
 192 780.42-2553.96 mt) of spawning biomass in the base model (Table e). Unfished age 1+
 193 biomass was estimated to be 2,523 mt in the base case model. The target spawning biomass
 194 ($SB_{40\%}$) is 890 mt, which corresponds with an equilibrium yield of 602 mt. Equilibrium yield
 195 at the proxy F_{MSY} harvest rate corresponding to $SPR_{50\%}$ is 507 mt (Figure f).

Table e: Summary of reference points and management quantities for the base case model.

Quantity	Estimate	Low 2.5% limit	High 2.5% limit
Unfished spawning output (mt)	2,224	1,246	3,202
Unfished age 1+ biomass (mt)	2,523	1,705	3,341
Unfished recruitment (R_0)	6,176	2,760	9,592
Spawning output(2018 mt)	1,655	771	2,540
Depletion (2018)	0.744	0.632	0.857
Reference points based on $SB_{40\%}$			
Proxy spawning output ($B_{40\%}$)	890	498	1,281
SPR resulting in $B_{40\%}$ ($SPR_{B40\%}$)	0.625	0.625	0.625
Exploitation rate resulting in $B_{40\%}$	0.048	0.042	0.055
Yield with $SPR_{B40\%}$ at $B_{40\%}$ (mt)	602	395	810
Reference points based on SPR proxy for MSY			
Spawning output	445	249	640
SPR_{proxy}	0.5		
Exploitation rate corresponding to SPR_{proxy}	0.071	0.061	0.08
Yield with SPR_{proxy} at SB_{SPR} (mt)	507	333	681
Reference points based on estimated MSY values			
Spawning output at MSY (SB_{MSY})	833	458	1,207
SPR_{MSY}	0.609	0.604	0.614
Exploitation rate at MSY	0.051	0.045	0.057
Dead Catch MSY (mt)	604	396	812
Retained Catch MSY (mt)	559	367	750

¹⁹⁶ **Ecosystem Considerations**

¹⁹⁷ In this assessment, neither environmental nor ecosystem considerations were explicitly included in the analysis. This is primarily due to a lack of relevant data or results of analyses
¹⁹⁸ that could contribute ecosystem-related quantitative information for the assessment.
¹⁹⁹

²⁰⁰ **Management Performance**

Table f: Recent trend in total catch (mt) relative to the management guidelines. Big skate was managed in the Other Species complex in 2013 and 2014, designated an Ecosystem Component species in 2015 and 2016, and managed with stock-specific harvest specifications since 2017.

Year	OFL (mt; ABC prior to 2011)	ABC (mt)	ACL (mt; OY prior to 2011)	Estimated total catch (mt)
2009				205.70
2010				196.20
2011				268.40
2012				269.60
2013	458.00	317.90	317.90	135.00
2014	458.00	317.90	317.90	372.40
2015				331.50
2016				411.50
2017	541.00	494.00	494.00	277.60
2018	541.00	494.00	494.00	172.60
2019	541.00	494.00	494.00	
2020	541.00	494.00	494.00	

²⁰¹ **Unresolved Problems and Major Uncertainties**

²⁰² **To be added**

²⁰³ **Decision Table**

²⁰⁴ **Template in Table h and associated discussion to be filled in later**

205 **Projected Landings, OFLs and Time-varying ACLs**

206 Potential OFLs projected by the model are shown in Table g. These values are based on an
207 SPR target of 50%, a P* of 0.45, and a time-varying Category 2 Sigma which creates the
208 buffer shown in the right-hand column.

Table g: Projections of landings, total mortality, OFL, and ACL values.

Year	Landings (mt)	Estimated total mortality (mt)	OFL (mt)	ACL (mt)	Buffer
2019	313.16	336.35	541.00	494.00	1.00
2020	313.16	336.32	541.00	494.00	1.00
2021	1042.23	1119.74	1275.51	1119.75	0.87
2022	987.51	1062.58	1222.62	1062.58	0.86
2023	942.80	1015.91	1179.51	1015.91	0.86
2024	906.41	977.59	1145.41	977.59	0.85
2025	876.49	945.64	1118.21	945.64	0.84
2026	850.59	917.76	1095.36	917.76	0.83
2027	828.05	893.39	1075.04	893.39	0.83
2028	805.87	869.37	1056.06	869.37	0.82
2029	784.60	846.33	1037.94	846.33	0.81
2030	764.95	825.07	1020.44	825.07	0.80

Table h: Summary of 10-year projections beginning in 2020 for alternate states of nature based on an axis of uncertainty for the model. Columns range over low, mid, and high states of nature, and rows range over different assumptions of catch levels. An entry of “-” indicates that the stock is driven to very low abundance under the particular scenario.

		States of nature					
		Low State		Base State		High State	
	Year	Catch	Spawning Output	Depletion	Spawning Output	Depletion	Spawning Output
Default harvest, for Low State	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
Default harvest, for Base State	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
Default harvest, for High State	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-
Average Catch	2019	-	-	-	-	-	-
	2020	-	-	-	-	-	-
	2021	-	-	-	-	-	-
	2022	-	-	-	-	-	-
	2023	-	-	-	-	-	-
	2024	-	-	-	-	-	-
	2025	-	-	-	-	-	-
	2026	-	-	-	-	-	-
	2027	-	-	-	-	-	-
	2028	-	-	-	-	-	-

Table i: Base case results summary.

Quantity	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019
Landings (mt)	313.160	313.160	1042.228	987.509	942.796	906.409	876.485	850.594	828.055	805.865
Total Est. Catch (mt)	336.345	336.325	1119.745	1062.581	1015.907	977.592	945.644	917.761	893.393	869.365
OFL (mt)	541.00	541.00	1275.51	1222.62	1179.51	1145.41	1118.21	1095.36	1075.04	1056.06
ACL (mt)	494.000	494.000	1119.750	1062.580	1015.910	977.592	945.643	917.762	893.393	869.365
(1-SPR)(1-SPR _{50%})	0.20	0.26	0.26	0.14	0.36	0.32	0.39	0.28	0.18	
Exploitation rate	0.01	0.01	0.01	0.01	0.02	0.02	0.02	0.02	0.01	
Age 1+ biomass (mt)	18810.9	18968.2	19113.5	19171.0	19221.9	19394.4	19315.6	19300.1	19211.4	19275.8
Spawning Output	1603.7	1617.6	1625.6	1634.8	1657.3	1657.0	1659.8	1652.2	1655.4	1667.2
95% CI	(726.67- 2480.71)	(738.18- 2496.94)	(745.16- 2506.06)	(753.17- 2516.41)	(772.22- 2542.46)	(772.35- 2541.69)	(774.79- 2544.85)	(768.4-2535.96)	(770.86- 2539.94)	(780.42- 2553.96)
Depletion	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7	0.7
95% CI	(0.6-0.842)	(0.608-0.847)	(0.613-0.849)	(0.618-0.852)	(0.631-0.859)	(0.632-0.859)	(0.634-0.859)	(0.63-0.856)	(0.632-0.857)	(0.639-0.86)
Recruits	5393.54	5414.62	5426.77	5440.55	5474.01	5473.54	5477.66	5466.40	5471.15	5488.48
95% CI	(2966.19 - 9807.3)	(2982.16 - 9831.17)	(2991.46 - 9844.65)	(3002 - 9859.94)	(3027.33 - 9898.09)	(3027.82 - 9894.77)	(3031.98 - 9896.09)	(3024.88 - 9878.58)	(3029.62 - 9880.26)	(3043.55 - 9897.46)




Figure f: Equilibrium yield curve for the base case model. Values are based on the 2018 fishery selectivity and with steepness fixed at 0.718.

²⁰⁹ **Research and Data Needs**

²¹⁰ We recommend the following research be conducted before the next assessment:

²¹¹ 1. **Data!:**

²¹² 2. **xxxx:**

²¹³ 3. **xxxx:**

²¹⁴ 4. **xxxx:**

²¹⁵ 5. **xxxx:**

²¹⁶ **To be continued**

217 **1 Introduction**

218 Skates are the largest and most widely distributed group of batoid fish with approximately
219 245 species ascribed to two families (Ebert and Compagno (2007), McEachran and Miyake
220 (1990)). Skates are benthic fish that are found in all coastal waters but are most common
221 in cold temperatures and polar waters (Ebert and Compagno 2007).

222 There are eleven species of skates in three genera (*Amblyraja*, *Bathyraja*, and *Raja*) present
223 in the Northeast Pacific Ocean off California, Oregon and Washington (Ebert 2003). Of that
224 number, just three species (Longnose Skate, *Raja rhina*; Big Skate, *Raja binoculata*; and
225 Sandpaper Skate, *Bathyraja interrupta*) make up over 95 percent of West Coast Ground-
226 fish Bottom Trawl Survey (WCGBTS) catches in terms of biomass and numbers, with the
227 Longnose Skate leading in both categories (with 62 percent of biomass and 56 percent of
228 numbers).

229 Big Skate (*Raja binoculata*) is the largest of the skate species in North America with a docu-
230 mented maximum length of 244 cm total length and a maximum weight of 91 kg (Eschmeyer
231 and Herald 1983). The species name “binoculata” (two-eyed) refers to the prominent ocellus
232 at the base of each pectoral fin. Big Skates are usually seen buried in sediment with only
233 their eyes showing.

234 **1.1 Biology**

235 Big Skate is oviparous, and is one of two skate species that have multiple embryos per
236 egg case (Ebert et al. 2008). From 1–8 embryos can be contained in a single, large egg
237 capsule, but most have 3–4 (DeLacy and Chapman 1935, Hitz 1964, Ford 1971). Eggs
238 are deposited year-round on sand or mud substrates at depths of ~50–150 m (Hitz 1964,
239 Ebert and Compagno 2007). Embryos hatch from eggs after 6–20 months, with shorter
240 developmental periods associated with warmer temperatures (Hoff, GR 2009). In captivity,
241 Big Skate females may produce > 350 eggs/year (average of 2 embryos/egg case; Chiquillo,
242 Kelcie L and Ebert, David A and Slager, Christina J and Crow, Karen D (2014)) from long-
243 term sperm storage (???). Size at birth is 18–23 cm TL (Ebert 2003). Maximum size is 244
244 cm TL [Eschmeyer and Herald (1983), with females growing to larger sizes].

245 Size at maturity has been variably estimated for Big Skate populations off California, British
246 Columbia, and Alaska. Off central California, Zeiner and Wolf (1993) reported sizes at first
247 maturity of ~129 cm TL (females) and ~100 cm TL (males). A similar size at maturity was
248 estimated for females from the Gulf of Alaska (first = 126 cm TL, 50% = 149 cm TL), but
249 male estimates were considerably greater (first = 124 cm TL, 50% = 119 cm TL; Ebert et
250 al. (2008)). Much smaller sizes at first (female = 60 cm TL, male = 50 cm TL) and 50%
251 (female = 90 cm TL, male = 72 cm TL) maturity were generated for the Longnose Skate
252 populations off British Columbia (McFarlane GA and King JR 2006); however, maturity

253 evaluation criteria were flawed (subadults were considered to be mature), and these results
254 are therefore not considered valid.

255 Age and growth parameters have been established from California, British Columbia, and
256 the Gulf of Alaska. Maximum ages off central California (females = 12, males = 11; Zeiner,
257 S.J. and P. Wolf. (1993)) and in the Gulf of Alaska (females = 14, males = 15; Gburski et
258 al. 2007) were similar, but estimates off British Columbia were much greater (females = 26,
259 males = 25; McFarlane and King 2006). It is important to note that age estimates are based
260 on an unvalidated method and geographic differences in size or age may reflect differences
261 in sampling or ageing criteria. In the Gulf of Alaska, Big Skates reach 50% maturity at 10
262 years and 7 years for females and males, respectively (Gburski, C.M. and Gaichas, S.K. and
263 Kimura, D.K. (2007), Ebert et al. (2008)). Generation length estimates range from 11.5
264 (Zeiner, S.J. and P. Wolf. 1993) to 17 years (McFarlane GA and King JR 2006).

Table 1: Regional comparison of life history parameter estimates.

	California		British Columbia		Gulf of Alaska	
	Female	Male	Female	Male	Female	Male
1st Maturity (TL cm)	129	100	60	50	126	124
50% Maturity (TL cm)			90	72	149	119
Max Age (year)	12	11	26	25	14	15
1st Maturity (year)	12	10	6	5	7	9
50% Maturity (year)			8	10	10	7

265 1.2 Distribution and Life History

266 The Big Skate is most common in soft-sediment habitats in coastal waters of the continental
267 shelf (Bizzarro, JJ and Broms, KM and Logsdon, MG and Ebert, DA and Yoklavich, MM
268 and Kuhnz, LA and Summers, AP (2014), Farrugia et al. (2016)). Use of mixed substrate
269 (e.g., mud with boulders) increases with ontogeny but hard substrates are largely avoided
270 (Bizzarro (2015)). In the GOA, the Big Skate is the most commonly encountered skate
271 species in continental shelf waters at 100–200 m depth, and is most abundant in the central
272 and western areas of the GOA (Stevenson, DE and Orr, JW and Hoff, GR and McEachran,
273 JD (2008); Bizzarro, JJ and Broms, KM and Logsdon, MG and Ebert, DA and Yoklavich,
274 MM and Kuhnz, LA and Summers, AP (2014)). Off the U.S. Pacific Coast, the Big Skate is
275 most densely distributed on the inner continental shelf (< 100 m; Bizzarro, JJ and Broms,
276 KM and Logsdon, MG and Ebert, DA and Yoklavich, MM and Kuhnz, LA and Summers,
277 AP (2014)). Eggs are mainly deposited between 70–90 m on sand or mud substrates (Hitz
278 (1964); NMFS-NWFSC-FRAM, unpub. data). Juveniles typically occur in shallower waters
279 than adults (Bizzarro (2015)). Core habitat regions of Big Skate off the U.S. Pacific Coast
280 and in the Gulf of Alaska are spatially segregated from those of other species (Bizzarro, JJ

²⁸¹ and Broms, KM and Logsdon, MG and Ebert, DA and Yoklavich, MM and Kuhnz, LA and
²⁸² Summers, AP ([2014](#))).

²⁸³ Big Skates are highly mobile and capable of long range (> 2000 km) movements
²⁸⁴ (KingandMcF2010; Farrugia et al. ([2016](#))). For example, in British Columbia, a study
²⁸⁵ revealed that ~75% of tagged individuals were recaptured within 21 km of the tagging
²⁸⁶ locations, but 15 of the tagged individuals (0.1%) moved over 1,000 km (max = 2340 km;
²⁸⁷ King, JR and McFarlane, GA ([2010](#))). In the Gulf of Alaska, a year of satellite tag data
²⁸⁸ showed that six of twelve tagged individuals moved over 100 km, with one skate moving >
²⁸⁹ 2,000 km (Farrugia et al. [2016](#)). Although primarily benthic, Big Skates utilize the entire
²⁹⁰ water column including surface waters (Farrugia et al. ([2016](#))). They have broad thermal
²⁹¹ tolerances 2–19° C that enable their occurrence from boreal to subtropical latitudes (Love,
²⁹² Milton S ([2011](#)); Farrugia et al. ([2016](#))).

²⁹³ The Big Skate is broadly distributed, occurring from the southeastern Bering Sea (Mecklen-
²⁹⁴ burg, CW and Mecklenburg, TA and Thorsteinson, LK [2002](#)) to southern Baja California
²⁹⁵ (22.90° N, 110.03° W; (Castro-Aguirre et al. [1993](#))) and the Gulf of California (Castro-
²⁹⁶ Aguirre and Pérez [1996](#)). It has been reported at depths of 2–501 m (min: Miller et al.
²⁹⁷ ([1980](#)); max: Farrugia et al. ([2016](#))) but is most common on the inner continental shelf (<
²⁹⁸ 100 m; (Love, Milton S [2011](#)); (Bizzarro [2015](#))). Big Skates are highly mobile and capable
²⁹⁹ of long range (> 2000 km) movements ((King and McFarlane [2009](#)); (Farrugia et al. [2016](#))).

³⁰⁰ In 2012, the Big Skate was moved from genus *Raja* to the new genus *Beringraja* together
³⁰¹ with the Mottled Skate (*B. pulchra*) (Ishihara et al. [2012](#)). These are the only two skates
³⁰² with multiple embryos per egg case, and they are very similar morphologically and genetically
³⁰³ (Bizzarro, J. [2019](#)).

³⁰⁴ 1.3 Ecosystem Considerations

³⁰⁵ Big Skates are opportunistic, generalist mesopredators with highly variable spatio-temporal
³⁰⁶ trophic roles (Ebert and Compagno ([2007](#)); Bizzarro ([2015](#))). Off central California, diet of
³⁰⁷ Big Skates is composed mainly of fishes, shrimps, and crabs (in descending order), with larger
³⁰⁸ skates incorporating more fishes (Bizzarro et al. ([2007](#))); however, in the Gulf of Alaska, Big
³⁰⁹ Skate diet consists mainly of crabs (esp. Tanner Crabs) throughout ontogeny, with relatively
³¹⁰ small portions of fishes and shrimps (Bizzarro ([2015](#))). Correspondingly, trophic level and
³¹¹ general diet composition estimates differ significantly between California and Gulf of Alaska
³¹² Big Skate populations (Bizzarro ([2015](#))).

³¹³ Big Skates and their egg cases are preyed upon by a variety of vertebrates and invertebrates.
³¹⁴ Snails and other molluscs bore holes in egg cases to feed on developing embryos and especially
³¹⁵ their protein rich yolk-sacs (Bizzarro, pers. obs; Hoff, GR ([2009](#))). Sevengill Sharks, Brown
³¹⁶ Rockfish, and Stellar Sea Lions are known predators of juvenile and adult Big Skates (Ebert
³¹⁷ ([2003](#)), Love, Milton S ([2011](#))). Northern Sea Lions consume free-living Big Skates and their
³¹⁸ egg cases (Ebert ([2003](#)), Love, Milton S ([2011](#))).

³¹⁹ In this assessment, ecosystem considerations were not explicitly included in the analysis.
³²⁰ This is primarily due to a lack of relevant data and results of analyses (conducted elsewhere)
³²¹ that could contribute ecosystem-related quantitative information for the assessment.

³²² 1.4 Fishery Information

³²³ Big Skate are caught in commercial and recreational fisheries on the West Coast using line
³²⁴ and trawl gears. There is a limited market for pectoral fins (skate wings).

³²⁵ The history of Big Skate is not well documented. They were used as a food source by the
³²⁶ native Coastal and Salish Tribes (Batdorf, C [1990](#)) long before Europeans settled in the
³²⁷ Pacific Northwest and then as fertilizer by the settlers (Bowers, G. M. [1909](#)). No directed
³²⁸ fishery for Big Skate has been documented; rather, they were taken along with other skates
³²⁹ and rays as “scrap fish” and used for fertilizer, fish meal and oil (Lippert [2019](#)).

³³⁰ Skates have been regarded as a predator on desirable market species such as Dungeness
³³¹ crab, and were thought of as nuisance fish with no appeal as a food item save for small local
³³² markets. They had been discarded or harvested at a minimal level until their livers became
³³³ valued along with those of other cartilaginous fishes for the extraction of vitamin A in the
³³⁴ 1940s. Chapman (Chapman, W.M. [1944](#)) recorded that “At present they are being fished
³³⁵ heavily, in common with the other elasmobranchs of the coast, for the vitamins in their livers.
³³⁶ The carcasses are either thrown away at sea or made into fish meal. Little use is made of
³³⁷ the excellent meat of the wings”.

³³⁸ Little information is available about the historic Washington fishery for Big Skate. In records
³³⁹ before 2000, they are lumped together with other skates or in market categories (Lippert
³⁴⁰ [2019](#)); this necessitates considerable attention to reconstructing the fishery by observing
³⁴¹ the composition of skate catches in the modern fishery and applying those to the recently
³⁴² reconstructed historical records.

³⁴³ Very little information is known about the Big Skate historical fishery in Oregon. The infor-
³⁴⁴ mation we do have is mainly from historical landing data and species composition samples
³⁴⁵ starting in the mid-nineties. The bulk of the catch is from the bottom trawl and longline
³⁴⁶ fisheries, with smaller amounts as by-catch in mid-water trawl and the shrimp trawl fishery.
³⁴⁷ Big Skate was lumped into the nominal “Skate” category until 2015 when it was separated
³⁴⁸ into its own market category. Species composition data have been vitally important in
³⁴⁹ reconstructing the pre-2015 historical catch (Calavan [2019](#)).

³⁵⁰ 1.5 Stock Status and Management History

³⁵¹ Big Skate were managed in the “Other Fish” complex until 2015 when they were designated
³⁵² an Ecosystem Component (EC) species. Catches of Big Skate are estimated to have averaged

³⁵³ 95 mt from 2007–2011, along with large landings of “Unspecified Skate”. Analysis of Oregon
³⁵⁴ port-sampling data indicates that about 98 percent of the recent Unspecified Skate landings
³⁵⁵ in Oregon were comprised of Big Skate. Such large landings indicates targeting of Big Skate
³⁵⁶ has occurred and an EC designation was not warranted. Based on this evidence, Big Skate
³⁵⁷ was redesignated as an actively-managed species in the fishery. Big skate have been managed
³⁵⁸ with stock-specific harvest specifications since 2017.

³⁵⁹ The recent OFL of 541 mt was calculated by applying approximate MSY harvest rates to
³⁶⁰ estimates of stock biomass from the Northwest Fisheries Science Center (NWFSC) West
³⁶¹ Coast Groundfish Bottom Trawl Survey. This survey-based biomass estimate is likely un-
³⁶² derestimated since Big Skate are distributed all the way to the shoreline and no West Coast
³⁶³ trawl surveys have been conducted in water shallower than 55 meters. This introduces an
³⁶⁴ extra source of uncertainty to management and suggests that increased precaution is needed
³⁶⁵ to reduce the risk of overfishing the stock.

³⁶⁶ There has been consideration for managing Big Skate in a complex with Longnose Skate,
³⁶⁷ the other actively-managed West Coast skate species, but the two species have disparate
³⁶⁸ distributions and fishery interactions (Longnose Skate is much more deeply distributed than
³⁶⁹ Big Skate) and that option was not endorsed. The Pacific Fishery Management Council has
³⁷⁰ chosen to set the Annual Catch Limit (ACL) equal to the Allowable Biological Catch (ABC)
³⁷¹ with a buffer for management uncertainty (P^*) of 0.45.

³⁷² 1.6 Fisheries Off Alaska, Canada and Mexico

³⁷³ 1.6.1 Alaska

³⁷⁴ In Alaska, skates were primarily taken as bycatch in both longline and trawl fisheries until
³⁷⁵ 2003, when a directed skate fishery developed in the Gulf of Alaska, where Longnose and
³⁷⁶ Big skates comprise the majority of the skate biomass.

³⁷⁷ The Gulf of Alaska (GOA) skate complex is managed as three units. Big skates and Longnose
³⁷⁸ Skates each have separate harvest specifications, with acceptable biological catches (ABCs)
³⁷⁹ specified for each GOA regulatory area (western, central, and eastern). A single gulfwide
³⁸⁰ overfishing level (OFL) is specified for each stock. All remaining skate species are managed as
³⁸¹ an “Other Skates” group with gulfwide harvest specifications. All GOA skates are managed
³⁸² as Tier 5 stocks, where OFL and ABC are based on survey biomass estimates and natural
³⁸³ mortality rate (Alaska Fisheries Science Center [2018](#)).

³⁸⁴ In the Bering Sea and Aleutian Islands, skates are assessed as a group rather than as separate
³⁸⁵ species.

³⁸⁶ **1.6.2 Canada**

³⁸⁷ In Canada historic information regarding skate catches goes back to the 1950's. Prior to
³⁸⁸ 1990's skates were taken mostly as bycatch and landings were reported as part of a skate
³⁸⁹ complex (not by species). As with the West Coast, the trawl fishery is responsible for the
³⁹⁰ largest amount of bycatch. Skate catches off British Columbia accelerated in the early 1990's,
³⁹¹ partly due to emerging Asian markets. Since 1996, longnose skate has been targeted by the
³⁹² B.C. trawl fishery and, as a result, catches have been more accurately reported.

³⁹³ Assessments of Longnose Skate and Big Skate were conducted by Canada's Division of Fish-
³⁹⁴ eries and Oceans in 2015(King, J.R., Surry, A.M., Garcia, S., and P.J. Starr [2015](#)). For Big
³⁹⁵ Skate, a Bayesian surplus production model failed to provide plausible results, and two data-
³⁹⁶ limited approaches were investigated: Depletion-Corrected Average Catch Analysis (DCAC),
³⁹⁷ and a Catch-MSY (maximum sustainable yield) Approach.

³⁹⁸ DCAC produced a range of potential yield estimates that were above the long-term average
³⁹⁹ catch, with an upper bound that was three orders of magnitude larger than the long-term
⁴⁰⁰ average catch. The Catch-MSY approach was found to be quite sensitive to assumptions
⁴⁰¹ and was not recommended as the sole basis of advice to managers.

⁴⁰² The recommendation for management for both skate species was that they should be man-
⁴⁰³ aged with harvest yields based on mean historic catch, with consideration given to survey
⁴⁰⁴ trends and to the ranges of maximum sustainable yield estimates identified by the Catch-
⁴⁰⁵ MSY Approach. However, the analysis found no significant trends in abundance indices for
⁴⁰⁶ Big Skate, and mean historical catches were below the maximum MSY estimate from the
⁴⁰⁷ catch-MSY results.

⁴⁰⁸ **1.6.3 Mexico**

⁴⁰⁹ No information is available on any fishery for Big Skate in Mexican waters, where they rarely
⁴¹⁰ occur, however they may be taken in the artisanal fishery.

411 **2 Fishery Data**

412 **2.1 Data**

413 Data used in the Big Skate assessment are summarized in Figure 1. Descriptions of the data
414 sources are in the following sections.

415 **2.2 Fishery Landings and Discards**

416 Catch information for Big Skate is very limited, in part because the requirement to sort
417 landings of Big Skate in the shore-based Individual Fishing Quota fishery from landings in
418 the “Unidentified Skate” category was not implemented until June 2015. The historical catch
419 of Big Skate therefore relies on the historical reconstruction of the landings of all skates as
420 well as an analysis of discards of Longnose Skate. The estimated landings for each state and
421 the tribal fishery are provided in Table 2 and shown in Figure 3.

422 **2.2.1 Washington Commercial Skate Landings Reconstruction**

423 Estimates of landings of Big Skate in Washington state were estimated as a fraction of total
424 skate landings as described in (Gertseva, V. 2019). The approach relied on trawl survey
425 estimates of depth distributions for each species, combined with logbook estimates of fishing
426 depths in each year.

427 The WCGBT Survey data was used to estimate proportions of longnose and big skates by
428 depth (aggregated into 100m bins) and year for the period of the survey (between 2003 and
429 2018). Big Skate were primarily found in the 0–100m and 100–200m. Trawl logbook data
430 include information on the amount of retained catch of skate (all species combined) within
431 each haul as well depth of catch. The proportion of Big Skate for each depth bin was assigned
432 to the skate catch for each haul within those depth bins and summed to get a total for each
433 year. When survey skate information was available (2003-2018), survey skate proportions
434 were applied by depth and year to account for inter-annual variability in those proportions.
435 Prior to 2003, average proportions from 2003-2007 within each depth bin were applied.

436 These estimated annual proportion of Big Skate relative to all skates from the logbook
437 analysis was then applied to total Washington skate landings by year (provided by WDFW)
438 to account for landings that weren’t included in the available logbook data. Prior to 1987
439 (when no logbook data were available), the average proportion Big Skate within the combined
440 skate category, calculated from 1987-1992 logbook data, was applied to total skate landings
441 in Washington. Estimated Big Skate landings provided by WDFW were used for the period
442 from 2004 forward.

443 2.2.2 Oregon Commercial Skate Landings Reconstruction

444 Oregon Department of Fish and Wildlife (ODFW) provided newly reconstructed commercial
445 landings for all observed skate species for the 2019 assessment cycle (1978 – 2018). In
446 addition, the methods were reviewed at a pre-assessment workshop. Historically, skates were
447 landed as a single skate complex in Oregon. In 2009, longnose skates were separated into
448 their own single-species landing category, and in 2014, big skates were also separated. The
449 reconstruction methodology differed by these three time blocks in which species composition
450 collections diverged (1978 – 2008; 2009 – 2014; 2015 – 2018).

451 Species compositions of skate complexes from commercial port sampling are available
452 throughout this time period but are generally limited, which precluded the use of all strata
453 for reconstructing landings. Quarter and port were excluded, retaining gear type, PMFC
454 area, and market category for stratifying reconstructed landings within the three time
455 blocks. Bottom trawl gear types include multiple bottom trawl gears, and account for
456 greater than 98% of skate landings . Minor gear types include primarily bottom longline
457 gear, but also include mid-water trawl, hook and line, shrimp trawl, pot gear and scallop
458 dredge.

459 For bottom trawl gears, trawl logbook areas and adjusted skate catches were matched with
460 strata-specific species compositions. In Time Block 1 (1978 – 2008), all bottom trawl gear
461 types were aggregated due to a lack of specificity in the gear recorded on the fish tickets.
462 However, in Time Blocks 2 and 3, individual bottom trawl gear types were retained. Some
463 borrowing of species compositions was required (31% of strata) and when necessary, borrowed
464 from the closest area or from the most similar gear type . Longline gear landings were
465 reconstructed in a similar fashion as to bottom trawl and required some borrowing among
466 strata as well (25%).

467 Due to insufficient species compositions, mid-water trawl landings were reconstructed using a
468 novel depth-based approach. Available compositions indicate that the proportion by weight
469 of big skates within a composition drops to zero at approximately 100 fathoms, and an inverse
470 relationship is observed for longnose skate, where the proportion by weight is consistently
471 one beyond 100 – 150 fathoms . Complex-level landings were assigned a depth from logbook
472 entries and these species specific depth associations were used to parse out landings by
473 species. The approach differed somewhat by time block . Landings from shrimp trawls were
474 handled using a similar methodology. Finally, very minor landings from hook and line, pot
475 gear and scallop dredges were assigned a single aggregated species composition, as they lack
476 any gear-specific composition samples. Landings from within a time block were apportioned
477 by year using the proportion of the annual ticket landings.

478 Results indicate that the species-specific landings from this reconstruction are very similar
479 to those from Oregon's commercial catch reconstruction (Karnowski et al. 2014) during the
480 overlapping years but cover a greater time period with methodology more applicable to skates
481 in particular. ODFW intends to incorporate reconstructed skate landings into PacFIN in
482 the future (A. Whitman, ODFW; pers. comm.).

483 **2.2.3 California Catch Reconstruction**

484 A reconstruction of historical skate landings from California waters was developed for the
485 1916–2017 time period using a combination of commercial catch data (spatially explicit block
486 summary catches and port sample data from 2009-2017) and fishery-independent survey data
487 (Bizzarro, J. 2019). Virtually all landings in California were of “unspecified skate” until
488 species-composition sampling of skate market categories began in 2009.

489 From 2009 through 2017, catch estimates were based on these market category species-
490 composition samples, and the average of those species-compositions was hindcast to 2002,
491 based on the assumption that those data were representative of the era of large area closures
492 in the post-2000 period.

493 For the period from 1936-1980, spatially explicit landings data (the California Department
494 of Fisheries and Wildlife (CDFW) block summary data) were merged with survey data to
495 provide species-specific estimates.

496 For years 1981-2001, a “blended” product of these two approaches was taken, in which
497 a linear weighting scheme blended the two sets of catch estimates through that period.
498 Landings estimates were also scaled upwards by an expansion factor for skates landed as
499 “dressed” based on fish ticket data. Prior to 1981 these data had not been reported and
500 skate landings were scaled by the “average” percentage landed as dressed in the 1981-1985
501 time period, but by the late 1980s nearly all skates were landed round.

502 As no spatial information on catch is available from 1916-1930, and the block summary
503 data were very sparse in the first few years of the CDFW fish ticket program (1931-1934),
504 spatial information from the late 1930’s was used to hindcast to the 1916–1935 time period.
505 However, since Washington and Oregon did not have catch estimates for this year period,
506 the California estimates of catch prior to 1938 were not used as they were subsumed into an
507 estimated of the total catch across all states increasing linearly from 1916 to 1950.

508 **2.2.4 Tribal Catch in Washington**

509 Tribal catch of Big Skate was provided by WDFW as all landings took place in Washington
510 State. The landings were estimated from limited state sampling of species compositions in
511 combined skate category. Anecdotal evidence suggest that most of the catch in tribal fishery
512 is retained, and discard is minimal.

513 **2.2.5 Fishery Discards**

514 Fishery discards of Big Skate are highly uncertain. The method used to estimate discards for
515 Longnose Skate was based on a strong correlation ($R^2 = 95.7\%$) between total mortality of

516 that species, and total mortality of Dover Sole for the years 2009–2017 during which Longnose
517 were landed separately from other skates. In contrast, the sorting requirement for Big Skate
518 occurred too recently to provide an adequate range of years for this type of correlation.
519 Furthermore, there is greater uncertainty in the total mortality for the shallow-water species
520 with which Big Skate most often co-occurs, such as Sand Sole and Starry Flounder, than
521 there is for Dover Sole, which has been the subject of recurring stock assessments.

522 Both what discard rate information is available and anecdotal information from those in-
523 volved in the fishery for both skate species indicate that discarding for Big Skate and Long-
524 nose Skate in the years prior to 1995 were driven by the same market forced and the discard
525 rates were similar. Therefore, the discard rate for Longnose Skate was used as a proxy for
526 the discards of Big Skate in order to estimate Big Skate discards.

527 The reconstructed landings of Big Skate for the period 1950–1995 had a mean of 63.1 t with
528 no significant trend (a linear model fit to the data increased from 62.8 t in 1950 to 63.5 in
529 1995. The estimated tribal catch prior to 1995 averaged less than 1 t and was not included
530 in this analysis of Big Skate discards for the years prior to 1995.

531 The mean discard rate for LN was 92.46%, also with no significant linear trend (the linear
532 fit decreased from 92.8% in 1950 to 92.1% in 1995). An estimate of the mean annual discard
533 amount can therefore be calculated as from the mean discard rate and the mean landings as
534 $\bar{L}/(1 - \bar{d})$ where \bar{L} is the mean landings across that time period and \bar{d} is the mean discards
535 (Figure ??).

536 Two alternative methods were used to estimate the mean annual discard amount: applying
537 the annual LN discard rates to the annual BS catch, and applying 3-year moving averages
538 of these two quantities. The use of the annual values resulted in an implausibly high degree
539 of annual variability among the estimates, with the most extreme being a spike of 2146.4 in
540 1979 compared to 1032.7 t the year before and 654.0 the year after. The use of the 3-year
541 moving average dampened this variability and these estimates were retained for a sensitivity
542 analysis (Figure 4).

543 A discard mortality rate of 50 percent was assumed for all discards, following the assumption
544 used for the Longnose Skate assessment conducted for the U.S. West Coast in 2007 (Gertseva,
545 V and Schrippa, MJ 2007) The same rate has been used for skates in the trawl fishery in
546 British Columbia, based on an approximate average of these reported rates. In 2015, PFMC's
547 Groundfish Management Team (GMT) conducted a comprehensive literature review of skate
548 discard mortality, and concluded that the current assumption regarding Big Skate discard
549 mortality is consistent with existing reported rates for other similar species.

550 Estimation of discard rates (discards amount relative to total catch) during the period of the
551 West Coast Groundfish Observer Program (WCGOP), which began in 2002, was hindered
552 by the landings of Big Skate primarily occurring in the “unspecified skate” category prior
553 to 2015. Therefore, a discard rate was computed using the combination of Big Skate and
554 unspecified skate under the assumption that the vast majority of the unspecified skates were

- 555 Big Skate. A coefficient of variation was calculated for the by bootstrapping vessels within
556 ports because the observer program randomly chooses vessels within ports to be observed.
557 For the years after the catch share program was implemented in 2011, the trawl fishery was
558 subject to 100% observer coverage and discarding is assumed to be known with minimal
559 error (CV = 0.01).
- 560 The mean body weight of discarded Big Skates, calculated from the weight and count of
561 baskets of discarded Big Skate, was available for the years 2002–2017.

562 **3 Fishery-Independent Data Sources**

563 **3.1 Indices of abundance**

564 **3.1.1 Alaska Fisheries Science Center (AFSC) Triennial Shelf Survey**

565 Research surveys have been used since the 1970s to provide fishery-independent information
566 about the abundance, distribution, and biological characteristics of Big Skate. A coast-
567 wide survey was conducted in 1977 (Gunderson, Donald Raymond and Sample, Terrance M.
568 1980) by the Alaska Fisheries Science Center, and repeated every three years through 2001.
569 The final year of this survey, 2004, was conducted by the NWFSC according to the AFSC
570 protocol. We refer to this as the **Triennial Survey**.

571 The survey design used equally-spaced transects from which searches for tows in a specific
572 depth range were initiated. The depth range and latitudinal range was not consistent across
573 years, but all years in the period 1980–2004 included the area from 40° 10'N north to the
574 Canadian border and a depth range that included 55–366 meters, which spans the range
575 where the vast majority of Big Skate encountered in all trawl surveys. Therefore the index
576 was based on this depth range. The survey as conducted in 1977 had incomplete coverage
577 and is not believed to be comparable to the later years, and is not used in the index.

578 **3.1.2 Northwest Fisheries Science Center West Coast Groundfish Bottom Trawl 579 Survey**

580 In 2003, the NWFSC took over an ongoing slope survey the AFSC had been conducting,
581 and expanded it spatially to include the continental shelf. This survey, referred to in this
582 document as the “WCGBT Survey” or “WCGBTS”, is conducted annually. It uses a random-
583 grid design covering the coastal waters from a depth of 55 m to 1,280 m from late-May
584 to early-October (Bradburn, M.J. and Keller, A.A and Horness, B.H. 2011 , Keller, A.A.
585 and Wallace, J.R. and Methot, R.D. 2017). Four chartered industry vessels are used each
586 year (with the exception of 2013 when the U.S. federal-government shutdown curtailed the
587 survey).

588 **3.1.3 Index Standardization**

589 The index standardization methods for the two bottom trawl surveys matched that used for
590 Longnose Skate and additional detail is provided in (Gertseva, V. 2019). The data from both
591 surveys was analyzed using a spatio-temporal delta-model (Thorson, J. T. and Shelton, A.
592 O. and Ward, E. J. and Skaug, H. J. 2015), implemented as an R package VAST (Thorson,
593 James T. and Barnett, Lewis A. K. 2017) and publicly available online (<https://github.com/James-Thorson/VAST>). Spatial and spatio-temporal variation is specifically included
594 in both encounter probability and positive catch rates, a logit-link for encounter probability,
595 and a log-link for positive catch rates. Vessel-year effects were included for each unique
596 combination of vessel and year in the database for the WCGBT Survey but not the Triennial
597 survey. Further details regarding model structure are available in the user manual (https://github.com/James-Thorson/VAST/blob/master/examples/VAST_user_manual.pdf).
598
600 Spatial patterns in the survey estimates show Big Skate widely distributed along the coast,
601 with higher densities in the central and more northern areas and closer to shore 7.

602 **3.1.4 Internation Pacific Halibut Commission Longline Survey**

603 The IPHC has conducted an annual longline survey for Pacific Halibut off the coast of Oregon
604 and Washington since 1997 (no surveys were performed in 1998 or 2000). Beginning in 1999,
605 this has been a fixed station design, with 84 locations in this area (station locations differed
606 in 1997, and are therefore not comparable with subsequent surveys). 400 to 800 hooks have
607 been deployed at each station in 100-hook groups (typically called “skates” although that
608 term will be avoided here to avoid confusion). The gear used to conduct the survey was
609 designed to efficiently sample Pacific Halibut and used 16/0 (#3) circle hooks baited with
610 Chum Salmon.

611 In some years from 2011 onward, additional stations were added to the survey to sample
612 Yelloweye Rockfish. These stations were excluded from the analysis, as were additional
613 stations added in 2013, 2014, and 2017, off the coast of California (south of 42 degrees
614 latitude). Some variability in exact sampling location is practically unavoidable, and leeway
615 is given in the IPHC methods to center the set on the target coordinates while allowing wind
616 and currents to dictate the actual direction in which the gear is deployed. This can result in
617 different habitats being accessed at each fixed deployment location across years. One station
618 that was very close to the U.S. Canada border had the mid-point of the set in Canada in 2
619 out of the 19 years of the survey. For consistency among years, all samples from this station
620 were included in the analysis, including those in Canada.

621 In most years, bycatch of non-halibut species has been recorded during this survey on the
622 first 20 hooks of each 100-hook group, although in 2003 only 10% of the hooks were observed
623 for bycatch, and starting in 2012, some stations had 100% of the hooks observed for bycatch.
624 Combining these observation pattern with the number of hooks deployed each year, resulted

in most stations having 80, 100, 120, 140, or 160 hooks observed, with a mean of 144 hooks and a maximum of 800 hooks observed. The depth range of the 84 stations considered was 42–530 m, thus extending beyond the range of Big Skate, but 74% of the stations were shallower than 200 m. Big Skate have been observed at 51 of the 84 the standard stations that were retained for this analysis, but no station had Big Skates observed in more than 12 out of the 19 years of survey data, and only 10% of the station/year combinations had at least one observed Big Skate (Figure X). Of those station/year combinations with at least one Big Skate observed, the Big Skates were observed on an average of 1.3% of the hooks observed. The highest proportion was 10 Big Skates out of 81 hooks observed at one station.

The IPHC longline survey catch data were standardized using a Generalized Linear Model (GLM) with binomial error structure. Catch-per-hook was modeled, rather than catch per station due to the variability in the number of hooks deployed and observed each year. The binomial error structure was considered logical, given the binary nature of capturing (or not) a Longnose Skate on each longline hook. The modeling approach is identical to that which has been applied in the past for Yelloweye Rockfish (Stewart et al. 2009), and Spiny Dogfish (Gertseva and Taylor 2011). MCMC sampling of the GLM parameters was used to estimate the variability around each index estimate. The median index estimates themselves were approximately equal to the observed mean catch rate in each year (Figure Y). In recent years, the IPHC standardization of the index of halibut abundance has included an adjustment to account for missing baits on hooks returned empty in an effort to account for reduced catchability of the gear that may result from the lost bait. This adjustment was not included in the analysis for Big Skate although it could be considered in future years.

647 **4 Biological Parameters and Data**

648 **4.1 Measurement Details and Conversion Factors**

649 Some size measurements were taken as either disc width or inter-spiracle width rather than
650 total length. A conversion from disc width to total length was estimated as $L = 1.3399 * W$
651 based on from 95 samples from WCGBT Survey where both measurements collected (R-
652 squared = 0.9983). Little sex difference observed, so using single relationship for both sexes
653 (Figure 15). This estimate is similar to the conversion estimated by Ebert (2008) for Big
654 Skate in Alaska. The inter-spiracle width to total length was converted based on estimates
655 from Downs & Cheng (2013):

656
$$L = 12.111 + 9.761 * ISW \text{ (females),}$$

657
$$L = 3.824 + 10.927 * ISW \text{ (males).}$$

658 **4.2 Fishery dependent length and age composition data**

659 Fishery length composition data was available from PacFIN were available for the years
660 1995–2018 (with the exception of 2000) as shown in Table 4. Ages were available from only
661 2004, 2008–2012, and 2018. These were all represented as conditioned on length in order to
662 provide more detailed information about the relationship between age and length, to reduce
663 any influence of size-based selectivity on the age composition, and to ensure independence
664 from the length samples. Furthermore, the samples from Washington in 2009 were sampled
665 using a length-stratified system, so should only be treated as conditioned on length.

666 Length compositions of Big Skate discarded in commercial fisheries measured by the West
667 Coast Groundfish Observer program were available for the years 2010–2017.

668 The input sample sizes for the length compositions were calculated via the Stewart Method
669 (Ian Stewart, personal communication, IPHC):

670
$$\text{Input } N = N_{\text{hauls}} + 0.138 * N_{\text{fish}} \text{ if } N_{\text{fish}}/N_{\text{hauls}} \text{ is } < 44,$$

671
$$\text{Input } N = 7.06 * N_{\text{hauls}} \text{ if } N_{\text{fish}}/N_{\text{hauls}} \text{ is } \geq 44.$$

672 However, no haul had greater than 44 Big Skate sampled, so only the first formula was used.

673 **4.3 Survey length and age composition data**

674 Lengths of Big Skate were only collected form the Triennial survey in 1998, 2001, and 2004,
675 but 1998 had only 3 samples and were excluded from this analysis. Length compositions were

available for all years of the WCGBT Survey. Sample sizes for both surveys are provided in Table 5. The WCGBT Survey used disc width for the years 2006 and 2007 and total length in all other years. Those samples where only disc width was measured were converted to total length using the formula above.

The length compositions from the fishery and each of the two surveys aggregated across all years is shown in Figure 9.

Ages were available from the WCGBT Survey in the years 2009, 2010, 2016, 2017, and 2018. No ages were available from the Triennial Survey.

Ageing Precision and Bias

Ages of Big Skate were all estimated based on growth band counts of sectioned vertebrae. Ageing precision and bias were estimated using double-reads of 518 Big Skate vertebrae using the approach of Punt et al. (2008). The results showed strong agreement among readers (Figure 13), with a standard deviation of the ageing error increasing from about 0.4 at age 0 to 1.6 years at age 15 (Figure 14).

Weight-Length

The mean weight as a function of length was estimated from 1159 samples from the WCGBT Survey using a linear regression on a log-log scale. Sex was not found to be a significant predictor, so a single relationship was estimated: $Weight = 0.0000074924 * Length^{2.9925}$ (Figure 15).

Sex Ratio, Maturity, and Fecundity

The female maturity relationship was based on visual maturity estimates from port samplers ($n = 278$, of which 241 were from Oregon and 37 from Washington, with 24 mature specimens) as well as 55 samples from the WCGBT Survey (of which 4 were mature). The resulting relationship was $L_{50\%} = 148.2453$ with a slope parameter of $Beta = -0.13155$ in the relationship $M = (1 + Beta(L - L_{50\%}))^{-1}$ (Figure 16).

4.4 Environmental or Ecosystem Data Included in the Assessment

In this assessment, neither environmental nor ecosystem considerations were explicitly included in the analysis. This is primarily due to a lack of relevant data or results of analyses that could contribute ecosystem-related quantitative information for the assessment.

705 **5 Assessment**

706 **5.1 Previous Assessments**

707 No previous stock assessment has been conducted for Big Skate. The current management
708 is based on an OFL estimate calculated from a proxy for F_{MSY} and average survey biomass
709 from the WCGBT Survey during the years 2010–2012 (Taylor IG and Cope, J and Hamel
710 O and Thorson, J 2013). The F_{MSY} estimate was based on the product of an assumed
711 F_{MSY}/M ratio and an M estimate of 0.162 based on the maximum age of 26 reported by
712 McFarlane and King (McFarlane GA and King JR 2006). Values were sampled from an
713 assumed distribution around all these quantities to develop a measure of uncertainty around
714 the OFL estimate.

715 **5.2 Model Description**

716 **5.2.1 Modeling Software**

717 The STAT team used Stock Synthesis version 3.30.13 (Methot, Richard D. and Wetzel,
718 Chantell R. (2013), Methot, RD Jr. and Wetzel, CR and Taylor, IG (2019)). The r4ss
719 package version 1.35.1 (Taylor et al. 2019) was used to post-process the output data from
720 Stock Synthesis.

721 **5.2.2 Summary of Data for Fleets and Areas**

722 Catch is divided among 4 fleets in the base model:

- 723 • *Fishery (current)* combines all non-tribal sources of catch for the years 1995 onward,
- 724 • *Discard (historical)* includes the estimated discard amount calculated from the esti-
725 mated Longnose Skate discard rate as described above. The input catch for this fleet
726 was 50
- 727 • *Fishery (historical)* includes the reconstructed landings estimates from each of the
728 three states for 1916–1994.
- 729 • *Tribal* includes the estimates of catch of Big Skate by treaty tribes.

⁷³⁰ **5.2.3 Other Specifications**

⁷³¹ This assessment covers the U.S. West Coast stock of Big Skate in off the coasts of Washington,
⁷³² Oregon and California, the area bounded by the U.S.-Canada border to the north, and the
⁷³³ U.S.-Mexico border to the south. The population is treated as a single coastwide stock
⁷³⁴ with no net movement in or out of the area. Females and males are modeled separately as
⁷³⁵ there is evidence for differences in growth based on both the age and length data, as well as
⁷³⁶ patterns in the sex ratios associated with the length composition data. Natural Mortality
⁷³⁷ is estimated within the model using natural mortality prior developed by Hamel (2015). A
⁷³⁸ Beverton-Holt stock-recruit function is assumed with no deviations from the spawner-recruit
⁷³⁹ curve estimated.

⁷⁴⁰ The length composition data are stratified into 37 5-cm bins, ranging between 20 and 200
⁷⁴¹ cm and the age data are stratified into ages 0–15+, conditioned on the same length bin
⁷⁴² structure. The population dynamics are computed over a larger range of lengths age ages,
⁷⁴³ with the 5-cm length bins extending up to 250 cm and the numbers-at-age computed up to
⁷⁴⁴ age 20.

⁷⁴⁵ **5.2.4 Data Weighting**

⁷⁴⁶ The Francis (2011) data weighting method “TA1.8” as implemented in the r4ss package was
⁷⁴⁷ used for all length and age composition data.

⁷⁴⁸ **5.2.5 Priors**

⁷⁴⁹ *Natural Mortality* A log-normal prior for natural mortality was based on a meta-analysis
⁷⁵⁰ completed by Hamel (2015). The Hamel prior for M is $\text{lognormal}(\ln(5.4/\text{max age}), .438)$,
⁷⁵¹ which based on the single 15-year-old fish observed out of 1034 ages from the WCGBT
⁷⁵² Survey. This results in $\text{lognormal}(\log(0.36) = -1.021651, 0.438)$ prior.

⁷⁵³ *Survey Catchability* The lack of contrast in the data resulted in unstable model results
⁷⁵⁴ under a variety of configurations. To keep biomass estimates within a plausible range,
⁷⁵⁵ the assessment uses a prior on the WCGBTS survey catchability parameter (q) that was
⁷⁵⁶ originally developed for the 2007 Longnose Skate assessment (Gertseva, V and Schrippa, MJ
⁷⁵⁷ 2007), and is being used for the concurrent Longnose Skate assessment (Gertseva, V. 2019).
⁷⁵⁸ The prior for the WCGBT Survey was derived as follows.

⁷⁵⁹ Several factors inform catchability in the survey. The WCGBT Survey covers the full latitudi-
⁷⁶⁰ nal range of Longnose Skate modeled in the assessment, and thus, the latitudinal availability
⁷⁶¹ factor was assumed to be one (complete latitudinal coverage). The survey coverage exceeds

762 the maximum depth distribution of Longnose Skates but doesn't fully cover the shallow end
763 of the skate distribution. A range of 95 to 100 percent was assumed for the depth availability.
764 A range of 75 to 95 percent was assumed for vertical availability on the basis that skates are
765 known to bury in the mud, and therefore some may be unavailable to the bottom trawl gear.

766 The largest bounds were placed on the probability of capture, given that a fish is in the net
767 path. It is known that flatfish can be herded by trawl gear, and it is possible that this could
768 also occur for skates. However, it is also possible that skate could avoid the trawl nets. For
769 capture probability, a range of 75 to 150 percent was assumed. The best estimates for each
770 of these factors were set at the midpoint of the range for individual factors, except for the
771 probability of capture, which was given a value of one. The overall estimate for the survey
772 catchability was thus estimated to be 0.83 with bounds of (0.53, 1.43). The best estimate
773 was converted to the median of a lognormal distribution and the bounds to 99 percent of
774 that distribution. This resulted in a normal prior on $\log(q)$, with a mean of -0.188, and
775 standard deviation of 0.187.

776 5.2.6 Estimated Parameters

777 A full list of all estimated and fixed parameters is provided in Tables 7.

778 The base model has a total of 44 estimated parameters in the following categories:

- 779 • 1 natural mortality parameter applied to both sexes,
- 780 • 6 parameters related to female growth and the variability in **length age age,**
- 781 • 2 parameters related to male growth relative to female growth,
- 782 • 1 stock-recruit parameter ($\log(R_0)$ controlling equilibrium recruitment)
- 783 • 3 catchability parameters (1 for the WCGBT Survey and 1 each for the early and late
784 periods of the Triennial Survey)
- 785 • 2 extra standard deviation parameters (1 for each survey)
- 786 • 29 selectivity parameters, including 16 related to time-varying retention rate

787 The estimated parameters are described in greater detail below and a full list of all estimated
788 and parameters is provided in Table 7.

789 *Growth.* Examination of patterns of age-at-length and length-at-age indicated unusual pat-
790 terns of growth for Big Skate. The youngest fish show near-linear growth, and average size
791 for both sexes is similar. However, older fish show considerable sex-based differences in size.

792 This led to the choice to model growth using the “growth cessation model” recently developed
793 by Maunder et al. (2018). The estimated growth curves are shown in Figure 17. This
794 model provided two key advantages over the more common von Bertalanffy growth model
795 in the case of Big Skate: it allowed essentially linear growth for the early years and it al-
796 lowed growth for the earlier ages to be similar between females and males while diverging at
797 older ages. The growth cessation model also improve the negative log likelihood by 45 units
798 relative to the von Bertalanffy growth model.

799 *Natural Mortality.* Male natural mortality was assumed equal to the value estimated for
800 females. Sensitivity analyses were used to test the impact of both the prior on natural
801 mortality and the assumption of equal natural mortality for both sexes.

802 *Selectivity.*

803 A double-normal selectivity function was used for all fleets to allow consideration of both
804 asymptotic and dome-shaped patterns. For the fishery and the Triennial survey, the dif-
805 ference in likelihood between dome-shaped and asymptotic patterns was very small and in
806 the case of the Triennial survey, the dome-shape occurred at a length beyond almost all
807 observations, indicating that this shape was likely driven by fit to other data sources, such
808 as the index, rather than the length composition data. The WCGBT Survey was allowed
809 to remain dome-shaped as this survey had the selectivity peak at a smaller length than the
810 other fleets and the likelihood was improved by the dome-shape. The WCGBT Survey also
811 has the shortest hauls, with 15 minutes or less of bottom contact, so larger skates may be
812 better able to escape the net.

813 In order to fit a strong skew in the sex ratios toward males for the length bins in which
814 the majority of the samples were found, it was necessary to estimate a sex-specific offset
815 of selectivity. Two offset parameters were estimated for all fleets, one for the difference in
816 length at peak selectivity and another for the maximum selectivity. The ascending slope was
817 assumed equal in all cases, as was the descending slope for the WCGBT Survey.

818 **5.2.7 Fixed Parameters**

819 The steepness of the Beverton-Holt stock-recruit curve was fixed at 0.4. The same value
820 was used in the 2007 Longnose Skate assessment (Gertseva, V and Schrippa, MJ 2007) and
821 is being considered for the ongoing 2019 Longnose Skate assessment. This value reflects a
822 K-type reproductive strategy associated with elasmobranchs in general. The influence of the
823 assumption of $h = 0.4$ on model output was explored via a likelihood profile analysis.

824 **5.3 Model Selection and Evaluation**

825 **5.3.1 Key Assumptions and Structural Choices**

826 **To be added prior to May 20 CIE pre-review deadline.**

827 **5.3.2 Alternate Models Considered**

828 **To be added prior to May 20 CIE pre-review deadline.**

829 **5.3.3 Convergence**

830 One hundred sets of jittered starting values were generated using the jitter function built
831 into Stock Synthesis, with used with jitter input = 0.1. The same likelihood as the base
832 model was returned by 54 out of the 100 runs, while the others all had worse total likelihood.

833 **5.4 Response to the Current STAR Panel Requests**

834 **Request No. 1:**

835

836 **Rationale:** xxx

837 **STAT Response:** xxx

838 **Request No. 2:**

839

840 **Rationale:** xxx

841 **STAT Response:** xxx

842 **Request No. 3:**

843

844 **Rationale:** x.

845 **STAT Response:** xxx

846 **Request No. 4:**

847

848 **Rationale:** xxx

849 **STAT Response:** xxx

850 **Request No. 5:**

851

852 **Rationale:** xxx

853 **STAT Response:** xxx

854 **5.5 Base Case Model Results**

855 The following description of the model results reflects a base model that incorporates all of
856 the changes made during the STAR panel (see previous section). The base model parameter
857 estimates and their approximate asymptotic standard errors are shown in Table 7 and the
858 likelihood components are in Table 8. Estimates of derived reference points and approximate
859 95% asymptotic confidence intervals are shown in Table e. Time-series of estimated stock
860 size over time are shown in Table 9.

861 **5.5.1 Parameter Estimates**

862 The additional survey variability (process error added directly to each year's input variabil-
863 ity) for all surveys was estimated within the model.

864 (Figure 31).

865 The stock-recruit curve ... Figure 32 with estimated recruitments also shown. } } \$**

866 **5.5.2 Fits to the Data**

867 *Indices.* The observed indices show much more variability than the model expectation, with
868 the fit to the WCGBT Survey essentially a flat line (Figure 19) and the fit to the Triennial
869 Survey only showing a noticeable change over time due to the separate catchability parameter
870 estimated for the early and late periods (Figure 20).

871 *Length Data.* The fits to the length data are much better thanks to the combination of the
872 growth cessation model and the sex-specific offsets to selectivity (Figures 21–22).

873 *Conditional Age-at-Length.* The conditional age-at-length data is likewise fit reasonably well,
874 with some patterns in residuals showing variability among years, but no clear pattern that
875 is consistent across years (Figures 23 and 24).

876 *Sex Ratios.* Sex ratio data is not included in the likelihood as such, but as a part of the
877 length composition likelihood. The proportions of females and males are compiled into a

single vector that is compared to the model expectations in the multinomial likelihood. The patterns in sex ratio by length bin show fewer females than males for the middle range of sizes (70–120 cm), with a shift to almost 100% females for the largest size bins (over 130 cm). The use of sex-specific growth curves was adequate to fit the ratios for the largest bins, but ratio skews toward males at lengths where the mean ages are similar for females and males. The fit to this part of the sex ratio pattern required an offset in selectivity.

5.5.3 Uncertainty and Sensitivity Analyses

A number of sensitivity analyses were conducted, including:

1. Setting all selectivity curves to be asymptotic
2. Setting all selectivity curves to be dome-shaped
3. Removing the sex-specific offset on the selectivity curves
4. Removing the prior on catchability for the WCGBT Survey
5. Estimating a single catchability for all years in the Triennial Survey
6. Estimating separate natural mortality parameters for males and females
7. Removing the prior on natural mortality
8. Using the von Bertalanffy growth model
9. Using the Richards growth model
10. Tuning the sample sizes using the McAllister-Ianelli method
11. Estimating historic discards based on 3yr average of discard rates and landings
12. Changeing discard mortality from 0.5 to 0.4
13. Changeing discard mortality from 0.5 to 0.6

Results of these sensitivities are shown in Figures 34 to 36, and Tables 10 to 12.

Additional text to be added

5.5.4 Retrospective Analysis

To be added

903 **5.5.5 Likelihood Profiles**

904 Likelihood profiles were conducted over $\log(R_0)$, stock-recruit steepness (h) and natural
905 mortality (M).

906 Results of these profiles are shown in Figures 39 to 44.

907 **Additional text to be added**

908 **5.5.6 Reference Points**

909 Reference points were calculated using the estimated selectivities and catch distribution
910 among fleets in the most recent year of the model, (2017). Sustainable total yield (landings
911 plus discards) were 507 mt when using an $SPR_{50\%}$ reference harvest rate and with a 95%
912 confidence interval of 333 mt based on estimates of uncertainty. The spawning biomass
913 equivalent to 40% of the unfished level ($SB_{40\%}$) was 890 mt.

914 (Figure 29

915 The 2018 spawning biomass relative to unfished equilibrium spawning biomass is
916 above/below the target of 40% of unfished levels (Figure ??). The relative fishing intensity,
917 $(1 - SPR)/(1 - SPR_{50\%})$, has been xxx the management target for the entire time series
918 of the model.

919 Table e shows the full suite of estimated reference points for the base model and Figure 45
920 shows the equilibrium curve based on a steepness value xxx.

₉₂₁ **6 Harvest Projections and Decision Tables**

₉₂₂ The forecasts of stock abundance and yield were developed using the final base model, with
₉₂₃ the forecasted projections of the OFL presented in Table [g](#).

₉₂₄ The forecasted projections of the OFL for each model are presented in Table [h](#).

₉₂₅ 7 Regional Management Considerations

926 **8 Research Needs**

927 There are a number of areas of research that could improve the stock assessment for Big
928 Skate. Below are issues identified by the STAT team and the STAR panel:

929 1. Data!:

930 2. xxxx:

931 3. xxxx:

932 4. xxxx:

933 5. xxxx:

934 **9 Acknowledgments**

935 The authors gratefully acknowledge the time and effort reviewers Stacey Miller, Jim Hastie
936 and Owen Hamel put into making this a polished document.

⁹³⁷ 10 Tables

⁹³⁸ 10.1 Data Tables

Table 2: Landings by source. Landings are reconstructed histories 1916-1995.

Year	CA (mt)	OR (mt)	WA (mt)	Tribal (mt)	Total (mt)
1916	78.30	0.00	0.00	0.00	78.30
1917	80.10	0.00	0.00	0.00	80.10
1918	101.20	0.00	0.00	0.00	101.20
1919	75.20	0.00	0.00	0.00	75.20
1920	122.00	0.00	0.00	0.00	122.00
1921	17.80	0.00	0.00	0.00	17.80
1922	30.80	0.00	0.00	0.00	30.80
1923	34.20	0.00	0.00	0.00	34.20
1924	33.40	0.00	0.00	0.00	33.40
1925	46.70	0.00	0.00	0.00	46.70
1926	59.30	0.00	0.00	0.00	59.30
1927	67.10	0.00	0.00	0.00	67.10
1928	116.70	0.00	0.00	0.00	116.70
1929	107.50	0.00	0.00	0.00	107.50
1930	70.80	0.00	0.00	0.00	70.80
1931	43.60	0.00	0.00	0.00	43.60
1932	73.30	0.00	0.00	0.00	73.30
1933	46.50	0.00	0.00	0.00	46.50
1934	57.40	0.00	0.00	0.00	57.40
1935	70.60	0.00	0.00	0.00	70.60
1936	87.70	0.00	0.00	0.00	87.70
1937	115.40	0.00	0.00	0.00	115.40
1938	99.40	0.00	0.00	0.00	99.40
1939	90.90	0.00	0.00	0.00	90.90
1940	60.30	5.30	0.00	0.00	65.70
1941	53.10	56.40	0.00	0.00	109.40
1942	27.00	34.40	0.00	0.00	61.40
1943	20.40	0.90	0.00	0.00	21.30
1944	7.80	1.60	0.00	0.00	9.50
1945	13.30	0.30	0.00	0.00	13.50
1946	17.10	1.80	0.00	0.00	18.90
1947	24.10	0.00	0.00	0.00	24.10
1948	30.70	5.70	0.00	0.00	36.30
1949	31.90	0.00	7.20	0.00	39.10
1950	32.20	2.10	2.10	0.00	36.40
1951	21.70	4.70	3.90	0.00	30.30

Continued on next page

Table 2: Landings by source. Landings are reconstructed histories 1916-1995.

Year	CA (mt)	OR (mt)	WA (mt)	Tribal (mt)	Total (mt)
1952	39.10	0.10	7.80	0.00	46.90
1953	124.90	1.20	1.60	0.00	127.60
1954	38.80	2.30	1.20	0.00	42.40
1955	45.70	35.60	1.60	0.00	82.90
1956	40.40	2.60	3.10	0.00	46.10
1957	49.50	0.00	2.50	0.00	52.00
1958	38.80	0.00	0.20	0.00	38.90
1959	46.50	0.00	0.80	0.00	47.30
1960	39.20	0.00	0.70	0.00	39.80
1961	54.40	40.90	4.60	0.00	99.80
1962	44.40	27.90	5.20	0.00	77.60
1963	53.20	30.40	2.10	0.00	85.70
1964	49.90	28.30	2.70	0.00	80.90
1965	34.30	12.80	3.50	0.00	50.60
1966	36.40	20.10	0.60	0.00	57.00
1967	53.30	15.60	6.60	0.00	75.50
1968	55.30	45.40	8.80	0.00	109.50
1969	32.50	33.80	6.60	0.00	72.90
1970	16.30	11.90	0.10	0.00	28.20
1971	18.50	3.10	0.00	0.00	21.60
1972	33.50	2.00	0.10	0.00	35.60
1973	40.70	0.90	0.00	0.00	41.70
1974	21.90	5.90	0.10	0.00	27.80
1975	39.80	2.00	0.00	0.00	41.80
1976	20.70	31.30	0.20	0.00	52.20
1977	32.80	31.50	0.60	0.00	64.90
1978	67.70	77.30	4.00	0.00	149.10
1979	90.50	75.50	30.40	0.00	196.40
1980	17.60	34.10	5.20	0.00	56.90
1981	138.00	14.80	6.50	0.00	159.30
1982	78.30	5.20	14.60	0.00	98.10
1983	55.30	14.20	8.90	0.00	78.40
1984	26.20	4.90	1.60	0.00	32.70
1985	60.30	0.40	4.90	0.00	65.60
1986	27.20	1.60	8.90	0.00	37.80
1987	22.60	1.90	18.40	1.00	43.90
1988	15.30	0.30	10.90	1.20	27.60
1989	18.90	0.20	6.20	0.00	25.30
1990	25.10	0.00	9.60	0.10	34.90
1991	22.80	0.20	21.50	0.10	44.60
1992	24.60	0.30	11.20	0.00	36.10

Continued on next page

Table 2: Landings by source. Landings are reconstructed histories 1916-1995.

Year	CA (mt)	OR (mt)	WA (mt)	Tribal (mt)	Total (mt)
1993	29.00	0.20	21.00	0.60	50.70
1994	27.70	2.50	20.50	0.10	50.70
1995	43.00	41.20	21.80	0.10	106.00
1996	146.70	138.50	22.80	0.10	308.10
1997	228.40	215.40	84.00	0.20	528.00
1998	120.50	51.40	22.70	0.20	194.90
1999	109.50	131.30	41.40	0.40	282.60
2000	69.40	193.60	97.70	0.30	361.00
2001	75.30	115.10	26.70	0.40	217.50
2002	34.70	102.80	70.80	4.80	213.10
2003	48.80	223.00	65.70	5.40	342.80
2004	45.20	105.90	98.00	4.60	253.80
2005	33.40	151.30	113.10	15.70	313.40
2006	102.40	206.60	66.20	24.90	400.00
2007	35.50	190.40	29.10	19.90	274.90
2008	46.00	280.10	36.80	3.20	366.00
2009	9.60	162.00	16.50	17.50	205.70
2010	1.20	157.50	25.00	12.50	196.20
2011	0.50	231.50	10.00	26.40	268.40
2012	6.80	216.30	5.00	41.60	269.60
2013	20.90	92.30	13.00	8.80	135.00
2014	41.00	286.00	16.80	28.60	372.40
2015	35.20	218.80	1.00	76.60	331.50
2016	15.00	317.50	1.20	77.80	411.50
2017	28.00	188.00	1.40	60.20	277.60
2018	23.80	115.80	2.40	30.60	172.60

Table 3: Index inputs.

Year	WCGBTS		Triennial		IPHC	
	Obs	se_log	Obs	se_log	Obs	se_log
1980			467.83	0.53		
1983			911.85	0.30		
1986			996.75	0.29		
1989			1431.65	0.22		
1992			2426.18	0.20		
1995			497.24	0.26		
1998			2437.75	0.20		
1999					0.00	0.17
2001			1669.73	0.23	0.00	0.29
2002					0.00	0.53
2003	8170.51	0.20			0.00	0.43
2004	14349.00	0.18	3674.14	0.19	0.00	0.20
2005	12122.52	0.16			0.00	0.18
2006	9273.79	0.18			0.00	0.64
2007	8137.47	0.18			0.00	0.34
2008	5494.76	0.21			0.00	0.81
2009	10721.30	0.17			0.00	0.48
2010	11475.29	0.14			0.00	0.24
2011	8029.69	0.16			0.00	0.20
2012	11593.79	0.16			0.00	0.61
2013	11521.85	0.17			0.00	0.20
2014	19855.79	0.13			0.00	0.19
2015	19251.41	0.13			0.00	0.16
2016	17141.95	0.15			0.00	0.17
2017	13237.37	0.14			0.00	0.18
2018	14568.79	0.14			0.00	0.26

Table 4: PacFIN Samples.

Year	CA		OR		WA		All Landings		Discards	
	Ntows	Nfish	Ntows	Nfish	Ntows	Nfish	Ntows	Nfish	Ntows	Nfish
Lengths										
1995			6	55			6	55		
1996			3	8			3	8		
1997			1	14			1	14		
1998			1	2			1	2		
1999			1	8			1	8		
2000										
2001			3	43			3	43		
2002			6	199			6	199		
2003			9	202			9	202		
2004			2	27	2	12	4	39		
2005			7	123	6	87	13	210		
2006			13	310	15	191	28	501		
2007	1	1	10	128	9	172	20	301		
2008			10	94	8	94	18	188		
2009	8	32	17	234	1	18	26	284		
2010	2	8	15	186			17	194	149	349
2011	2	2	29	418	4	9	35	429	554	1518
2012	3	43	24	477	3	38	30	558	544	1405
2013	11	201	11	252	8	168	30	621	443	987
2014	15	217	11	237	5	249	31	703	676	1625
2015	25	237	21	411	2	5	48	653	688	1557
2016	14	181	34	444	7	98	55	723	652	1456
2017	14	239	50	668	12	47	76	954	508	1248
2018	15	133	46	552	14	98	75	783		
Ages										
2004					2	11	2	11		
2008			8	80			8	80		
2009			10	87	8	65	18	152		
2010			10	102			10	102		
2011			21	202			21	202		
2012			12	120			12	120		
2018			6	39	13	93	19	132		

Table 5: Samples from the surveys.

NA.	Triennial		WCGBTS		IPHC	
	Triennial	NA..1	WCGBTS	NA..2	IPHC	NA..3
Year	Ntows	Nfish	Ntows	Nfish	Nsets	Nfish
Lengths						
2001	41	81				
2003			60	197		
2004	39	100	81	262		
2005			99	328		
2006			67	154		
2007			76	192		
2008			53	159		
2009			82	305		
2010			130	466		
2011			99	360		
2012			104	395		
2013			84	316		
2014			149	552	14	54
2015			134	546		
2016			105	422		
2017			125	496		
2018			123	331		
Ages						
2009			77	230		
2010			124	333		
2016			100	138		
2017			110	164		
2018			118	169		

940 **10.2 Model Results Tables**

Table 6: Results from 100 jitters from the base case model.

Description	Value	NA	NA
Returned to base case	-	-	-
Found local minimum	-	-	-
Found better solution	-	-	-
Error in likelihood	-	-	-
Total	100	100	100

Table 7: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
1	NatM_p_1_Fem_GP_1	0.445	3	(0.1, 0.6)	OK	0.030	Log_Norm (-1.02165, 0.438)
2	L_at_Amin_Fem_GP_1	20.094	2	(10, 40)	OK	1.033	None
3	Linf_Fem_GP_1	175.671	2	(100, 300)	OK	4.012	None
4	VonBert_K_Fem_GP_1	12.137	1	(0.005, 30)	OK	0.359	None
5	Cessation_Fem_GP_1	5.652	3	(0.1, 10)	OK	12.041	None
6	SD_young_Fem_GP_1	5.706	5	(1, 20)	OK	0.903	None
7	SD_old_Fem_GP_1	7.085	5	(1, 20)	OK	0.921	None
8	Wtlen_1_Fem_GP_1	0.000	-3	(0, 3)			None
9	Wtlen_2_Fem_GP_1	2.993	-3	(2, 4)			None
10	Mat50%_Fem_GP_1	148.245	-3	(10, 140)			None
11	Mat_slope_Fem_GP_1	-0.132	-3	(-0.09, -0.05)			None
12	Eggs/kg_inter_Fem_GP_1	1.000	-3	(-3, 3)			None
13	Eggs/kg_slope_wt_Fem_GP_1	0.000	-3	(-3, 3)			None
14	NatM_p_1_Mal_GP_1	0.000	-2	(-3, 3)			None
15	L_at_Amin_Mal_GP_1	0.000	-2	(-1, 1)			None
16	Linf_Mal_GP_1	-0.373	2	(-1, 1)	OK	0.025	None
17	VonBert_K_Mal_GP_1	0.101	3	(-10, 20)	OK	0.034	None
18	Cessation_Mal_GP_1	0.200	-3	(-3, 3)			None
19	SD_young_Mal_GP_1	0.000	-5	(-1, 1)			None
20	SD_old_Mal_GP_1	0.000	-5	(-1, 1)			None
21	Wtlen_1_Mal_GP_1	0.000	-3	(0, 3)			None
22	Wtlen_2_Mal_GP_1	2.993	-3	(2, 4)			None
23	CohortGrowDev	1.000	-5	(0, 2)			None
24	FracFemale_GP_1	0.500	-99	(0.001, 0.999)			None
25	SR_LL(R0)	8.728	3	(5, 15)	OK	0.282	None
26	SR_BH_stEEP	0.400	-3	(0.2, 1)			None

Continued on next page

Table 7: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
27	SR_sigmaR	0.300	-2	(0, 0.4)			None
28	SR_regime	0.000	-1	(-2, 2)			None
29	SR_autocorr	0.000	-99	(0, 0)			None
78	LnQ_base_WCGBT5(5)	-0.209	1	(-2, 2)	OK	0.184	Normal (-0.188, 0.187)
79	Q_extraSD_WCGBT5(5)	0.162	1	(0, 2)	OK	0.057	None
80	LnQ_base_Triennial(6)	-1.046	1	(-10, 2)	OK	0.694	None
81	Q_extraSD_Triennial(6)	0.365	1	(0, 2)	OK	0.146	None
82	LnQ_base_Triennial(6)_1995	-0.731	1	(-7, 0)	OK	0.693	None
83	Size_DblN_peak_(1)	94.092	4	(80, 150)	OK	4.912	None
84	Size_DblN_top_logit_(1)	-15.000	-5	(-15, 4)			None
85	Size_DblN_ascend_se_(1)	7.156	4	(-1, 9)	OK	0.118	None
86	Size_DblN_descend_se_(1)	20.000	-5	(-1, 20)			None
87	Size_DblN_start_logit_(1)	-999.000	-4	(-999, 9)			None
88	Size_DblN_end_logit_(1)	-999.000	-5	(-999, 9)			None
89	Retain_L_infl_(1)	66.219	2	(15, 150)	OK	0.671	None
90	Retain_L_width_(1)	4.876	2	(0.1, 10)	OK	0.354	None
91	Retain_L_asymptote_logit_(1)	2.048	3	(-10, 20)	OK	0.359	None
92	Retain_L_maleoffset_(1)	0.000	-3	(0, 0)			None
93	DiscMort_L_infl_(1)	5.000	-4	(5, 15)			None
94	DiscMort_L_width_(1)	0.000	-4	(0.001, 10)			None
95	DiscMort_L_level_old_(1)	0.500	-5	(0, 1)			None
96	DiscMort_L_male_offset_(1)	0.000	-5	(0, 0)			None
97	SzSel_Fem_Peak_(1)	-5.537	4	(-50, 50)	OK	2.174	None
98	SzSel_Fem_Ascend_(1)	0.000	-4	(-5, 5)			None
99	SzSel_Fem_Descend_(1)	0.000	-4	(-5, 5)			None
100	SzSel_Fem_Final_(1)	0.000	-4	(-5, 5)			None
101	SzSel_Fem_Scale_(1)	0.744	4	(0.5, 1.5)	OK	0.095	None

Continued on next page

Table 7: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior	(Exp.Val, SD)
102	Size_DblN_peak_WCGBT5(5)	76.187	4	(50, 150)	OK	6.668	None	
103	Size_DblN_top_logit_WCGBT5(5)	-15.000	-5	(-15, 4)			None	
104	Size_DblN_ascend_se_WCGBT5(5)	6.503	4	(-1, 9)	OK	0.371	None	
105	Size_DblN_descend_se_WCGBT5(5)	16.488	5	(-1, 20)	OK	56.568	None	
106	Size_DblN_start_logit_WCGBT5(5)	-5.000	-4	(-999, 9)			None	
107	Size_DblN_end_logit_WCGBT5(5)	-999.000	-5	(-999, 9)			None	
108	SzSel_Fem_Peak_WCGBT5(5)	-8.052	4	(-50, 50)	OK	4.166	None	
109	SzSel_Fem_Ascend_WCGBT5(5)	0.000	-4	(-5, 5)			None	
110	SzSel_Fem_Descend_WCGBT5(5)	0.000	-4	(-5, 5)			None	
111	SzSel_Fem_Final_WCGBT5(5)	0.000	-4	(-5, 5)			None	
112	SzSel_Fem_Scale_WCGBT5(5)	0.696	4	(0.5, 1.5)	OK	0.125	None	
113	Size_DblN_peak_Triennial(6)	187.722	4	(50, 200)	OK	34.761	None	
114	Size_DblN_top_logit_Triennial(6)	-15.000	-5	(-15, 4)			None	
115	Size_DblN_ascend_se_Triennial(6)	8.474	4	(-1, 9)	OK	0.422	None	
116	Size_DblN_descend_se_Triennial(6)	20.000	-5	(-1, 20)			None	
117	Size_DblN_start_logit_Triennial(6)	-4.789	4	(-15, 9)	OK	0.786	None	
118	Size_DblN_end_logit_Triennial(6)	-999.000	-5	(-999, 9)			None	
119	SzSel_Fem_Peak_Triennial(6)	0.000	-4	(-50, 50)			None	
120	SzSel_Fem_Ascend_Triennial(6)	0.000	-4	(-5, 5)			None	
121	SzSel_Fem_Descend_Triennial(6)	0.000	-4	(-5, 5)			None	
122	SzSel_Fem_Final_Triennial(6)	0.000	-4	(-5, 5)			None	
123	SzSel_Fem_Scale_Triennial(6)	0.604	4	(0.5, 1.5)	OK	0.130	None	
124	Retain_L_asymptote_logit_2005	2.299	4	(-10, 20)	OK	0.566	None	
125	Retain_L_asymptote_logit_2006	3.304	4	(-10, 20)	OK	1.305	None	
126	Retain_L_asymptote_logit_2007	3.962	4	(-10, 20)	OK	1.982	None	
127	Retain_L_asymptote_logit_2008	11.091	4	(-10, 20)	OK	111.895	None	

Continued on next page

Table 7: List of parameters used in the base model, including estimated values and standard deviations (SD), bounds (minimum and maximum), estimation phase (negative values indicate not estimated), status (indicates if parameters are near bounds, and prior type information (mean, SD)).

No.	Parameter	Value	Phase	Bounds	Status	SD	Prior (Exp.Val, SD)
128	Retain_L_asymptote_logit_2009	4.917	4	(-10, 20)	OK	3.735	None
129	Retain_L_asymptote_logit_2010	13.242	4	(-10, 20)	OK	88.124	None
130	Retain_L_asymptote_logit_2011	14.640	4	(-10, 20)	OK	74.025	None
131	Retain_L_asymptote_logit_2012	13.890	4	(-10, 20)	OK	81.550	None
132	Retain_L_asymptote_logit_2013	3.454	4	(-10, 20)	OK	0.333	None
133	Retain_L_asymptote_logit_2014	3.619	4	(-10, 20)	OK	0.276	None
134	Retain_L_asymptote_logit_2015	3.404	4	(-10, 20)	OK	0.261	None
135	Retain_L_asymptote_logit_2016	2.885	4	(-10, 20)	OK	0.192	None
136	Retain_L_asymptote_logit_2017	2.819	4	(-10, 20)	OK	0.193	None

Table 8: Likelihood components from the base model.

Likelihood component	Value
TOTAL	1097.30
Catch	0.00
Survey	-98.12
Length composition	763.02
Age composition	421.52
Recruitment	10.88
Forecast recruitment	0.00
Parameter priors	0.00
Parameter soft bounds	0.01

Table 9: Time-series of population estimates from the base-case model. Relative exploitation rate is $(1 - SPR)/(1 - SPR_{50\%})$.

Year	Total biomass (mt)	Spawning biomass (mt)	Depletion	Age-0 recruits	Total catch (mt)	Relative exploitation rate	SPR
1916	25232	2224	0.000	6176	0	0.00	1.00
1917	25232	2224	0.000	6176	12	0.00	0.99
1918	25221	2223	0.999	6175	25	0.00	0.99
1919	25199	2220	0.998	6172	37	0.00	0.98
1920	25169	2217	0.997	6168	49	0.00	0.98
1921	25131	2212	0.995	6164	62	0.00	0.97
1922	25087	2206	0.992	6157	74	0.00	0.97
1923	25037	2200	0.989	6150	86	0.00	0.96
1924	24981	2192	0.985	6142	99	0.00	0.96
1925	24920	2183	0.981	6132	111	0.00	0.96
1926	24854	2173	0.977	6122	123	0.01	0.95
1927	24783	2163	0.973	6111	136	0.01	0.94
1928	24707	2153	0.968	6100	148	0.01	0.94
1929	24627	2142	0.963	6088	160	0.01	0.93
1930	24544	2130	0.958	6076	172	0.01	0.93
1931	24456	2118	0.953	6063	185	0.01	0.92
1932	24365	2106	0.947	6049	197	0.01	0.92
1933	24271	2094	0.941	6035	210	0.01	0.91
1934	24174	2081	0.936	6020	222	0.01	0.91
1935	24074	2067	0.929	6005	234	0.01	0.90
1936	23971	2053	0.923	5989	246	0.01	0.90
1937	23866	2039	0.917	5973	259	0.01	0.89
1938	23758	2025	0.910	5956	271	0.01	0.89
1939	23648	2010	0.904	5939	329	0.01	0.87
1940	23494	1991	0.895	5916	329	0.02	0.86
1941	23353	1972	0.887	5894	363	0.02	0.85
1942	23193	1952	0.878	5869	351	0.02	0.85
1943	23059	1933	0.869	5846	343	0.02	0.86
1944	22943	1917	0.862	5826	350	0.02	0.85
1945	22829	1900	0.854	5805	364	0.02	0.85
1946	22708	1884	0.847	5784	379	0.02	0.84
1947	22581	1868	0.840	5763	394	0.02	0.83
1948	22447	1851	0.832	5742	412	0.02	0.83
1949	22306	1834	0.825	5720	426	0.02	0.82
1950	22162	1818	0.817	5698	424	0.02	0.82
1951	22032	1801	0.810	5677	418	0.02	0.82
1952	21917	1786	0.803	5656	434	0.02	0.81

Continues next page

Table 9: Time-series of population estimates from the base-case model. Relative exploitation rate is $(1 - SPR)/(1 - SPR_{50\%})$.

Year	Total biomass (mt)	Spawning biomass (mt)	Depletion	Age-0 recruits	Total catch (mt)	Relative exploitation rate	SPR
1953	21794	1771	0.796	5635	515	0.03	0.78
1954	21603	1748	0.786	5604	430	0.02	0.81
1955	21507	1734	0.780	5584	470	0.02	0.80
1956	21377	1718	0.772	5561	434	0.02	0.81
1957	21290	1706	0.767	5544	439	0.02	0.81
1958	21201	1694	0.762	5527	426	0.02	0.81
1959	21126	1685	0.757	5514	435	0.02	0.81
1960	21045	1675	0.753	5500	427	0.02	0.81
1961	20974	1667	0.750	5489	487	0.03	0.78
1962	20849	1655	0.744	5471	465	0.02	0.79
1963	20754	1645	0.740	5456	473	0.02	0.79
1964	20658	1635	0.735	5440	468	0.02	0.79
1965	20575	1624	0.730	5425	438	0.02	0.80
1966	20525	1616	0.727	5413	444	0.02	0.80
1967	20470	1608	0.723	5401	463	0.02	0.79
1968	20399	1599	0.719	5387	497	0.03	0.78
1969	20299	1588	0.714	5369	460	0.02	0.79
1970	20238	1581	0.711	5358	416	0.02	0.81
1971	20223	1578	0.710	5354	409	0.02	0.81
1972	20211	1577	0.709	5352	423	0.02	0.80
1973	20184	1574	0.708	5348	429	0.02	0.80
1974	20150	1571	0.706	5343	415	0.02	0.81
1975	20130	1570	0.706	5341	429	0.02	0.80
1976	20097	1567	0.705	5337	440	0.02	0.80
1977	20057	1564	0.703	5331	452	0.02	0.79
1978	20010	1559	0.701	5324	536	0.03	0.76
1979	19887	1546	0.695	5304	584	0.03	0.74
1980	19732	1529	0.688	5277	444	0.02	0.79
1981	19724	1524	0.685	5268	547	0.03	0.75
1982	19618	1510	0.679	5246	486	0.03	0.77
1983	19576	1502	0.676	5233	466	0.03	0.78
1984	19551	1497	0.673	5224	420	0.02	0.80
1985	19565	1497	0.673	5224	453	0.03	0.79
1986	19541	1495	0.672	5221	425	0.02	0.80
1987	19539	1497	0.673	5224	431	0.02	0.79
1988	19529	1499	0.674	5228	415	0.02	0.80
1989	19534	1502	0.676	5233	413	0.02	0.80

Continues next page

Table 9: Time-series of population estimates from the base-case model. Relative exploitation rate is $(1 - SPR)/(1 - SPR_{50\%})$.

Year	Total biomass (mt)	Spawning biomass (mt)	Depletion	Age-0 recruits	Total catch (mt)	Relative exploitation rate	SPR
1990	19541	1506	0.677	5238	422	0.02	0.80
1991	19540	1507	0.678	5240	432	0.02	0.79
1992	19531	1506	0.677	5239	424	0.02	0.80
1993	19534	1505	0.677	5238	438	0.02	0.79
1994	19524	1503	0.676	5234	438	0.02	0.79
1995	19515	1500	0.675	5230	120	0.01	0.94
1996	19808	1525	0.686	5269	348	0.02	0.83
1997	19858	1529	0.688	5277	596	0.03	0.73
1998	19673	1512	0.680	5250	220	0.01	0.89
1999	19862	1529	0.688	5277	319	0.02	0.85
2000	19941	1538	0.692	5291	408	0.02	0.81
2001	19931	1539	0.692	5292	245	0.01	0.88
2002	20076	1554	0.699	5316	240	0.01	0.88
2003	20212	1569	0.706	5340	386	0.02	0.82
2004	20197	1571	0.707	5344	286	0.02	0.86
2005	20281	1582	0.711	5361	347	0.02	0.84
2006	20304	1588	0.714	5369	429	0.02	0.80
2007	20254	1585	0.713	5365	292	0.02	0.86
2008	20344	1593	0.716	5377	387	0.02	0.82
2009	20342	1591	0.715	5374	217	0.01	0.90
2010	20501	1604	0.721	5394	207	0.01	0.90
2011	20652	1618	0.727	5415	282	0.01	0.87
2012	20714	1626	0.731	5427	282	0.01	0.87
2013	20769	1635	0.735	5441	144	0.01	0.93
2014	20947	1657	0.745	5474	397	0.02	0.82
2015	20874	1657	0.745	5474	351	0.02	0.84
2016	20859	1660	0.746	5478	441	0.02	0.80
2017	20770	1652	0.743	5466	297	0.02	0.86
2018	20833	1655	0.744	5471	185	0.01	0.91
2019	0	1667	0.750	5488			

Table 10: Sensitivity of the base model to assumptions about selectivity and catchability.

Label	Base model	Sel all asymptotic	Sel all domed	Sel no sex offset	Q no prior on WCGBTS	Q no offset on triennial
TOTAL likelihood	402.12	402.10	402.12	402.11	401.67	402.95
Survey likelihood	-9.72	-9.71	-9.72	-9.72	-9.31	-9.38
Length comp likelihood	341.44	341.39	341.44	341.44	340.46	342.01
Age comp likelihood	97.14	97.16	97.14	97.14	97.08	96.99
Discard likelihood	-22.45	-22.42	-22.45	-22.45	-22.14	-22.64
Mean body wt likelihood	-4.42	-4.45	-4.42	-4.42	-4.60	-4.27
Parm priors likelihood	0.12	0.12	0.12	0.12	0.17	0.23
Recr Virgin millions	6.18	6.13	6.18	6.18	34.78	5.94
log(R0)	8.73	8.72	8.73	8.73	10.46	8.69
NatM Female	0.45	0.44	0.45	0.45	0.46	0.45
NatM Male	0.45	0.44	0.45	0.45	0.46	0.45
Linf Female	175.67	175.62	175.67	175.67	175.61	175.40
Linf Male	120.97	120.94	120.97	120.97	120.95	121.01
Q WCGBTS	0.81	0.81	0.81	0.81	0.14	0.90
SSB Virgin thousand mt	2.22	2.27	2.22	2.22	9.93	1.91
SSB 2019 thousand mt	1.67	1.71	1.67	1.67	9.50	1.37
Bratio 2019	0.75	0.75	0.75	0.75	0.96	0.72
SPRratio 2018	0.18	0.17	0.18	0.18	0.03	0.20
Retained Catch MSY	558.67	562.75	558.67	558.67	2793.89	510.57
Dead Catch MSY	603.92	608.37	603.92	603.92	3030.18	551.56

Table 11: Sensitivity of the base model to assumptions about catches.

Label	Base model	Discards based on 3yr averages	Discard mortality 0 4	Discard mortality 0 6
TOTAL likelihood	402.12	401.58	401.85	402.36
Survey likelihood	-9.72	-9.92	-9.98	-9.49
Length comp likelihood	341.44	341.12	341.61	341.28
Age comp likelihood	97.14	97.24	97.14	97.13
Discard likelihood	-22.45	-22.51	-22.66	-22.26
Mean body wt likelihood	-4.42	-4.46	-4.39	-4.45
Parm priors likelihood	0.12	0.11	0.11	0.14
Recr Virgin millions	6.18	6.02	6.19	6.19
log(R0)	8.73	8.70	8.73	8.73
NatM Female	0.45	0.44	0.44	0.45
NatM Male	0.45	0.44	0.44	0.45
Linf Female	175.67	175.76	175.68	175.66
Linf Male	120.97	120.95	120.96	120.98
Q WCGBTS	0.81	0.83	0.82	0.80
SSB Virgin thousand mt	2.22	2.23	2.29	2.17
SSB 2019 thousand mt	1.67	1.62	1.67	1.66
Bratio 2019	0.75	0.73	0.73	0.77
SPRratio 2018	0.18	0.18	0.18	0.18
Retained Catch MSY	558.67	551.42	567.17	552.69
Dead Catch MSY	603.92	595.86	612.92	597.60

Table 12: Sensitivity of the base model to assumptions about biology and misc.

Label	Base model	Bio separate M by sex	Bio no M prior	Bio von Bertalanffy growth	Bio Richards growth	Misc McAllister Ianelli tuning
TOTAL likelihood	402.12	399.94	402.00	445.19	456.54	1116.89
Survey likelihood	-9.72	-9.88	-9.72	-9.54	-9.73	-9.66
Length comp likelihood	341.44	338.79	341.48	387.56	362.67	564.52
Age comp likelihood	97.14	97.53	97.09	94.06	129.88	591.26
Discard likelihood	-22.45	-22.79	-22.47	-22.39	-21.98	-22.34
Mean body wt likelihood	-4.42	-3.92	-4.41	-5.05	-4.33	-7.13
Parm priors likelihood	0.12	0.21	0.01	0.53	0.01	0.24
Recr Virgin millions	6.18	5.19	6.29	17.80	0.00	7.26
log(R0)	8.73	8.55	8.75	9.79	8.03	8.89
NatM Female	0.45	0.47	0.45	0.57	0.36	0.46
NatM Male	0.45	0.40	0.45	0.57	0.36	0.46
Linf Female	175.67	175.53	175.65	587.20	2595.92	176.97
Linf Male	120.97	120.15	120.99	236.34	136.91	120.50
Q WCGBTS	0.81	0.81	0.81	0.84	0.85	0.77
SSB Virgin thousand mt	2.22	1.37	2.20	1.25	0.00	2.37
SSB 2019 thousand mt	1.67	0.87	1.65	1.02	0.00	1.83
Bratio 2019	0.75	0.63	0.75	0.82	0.00	0.77
SPRratio 2018	0.18	0.26	0.18	0.13	0.89	0.16
Retained Catch MSY	558.67	432.06	561.23	751.54	0.00	601.43
Dead Catch MSY	603.92	465.72	606.68	812.55	0.00	650.09

Table 13: Summaries of key assessment outputs and likelihood values from the retrospective analysis. Note that male growth parameters are exponential offsets from female parameters, and depletion and SPR ratio are for the year of 2017. The base model includes all of the data. Retro1 removes the last year of data (2016), Retro2 removes the last two years of data, Retro3 removes three years and Retro4 removes four years.

Label	Base	Retro1	Retro2	Retro3	Retro4
Female natural mortality	0.26	0.26	0.26	0.26	0.26
Steepness	0.72	0.72	0.72	0.72	0.72
lnR0	8.16	8.09	8.07	8.04	8.08
Total Biomass (mt)	2796.86	2593.78	2568.77	2498.07	2650.36
Depletion	57.41	53.57	50.74	50.72	54.78
SPR ratio	0.72	0.76	0.79	0.80	0.74
Female Lmin	12.43	12.45	12.90	12.63	13.03
Female Lmax	33.31	33.50	33.39	33.37	33.46
Female K	0.25	0.24	0.24	0.25	0.23
Male Lmin (offset)	0.00	0.00	0.00	0.00	0.00
Male Lmax (offset)	-0.16	-0.16	-0.15	-0.16	-0.15
Male K (offset)	-0.29	-0.30	-0.43	-0.41	-0.56
Negative log-likelihood	1097.30	1047.56	1009.37	961.81	897.04
No. parameters	0.00	0.00	0.00	0.00	0.00
TOTAL	0.00	0.00	0.00	0.00	0.00
Equilibrium catch	-98.12	-92.00	-89.12	-81.75	-80.59
Survey	763.02	739.90	720.39	700.10	670.66
Length composition	421.52	390.56	369.97	336.26	299.84
Age composition	10.88	9.09	8.12	7.20	7.12
Recruitment	0.00	0.00	0.00	0.00	0.00
Forecast Recruitment	0.00	0.00	0.00	0.00	0.00
Parameter priors	0.01	0.01	0.01	0.01	0.01

Table 14: Summaries of key assessment outputs and likelihood values from selected likelihood profile runs on virgin recruitment ($\ln R_0$) and steepness. Note that male growth parameters are exponential offsets from female parameters, and depletion and SPR ratio are for the year of 2017.

Label	R07400	R07800	R08200	R08600	R09000	h0410	h0570	h0710	h0870	h0990
Female M	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26	0.26
Steepness	0.72	0.72	0.72	0.72	0.72	0.41	0.57	0.71	0.87	0.99
lnR0	7.40	7.80	8.20	8.60	9.00	8.34	8.21	8.16	8.13	8.11
Total biomass (m)	1623.19	2113.03	2894.72	4173.95	6142.97	3313.42	2943.85	2802.69	2712.12	2667.97
Depletion (%)	46.83	49.83	58.31	66.23	71.80	51.20	55.27	57.32	58.81	59.60
SPR ratio	1.05	0.91	0.70	0.49	0.34	0.68	0.71	0.72	0.72	0.73
Female Lmin	12.16	12.41	12.43	12.39	12.36	12.43	12.44	12.43	12.43	12.43
Female Lmax	34.29	33.83	33.26	32.76	32.42	33.19	33.28	33.31	33.33	33.34
Female K	0.24	0.25	0.25	0.26	0.26	0.25	0.25	0.25	0.25	0.25
Male Lmin (offset)	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Male Lmax (offset)	-0.18	-0.17	-0.16	-0.15	-0.15	-0.16	-0.16	-0.16	-0.16	-0.16
Male K (offset)	-0.22	-0.31	-0.29	-0.24	-0.21	-0.27	-0.29	-0.29	-0.30	-0.30
Negative log-likelihood										
TOTAL	1117.15	1101.02	1097.33	1099.69	1102.95	1101.35	1098.58	1097.35	1096.72	1100.21
Catch	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Equil_catch	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Survey	-100.10	-99.20	-97.99	-97.00	-96.37	-98.27	-98.18	-98.12	-98.06	-98.03
Length_comp	761.18	760.12	763.44	767.61	770.76	765.11	763.69	763.05	762.58	762.33
Age_comp	437.32	427.37	421.09	418.57	417.98	420.58	421.24	421.51	421.68	421.77
Recruitment	18.74	12.72	10.80	10.50	10.58	12.55	11.40	10.90	10.56	10.38
Forecast_Recruitment	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Parm_priors	0.00	0.00	0.00	0.00	0.00	1.38	0.42	0.01	-0.04	3.76
Parm_softbounds	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01	0.01
Parm_devs	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00
Crash_Pen	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00

Table 15: Summaries of key assessment outputs and likelihood values from selected likelihood profile runs on female natural mortality. Note that male growth parameters are exponential offsets from female parameters, and depletion and SPR ratio are for the year of 2017.

Label	M0220	M0260	M0300	M0350	M0400
Female M	0.22	0.26	0.30	0.35	0.40
Steepness	0.72	0.72	0.72	0.72	0.72
lnR0	7.67	8.20	8.95	12.21	31.00
Total biomass (m)	2259.39	2861.79	4632.81	89473.50	9753570000000.00
Depletion (%)	47.72	58.15	68.08	79.27	79.74
SPR ratio	0.97	0.70	0.41	0.02	0.00
Female Lmin	12.39	12.44	12.43	12.39	12.24
Female Lmax	33.23	33.31	33.31	33.25	33.73
Female K	0.25	0.25	0.25	0.25	0.24
Male Lmin (offset)	0.00	0.00	0.00	0.00	0.00
Male Lmax (offset)	-0.16	-0.16	-0.15	-0.15	-0.15
Male K (offset)	-0.27	-0.30	-0.31	-0.32	-0.36
Negative log-likelihood					
TOTAL	1102.66	1096.96	1092.96	1089.92	1091.52
Catch	0.00	0.00	0.00	0.00	0.00
Equil_catch	0.00	0.00	0.00	0.00	0.00
Survey	-97.79	-98.14	-98.33	-98.33	-98.95
Length_comp	765.50	762.85	760.88	759.19	755.26
Age_comp	422.97	421.41	420.05	418.75	425.16
Recruitment	11.91	10.82	10.30	10.05	9.54
Forecast_Recruitment	0.00	0.00	0.00	0.00	0.00
Parm_priors	0.06	0.00	0.06	0.25	0.51
Parm_softbounds	0.01	0.01	0.01	0.00	0.00
Parm_devs	0.00	0.00	0.00	0.00	0.00
Crash_Pen	0.00	0.00	0.00	0.00	0.00

Table 16: Projection of potential OFL, spawning biomass, and depletion for the base case model.

Yr	OFL contribution (mt)	ACL landings (mt)	Age 5+ biomass (mt)	Spawning Biomass (mt)	Depletion
2019	1389.940	313.160	0.000	1667.190	0.750
2020	1390.490	313.160	0.000	1664.770	0.749
2021	1390.540	1136.647	0.000	1662.950	0.748
2022	1327.210	1072.121	0.000	1581.990	0.711
2023	1278.000	1021.539	0.000	1507.590	0.678
2024	1241.120	982.221	0.000	1438.770	0.647
2025	1212.850	950.914	0.000	1374.480	0.618
2026	1189.120	923.817	0.000	1314.410	0.591
2027	1167.280	899.641	0.000	1259.890	0.566
2028	1145.980	875.107	0.000	1213.480	0.546
2029	1124.960	851.041	0.000	1177.730	0.530
2030	1104.190	828.385	0.000	1152.760	0.518

941 11 Figures

942 11.1 Data Figures




Figure 1: Summary of data sources used in the model.




Figure 2: Map showing the distribution of Big Skate within the area covered by the West Coast Groundfish Bottom Trawl Survey aggregated over the years 2003–2018.




Figure 3: Reconstructed landings by area. Tribal catch was all landed in Washington.




Figure 4: Estimated total catch using different assumptions for discards. The discard rates shown in green lines are relative to the right-hand axis while all other values are relative to the left-hand axis.




Figure 5: Catch data input to the model under assumed fleet structure. The historical discards shown in green have been scaled to account for an assumed 50% discard mortality. Discards during the period from 1995 onward are not represented here as they are estimated within the model.




Figure 6: Map of estimated density by year for Big Skate in the Triennial survey.
71




Figure 7: Map of estimated density by year for Big Skate in the WCGBT Survey.

Big Skate per 100 observed hooks in IPHC longline survey




Figure 8: Map of catch rates by year for Big Skate in the International Pacific Halibut Commission longline survey.

943 11.2 Biology Figures




Figure 9: Length comp data, aggregated across time by fleet.




Figure 10: Length comp data for all years and fleets. Bubble size indicates the observed proportions, with females in red and males in blue.




Figure 11: Conditional age-at-length data from the fishery.




Figure 12: Conditional age-at-length data from the WCGBT Survey.




Figure 13: Comparison of reads from each of two age readers for Big Skate. Sample sizes associated with each combination of ages are shown by the size circles and the numbers within them. The blue histograms show the distribution of ages estimated by each reader.




Figure 14: Estimated ageing imprecision.




Figure 15: Estimated relationship between length and weight (left) and disc-width and length (right) for Big Skate. Colored points show observed values and the black line indicates the estimated relationship $W = 0.0000074924L^{2.9925}$.




Figure 16: Estimated maturity relationship for female Big Skate. Gray points indicate average observed functional maturity within each length bin with point size proportional to the number of samples (indicated by text within each point).

⁹⁴⁴ **11.3 Model Results Figures**

⁹⁴⁵ **11.3.1 Growth and Selectivity**




Figure 17: Estimated length-at-age for female and male Big Skate (top left panel). Shaded areas indicate 95% intervals for distribution of lengths at each age. Values represent beginning-of-year growth. Weight (thick line) and maturity (thin line) are shown in the top-right and lower-left panels as a function of length and age, respectively, where the values-at-age are calculated by mapping the length-based relationships through the estimated distribution of length at each age.




Figure 18: Selectivity at length for all of the fleets in the base model.

946 **11.3.2 Fits to the Data**




Figure 19: Fit to index data for WCGBT Survey. Lines indicate 95% uncertainty interval around index values. Thicker lines indicate input uncertainty before addition of estimated additional uncertainty parameter. The blue line indicates the model estimate.




Figure 20: Fit to index data for Triennial Survey. Lines indicate 95% uncertainty interval around index values. Thicker lines indicate input uncertainty before addition of estimated additional uncertainty parameter. The blue line indicates the model estimate with a change between 1992 and 1995 associated with the estimated change in catchability.




Figure 21: Fits to length comp data, aggregated across time by fleet.




Figure 22: Pearson residuals for length comp data for all years and fleets, with females in red and males in blue.




Figure 23: Pearson residuals for the fit to conditional age-at-length data from the fishery.




Figure 24: Pearson residuals for the fit to conditional age-at-length data from the WCGBT Survey.




Figure 25: Observed sex ratios (points) from the fishery length comp data with 75% intervals (vertical lines) calculated as a Jeffreys interval based on the adjusted input sample size. The model expectation is shown in the blue line.




Figure 26: Observed sex ratios (points) from the WCGBT Survey length comp data with 75% intervals (vertical lines) calculated as a Jeffreys interval based on the adjusted input sample size. The model expectation is shown in the blue line.




Figure 27: Fit to the discard fraction estimates. Points are model estimates with 95% uncertainty intervals. The model estimate is shown in the blue lines.




Figure 28: Fit to the mean weight of the discards. Points are model estimates with 95% uncertainty intervals. The model estimate is shown in the blue lines.

947 **11.3.3 Time Series Figures**

Spawning biomass (mt) with ~95% asymptotic intervals




Figure 29: Estimated spawning biomass (mt) with approximate 95% asymptotic intervals.




Figure 30: Estimated spawning depletion with approximate 95% asymptotic intervals.



Figure 31: Estimated time-series of recruitment for Big Skate.




Figure 32: Estimated recruitment and the assumed stock-recruit relationship.




Figure 33: Estimated total catch including discards estimated within the model. The historical discards shown in green have been scaled to account for an assumed 50% discard mortality but the discards in the recent period show both live and dead discards.

⁹⁴⁸ 11.3.4 Sensitivity Analyses and Retrospectives




Figure 34: Time series of spawning biomass (mt) estimated in sensitivity analyses related to selectivity and catchability.




Figure 35: Time series of spawning biomass (mt) estimated in sensitivity analyses related to historic catch and discards.




Figure 36: Time series of spawning biomass (mt) estimated in sensitivity analyses related to biology and other assumptions.




Figure 37: Comparison of the estimated growth curves from the sensitivities analyses.

⁹⁴⁹ 11.3.5 Likelihood Profiles




Figure 38: Likelihood profile over the log of equilibrium recruitment (R_0).




Figure 39: Likelihood profile over the log of equilibrium recruitment (R_0).




Figure 40: Time series of spawning biomass (mt) estimated for the models included in the profile over the log of equilibrium recruitment (R_0).




Figure 41: Likelihood profile over stock-recruit steepness (h).




Figure 42: Time series of spawning biomass (mt) estimated for the models included in the profile over stock-recruit steepness (h).




Figure 43: Likelihood profile over natural mortality (M).




Figure 44: Time series of spawning biomass (mt) estimated for the models included in the profile over natural mortality (M).

⁹⁵⁰ 11.3.6 Reference Points and Forecasts




Figure 45: Equilibrium yield curve for the base case model. Values are based on the fishery selectivity and with steepness fixed at 0.4.

⁹⁵¹ Appendix A. Detailed fits to length composition data




Figure A46: Length comps, retained, Fishery. ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAlister_Iannelli tuning method.




Figure A47: Length comps, discard, Fishery. ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAlister_Iannelli tuning method.




Figure A48: Length comps, whole catch, WCGBT Survey. 'N adj.' is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method.




Figure A49: Length comps, whole catch, Triennial Survey. ‘N adj.’ is the input sample size after data_weighting adjustment. N eff. is the calculated effective sample size used in the McAllister_Iannelli tuning method.

952 **References**

- 953 Alaska Fisheries Science Center. 2018. Assessment of the skate stock complex in the Gulf of
954 Alaska. Available from {<https://www.afsc.noaa.gov/REFM/Docs/2018/GOA/GOAskate.pdf>}.
- 955 Batdorf, C. 1990. Northwest Native Harvest. Hancock House Publishers Ltd.; Surrey, B.C.,
956 Canada.
- 957 Bizzarro, J. 2015. Comparative resource utilization of eastern north pacific skates (raji-
958 formes: Rajidae) with applications for ecosystem-based fisheries management. WA: Univer-
959 sity of Washington.
- 960 Bizzarro, J. 2019. Manuscript in preparation.
- 961 Bizzarro, JJ and Broms, KM and Logsdon, MG and Ebert, DA and Yoklavich, MM and
962 Kuhnz, LA and Summers, AP. 2014. Spatial segregation in eastern north Pacific skate
963 assemblages. PloS one **9**(10).
- 964 Bizzarro, J., Robinson, H., Rinewalt, C., and Ebert, D. 2007. Comparative feeding ecology
965 of four sympatric skate species off central California, USA. In Biology of skates. Springer.
966 pp. 91–114.
- 967 Bowers, G. M. 1909. Report of The Commissioner For the Year Ending June 30, 1909. Part
968 XXVIII. Washington Printing Office.
- 969 Bradburn, M.J. and Keller, A.A and Horness, B.H. 2011. The 2003 to 2008 US West
970 Coast bottom trawl surveys of groundfish resources off Washington, Oregon, and Califor-
971 nia: estimates of distribution, abundance, length, and age composition. NOAA Technical
972 Memorandum NMFS NOAA-TM-NMFS-NWFSC-114: 323 pp.
- 973 Calavan, T. 2019. Oregon Department of Fisheries; Wildlife; Personal Communication,
974 Newport, OR, USA.
- 975 Castro-Aguirre, J.L., and Pérez, H.E. 1996. Catálogo sistemático de las rayas y especies
976 afines de méxico: Chondrichthyes: Elasmobranchii: Rajiformes: Batoideiomorpha. Unam.
- 977 Castro-Aguirre, J., Schmitter, J., Balart, E., and Torres-Orozco, R. 1993. Sobre la dis-
978 tribución geográfica de algunos peces bentónicos de la costa oeste de baja california sur,
979 méxico, con consideraciones ecológicas y evolutivas. In Anales de la escuela nacional de
980 ciencias biológicas, méxico. pp. 75–102.
- 981 Chapman, W.M. 1944. The Latent Fisheries of Washington and Alaska. Washington State
982 Department of Fisheries.

- 983 Chiquillo, Kelcie L and Ebert, David A and Slager, Christina J and Crow, Karen D. 2014.
984 The secret of the mermaid's purse: Phylogenetic affinities within the Rajidae and the evolution
985 of a novel reproductive strategy in skates. *Molecular Phylogenetics and Evolution* **75**:
986 245–251. Elsevier.
- 987 DeLacy, A.C., and Chapman, W.M. 1935. Notes on some elasmobranchs of puget sound,
988 with descriptions of their egg cases. *Copeia* **1935**(2): 63–67. JSTOR.
- 989 Downs, D.E., and Cheng, Y.W. 2013. Length-length and width-length conversion of long-
990 nose skate and big skate off the pacific coast: Implications for the choice of alternative
991 measurement units in fisheries stock assessment. *North American journal of fisheries man-
992 agement* **33**(5): 887–893. Taylor & Francis.
- 993 Ebert, D. 2003. Sharks, rays, and chimaeras of california. Univ of California Press.
- 994 Ebert, D.A., and Compagno, L.J. 2007. Biodiversity and systematics of skates (chon-
995 drichthyes: Rajiformes: Rajoidei). In *Biology of skates*. Springer. pp. 5–18.
- 996 Ebert, D.A., Smith, W.D., and Cailliet, G.M. 2008. Reproductive biology of two commer-
997 cially exploited skates, raja binoculata and r. Rhina, in the western gulf of alaska. *Fisheries
998 Research* **94**(1): 48–57. Elsevier.
- 999 Eschmeyer, W.N., and Herald, E.S. 1983. A field guide to pacific coast fishes: North america.
1000 Houghton Mifflin Harcourt.
- 1001 Farrugia, T.J., Goldman, K.J., Tribuzio, C., and Seitz, A.C. 2016. First use of satellite
1002 tags to examine movement and habitat use of big skates beringraja binoculata in the gulf of
1003 alaska. *Marine Ecology Progress Series* **556**: 209–221.
- 1004 Ford, P. 1971. Differential growth rate in the tail of the pacific big skate, (*Raja binoculata*).
1005 *Journal of the Fisheries Board of Canada* **28**(1): 95–98. NRC Research Press.
- 1006 Francis, R.I.C.C. 2011. Data weighting in statistical fisheries stock assessment models. *Canan-
1007 dian Journal of Fisheries and Aquatic Sciencies* **68**: 1124–1138.
- 1008 Gburski, C.M. and Gaichas, S.K. and Kimura, D.K. 2007. Age and growth of big skate
1009 (*Raja binoculata*) and longnose skate (*Raja rhina*) in the Gulf of Alaska. In *Biology of
1010 Skates*. Springer, Dordrecht.
- 1011 Gertseva, V. 2019. Manuscript in preparation.
- 1012 Gertseva, V and Schrippa, MJ. 2007. Status of the Longnose Skate (*Raja rhina*) off the
1013 continental US Pacific Coast in 2007. Pacific Fishery Management Council, Portland, OR.
1014 Available from <http://www.pcouncil.org/groundfish/stock-assessments/>.

- 1015 Gertseva, V., and Taylor, I. 2011. Status of spiny dogfish shark resource off the continental
1016 us pacific coast in 2011. PFMC. 2011. Pacific Fishery Management Council, Portland, OR.
1017 Available from [\](http://www.pcouncil.org/groundfish/stock-assessments/).
- 1018 Gunderson, Donald Raymond and Sample, Terrance M. 1980. Distribution and
1019 abundance of rockfish off Washington, Oregon and California during 1977. Northwest
1020 and Alaska Fisheries Center, National Marine Fisheries Service. Available from
1021 [\](http://spo.nmfs.noaa.gov/mfr423-4/mfr423-42.pdf).
- 1022 Hamel, Owen S. 2015. A method for calculating a meta-analytical prior for the natural
1023 mortality rate using multiple life history correlates. ICES Journal of Marine Science: Journal
1024 du Conseil **72**(1): 62–69. doi: [10.1093/icesjms/fsu131](https://doi.org/10.1093/icesjms/fsu131).
- 1025 Hitz, C.R. 1964. Observations on egg cases of the big skate (*raja binoculata girard*) found
1026 in oregon coastal waters. Journal of the Fisheries Board of Canada **21**(4): 851–854. NRC
1027 Research Press.
- 1028 Hoff, GR. 2009. Skate *Bathyraja* spp. egg predation in the eastern Bering Sea. J. Fish.
1029 Biol. **74**: 250–269.
- 1030 Ishihara, H., Treloar, M., Bor, P., Senou, H., and Jeong, C. 2012. The comparative mor-
1031 phology of skate egg capsules (Chondrichthyes: Elasmobranchii: Rajiformes). Bulletin of
1032 the Kanagawa Prefectural Museum (Natural Science) **41**: 9–25.
- 1033 Keller, A.A. and Wallace, J.R. and Methot, R.D. 2017. The Northwest Fisheries Science
1034 Center's West Coast Groundfish Bottom Trawl Survey: History, Design, and Description.
1035 NOAA Technical Memorandum NMFS NOAA-TM-NMFS-NWFSC-136: 38 pp.
- 1036 King, J., and McFarlane, G. 2009. Biological results of the strait of georgia spiny dogfish
1037 (*squalus acanthias*) longline survey, october 10-22, 2008. Fisheries; Oceans Canada, Science
1038 Branch, Pacific Region.
- 1039 King, J.R., Surry, A.M., Garcia, S., and P.J. Starr. 2015. Big skate (*Raja binoculata*)
1040 and longnose skate (*R. rhina*) stock assessments for British Columbia. Ottawa : Canadian
1041 Science Advisory Secretariat.
- 1042 King, JR and McFarlane, GA. 2010. Movement patterns and growth estimates of big skate
1043 (*Raja binoculata*) based on tag-recapture data. Fish. Res. **101**: 50–59.
- 1044 Lippert, G. 2019. Washington Department of Fisheries; Wildlife; Personal Communication,
1045 Olympia, Washington, USA.
- 1046 Love, Milton S. 2011. Certainly more than you want to know about the fishes of the Pacific
1047 Coast: a postmodern experience. Really Big Press.

- 1048 Maunder, M.N., Deriso, R.B., Schaefer, K.M., Fuller, D.W., Aires-da-Silva, A.M., Minte-
1049 Vera, C.V., and Campana, S.E. 2018. The growth cessation model: A growth model for
1050 species showing a near cessation in growth with application to bigeye tuna (*thunnus obesus*).
1051 *Marine biology* **165**(4): 76. Springer.
- 1052 McEachran, J., and Miyake, T. 1990. Zoogeography and bathymetry of skates (chon-
1053 drichthyes, rajidae). Elasmobranchs as living resources. *Advances in biology, Ecology, Sys-*
1054 *tematics and the status of the fisheries:* 305–326.
- 1055 McFarlane GA and King JR. 2006. Age and growth of big skate (*Raja binoculata*) and
1056 longnose skate (*Raja rhina*) in British Columbia waters. *Fisheries Research* **May 1 (2-3)**:
1057 169–78.
- 1058 Mecklenburg, CW and Mecklenburg, TA and Thorsteinson, LK. 2002. *Fishes of Alaska.*
1059 American Fisheries Society, Bethesda, Maryland.
- 1060 Methot, RD Jr. and Wetzel, CR and Taylor, IG. 2019. Stock Synthesis User Manual Version
1061 3.30.13. NOAA Fisheries. Seattle, WA. Available from <https://vlab.ncep.noaa.gov/web/stock-synthesis>.
- 1063 Methot, Richard D. and Wetzel, Chantell R. 2013. Stock synthesis: A biological and statis-
1064 tical framework for fish stock assessment and fishery management. *Fisheries Research* **142**:
1065 86–99.
- 1066 Miller, B.S., Cross, J.N., Steinfort, S.N., Fresh, K.L., and Simenstad, C.A. 1980. Nearshore
1067 fish and macroinvertebrate assemblages along the strait of juan de fuca including food habits
1068 of the common nearshore fish.
- 1069 Punt AE and Smith DC and KrusicGolub K and Robertson S. 2008. Quantifying age-reading
1070 error for use in fisheries stock assessments, with application to species in Australia's southern
1071 and eastern scalefish and shark fishery. *Canadian Journal of Fisheries and Aquatic Sciences*.
- 1072 Stevenson, DE and Orr, JW and Hoff, GR and McEachran, JD. 2008. Emerging patterns
1073 of species richness, diversity, population density, and distribution in the skates (Rajidae) of
1074 Alaska. *Fish Bull* **106**: 24–39.
- 1075 Stewart, I.J., Wallace, J.R., and McGilliard, C. 2009. Status of the us yelloweye rockfish
1076 resource in 2009. *In Pacific Fishery Management Council, Portland, OR. Available from*
1077 <http://www.pcouncil.org/groundfish/stock-assessments/>.
- 1078 Taylor, I.G., Stewart, I.J., Hicks, A.C., Garrison, T.M., Punt, A.E., Wallace, J.R., Wetzel,
1079 C.R., Thorson, J.T., Takeuchi, Y., Ono, K., Monnahana, C.C., Stawitz, C.C., A'mar, Z.T.,
1080 Whitten, A.R., Johnson, K.F., Emmet, R.L., Anderson, S.C., Lambert, G.I., Stachura,
1081 M.M., Cooper, A.B., Stephens, A., Klaer, N.L., McGilliard, C.R., Iwasaki, W.M., Doering,
1082 K., and Havron, A.M. 2019. R4ss: R code for stock synthesis. Available from <https://github.com/r4ss>.

1084 Taylor IG and Cope, J and Hamel O and Thorson, J. 2013. Deriving estimates of OFL for
1085 species in the “Other Fish” complex or potential alternative complexes. Pacific Fishery Man-
1086 agement Council, Portland, OR. Available from <http://www.pcouncil.org/groundfish/stock->
1087 [assessments/](#).

1088 Thorson, James T. and Barnett, Lewis A. K. 2017. Comparing estimates of abun-
1089 dance trends and distribution shifts using single- and multispecies models of fishes and
1090 biogenic habitat. ICES Journal of Marine Science: Journal du Conseil: fsw193. doi:
1091 [{10.1093/icesjms/fsw193}](https://doi.org/10.1093/icesjms/fsw193).

1092 Thorson, J. T. and Shelton, A. O. and Ward, E. J. and Skaug, H. J. 2015. Geostatistical
1093 delta-generalized linear mixed models improve precision for estimated abundance indices
1094 for West Coast groundfishes. ICES Journal of Marine Science **72**(5): 1297–1310. doi:
1095 [{10.1093/icesjms/fsu243}](https://doi.org/10.1093/icesjms/fsu243).

1096 Zeiner, S.J. and P. Wolf. 1993. Growth characteristics and estimates of age at maturity of
1097 two species of skates (*Raja binoculata*) and (*Raja rhina*) from Monterey Bay, California.