


Modelling the orientation dynamics of “chaotic mixing”/ turbulence.

Ian Towey
04128591

Msc Data and Computational Science


July 27, 2018

Introduction : Why study turbulence




Introduction : Why study turbulence

- It is a common phenomenon encountered in science and engineering.




Introduction : Why study turbulence

- It is a common phenomenon encountered in science and engineering.
- No analytical theory to predict the evolution of turbulent flows, closure problem.



Introduction : Why study turbulence

- It is a common phenomenon encountered in science and engineering.
- No analytical theory to predict the evolution of turbulent flows, closure problem.
- Large scale direct numerical simulations of turbulence are computationally intractable.




Introduction : Why study turbulence

- It is a common phenomenon encountered in science and engineering.
- No analytical theory to predict the evolution of turbulent flows, closure problem.
- Large scale direct numerical simulations of turbulence are computationally intractable.

“Turbulence is the most important unsolved problem of classical physics.”

- Richard P. Feynman
In The Feynman Lectures on
Physics (1964).



Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.
- Define Orientation dynamics model.

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.
- Define Orientation dynamics model.
- Define the model of the orientation dynamics using a system of SDEs. Derive the corresponding Fokker-Planck PDE

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.
- Define Orientation dynamics model.
- Define the model of the orientation dynamics using a system of SDEs. Derive the corresponding Fokker-Planck PDE
- Numerical Simulation of two-dimensional turbulence.

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.
- Define Orientation dynamics model.
- Define the model of the orientation dynamics using a system of SDEs. Derive the corresponding Fokker-Planck PDE
- Numerical Simulation of two-dimensional turbulence.
- Solve Fokker-Planck numerically to find the PDF of the orientation angle and growth rate.

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.
- Define Orientation dynamics model.
- Define the model of the orientation dynamics using a system of SDEs. Derive the corresponding Fokker-Planck PDE
- Numerical Simulation of two-dimensional turbulence.
- Solve Fokker-Planck numerically to find the PDF of the orientation angle and growth rate.
- Compare the PDF of the orientation angle from the vorticity simulation vs and FP model.

Context of work

Modelling orientation angle and growth rate of a tracer gradient in two-dimensional turbulence:

- Define 2-d turbulence model.
- Define Orientation dynamics model.
- Define the model of the orientation dynamics using a system of SDEs. Derive the corresponding Fokker-Planck PDE
- Numerical Simulation of two-dimensional turbulence.
- Solve Fokker-Planck numerically to find the PDF of the orientation angle and growth rate.
- Compare the PDF of the orientation angle from the vorticity simulation vs and FP model.
- Use uncertainty quantification methods to fit angle and growth rate PDF parameters to the simulated data.

Advection-Diffusion equation

$$\frac{\partial \theta}{\partial t} + \mathbf{u} \cdot \nabla \theta = \kappa \nabla^2 \theta$$

Advection-Diffusion equation

$$\frac{\partial \theta}{\partial t} + \mathbf{u} \cdot \nabla \theta = \kappa \nabla^2 \theta$$

- $\mathbf{u}(x, t)$ is the velocity field
- θ is the concentration of some passive scalar (tracer)
- κ is the diffusion coefficient (in this context $\kappa = 0$)
- $\nabla \cdot \mathbf{u} = 0$ (incompressible flow)

Advection-Diffusion equation

$$\frac{\partial \theta}{\partial t} + \mathbf{u} \cdot \nabla \theta = \kappa \nabla^2 \theta$$

- $\mathbf{u}(x, t)$ is the velocity field
- θ is the concentration of some passive scalar (tracer)
- κ is the diffusion coefficient (in this context $\kappa = 0$)
- $\nabla \cdot \mathbf{u} = 0$ (incompressible flow)

The model for the orientation dynamics will be derived from the Advection-Diffusion equation

Vorticity Equation

$$\frac{\partial \omega}{\partial t} + \mathbf{u} \cdot \nabla \omega = (-1)^p \nu_p \nabla^{2p} \omega + \nabla \times \mathbf{F} - \nabla \times \mathbf{D}$$

Vorticity Equation

$$\frac{\partial \omega}{\partial t} + \mathbf{u} \cdot \nabla \omega = (-1)^p \nu_p \nabla^{2p} \omega + \nabla \times \mathbf{F} - \nabla \times \mathbf{D}$$

- ω is the vorticity, describes the rotation of a fluid particle some point
- $\mathbf{u}(x, t)$ is the velocity
- ν is the viscosity
- \mathbf{F} is a forcing term
- \mathbf{D} is a damping term

Orientation Dynamics

Orientation Dynamics

$$\frac{\partial \mathcal{B}}{\partial t} + \mathbf{u} \cdot \nabla \mathcal{B} = \mathcal{B} \cdot \nabla \mathbf{u}$$

$$\mathcal{B} = (-\theta_y, \theta_x)$$

Orientation Dynamics

$$\frac{\partial \mathcal{B}}{\partial t} + \mathbf{u} \cdot \nabla \mathcal{B} = \mathcal{B} \cdot \nabla \mathbf{u}$$

$$\mathcal{B} = (-\theta_y, \theta_x)$$

$$\frac{d}{dt} |\mathcal{B}|^2 = -2\mu \sin \zeta |\mathcal{B}|^2$$

Orientation Dynamics

$$\frac{\partial \mathcal{B}}{\partial t} + \mathbf{u} \cdot \nabla \mathcal{B} = \mathcal{B} \cdot \nabla \mathbf{u}$$

$$\mathcal{B} = (-\theta_y, \theta_x)$$

$$\frac{d}{dt} |\mathcal{B}|^2 = -2\mu \sin \zeta |\mathcal{B}|^2$$

The growth rate of the tracer gradient is defined as

$$\Lambda = -2\mu \sin \zeta$$

- μ is the strain rate
- ζ orientation angle

Stochastic Model Equation Model

Stochastic Model Equation Model

$$\begin{aligned}\gamma \frac{dX}{dt} &= w + (-\cos(X) + k\sqrt{\delta})Y + Z \\ \frac{dY}{dt} &= -\frac{Y}{\tau} + \frac{\sqrt{D_Y}}{\tau} \xi_Y \\ \frac{dZ}{dt} &= -\frac{Z}{\tau} + \frac{\sqrt{D_Z(1-k^2)}}{\tau} \xi_Z\end{aligned}$$

Stochastic Model Equation Model

$$\begin{aligned}\gamma \frac{dX}{dt} &= w + (-\cos(X) + k\sqrt{\delta})Y + Z \\ \frac{dY}{dt} &= -\frac{Y}{\tau} + \frac{\sqrt{D_Y}}{\tau} \xi_Y \\ \frac{dZ}{dt} &= -\frac{Z}{\tau} + \frac{\sqrt{D_Z(1-k^2)}}{\tau} \xi_Z\end{aligned}$$

Corresponding Fokker-Planck equation



$$\frac{\partial P}{\partial t} = \mathcal{L}_{OU} P - \frac{\partial}{\partial t}(VP)$$

where

$$\mathcal{L}_{OU} P = \frac{1}{\tau_Y} \frac{\partial}{\partial t} (y \circ) + \frac{1}{\tau_Y^2} \frac{\partial^2}{\partial y^2} + \frac{1}{\tau_Z} \frac{\partial}{\partial Z} (z \circ) + \frac{\rho}{\tau_Z^2} \frac{\partial^2}{\partial z^2} + \frac{2c\sqrt{\rho}}{\tau_Y \tau_Z} \frac{\partial^2}{\partial y \partial z}$$



$$V = 2(w + y \cos x + z)$$

Numerical Simulation : Vorticity Equation





Numerical Simulation : Vorticity Equation

- Periodic Boundary Conditions





Numerical Simulation : Vorticity Equation

- Periodic Boundary Conditions
- Discretise the Vorticity equation





Numerical Simulation : Vorticity Equation

- Periodic Boundary Conditions
- Discretise the Vorticity equation
- Convert discretised vorticity equation to fourier space





Numerical Simulation : Vorticity Equation

- Periodic Boundary Conditions
- Discretise the Vorticity equation
- Convert discretised vorticity equation to fourier space
- Save snapshots of ω at regular intervals



Numerical Simulation : Vorticity Equation

- Periodic Boundary Conditions
- Discretise the Vorticity equation
- Convert discretised vorticity equation to fourier space
- Save snapshots of ω at regular intervals
- Extract empirical PDF of angle X statistically stable dataset



Numerical Simulation : Fokker Planck

Numerical Simulation : Fokker Planck

- Periodic Boundary Conditions

Numerical Simulation : Fokker Planck

- Periodic Boundary Conditions
- Solve for PDF of the Fokker-Planck

Numerical Simulation : Fokker Planck


- Periodic Boundary Conditions
- Solve for PDF of the Fokker-Plank
- Extract marginal probability of X for the PDF of the angle X

Numerical Simulation : Fokker Planck

- Periodic Boundary Conditions
- Solve for PDF of the Fokker-Plank
- Extract marginal probability of X for the PDF of the angle X
- Compute the PDF of the growth rate Λ using the joint PDF of XY


Analysis to date

- Verify simulation of vorticity has reached a steady state




Analysis to date

- Verify simulation of vorticity has reached a steady state
- Compare the PDF of the angle X from the SIM vs the SDE model



Analysis to date

- Verify simulation of vorticity has reached a steady state
- Compare the PDF of the angle X from the SIM vs the SDE model
- Compare the PDF of the growth rate from th SIM vs the SDE model



Outstanding Work

Outstanding Work

- Apply uncertainty quantification methods to the SDE/Fokker-Planck equation to fit the model parameters

Conclusions

The End