

Notes on Topology

January 13, 2020

Contents

1	Topological Spaces	2
1.1	Topology	2
1.2	Interior, exterior, boundary and closure	2
1.3	Neighborhoods	3
2	Continuity and Limits	5
2.1	Continuity	5
2.2	Limits	6
2.3	Sequences	6
3	Metric Spaces	7
3.1	Metrics	7
3.2	The Topology of a Metric Space	7
3.3	Continuity	8

Some conventions:

- $\mathbb{N} = \{0, 1, 2, \dots\}$,
- A_+ with $A = \mathbb{R}, \mathbb{Q}, \mathbb{N}, \mathbb{Z}$, refers to the respective subset of positive elements,
- A_- is the same as above but for negative elements.

1 Topological Spaces

1.1 Topology

Definition 1.1. Let E be a set. The set $\tau \subseteq \mathcal{P}(E)$ is said to be a *topology on E* if it satisfies the following axioms:

1. $\emptyset, E \in \tau$,
2. $O_1, O_2 \in \tau \Rightarrow (O_1 \cap O_2) \in \tau$,
3. $(O_j \in \tau : j \in J) \Rightarrow (\cup_{j \in J} O_j) \in \tau$.

A *topological space* is an ordered pair (E, τ) where E is a set and τ a topology on E . An element of $X \in \mathcal{P}(E)$ is said to be an *open set* if $X \in \tau$, and said to be a *closed set* if $E \setminus X \in \tau$.

Definition 1.2. A topological space (E, τ) is said to be *Hausdorff* if

$$\forall a, b \in E \text{ such that } a \neq b, \text{ there exists } O_a, O_b \in \tau \text{ and } O_a \cap O_b = \emptyset$$

Definition 1.3. Let τ_1, τ_2 be two topologies on a set E . The topology τ_1 is said to be *finer* than τ_2 if $\tau_2 \subseteq \tau_1$. If in addition $\tau_1 \neq \tau_2$, then τ_1 is said to be *strictly finer* than τ_2 .

Two topologies are said to be *comparable* if one is finer than the other.

1.2 Interior, exterior, boundary and closure

Definition 1.4. Let (E, τ) be a topological space, a an element of E , and $X \in \mathcal{P}(E)$. Then:

$$\begin{aligned} a \text{ is an interior point of } X &\Leftrightarrow \exists O \in \tau, a \in O \text{ and } O \subseteq X, \\ a \text{ is an exterior point of } X &\Leftrightarrow \exists O \in \tau, a \in O \text{ and } O \subseteq E \setminus X, \\ a \text{ is a boundary point of } X &\Leftrightarrow \forall O \in \tau, a \in O, O \cap X \neq \emptyset \text{ and } O \cap (E \setminus X) \neq \emptyset \end{aligned}$$

Notation 1. Note that the following sets depends on the fixed topology.

- $\text{int } X$ is the set of all interior points of X ,

- $\text{ext } X$ is the set of all exterior points of X ,
- ∂X is the set of all boundary points of X ,
- The set $\overline{X} = \text{int } X \cup \partial X$ is called the *closure* of X .

Remark 1. The elements of \overline{X} are called *adherent* point of X . It follows that

$$a \text{ is an adherent point of } X \Leftrightarrow \forall O \in \tau, a \in O \Rightarrow (O \cap X) \neq \emptyset.$$

Definition 1.5. Let (E, τ) be a topological space, let X be a subset of E and $a \in E$. The element a is said to be an *accumulation* point of X if:

$$\forall O \in \tau, a \in O \Rightarrow \exists b, a \neq b.$$

The set of such points is called the *derivative* of X , denoted X' . It follows that $X \cup X' = \overline{X}$.

Proposition 1.1. Let (E, τ) be a topological space and $X \in \mathcal{P}(E)$. Then

1. $a \in \overline{X} \Leftrightarrow \forall O \in \tau, a \in O \Rightarrow (O \cap X) \neq \emptyset$,
2. $\text{int } X \subseteq X \subseteq \overline{X}$,
3. $X \in \tau \Leftrightarrow X = \text{int } X$,
4. $(E \setminus X) \in \tau \Leftrightarrow \overline{X} = X$,
5. $\overline{X} = X \cup \partial X$,
6. $X \subseteq Y \Rightarrow \text{int } X \subseteq \text{int } Y \text{ and } \overline{X} \subseteq \overline{Y}$,
7. $\overline{X \cup Y} = \overline{X} \cup \overline{Y} \text{ and } \overline{X \cap Y} \subseteq \overline{X} \cap \overline{Y}$,
8. $\overline{\overline{X}} = \overline{X}$.

Definition 1.6. Let (E, τ) be a topological space, and $X, Y \subseteq E$. The set X is said to be *dense* on Y if $Y \subseteq \overline{X}$.

Definition 1.7. Let (E, τ) be a topological space; it is said to be *separable* if there exists a subset X of E that is both dense and countable.

1.3 Neighborhoods

Definition 1.8. Let (E, τ) be a topological space and $a \in E$. A subset X of E is said to be a *neighborhood* of a if there exists $O \in \tau$, such that $a \in O$ and $O \subseteq X$.

The set X is said to be a neighborhood of $A \subseteq E$, if for every element $a \in A$, X is a neighborhood of a . The set of all neighborhoods of a point a is denoted by \mathcal{V}_a , and \mathcal{V}_A denotes the set of all neighborhoods of an arbitrary set A .

Proposition 1.2.

$$\begin{aligned}
X \in \tau &\Leftrightarrow X \text{ is a neighborhood of } x \in X, \\
V_1, V_2 \in \mathcal{V}_a &\Rightarrow (V_1 \cap V_2) \in \mathcal{V}_a, \\
V \in \mathcal{V}_a \text{ and } V \subseteq W &\Rightarrow W \in \mathcal{V}_a, \\
V \in \mathcal{V}_a &\Rightarrow \text{int}(V) \in \mathcal{V}_a.
\end{aligned}$$

Definition 1.9. Let \mathcal{W} be a class of neighborhoods of a given point a (or a given set X) on the topological space (E, τ) . We call \mathcal{W} a *fundamental neighborhood system* of a (or X) if for every neighborhood V of a (or X), there exists $W \in \mathcal{W}$ such that $W \subseteq V$.

Definition 1.10. A topological space (E, τ) is said to satisfy the *first axiom of countability* if every $a \in E$, has a countable fundamental neighborhood system.

Definition 1.11. Let (J, \leq) be a well-ordered set. A class of neighborhoods $(V_j)_{j \in J}$ of a , indexed on J , is said to be a *nested* fundamental system of neighborhoods of a , if $V_k \subseteq V_j$ whenever $j \leq k$.

Proposition 1.3. *Every topological space that satisfies the first axiom of countability has a nested fundamental neighborhood system.*

Proof. Let (E, τ) be a topological space, and let $(W_n)_{n \in \mathbb{N}}$ be a fundamental neighborhood system of $a \in E$. For each $n \in \mathbb{N}$, define $V_n = \bigcap_{k=0}^n W_k$. Then $(V_n)_{n \in \mathbb{N}}$ is a fundamental neighborhood system of a , and $V_k \subseteq V_i$ whenever $i \leq k$. Therefore $(V_n)_{n \in \mathbb{N}}$ is a nested fundamental neighborhood system of a . \square

Theorem 1.4. *Let (E, τ) be a topological space; (E, τ) is separated if and only if, for every $a \in E$, the intersection of closed neighborhood of a is equal to $\{a\}$.*

Proof. Exercise! \square

Corollary 1.4.1. *For every Hausdorff (separated) topological space E and $a \in E$, the singleton $\{a\}$ is a closed set.*

Definition 1.12. In a topological space E , a point $a \in E$ is said to be *isolated* if $\{a\}$ is open, or equivalently, if $\{a\}$ is a neighborhood of a .

If E is a topological space, the class of all open sets of E , such that $a \in E$ is a fundamental neighborhood system of a . Therefore every point has a fundamental system of open neighborhoods. It is not true that every point has a fundamental system of closed neighborhoods. Hence the following definition.

Definition 1.13. A topological space E is said to be *regular* if it is Hausdorff and if every point $a \in E$ has a fundamental system of closed neighborhoods.

Definition 1.14. The class $\mathcal{B} \subseteq \mathcal{P}(E)$ is said to be a *basis* of a topological space (E, τ) (or a basis of τ) if, $\mathcal{B} \subseteq \tau$ and every open set of E is the intersection of elements of \mathcal{B} .

Remark 2. A basis for a topology τ determines τ in its entirety, since every open is the intersection of elements from the basis. If a topological space has a countable basis, then it is said to satisfy the *second axiom of countability*.

Theorem 1.5. Let (E, τ) be a topological space and $\mathcal{B} \in \tau$. The class \mathcal{B} is a basis of E if and only if, for every $a \in E$, the class $\{X \in \mathcal{B} : a \in X\}$ is a fundamental neighborhood system of a .

Proof. Exercise! □

Theorem 1.6. Let $\mathcal{B} \subseteq \mathcal{P}(E)$. Then there exists a unique topology τ in E , for which \mathcal{B} is a basis of the topological space (E, τ) if and only if, \mathcal{B} satisfies the following two conditions:

1.

$$\left(\bigcup_{A \in \mathcal{B}} A \right) = E.$$

2. For all $A, B \in \mathcal{B}$ and $x \in A \cap B$, there exists $C \in \mathcal{B}$, such that $x \in C$ and $C \subseteq (A \cap B)$.

Proof. □

2 Continuity and Limits

2.1 Continuity

Definition 2.1. Let (E_1, τ_1) and (E_2, τ_2) be topological spaces; let $f : (E_1, \tau_1) \rightarrow (E_2, \tau_2)$ be a function and a a point of E_1 . The function f is *continuous* on a if:

$$\forall O_2 \in \tau_2, f(a) \in O_2 : \exists O_1 \in \tau_1, a \in O_1 \Rightarrow f(O_1) \subseteq O_2.$$

This may be equivalently stated as:

$$\forall V \in \mathcal{V}_{f(a)}, \exists U \in \mathcal{V}_a : f(U) \subseteq V.$$

We'll prove that these two definitions are indeed equivalent.

Proof. □

Theorem 2.1. Let E_1, E_2 be topological spaces and $f : E_1 \rightarrow E_2$ a function. The following conditions are equivalent:

1. f is continuous;
2. If O is an open set of E_2 then its pre-image $f^{-1}(O)$ is an open set of E_1 ;
3. If C is a closed set of E_2 , then $f^{-1}(C)$ is closed in E_1 .

Proof. □

Definition 2.2. Let (E_1, τ_1) and (E_2, τ_2) be two topological spaces and $f : (E_1, \tau_1) \rightarrow (E_2, \tau_2)$ a function. The function f is said to be *open* (respectively *closed*) if $f(O) \in \tau_2$ (respectively $E_2 \setminus f(O) \in \tau_2$) whenever $O \in \tau_1$ (respectively $E_1 \setminus O \in \tau_1$).

Theorem 2.2. Let E_1, E_2, E_3 be topological spaces and $f : E_1 \rightarrow E_2$, $g : E_2 \rightarrow E_3$ two given functions. If f is continuous at $a \in E_1$ and g is continuous at $b = f(a) \in E_2$, then $g \circ f$ is continuous at $a \in E_1$.

Proof. □

Definition 2.3. If $f : E_1 \rightarrow E_2$ is continuous and bijective, then it is called a *homeomorphism*. The topological space E_1 is said to be *homeomorphic* to E_2 if such a function exists.

2.2 Limits

Definition 2.4. Let (E_1, τ_1) and (E_2, τ_2) be topological spaces. Let $f : X \subseteq E_1 \rightarrow E_2$ be a function, let $a \in \overline{X}$ and $\lambda \in E_2$.

- We say that $f(x)$ tends to λ when x tends to a , if:

$$\forall O_2 \in \tau_2, \lambda \in O_2 : \exists O_1 \in \tau_1, a \in O_1 : x \in (O_1 \cap X) \Rightarrow f(x) \in O_2.$$

- If (E_2, τ_2) is a Hausdorff space, then λ is said to be the *limit* of $f(x)$ as x tends to a , and write:

$$\lim_{x \rightarrow a} f(x) = \lambda.$$

- Let $D \subseteq X$, suppose $a \in \overline{D}$. We say that $f(x)$ tends to λ when x tends to a by elements of D if f restricted to D tends to λ when x tends to a . In this case we write:

$$\lambda = \lim_{x \rightarrow a, x \in D} f(x), \text{ which is logically equivalent to:}$$

$$\forall O_2 \in \tau_2, \lambda \in O_2 : \exists O_1 \in \tau_1, a \in O_1 : x \in (O_1 \cap D) \Rightarrow f(x) \in O_2.$$

2.3 Sequences

Definition 2.5. Let $(x_n)_{n \in \mathbb{N}}$ be a sequence defined on a topological space (E, τ) . The sequence (x_n) is said to converge to $a \in E$ as n tends to $+\infty$, written as $x_n \rightarrow a$ (meaning $a = \lim_{n \rightarrow +\infty} x_n$), if:

$$\forall O \in \tau, a \in O : \exists p \in \mathbb{N}, n \geq p \Rightarrow x_n \in O.$$

Theorem 2.3. Let (E, τ) a topological space, $X \subseteq E$, and $a \in E$.

3 Metric Spaces

3.1 Metrics

Definition 3.1. A function $d : E \times E \rightarrow \mathbb{R}$ is said to be a *metric* on a set E if, for any $x, y, z \in E$:

1. $d(x, y) \geq 0$;
2. $d(x, y) = 0 \Leftrightarrow x = y$;
3. $d(x, y) = d(y, x)$;
4. $d(x, y) \leq d(x, z) + d(z, y)$.

The pair (E, d) is called a *metric space* if d is a metric on E .

TO-DO: Insert Examples

3.2 The Topology of a Metric Space

Given any metric space (E, d) it is possible to define a topology on it. Consider the sets

$$\begin{aligned} B_{op}(a; R) &:= \{x \in E : d(x, a) < R\}, \\ B_{cl}(a; R) &:= \{x \in E : d(x, a) \leq R\} \end{aligned}$$

The former is called the *open ball* (the latter, *closed ball*) with center a and radius R . We also define the *sphere* centered at a with radius R to be the set

$$S(a; R) := \{x \in E : d(x, a) = R\}.$$

By convention $B(a; R) = B_{op}(a; R)$.

The geometric motivation for these definitions becomes clear once the \mathbb{R}^2 case is considered.

TO-DO: Insert pictures

From these definitions we may easily deduce that:

$$\begin{aligned} B_{cl}(a; R) &= B_{op}(a; R) \cup S(a; R); \\ B_{op}(a; R) &\subset B_{cl}(a; R); \\ R < R' &\Rightarrow B_{cl}(a; R) \subset B_{op}(a; R'). \end{aligned}$$

Definition 3.2. Let (E, d) be a metric space and $X \subset E$. The set X is said to be *bounded* if $\exists a \in E$ and $\exists R \in \mathbb{R}^+$, such that $X \subseteq B_{op}(a; R)$.

Analogously, a function $f : I \rightarrow E$ (with I an arbitrary set) is bounded if $f(I)$ is a bounded set. In particular, a sequence $(x_n)_{n \in \mathbb{N}}$ on E is bounded if:

$$\exists a \in E, \exists R \in \mathbb{R}^+ : \forall n \in \mathbb{N} : d(x_n, a) \leq R.$$

Definition 3.3. Let (E, d) be a metric space. Consider the following set:

$$\tau_d := \{O \subset E : \forall a \in O, \exists R \in \mathbb{R}^+ : B(a; R) \subset O\}.$$

This set will be called the *generated topology by the metric d on E* .

Theorem 3.1. *Every metric space is Hausdorff.*

Apply concepts from topology to derive a "definition" of convergence for metric spaces.

Definition 3.4. Let d_1 and d_2 be two metrics on E ; they are said to be *equivalent* if $\tau_{d_1} = \tau_{d_2}$.

Proposition 3.2. *Let d_1, d_2 be metrics on E . If there exists $\alpha, \beta \in \mathbb{R}^+$ such that, for every $x, y \in E$:*

$$\alpha d_1(x, y) \leq d_2(x, y) \leq \beta d_1(x, y)$$

3.3 Continuity

Definition 3.5. continuity at a point

Definition 3.6. Continuous on a set

Definition 3.7. Uniformly continuous

Definition 3.8. lipschitzian

Definition 3.9. Lipschitz's constant

Definition 3.10. function is a contraction

Theorem 3.3. *Let I be a non-degenerate interval of \mathbb{R} and let $f : I \rightarrow \mathbb{R}$ be a continuous function on I and differentiable on (I) . Then f is lipschitzian, if and only if, f' is bounded. In this case $L_f = \sup_{y \in \text{int}(I)} |f'(y)|$.*