

Conformational Analysis

Ingyu Bahng

May 2026

Contents

1	Rotamers	2
2	Cycloalkane & Ring Strain	4

1 Rotamers

Core Concept 1.1: Rotamers

Rotamers, or rotational isomers, represent a specific class of conformational isomers characterized by distinct spatial arrangements arising exclusively from rotation about single (σ) bonds. Consistent with the fundamental principle established throughout Thermodynamics—that matter adopts configurations minimizing potential energy (PE)—molecular conformations similarly favor energetically favorable arrangements.

Core Concept 1.2: Rotamer Example (Ethane & Propane)

The conformational behavior of ethane can be effectively visualized through **Newman projections**, which reveal distinct rotameric states.




Figure 1: Newman projections of ethane

Rotation about the C-C σ bond generates two principal conformations: **eclipsed** and **staggered**. The staggered conformation represents the global energy minimum due to minimized $e^- - e^-$ repulsions between C-H σ bonds. Conversely, the eclipsed conformation constitutes a local energy maximum, characterized by maximized $e^- - e^-$ repulsions between adjacent C-H bonds. The PE profile for complete rotation about the C-C σ bond is depicted below.




Figure 2: Potential energy diagram of ethane Newman projections

The barrier to rotation ($3.0 \text{ kcal mol}^{-1}$) represents the activation energy (E_a) required for interconversion between staggered and eclipsed conformations. Notably, all staggered conformations of ethane are energetically degenerate, resulting in no net energy change upon rotation—the process is neither endothermic nor exothermic. This general pattern persists in propane, with a predictable modification.




Figure 3: Potential energy diagram of propane Newman projections

The rotational barrier in propane increases to $3.4 \text{ kcal mol}^{-1}$, reflecting the enhanced e^- density of the methyl substituent relative to hydrogen in ethane. This increase in electron density amplifies e^- - e^- repulsive interactions, thereby elevating the energy barrier.

Core Concept 1.3: Rotamer Example (Butane)

Butane exhibits a more complex conformational landscape due to the presence of two methyl groups.




Figure 4: Newman projections of butane

Applying principles derived from propane's conformational analysis, the **anti** staggered conformation achieves maximum stability by positioning methyl groups at the greatest possible separation ($\theta = 180^\circ$). The **gauche** staggered conformation ($\theta = 60^\circ, 300^\circ$) represents the next most stable arrangement. Among eclipsed conformations, the fully eclipsed state ($\theta = 0^\circ$) exhibits minimum stability due to direct overlap of electron-dense methyl groups, while the half-eclipsed conformation ($\theta = 120^\circ, 240^\circ$) shows intermediate instability.




Figure 5: Potential energy diagram of butane Newman projections

Core Concept 1.4: Stereoisomeric Nomenclature

Stereoisomers are molecules with identical connectivity but differing three-dimensional spatial arrangements. This broad classification encompasses the following two categories.

- Conformational isomers** arise from bond rotations that occur freely at room temperature, with rotamers serving as the archetypal example. These isomers interconvert rapidly without bond cleavage.
- Configurational isomers**, in contrast, possess permanent stereochemical arrangements that can only be altered through bond-breaking and bond-forming processes. This category includes geometric isomers (*cis/trans*) and enantiomers (*R/S* designations).

2 Cycloalkane & Ring Strain

Core Concept 2.1: Cycloalkane Ring Strain

Cycloalkanes exhibit ring strain (or *elevated potential energy*) arising from the following three factors.

- Bond angle strain**, particularly pronounced in smaller rings. Carbon atoms preferentially adopt sp^3 hybridization with ideal tetrahedral bond angles of 109.5° . However, geometric constraints imposed by ring closure force deviations from this optimal geometry.

For instance, cyclopropane exhibits internal bond angles of approximately 60° , representing a 49.5° compression from the ideal value. This substantial angular distortion destabilizes the molecule by increasing orbital overlap inefficiency and raising the overall potential energy.

- (b) **Eclipsing strain** results from unfavorable torsional interactions between adjacent C–H bonds. As seen in Core Concept 1.2, eclipsed conformations maximize $e^- - e^-$ repulsions between proximal σ bonds, elevating potential energy relative to staggered arrangements.
- (c) **Transannular strain** is characterized by steric repulsions between atoms positioned across the ring interior. This strain component becomes significant in medium-sized rings ($C_8 - C_{11}$) where the ring cavity is sufficiently small that non-bonded atoms on opposite sides of the ring approach at shorter distances.

These strain factors collectively drive cycloalkanes to adopt **non-planar conformations** that deviate from the intuitively expected flat ring geometry.




Figure 6: Non-planar cycloalkane conformations

Core Concept 2.2: Cyclohexane Conformational Analysis

Cyclohexane preferentially adopts the chair conformation (Figure 7) as its global energy minimum primarily due to the following reasons.




Figure 7: Cyclohexane chair conformation

- (a) **Minimization of eclipsing strain:** The chair conformation's stability comes from the staggered orientation observed along all C–C bonds. Newman projection analysis reveals that each C–C bond maintains a **staggered** arrangement, thereby eliminating the eclipsing strain felt in planar conformations.




Figure 8: Newman projection analysis of cyclohexane

- (b) **Minimization of transannular strain:** The chair conformation additionally minimizes steric interactions across the ring. In contrast, the **boat conformation** (Figure 10) suffers from both heightened transannular strain and increased eclipsing strain.

As a result, the boat conformation represents instead a **transition state** in cyclohexane ring inversion and consequently possesses negligible presence at equilibrium. On the other hand, the **twist (skew) boat conformation**—a geometrically relaxed intermediate—exhibits lower energy while retaining higher energy than the chair form (Figure 10).




Figure 9: Cyclohexane twist (skew) boat conformation

Similarly to the boat conformation, the **half-chair conformation**—an additional high-energy transition state—arises from analogous strain considerations during the chair-to-chair interconversion pathway as shown in Figure 10.




Figure 10: Cyclohexane conformation potential energy diagram

Core Concept 2.3: Monosubstituted Cyclohexanes

Whereas unsubstituted cyclohexane exhibits energetic degeneracy between chair conformations, monosubstituted derivatives (*e.g.*, methylcyclohexane) demonstrate conformational preference driven by differential transannular strain.




Figure 11: Methylcyclohexane chair conformations

Specifically, the **equatorial** methyl conformer predominates over the **axial** methyl conformer due to reduced 1,3-diaxial interactions as shown in Figure 11.

Core Concept 2.4: Axial vs Equatorial

Cyclohexane substituents occupy one of two geometrically distinct positions, each with characteristic spatial orientations and steric consequences.

- Axial** bonds orient perpendicular to the approximate plane of the cyclohexane ring, extending parallel to the ring's pseudo-axis.
- Equatorial** bonds, on the other hand, radiate outward approximately within the ring's equatorial plane, directed toward the ring periphery.




Figure 12: Axial (A) and equatorial (E) bonds within cyclohexane