

# Ethers

Ingyu Bahng

June 2026

## Contents

|   |                                       |   |
|---|---------------------------------------|---|
| 1 | Ether Synthesis                       | 2 |
| 2 | Ether Reactions                       | 5 |
| 3 | Sulfur Analogs: <i>A Brief Lesson</i> | 7 |

# 1 Ether Synthesis

## Core Concept 1.1: Ethers

Ethers are a class of organic compounds characterized by an oxygen bonded to two alkyl or aryl groups. Their general formula is  $R-O-R'$ , where  $R = R'$  (symmetrical ether) or  $R \neq R'$  (asymmetrical ether).





Figure 1: Dimethyl ether structure

## Core Concept 1.2: Williamson Ether Synthesis

**Williamson ether synthesis** is the primary way chemists make asymmetrical ethers. This method uses an alkoxide  $R'O^-$  as its nucleophile, thus works best when attacking non- $\beta$ -branched primary  $R-X$ s because  $\beta$ -branched, secondary, or tertiary  $R-X$ s will unintentionally use the E2 mechanism.



It's important to note that if the desired product is asymmetrical, the choice of corresponding R groups of the haloalkane and alkoxide is important because a tertiary or secondary haloalkane paired with a primary alkoxide will lead to E2 whereas the opposite combination will undergo  $S_N2$  (*the desired method*).




Figure 2: Williamson ether retrosynthesis; *unsymmetrical ethers have two choices, one is usually favorable over the other*

## Core Concept 1.3: Alcoholysis of $R_{\text{sec/tert}}X$

For  $R_{\text{sec/tert}}X$  haloalkanes, chemists use the protonated version of alkoxides (alcohol) to make the nucleophile less basic and consequently undergo  $S_N1$  or E1 rather than E2.




Figure 3: Alcoholysis of  $R_{\text{sec/tert}}X$

While E1 reactions still remain an unintended pathway, chemists minimize this by carrying out the reaction in lower temperatures.

#### Core Concept 1.4: Intramolecular Williamson Ether Synthesis (Cyclic Ether Formation)

So far, we've looked at only **intermolecular** ether synthesis reactions. **Intramolecular** ether synthesis (or *cyclic ether formation*) works in the same way as the intermolecular equivalent does.




Figure 4: Intramolecular Williamson ether synthesis

**Intramolecular are particularly favorable over intermolecular ether synthesis** because [1] it does not lower entropy and [2] highly diluted concentrations of the relevant alcohol do not effect it as much because the reaction is occurring with itself.

Mechanistically, this reaction occurs via a backside attack with inversion, prepared by the rearrangement of a less stable rotamer if needed.




Figure 5: Backside attack and inversion in cyclic ether formation

The relative rates of ring closure depend on the trade off between  $\Delta H$  (*ring strain*) and  $\Delta S$  (*lining up*) in the various transition states. The following is the ranking of ether ring formation speeds by ring size.

$$3 > 5 \geq 6 > 4 \geq 7 > 8$$

Reaction speed does not directly correspond to ring size. Although one might expect a linear relationship, the trend is more nuanced: 3-membered rings react fastest owing to the close proximity of the bonding atoms, 4-membered rings are the slowest, and 6-membered rings are surprisingly fast again due to favorable atomic geometry. In summary, this ordering stems from competing  $\Delta H$  and  $\Delta S$  contributions across the transition states of each ring size, rather than any simple geometric trend.






Figure 6: Relative rates of ring closure

**Core Concept 1.5: Symmetrical Ether Synthesis via Alcohols**


Ethers can also be formed by **alcohols and  $H^+$**  through  $S_N2$  and/or  $S_N1$  depending on the R groups. Chemists specifically use this technique to form symmetrical ethers because an alcohol mixture would otherwise produce a slurry of different ether products.



For  $R_{\text{prim}}-OH$ , this mechanism works via  $S_N2$  due to minimal steric hindrance.

Figure 7: Symmetrical ether formation via  $S_N2$ 

When dealing with  $R_{\text{sec/tert}}-OH$ , this mechanism shifts to  $S_N1$ .

Figure 8: Symmetrical ether formation via  $S_N1$ **Core Concept 1.6: Asymmetrical Ether Synthesis (Tetrabutyl Scenario)**

However, a unique alcohol combination that preferentially forms an asymmetrical ether is an **unhindered ROH and tetrabutyl mixture**.




Figure 9: Asymmetrical ether formation with t-Butyl group

The ROH preferentially attacks readily available t-Bu cations (due to hyperconjugation) via  $S_N1$ . And while the t-Bu cation could be attacked by another t-Bu alcohol molecule, steric hindrance makes this interaction less likely to occur compared to the smaller, less sterically hindered ROH.

In the laboratory, t-Bu is used for "**protection**" of other alcohols by forming these t-Bu ethers.

## 2 Ether Reactions

Let's go through a collection of different ether reactions.

### Core Concept 2.1: $R_{\text{prim}}-O-R_{\text{prim}}$ Reaction

$R_{\text{prim}}-O-R_{\text{prim}}$  are quite unreactive, exhibiting stability to bases, RLi, RMgBr, and dilute aqueous  $H^+$ . However they are reactive under  $H^+X^-$  exposure.






Figure 10: Ether cleavage with hydrohalic acids

### Core Concept 2.2: $R_{\text{prim}}-O-R_{\text{sec}}$ Reaction

$R_{\text{prim}}-O-R_{\text{sec}}$  ethers react either through  $S_N1$  or  $S_N2$  depending on the nucleophile strength (under acidic conditions). When paired with a **good nucleophile**, the attack occurs on the  $R_{\text{prim}}$  side carbon via  $S_N2$  to avoid the steric hindrance of  $R_{\text{sec}}$ .

Figure 11: Mixed ether reaction with good nucleophile ( $S_N2$ )

When paired with a **poor nucleophile**, an  $S_N1$  reaction occurs at the carbocation of  $R_{\text{sec}}$  (stabilized by hyperconjugation).

Figure 12: Mixed ether reaction with poor nucleophile ( $S_N1$ )

### Core Concept 2.3: t-Bu Ether Hydrolysis

t-Bu ether hydrolysis is the reverse of t-Bu ether formation covered in Core Concept 1.6. This reverse process is the "deprotection" of alcohol ROH by the t-Bu cation via acid. The resulting t-Bu cation, when exposed to heat, undergoes E1 to exaporate into an alkene.




Figure 13: Hydrolysis and deprotection of t-Butyl ethers

### Core Concept 2.4: Ether Ring Opening

Ether ring openings release ring strain by a  $Nu^-$  attacking directly one of the two carbons adjacent to the oxygen. This is a regioselective process, meaning  $S_N2$  occurs at the less hindered relevant carbon.




Figure 14: Regioselective ether ring opening

Many  $Nu^-$  will work. The following example shows that ether ring openings are regioselective and also **stereocontrolled**.




Figure 15: Stereocontrolled ether ring opening

With **neutral nucleophiles**, acidic conditions are required to activate the ether toward nucleophilic attack. Without  $H^+$ , the reaction will not occur.




Figure 16: Acid-activated ether ring opening with neutral nucleophile

One important distinction in the neutral nucleophile mechanism is regioselectivity towards the more hindered carbon. In asymmetrical ethers, the nucleophile preferentially attacks the more substituted carbon, as it better stabilizes the partial positive charge ( $\delta^+$ ) that develops through hyperconjugation and inductive effects—causing the C-O bond on that side to stretch and weaken. This gives the mechanism a hybrid  $S_N1 / S_N2$  character: it resembles  $S_N2$  in that nucleophile attack and leaving group departure occur simultaneously, yet resembles  $S_N1$  in that the partial positive charge mimics a carbocation.




Figure 17: Regioselectivity of acid-catalyzed ring opening of asymmetrical ethers

### 3 Sulfur Analogs: A Brief Lesson

#### Core Concept 3.1: Sulfur Analogs

Sulfur is directly below oxygen in the periodic table, and thus behaves very similarly.


Thiols ( $RSH$ ) are **more acidic than alcohols** because the  $S-H$  bond is weaker due to sulfur's higher-energy molecular orbitals. The lower electronegativity difference in  $S-H$  compared to  $O-H$  also results in **weaker hydrogen bonding**, making thiols less polar overall. *This is illustrated by  $H_2S$ , which is a gas at  $25^\circ C$  with a boiling point of  $-60^\circ C$ , in stark contrast to  $H_2O$ .*

Despite being less basic than  $RO^-$ ,  $RS^-$  is a stronger nucleophile due to its greater **polarizability**—note that polarizability is distinct from polarity. Polarizability describes how easily an electron cloud can be distorted by an external force (Core Concept 2.4). Larger atoms like sulfur are more polarizable than oxygen because their diffuse, loosely-held electron clouds are more easily deformed. This allows  $RS^-$  to stretch and distort toward an electrophile, enhancing its nucleophilicity despite its lower basicity.

placeholder link

#### Core Concept 3.2: Sulfur Reactivity

The better nucleophilicity of  $RS^-$  eliminates the issue of  $E2$  reactions when paired with  $R_{sec}X$  like  $RO^-$  has.

Figure 18: Sulfur versus oxygen reaction with  $R_{sec}X$ 

Even neutral  $RSR'$  readily undergoes  $S_N2$ , whereas dimethyl ether ( $CH_3OCH_3$ ) does not react under the same conditions. When  $RSR'$  attacks an  $RX$ , the resulting sulfonium ion ( $R_3S^+$ ) becomes susceptible to attack by another nucleophile — the sulfide, once a good nucleophile, has now become a good leaving group, analogous to  $H_2O$ . This makes **sulfonium salts** ( $R_3S^+$ ) effective **alkylating agents**, meaning they can transfer an alkyl group to an incoming nucleophile.




Figure 19: Sulfide acting as a nucleophile to form a sulfonium ion (top), which then acts as an alkylating agent toward an incoming nucleophile (bottom)