

Conjugated Systems & Aromaticity

Ingyu Bahng

July 2026

Contents

1	Conjugated Systems	2
1.1	Organic Acids & Bases Review	2
1.2	Allylic Systems	4
1.3	Conjugated π Systems	5
2	Diel Alder Reactions	7
2.1	Rates of Diel Alder Reactions	8
3	Aromaticity & Aromatic Substitution	10
3.1	Aromatic Compounds	10
3.2	Electrophilic Aromatic Substitution	12
3.2.1	Reaction Types	12
3.2.2	Directing Groups	14
3.2.3	Aromatic Substituent Manipulations	16
3.3	Nucleophilic Aromatic Substitution	17

1 Conjugated Systems

1.1 Organic Acids & Bases Review

Core Concept 1.1: General Organic Compound Acidity

Committing the following pK_a values to memory as a baseline reference is essential, as they serve as structural anchors from which the acidities of related but distinct compounds can be systematically estimated by comparing functional group similarities, electron-withdrawing effects, and resonance stabilization of the conjugate base.












Compound	pK_a	Structure	Compound	pK_a	Structure
Alkane	50		Ammonium	10	
Alkene	45		α to Two Carbonyls	10	
Allylic	40		Carboxylic Acid	5	
Amine	35		Oxonium	0	
Alkyne	25		Trifluoro Acetic Acid	0	
α to an Ester	25		Hydrohalic Acid	<0	
α to a Ketone or Aldehyde	20		Sulfuric Acid	<0	
Alcohol (ROH)	15		Toluenesulfonic Acid	<0	
Phenol (PhOH)	10				

Table 1: Common acids and their pK_a values, ordered from least to most acidic

Core Concept 1.2: Acidity of Alkanes, Alkenes, and Alkynes

Let's take a look at the pK_a s of alkanes, alkenes, and alkynes.

alkane > alkene > alkyne

We can explain the observed pattern by going through the following factors

(a) **Electronegativity of atom attached to the acidic H**

This doesn't explain it because they are all carbons, so there is no difference

(b) **Inductive Effect**

This also doesn't explain it because only CHs exist nearby

(c) **Resonance**

Resonance is not possible, so no explanation

(d) **Charge**

All neutral, so no explanation

(e) **Hybridization**

There is a difference here, it is sp^3 vs sp^2 vs sp . The resulting lone pair e^- s on the relevant carbons as a result of dehydrogenation is stabilized best by the sp hybridized orbital because the higher percentage s characteristic allows e^- s to be held closer to the nucleus.

If you cannot explain acidity with the acid structure itself, then you look at conjugate bases

<figure of the conjugate bases>

Which is LEAST basic? alkyne conjugate base is the weakest conjugate base because the lone pair is least reactive and most stabilized. Why? $\rightarrow e^-$ s are gathered more to the opposite side of the bond so it attracts less H^+ in comparison to alkanes and alkenes. An energy diagram is shown below to easy visual comparison

<figure of energy diagram> caption: lone pair in $C(sp)$ orbital is most stabilized, thus least basic; increased s character of sp orbital increases probability of e^- s being closer to the positive charge of the nucleus

Core Concept 1.3: K_{eq} and pK_a s

Any given acid base chemical reaction has a $K_{eq} = 10(\text{conj Acid } pK_a - \text{acid } pK_a)$. Thus based on the pK_a values given in Table 1, we can determine the K_{eq} of any given acid and base reaction to a reasonable level of accuracy as shown in the following examples

(a) **chemical equation name placeholder**

<figure>

(b) **chemical equation name placeholder**

<figure>

(c) **chemical equation name placeholder**

<figure>

(d) **chemical equation name placeholder**

<figure>

Furthermore, pK_a values can also be predicted based on the reference pK_a s from Table 1, as shown in the following examples.

(a) placeholder

<figure>

explanation

(b) placeholder

<figure>

explanation

1.2 Allylic Systems

Core Concept 1.4: Allylic Carbons

Allylic carbons refer to sp^3 hybridized carbons attached to a C=C double bond. An allylic H refers to those that are attached to allylic carbons.

<figure>

Core Concept 1.5: Allylic System Resonance

An allylic hydrogen is even more acidic than an alkyne hydrogen due to resonance spreading out negative charge within the resulting allylic anion (making it a weaker base as a consequence).

<figure of the 2 reaction at the beginning showcasing the reaction within amine and then also the reaction below of resonance at the bottom of the same column in the rough notes>

Just as how negative charge (*filled octets*) is stabilized by resonance in allylic structures, radicals and positive charge (*unfilled octets*) are also stabilized by the same mechanism.

<figure>

The important thing to know is that if a lone pair can do resonance, it is in the p orbital.

Core Concept 1.6: Allylic System Bond Dissociation Energy

As a consequence of resonance induced stabilization, the bond dissociation energies of different hydrogens within an alkene follow a predictable pattern.

<triple reaction figure>

Above, the 3° carbon radical is lower in energy than the 3° carbon because of hyperconjugation between the radical carbon and neighboring σ bonds. The allylic, on the other hand, although it receives less hyperconjugation as a 2° carbon, it is able to perform resonance, making its bond dissociation consume the least amount of energy.

<energy diagram figure>

This leads to chemical consequences that play large roles in practice. Take a look at the reaction below for example.

<double reaction fig> explain things via the caption

Core Concept 1.7: Allylic System Molecular Orbital Structure

The relevant molecular orbitals of allylic structures are the p orbitals, which have the following energy levels in the comparative context of sp and sp² hybridized carbons as shown below. Notice that the energy level of sp² orbitals sits higher than that of sp orbitals.

<figure of sp vs sp² orbital energy levels>

The orbital structure on a skeletal diagram backbone would look like the following.

<figure> caption: σ bond orbitals overlap along internuclear axis (handshake) whereas π bond orbitals overlap above and below internuclear axes. All carbons are sp² hybridized

In the context of the entire allylic p orbital system, the energy diagram presents a unique 3 level pattern, going from bonding to non-bonding to anti-bonding.

<p orbital system figure>

The resulting HOMO-LUMOs of the allylic anion, radical, and cation types are as a result different.

<figure>

This is particularly relevant in the context of the **frontier MO theory**, which states that the nucleophile HOMO interacts with the electrophilic LUMO.

<figure>

1.3 Conjugated π Systems**Core Concept 1.8: Conjugated π System Stability**

A conjugated pi system refers to network of adjacent allylic structures that allow for resonance across a greater number of carbons. For instance, take a look at the hydrogenation enthalpies of the following alkenes.

<5 reactions figure>

Compound 5 is more stabilized than predicted based on intuitive addition of constituent enthalpies because of the ability of conjugated π systems to better stabilize excess or lacking charge.

<energy diagram figure>


When we look closer into the orbital energy diagrams of these conjugated π systems like we have done in REFERENCE TO 3 PI SYSTEM, we find that the conjugated π system orbital energy levels sit in an overall larger range but possesses smaller stepwise increases in energy, thus the HOMO LUMO gap is comparatively smaller compared to the non conjugated π system alkenes.

<pi orbital energy diagram figure> caption: source of stability in a conjugated π system are the lowest π MO e⁻s

The rules are the following.

- As conjugation increases (more p orbitals in a row), the energy gap between π MOs gets smaller
- As conjugation increases, the lowest energy π MO gets lower in energy and the highest energy π MO goes higher in energy

It's important to know that every high energy intermediate needs to be as stabilized as possible and that resonance from conjugated π systems trump stability originating from hyperconjugation.



This has particularly large consequences in the different reactions covered within Alkenes & Alkynes because many electrophilic additions progress in a Markovnikov or anti-Markovnikov manner, which takes into consideration intermediate state stability. Examples include alkene hydrohalogenation (*Core Concept 1.9*), which progresses in a Markovnikov manner, and hydroboration-oxidation (*Core Concept 1.13*), which progresses in an anti-Markovnikov manner.

Using our learned knowledge of conjugated π system stability, let's look at two examples.

Core Concept 1.9: Chlorination of Conjugated Cycloalkene

Our goal is to predict all of the products of the following reaction, including stereoisomers.

<figure of reaction>

First, we map out the chloronium intermediates.

<figure>

Use resonance to determine which carbon of halonium to attack with weak nucleophilic ROH.

<figure> caption: tertiary " σ bond resonance" vs 2° and allylic

Because the allylic side is more stable, the electrophilic carbons exist in two locations within the system

<figure> caption: the top resonance contributor is the Cl

Thus three possible attacks exist, giving us three distinct stereoisomeric products.

(a) **attack name placeholder**

<figure>

(b) **attack name placeholder**

<figure>

(c) **attack name placeholder**

<figure>

Core Concept 1.10: Radical Bromination of Terminal Alkene

The radical bromination of a terminal alkene works in a similar manner to the reaction shown in Core Concept 1.9, wherein resonance creates the possibility of multiple different types of attacks and consequently three distinct products.

We first create Br_2 through N-Bromo-succinimide, which allows for the production of **monobrominated products**. Realize that adding pure Br_2 to react with alkenes results in dihalogenation as mapped out in Core Concept 1.11.

<figure> caption: slow release of Br_2 to avoid alkene adding Br_2 directly

After the spontaneous dissociation of Br_2 into radical bromine atoms, the first propagation occurs, exposing a radical in the original alkene compound.

<figure of prop 1>

This radical alkene and the remaining bromine radical can carry out a second propagation interaction in three different manners.

(a) **propagation placeholder**

<figure>

(b) **propagation placeholder**

<figure>

(c) **propagation placeholder**

<figure>

To finalize, the full reaction looks like this.

<figure of full simplified reaction>

2 Diel Alder Reactions

Core Concept 2.1: Cycloadditions

Cycloadditions refer to 2 separate π systems reacting to form a new ring. There is no nucleophile nor an electrophile in this interaction.

We have already seen cycloadditions in two different reactions within Alkenes & Alkynes.

(a) **Ozonolysis** (*Core Concept 1.18*)

<figure>

(b) **OsO₄ Syn-Dihydroxylation** (*Core Concept 1.17*)

<figure>

Core Concept 2.2: Diel Alder Reactions

Diel alder reactions are a type of cycloaddition that combines a minimum 4π diene with a minimum 2π dienophile to form a cyclohexane.

<figure>

As such, diene and dienophile pairings that possess larger conjugated π systems than the stated minimums of 4π and 2π , respectively, can produce a massive diversity of stereoisomeric products

<figure>

Core Concept 2.3: Orbitals of Diel Alder Reactions

In diel alder reactions, π systems approach each other in a vertical manner to allow π systems to generate new σ overlaps

<figure> caption: p orbitals overlap to form σ interactions

Let's take a look at the energy diagrams of the respective conjugated π systems of the diene and dienophile.

<figure>

An important observation can be made in the context of which energy levels interact with which: HOMO of one reagent interacts with the LUMO of the other reagent.

<figure showing the two different homo lomo overlaps in diel alder reaction>

Core Concept 2.4: Diel Alder Stereochemistry

We can also categorize diel alder reactions by the different possible stereochemical approaches of the reagents.

- (a) **Top** and **bottom** diel alder reactions simply refer to when the dienophile approaches the diene from the **top** or **bottom**, respectively.

<figure>

- (b) **Endo** and **exo** distinctions refer to when the sterics of the dienophile are pointed **towards** or **away from** the corresponding diene

<figure>

Keep in mind that the bonds pointing towards the diene get flipped around the dienophile double bond in the direction opposite from where the diene approaches.

<figure>

As practice, consider the following diel alder reaction in which we need to predict all diel alder nondimer products.

<figure>

Two possible constant isomers can form from the two different dienophile positionings that can occur.

<figure of the products> caption: easier to think of the dienophile being flipped along a centered horizontal axis to produce the right product

And for each of these reactions, four different stereochemical approaches—endo top, endo bottom, exo top, exo bottom—exist.

- (a) **Isomer 1**

- **endo top**
- **endo bottom**
- **exo top**
- **exo bottom**

- (b) **Isomer 2**

- **endo top**
- **endo bottom**
- **exo top**
- **exo bottom**

2.1 Rates of Diel Alder Reactions

Core Concept 2.5: Diene Orientation

Diel alder reactions are more favorable when the resulting ring has minimal strain. The **s-cis** diene orientation does this better than the **s-trans** orientation as shown below. *The s in s-cis and s-trans stands for specifically along single bond as clarification because conventionally cis and trans in the context of alkenes look at the double bond as the reference bond.*

<figure of s-cis and s-trans> caption: a ring needs to be at least 8 atoms large to accommodate a trans π bond

Take the following dienes for instance.

<figure>

The relative diel alder reaction rates is $C > A > B$. C reigns superior because its structure restricts rotation along the relevant single bond, preventing any rearrangement to an s-trans arrangement. A and B, on the other hand, both can rotate along the relevant single bond, however B favors the s-trans arrangement because the s-cis arrangement has a comparatively higher level of steric hinderance between the alkenyl substituents to the relevant single bond.

Core Concept 2.6: Secondary Orbital Overlap

Under certain scenarios, orbitals not directly responsible for new bond formation between the diene and dienophile can undergo a secondary overlap to lower the overall E_a . Take a look at the following diel alder interaction between two cyclopentenenes.

<figure>

It's important to keep in mind that although favorable mixing of secondary orbital overlaps in the transition state will help stabilize that transition state, sterics still does not favor the endo approach. As a result, ambiguous diene dienophile pairs will usually need to undergo experiments to see which effect is more important in each case.

Core Concept 2.7: Electronic Effects

Diel alder reactions progress faster when the energies of the interacting orbitals are closer to one another. In other words, the small the HOMO-LUMO gap between reacting molecules, the faster the rate. This usually occurs when more conjugation occurs.

However, while more conjugation typically shrinks the HOMO-LUMO gap and speeds up a reaction, it also alters the wave topology (*the layout of the nodes*). If the orbital with the ideal energy gap happens to place a node on an essential bonding atom, that orbital becomes useless for the reaction, forcing the system to use a higher-energy, slower pathway. Symmetry and overlap must always be satisfied before energy gaps even matter.

<figure> caption: The left reagents actually go faster despite the small $\pi_3-\pi^*$ gap of the right reagents because π_3 is not favorable because there is a node of the atom that needs ot make a new σ bond

Core Concept 2.8: Groundstate Destabilization

Given that two sets of reagents that are similar will have similar transition states leading to the **same** product, the reactant pair in which the ground states of the initial diene and dienophile start at a higher energy levels (*due to something such as steric hinderance*) will have a faster diel alder reaction rate. Take the following diene dineophile pairs leading to the same product for instance.

<figure>

The energy level for the right diene dineophile pair possesses a greater level of steric hinderance between the isobutyl substituents wihtin the diene, placing it at a higher energy level relative to the left diene dienophile pair.

<figure>

As shown in the energy diagram above, the E_a is much lower for the more unstable reactant pair and thus progresses at a much faster rate because the total ΔH is lower.

3 Aromaticity & Aromatic Substitution

3.1 Aromatic Compounds

Core Concept 3.1: Aromatic Compounds

Name originates from these compounds' nice scent, but now it is a strictly structural definition covered in Core Concept 3.5.

While not limited to the base benzene structure, the extent of these notes will mostly refer to base benzene compounds such as the following.

<figure>

Core Concept 3.2: Aromatic Compound Stereochemistry

Additional substituent positioning to the the root aromatic compounds shown in FIG REFERENCE can be specified through the ortho, meta, and para keywords.

<figure>

The compound naming scheme follows a structure of *realtion-sub-root*. Take a look at the following examples.

<figure or ortho-nitro-aniline and para-sulfonate-tolvene and the three other examples>

Core Concept 3.3: Reactivity of Aromatic Compounds

Aromatic compounds are less reactive than predicted. For instance while a simple cyclohexene reacts with mCPBA to form a secondary ester ring, no reaction occurs when benzene and 3 mCPBA are mixed together.

<figure>

Similarly, a portion of the predicted energy drop during the hydrogenation of benzene is eliminated due to aromatic stabilization.

<figure>

In order to explain this, let's look at a top down view of the benzene molecular orbital diagram.

<figure>

explanation

Core Concept 3.4: Aromatic Compound Molecular Diagrams

The **Frost Diagram** is a method to quickly predict π molecular orbital energy levels for a cyclic conjugated system, following the following steps.

- (1) Compound must have acyclic array of p orbitals
- (2) Draw shape pointed down
- (3) Draw a circle around the shape
- (4) Intersections are energy levels
- (5) Middle of circle is nonbonding

(6) Add e^- s

Here are the respective orbital diagram examples for a pentene anion and a heptene cation. Benzene has already been shown in FIG REFERENCE ABOVE.

(a) **Pentene Anion**

<figure>

(b) **Heptene Cation**

<figure>

Core Concept 3.5: Aromaticity Rules

In order to a compound to be considered aromatic, it must follow a strict set of rules.

(1) Cyclic array of p orbitals

(2) e^- count in cyclic array fulfills the equation of $4n + 2$, where n is an integer (0, 1, 2, 3, 4, ...).

*An e^- count that fulfills the equation of $4n$ where n is an integer is much more reactive than predicted and is termed **anti-aromatic***

If the above criteria are not met, the compound is considered **nonaromatic**, such as the cyclohexene shown below.

<figure>

As practice, take a look at the follow two examples.

(a) **Pentene Anion**

<figure>

This is a cyclic array of p orbitals and the e^- count is 6, which fulfills the $4n + 2$ expression. Thus this compound is **considered aromatic**.

(b) **Pentene Cation**

<figure>

This compound is also a cyclic array of e^- s but its electron count is 4 which does not fulfill the expression $4n + 2$ but rather $4n$. Thus this compound is considered **anti-aromatic**.

Notice that the corresponding molecular orbital diagram results in two unpair e^- s, making it reactive.

<figure>

Core Concept 3.6: Aromatic Contributors

An **aromatic contributor** refers to a resonance structure of a molecule that possesses a fully conjugated, cyclic, planar π system satisfying the $4n + 2 \pi e^-$ count, even if the dominant or lowest-energy Lewis structure doesn't immediately look aromatic. Take the following molecule for instance.

<figure of the base non aromatic state to the left and the contributor state to the right>

Notice that while the base state is nonaromatic by definition because the conjugated π orbital system is not cyclic, an alternative state accessed through resonance is aromatic, making it an aromatic contributor.

3.2 Electrophilic Aromatic Substitution

Core Concept 3.7: Electrophilic Aromatic Substitution

Electrophilic aromatic substitution (EAS) refers to a reaction wherein an aromatic ring acts as a nucleophile. This reaction type follows the following general mechanism.

<figure of general mechanism>

3.2.1 Reaction Types

Core Concept 3.8: Halogenation

EAS halogenation effectively attaches a single halogen to an aromatic ring, generating the corresponding hydrohalogen as a byproduct. The general mechanism utilizes a **Lewis acid** (*any chemical species capable of accepting a pair of electrons*) as shown below.

<figure of general mechanism>

Analogously, here are examples showcasing chlorination and bromination of benzene.

(a) **Chlorination of Benzene**

<figure>

(b) **Bromination of Benzene**

<figure>

Core Concept 3.9: Nitration

Nitration effectively attaches an NO_2 nitro group onto an aromatic ring, acting primarily as a synthetic stepping stone in practice. First, let's look at the generation of an NO_2^+ electrophile through the exposure of nitric acid to sulfuric acid.

<figure>

With this in mind, the general nitration mechanism to a benzene follows the outline reaction mapped out in Core Concept 3.7.

<figure>

Core Concept 3.10: Sulfonation

Sulfonation entails the attachment of a SO_3H group onto an aromatic ring. The first step is a reaction between two H_2SO_4 molecules, a consequence of one of the reagents being **fumming H_2SO_4** .

<figure>

Then follows the same general EAS reaction mechanism covered in Core Concept 3.7.

<figure>

Core Concept 3.11: Alkylation

Alkylation is the process of adding an alkyl group onto an aromatic structure. In order for an alkane to become electrophilic, it must undergo a preparatory conversion to a carbocation state through the following reaction.

(a) **placeholder**

<figure>

This reaction is similar to the initial reaction shown in the chlorination pathway within Core Concept 3.8.

(b) **placeholder**

<BF₃ figure>

(c) **placeholder**

<figure>

(d) **placeholder**

<figure>

In certain instances, carbocation rearrangements can occur when the directly resulting carbocation is unstable.

<figure + mechanism>

Subsequently, these carbocation alkanes then proceed to perform EAS to aromatic structures.

<figure>

In the specific case above showcasing carbocation rearrangement of a cyclobutane to a cyclopentane, we can prevent rearrangement via a stepwise reaction pathway of **acylation** (Core Concept 3.12) and then reduction with Zn(Hg) and HCl.

(a) **Acylation**

<figure>

(b) **Reduction**

<figure>

(c)

Core Concept 3.12: Acylation

Acylation is the process of adding an acyl halide compound to an aromatic structure. We first need to make the acyl halide electrophilic through a lewis acid as shown in the following general reaction outline.

<figure of general acyl halid with lewis acid>

The resulting RCO⁺ molecule can then perform EAS with an aromatic ring. A full reaction using real compounds is shown below.

<figure of full reaction with FeCl₃>

3.2.2 Directing Groups

Core Concept 3.13: Directing Groups

So far, we have covered simply the EAS reaction types in Subsubsection 3.2.1 namely using benzene as our sole relevant aromatic structure. However, in scenarios where the aromatic ring is asymmetrical, chemists need a method to determine which carbon they want the EAS to occur on. Now we introduce **directing groups**, functional groups already attached to a benzene ring that dictate where (O, M, P) a new incoming electrophile will land.

<figure of the different directions on a benzene>

A summary table is provided below, but visit each Core Concept for a more detailed look into each directing group type.

Directing Group Type	Examples	Description
(1) Inductive O/P Director	$R = \text{alkyl [Me, Et, } i\text{Pr, ...]}$	Alkyl groups donate electron density inductively, directing substitution to the ortho and para positions.
(2) Inductive M Director	$R = \text{carbon w/ EN atoms [CCl}_3, \text{CF}_3, \text{...}]}$	Carbons bearing electronegative atoms withdraw electron density inductively, directing substitution to the meta position.
(3) Resonance O/P Director	i) R has lone pair adjacent to ring: [OMe, SR, NMe ₂] ii) $R = \text{carbon w/ } \pi \text{ system: [alkene, alkyne, benzene]}$	Substituents that donate electron density via resonance into the ring direct substitution to the ortho and para positions.
(4) Resonance M Director	[C=O, C $\equiv$ N, NO ₂]	Substituents that withdraw electron density via resonance from the ring accept electron density and direct substitution to the meta position.

Table 2: Classification of electrophilic aromatic substitution directing groups

Core Concept 3.14: Inductive O/P Directors

Inductive O/P directors are substituents that direct any EAS to the O and P positions within an aromatic ring.

<figure of first general reaction>

The substituents are generally alkyls, and the mechanism behind why they favor O/P over M positioning is due to the general stability of the different resonance structures that occur in each positioning scenario.

<triple figure>

As shown above, while nearly all of the different possible resonance structures place the transient carbocation charge on a 2° allylic position, O/P positioning allows for a 3° allylic carbocation position within 1 out of 3 resonance structures for each whereas M positioning does not. As such the energy levels of the resulting carbocation intermediaries follow the general ordering shown in the energy diagram below.

<energy diagram figure>

Core Concept 3.15: Inductive M Directors

Inductive M directors are substituents that direct any EAS to the M position within an aromatic ring. The substituents are carbons bearing highly electronegative atoms, making the carbon highly partially positive.

<figure>

The M position is favored because the O/P 3° allylic carbocation resonance structures have possess extra unstable repulsion between the carbocation and the partial positive carbon within the substituent. The M positioning, on the other hand, possess for a resonance structure that allows for maximal distance between the positive substituent carbon and the carbocation charge on the opposite side of the aromatic ring.

<figure>

Consequently, the resulting energy levels put O/P positioning energy above that of M positioning.

<energy diagram figure>

Core Concept 3.16: Resonance O/P Directors

Resonance O/P directors are substituents that direct any EAS to the O and P positions within an aromatic ring through donating e^- density. For instance, phenol possess resonance structures that place a carboanion negative charge on exclusively the O and P positions.

<figure>

These types of substituents specifically need to be lone pair adjacent to the aromatic ring. Another substituent type that allows for e^- donation is carbon with conjugated π systems. Take the following alkene for instance.

<figure>

Notice that the primary carbocation is able to form because it is allylic.

Here is a side note regarding the method by which inductive directors induce favorability of EAS positioning in comparison to resonance directors -> resonance directors affect the nucleophilicity capabilities of the relevant positions whereas the inductive versions manipulate favorability through transition state stability.

Core Concept 3.17: Resonance M Directors

Resonance M directors are substituents that direct any EAS to the M position within an aromatic ring through withdrawing e^- density. For instance, carbonyl substituents have the ability to withdraw e^- density in a manner that results in subsequent resonance structures to place a carbocation positive charge on the O/P positions, making these locations effectively less nucleophilic and the P position relatively more nucleophilic.

<figure>

As a result, when an electrophile approaches the transient aromatic ring, the nucleophilic attack occurs most favorably from the M position rather than the O/P position.

<figure>

Core Concept 3.18: Halogens as Directing Groups

Halogens can also act as directing groups, and based on data, they act as O/P directors. To understand the underlying basis of this observation, let's analyze monohalobenzene in the context of the inductive effect and

resonance.

<figure>

(a) **Inductive Effect**

<figure>

Under an O/P directing EAS interaction, inductive analysis would actually discourage O/P directing because the subsequent carbocation transition state places a carbocation that already has lowered e^- density even lower by residing next to the highly electronegative halogen atom. As a result, the dipole moment created destabilizes the structure.

(b) **Resonance**

<figure>

Resonance, on the other hand, allows the halogen to donate e^- density in order to fill all octet requirements throughout the entire molecule, which favors the O/P directing EAS interaction.

Although these two effects contradict one another, we can confidently say that the effects of resonance trump the inductive effect because real data in practice results in the O/P directing as favored.

3.2.3 Aromatic Substituent Manipulations

Core Concept 3.19: Reduction of Acyl Ketone

The reduction of an acyl ketone is a reduction reaction that was previously used in Core Concept 3.11 to replace a carbonyl oxygen with two hydrogen bonds.

<figure>

No mechanism is provided for this reaction.

Core Concept 3.20: Nitrobenzene Reduction to Aniline

The reduction of nitrobenzene work similarly to the reduction of acyl ketone (Core Concept 3.19) wherein oxygens are replaced by hydrogens. This reaction uses the same reactants to carry this replacement out.

<figure>

No mechanism is provided for this reaction.

Core Concept 3.21: Diazotization of Aniline

The diazotization of aniline is a potential subsequent reaction to nitrobenzene reduction as the product of the latter is the reactant of the former.

<figure>

The reaction mechanism occurs as follows, first starting with the conversion of NO_2^- to a nitrosyl cation.

<figure>

The highly reactive nitrozy cation then goes onto interact with the aniline in the following manner to produce **aryl diazonium**.

<figure>

3.3 Nucleophilic Aromatic Substitution

Core Concept 3.22: Nucleophilic Aromatic Substitution

Nucleophilic aromatic substitution (NAS or S_NAr) is an interaction in which the aromatic ring is the electrophile. NAS comes in two "flavors": **diazonium** (Core Concept 3.23) or **addition/elimination** (Core Concept 3.24) chemistry.

Recall that in electrophilic aromatic substitution (Subsection 3.2), the aromatic ring is the nucleophile.

Core Concept 3.23: Diazonium Chemistry

Diazonium chemistry begins by N_2 leaving a phenyldiazonium molecule—via breaking the bond between the σ $C(sp^2)-N(sp)$ bond—to produce a carbocation possessing benzene ring and an extremely stable N_2 gas molecule.

<figure>

It's important to notice that resonance is not possible in this specific carbocation state because the orbital that is empty is the $C(sp^2)$ orbital. We would need an unbalanced p orbital for resonance to occur. As a result, this phenyl cation is highly reactive and hungry for any nearby nucleophile to help stabilize its positive charge.

<figure>

As shown above, nearly all covered nucleophiles work in this interaction **with the exception of grignards and lithiates**. *The reasoning behind this will not be covered in the scope of these notes.*

Combining this diazonium chemistry with previously learned reactions, chemists can effectively attach nearly any nucleophile to a benzene molecule using the following reaction pathway.

<figure> caption: reference the core concepts for the reactions within the caption

Core Concept 3.24: Addition/Elimination

The NAS **addition/elimination** reaction occurs in the following general manner.

<figure>

Recall that we have seen resonance acceptors in resonance M directors (Core Concept 3.17), however, these resonance acceptors are not for the purpose of directing meta in the context of NAS chemistry. Their main purpose is to be able to accept e^- density after the nucleophile attacks the carbon with the leaving group.

Additionally, this reaction requires strong nucleophiles because it ultimately needs to break aromaticity as shown in the mechanism below. It should be noted that this step—the addition—is consequently the rate-determining step.

<figure>

As the mechanism showcases, the reaction is simply a relay of e^- density to the resonance acceptor, which then bounces back the e^- density in order to kick out the leaving group in the end. However, since the outcome of this interaction is ultimately the same as that of an S_N2 reaction, one may wonder why S_N2 is not a viable pathway.

Revisiting the S_N2 mechanism, recall that, in the lens of molecular orbitals, the nucleophile interacts with the σ^* anti-bonding orbital between the electrophile and leaving group in order to kick out the leaving group and create a bond simultaneously via a backside attack.

<figure>

In the aromatic ring scenario, this necessary σ^* interaction is extremely unlikely due to the physical barrier to entry because of the ring itself.

In the context of the frontier molecular orbital theory, the π^* orbital is lower in energy than σ^* , making it the LUMO and a much more accessible orbital to interact with, which is what happens in addition/elimination, the favorable pathway.

Let's go through two examples.

(a) Example 1: Predicting Major Kinetic Product

Consider the following reaction consisting of two possible leaving groups F and Cl with a NO_2 resonance acceptor.

<figure>

Both addition/elimination pathways are possible as shown below, ultimately leading to two distinct products.

<figure>

Since the nucleophile is held constant across both reaction pathways, the ability of the nucleophile to break aromaticity is not a factor. Instead, it is the greater dipole moment created by the more electronegative fluorine over the chlorine. Due to the greater partial positive charge of the fluorine carbon, MeO^- favors attacking that carbon because it is more electrophilic. Thus the reaction favors the substitution of the F over the Cl.

<figure>

(b) Example 2: Predicting Product of NAS Chemistry

Consider the following addition/elimination reaction below.

<figure>

In this case, simply only one of the leaving group substitutions work to donate e^- density into the resonance acceptor and the other doesn't.

<figure>

Thus the exclusive reaction outcome maps out as follows.

<figure>

(c) Example 3: Predicting Starting Material

Consider having to determine the starting aromatic compound to the following products based on the given nucleophile.

<figure>

Because we see two Cl^- and two $\text{H}_3\text{N}^+\text{R}$ on the product side, we know that the addition/elimination has occurred twice. Luckily the product quite literally has the nucleophile placed directly on center connected to the aromatic ring system via two bonds on the nitrogen. Based on this, we can reasonably predict the starting material to have had these two bonds substituted.

<figure>