

**Supplementary Information for "Causal motifs and existence of endogenous
cascades in directed networks with application to company defaults"**

I. TWO-EDGE VARIANCE CALCULATION

$$E[C^{(1,2)}] = N \sum_{k_i, k_o} P(k_i, k_o) c_{k_i, k_o}^{(1,2)}(t) \quad (1)$$

$$V[C^{(1,2)}] = N \sum_{k_i, k_o} P(k_i, k_o) \left[\left(c_{k_i, k_o}^{(1,2)}(t) - \frac{1}{N} E[C^{(1,2)}(t)] \right)^2 + c_{k_i, k_o}^{(1,2)COV}(t) \right] \quad (2)$$

$$E[\tilde{C}^{(1,2)}(t)] = N \sum_{k_i, k_o} P(k_i, k_o) \tilde{c}_{k_i, k_o}^{(1,2)}(t) \quad (3)$$

$$V[\tilde{C}^{(1,2)}(t)] = N \sum_{k_i, k_o} P(k_i, k_o) \left[\left(\tilde{c}_{k_i, k_o}^{(1,2)}(t) - \frac{1}{N} E[\tilde{C}^{(1,2)}(t)] \right)^2 + \tilde{c}_{k_i, k_o}^{(1,2)COV}(t) \right] \quad (4)$$

One-edge:

$$c_{k_i, k_o}^{(1)COV}(t) = 0 \quad (5)$$

$$\tilde{c}_{k_i, k_o}^{(1)COV}(t) = 0 \quad (6)$$

Two-edge:

$$\begin{aligned} c_{k_i, k_o}^{(2)COV}(t) &= 2 \cdot (c_{k_i, k_o}^{VI}(t) + c_{k_i, k_o}^{\Lambda I}(t) + c_{k_i, k_o}^{V\Lambda}(t)) + c_{k_i, k_o}^{VV}(t) + c_{k_i, k_o}^{\Lambda\Lambda}(t) \\ &\quad + c_{k_i, k_o}^I(t) + c_{k_i, k_o}^V(t) + c_{k_i, k_o}^\Lambda(t) \\ &= \sum_{z_i=0}^{k_i} \sum_{z_o=0}^{k_o} \left[2 \cdot \left(z_i \binom{z_o}{2} + \binom{z_i}{2} \binom{z_o}{2} + \binom{z_i}{2} z_o \right) + \binom{z_i}{3} + \binom{z_i}{4} + \binom{z_o}{3} + \binom{z_o}{4} \right. \\ &\quad \left. + \binom{z_i}{3} z_o + z_i \binom{z_o}{3} + z_i z_o + \binom{z_i}{2} + \binom{z_o}{2} \right] \times i_{k_i, k_o, z_i, z_o}^m(t) \end{aligned} \quad (7)$$

$$\begin{aligned} \tilde{c}_{k_i, k_o}^{(2)COV}(t) &= 2 \cdot (\tilde{c}_{k_i, k_o}^{VI}(t) + \tilde{c}_{k_i, k_o}^{\Lambda I}(t) + \tilde{c}_{k_i, k_o}^{V\Lambda}(t)) + \tilde{c}_{k_i, k_o}^{VV}(t) + \tilde{c}_{k_i, k_o}^{\Lambda\Lambda}(t) \\ &\quad + \tilde{c}_{k_i, k_o}^I(t) + \tilde{c}_{k_i, k_o}^V(t) + \tilde{c}_{k_i, k_o}^\Lambda(t) \\ &= \sum_{m_i=0}^{k_i} \sum_{m_o=0}^{k_o} \left[2 \cdot \left(\frac{1}{12} m_i \binom{m_o}{2} + \frac{1}{30} \binom{m_i}{2} \binom{m_o}{2} + \frac{1}{12} \binom{m_i}{2} m_o \right) \right. \\ &\quad \left. + \frac{1}{4} \binom{m_i}{3} + \frac{1}{5} \binom{m_i}{4} + \frac{1}{4} \binom{m_o}{3} + \frac{1}{5} \binom{m_o}{4} \right. \\ &\quad \left. + \frac{1}{20} \binom{m_i}{3} m_o + \frac{1}{20} m_i \binom{m_o}{3} + \frac{1}{6} m_i m_o + \frac{1}{3} \binom{m_i}{2} + \frac{1}{3} \binom{m_o}{2} \right] \times i_{k_i, k_o, m_i, m_o}^z(t) \end{aligned} \quad (8)$$

II. TEMPORAL EVOLUTION




FIG. 1: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ SI process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 2$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic














| Overlap | One node | Two nodes | Three nodes |
|---------------------|--|---|--|
| I-I |  $c^{V\Lambda}$ |  $c^{I\Lambda}$ |  c^{IV} |
| V-V |  c^{VV} |  c^{VV} |  c^V |
| $\Lambda - \Lambda$ |  $c^{\Lambda\Lambda}$ |  $c^{\Lambda\Lambda}$ |  c^Λ |
| I-V |  c^{IVV} |  c^{IV} | |
| I- Λ |  $c^{I\Lambda\Lambda}$ |  $c^{I\Lambda}$ | |
| V- Λ |  $c^{V\Lambda}$ | | |

TABLE I: Table with all possible causal motifs up to order 3.

III. Z-SCORE




FIG. 2: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ voter model process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 2$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic




FIG. 3: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ SI process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 4$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic




FIG. 4: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ voter model process on an Erdős-Rényi graph with $N = 1000$, $\langle k \rangle = 4$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic




FIG. 5: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ SI process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 6$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic




FIG. 6: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ voter model process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 6$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic




FIG. 7: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ SI process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 8$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic




FIG. 8: One-edge and two-edge statistics for $\zeta = 0.01, 0.1, 1, 10$ voter model process on an Erdős-Rényi graph with $N = 1000, \langle k \rangle = 8$. x-axis shows the time, while y-axis represents the relative frequency of a test statistic

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|--------------|--------------|--------------|--------------|--------------|
| 5.0 | 4.11 (1.22) | 4.66 (1.09) | 5.01 (1.04) | 5.12 (0.99) | 5.14 (0.92) |
| 10.0 | 6.35 (1.22) | 7.0 (1.07) | 7.15 (1.04) | 7.01 (0.99) | 6.9 (0.95) |
| 15.0 | 7.77 (1.22) | 8.5 (1.06) | 8.58 (1.05) | 8.04 (1.03) | 7.63 (1.01) |
| 20.0 | 8.8 (1.25) | 9.45 (1.16) | 9.41 (1.12) | 8.55 (1.06) | 8.14 (1.03) |
| 25.0 | 9.49 (1.31) | 10.11 (1.17) | 9.89 (1.12) | 8.88 (1.07) | 8.25 (1.07) |
| 30.0 | 9.9 (1.36) | 10.6 (1.22) | 10.09 (1.15) | 9.04 (1.11) | 8.39 (1.05) |
| 35.0 | 10.05 (1.38) | 10.95 (1.22) | 10.21 (1.13) | 9.08 (1.11) | 8.37 (1.1) |
| 40.0 | 9.93 (1.36) | 11.12 (1.25) | 10.28 (1.15) | 9.06 (1.13) | 8.23 (1.11) |
| 45.0 | 9.72 (1.39) | 11.15 (1.23) | 10.35 (1.18) | 9.06 (1.14) | 8.12 (1.09) |
| 50.0 | 9.4 (1.38) | 11.16 (1.27) | 10.16 (1.13) | 8.87 (1.11) | 7.83 (1.12) |
| 55.0 | 9.01 (1.38) | 11.09 (1.3) | 9.99 (1.16) | 8.56 (1.09) | 7.53 (1.09) |
| 60.0 | 8.5 (1.34) | 10.61 (1.21) | 9.69 (1.18) | 8.27 (1.08) | 7.27 (1.08) |
| 65.0 | 7.91 (1.28) | 10.12 (1.21) | 9.25 (1.13) | 7.93 (1.09) | 6.92 (1.07) |
| 70.0 | 7.2 (1.21) | 9.26 (1.34) | 8.77 (1.15) | 7.44 (1.05) | 6.5 (1.06) |
| 75.0 | 6.33 (1.2) | 8.05 (1.28) | 8.11 (1.09) | 6.85 (1.01) | 6.02 (1.07) |
| 80.0 | 5.33 (1.14) | 6.71 (1.17) | 7.32 (1.09) | 6.22 (1.03) | 5.46 (1.05) |
| 85.0 | 4.15 (1.11) | 5.29 (1.13) | 5.98 (1.2) | 5.29 (1.01) | 4.69 (1.05) |
| 90.0 | 2.85 (1.07) | 3.72 (1.11) | 4.16 (1.16) | 4.2 (1.02) | 3.7 (1.03) |
| 95.0 | 1.44 (1.04) | 1.94 (1.06) | 2.16 (1.14) | 2.57 (1.07) | 2.38 (1.03) |
| 100.0 | -0.08 (1.02) | 0.06 (1.04) | -0.03 (1.1) | -0.14 (1.05) | -0.05 (1.04) |

TABLE II: z-scores of one-edge statistic for an SI process with $\zeta = 0.01$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|-------------|-------------|-------------|--------------|-------------|
| 5.0 | 1.87 (1.31) | 2.2 (1.24) | 2.49 (1.22) | 2.8 (1.17) | 2.96 (1.15) |
| 10.0 | 3.25 (1.27) | 3.71 (1.18) | 4.1 (1.19) | 4.42 (1.07) | 4.49 (1.09) |
| 15.0 | 4.39 (1.23) | 4.88 (1.18) | 5.24 (1.12) | 5.46 (1.09) | 5.52 (1.07) |
| 20.0 | 5.23 (1.24) | 5.85 (1.15) | 6.06 (1.15) | 6.17 (1.11) | 6.13 (1.08) |
| 25.0 | 5.84 (1.22) | 6.44 (1.16) | 6.59 (1.16) | 6.64 (1.1) | 6.47 (1.06) |
| 30.0 | 6.4 (1.22) | 6.98 (1.18) | 7.08 (1.15) | 7.05 (1.14) | 6.69 (1.07) |
| 35.0 | 6.87 (1.16) | 7.43 (1.18) | 7.52 (1.18) | 7.24 (1.14) | 6.78 (1.07) |
| 40.0 | 7.15 (1.19) | 7.74 (1.15) | 7.71 (1.16) | 7.39 (1.12) | 6.89 (1.07) |
| 45.0 | 7.36 (1.18) | 7.85 (1.15) | 7.8 (1.15) | 7.45 (1.11) | 6.86 (1.04) |
| 50.0 | 7.4 (1.17) | 7.96 (1.19) | 7.82 (1.11) | 7.4 (1.13) | 6.7 (1.07) |
| 55.0 | 7.33 (1.16) | 7.92 (1.18) | 7.74 (1.11) | 7.31 (1.05) | 6.56 (1.06) |
| 60.0 | 7.24 (1.14) | 7.78 (1.15) | 7.64 (1.15) | 7.13 (1.04) | 6.41 (1.05) |
| 65.0 | 6.99 (1.2) | 7.59 (1.17) | 7.36 (1.07) | 6.86 (1.04) | 6.16 (1.07) |
| 70.0 | 6.51 (1.15) | 7.19 (1.11) | 7.03 (1.07) | 6.56 (1.04) | 5.83 (1.04) |
| 75.0 | 5.93 (1.17) | 6.73 (1.08) | 6.64 (1.08) | 6.12 (1.08) | 5.41 (1.03) |
| 80.0 | 5.1 (1.12) | 6.03 (1.1) | 5.99 (1.06) | 5.59 (1.03) | 4.96 (1.05) |
| 85.0 | 4.05 (1.1) | 5.02 (1.11) | 5.18 (1.05) | 4.83 (1.0) | 4.26 (0.99) |
| 90.0 | 2.85 (1.05) | 3.6 (1.06) | 4.0 (1.07) | 3.84 (0.99) | 3.43 (0.98) |
| 95.0 | 1.49 (1.02) | 1.91 (1.04) | 2.22 (1.07) | 2.4 (1.0) | 2.21 (0.97) |
| 100.0 | -0.05 (1.0) | 0.04 (1.0) | 0.06 (1.05) | -0.05 (1.01) | 0.01 (1.01) |

TABLE III: z-scores of one-edge statistic for an SI process with $\zeta = 0.1$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|-------------|--------------|--------------|-------------|-------------|
| 5.0 | 0.4 (1.15) | 0.55 (1.2) | 0.6 (1.15) | 0.71 (1.15) | 0.82 (1.15) |
| 10.0 | 0.78 (1.11) | 1.02 (1.17) | 1.13 (1.12) | 1.33 (1.12) | 1.46 (1.12) |
| 15.0 | 1.11 (1.13) | 1.43 (1.16) | 1.55 (1.11) | 1.86 (1.08) | 1.99 (1.1) |
| 20.0 | 1.49 (1.13) | 1.75 (1.14) | 1.93 (1.09) | 2.23 (1.1) | 2.43 (1.07) |
| 25.0 | 1.77 (1.07) | 2.05 (1.11) | 2.27 (1.1) | 2.57 (1.12) | 2.83 (1.05) |
| 30.0 | 2.05 (1.07) | 2.34 (1.1) | 2.55 (1.1) | 2.82 (1.1) | 3.1 (1.08) |
| 35.0 | 2.26 (1.1) | 2.61 (1.1) | 2.86 (1.12) | 3.07 (1.05) | 3.36 (1.05) |
| 40.0 | 2.45 (1.09) | 2.79 (1.1) | 3.1 (1.12) | 3.27 (1.09) | 3.52 (1.06) |
| 45.0 | 2.68 (1.08) | 3.0 (1.08) | 3.24 (1.08) | 3.45 (1.08) | 3.62 (1.05) |
| 50.0 | 2.88 (1.04) | 3.13 (1.08) | 3.4 (1.08) | 3.56 (1.04) | 3.72 (1.04) |
| 55.0 | 2.96 (1.04) | 3.26 (1.08) | 3.47 (1.07) | 3.65 (1.02) | 3.75 (1.06) |
| 60.0 | 3.02 (1.07) | 3.3 (1.07) | 3.48 (1.07) | 3.67 (1.05) | 3.7 (1.04) |
| 65.0 | 3.02 (1.05) | 3.32 (1.05) | 3.45 (1.05) | 3.68 (1.05) | 3.73 (1.03) |
| 70.0 | 2.98 (1.04) | 3.26 (1.02) | 3.42 (1.04) | 3.59 (1.05) | 3.61 (1.05) |
| 75.0 | 2.89 (1.06) | 3.15 (1.03) | 3.33 (1.06) | 3.43 (1.05) | 3.48 (1.06) |
| 80.0 | 2.68 (1.03) | 2.96 (1.02) | 3.13 (1.05) | 3.21 (1.05) | 3.23 (1.04) |
| 85.0 | 2.36 (1.04) | 2.61 (1.03) | 2.77 (1.04) | 2.84 (1.03) | 2.87 (1.05) |
| 90.0 | 1.9 (1.03) | 2.1 (1.02) | 2.23 (1.02) | 2.35 (1.02) | 2.35 (1.03) |
| 95.0 | 1.18 (1.01) | 1.33 (1.03) | 1.44 (1.03) | 1.56 (1.02) | 1.58 (1.04) |
| 100.0 | 0.02 (1.03) | -0.02 (1.04) | -0.04 (1.01) | 0.05 (0.99) | 0.07 (1.03) |

TABLE IV: z-scores of one-edge statistic for an SI process with $\zeta = 1.0$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|--------------|--------------|-------------|--------------|--------------|
| 5.0 | 0.0 (1.03) | 0.11 (1.04) | 0.03 (1.06) | 0.09 (1.01) | 0.07 (1.01) |
| 10.0 | 0.02 (1.03) | 0.16 (1.04) | 0.13 (1.04) | 0.21 (1.07) | 0.2 (1.05) |
| 15.0 | 0.09 (1.06) | 0.21 (1.05) | 0.17 (1.06) | 0.28 (1.05) | 0.29 (1.04) |
| 20.0 | 0.17 (1.04) | 0.27 (1.01) | 0.26 (1.05) | 0.35 (1.05) | 0.38 (1.04) |
| 25.0 | 0.19 (1.07) | 0.32 (1.02) | 0.31 (1.05) | 0.44 (1.01) | 0.48 (1.05) |
| 30.0 | 0.25 (1.06) | 0.36 (1.02) | 0.39 (1.05) | 0.5 (1.0) | 0.55 (1.08) |
| 35.0 | 0.29 (1.05) | 0.44 (1.0) | 0.45 (1.03) | 0.58 (1.02) | 0.61 (1.06) |
| 40.0 | 0.34 (1.06) | 0.46 (1.0) | 0.5 (1.03) | 0.62 (1.05) | 0.66 (1.03) |
| 45.0 | 0.36 (1.06) | 0.48 (1.0) | 0.54 (1.02) | 0.68 (1.04) | 0.71 (1.03) |
| 50.0 | 0.38 (1.09) | 0.51 (1.01) | 0.6 (1.03) | 0.69 (1.03) | 0.78 (1.02) |
| 55.0 | 0.4 (1.07) | 0.54 (0.99) | 0.64 (1.02) | 0.75 (1.03) | 0.83 (1.0) |
| 60.0 | 0.43 (1.06) | 0.59 (0.99) | 0.67 (1.04) | 0.77 (1.01) | 0.87 (1.0) |
| 65.0 | 0.45 (1.04) | 0.61 (0.98) | 0.67 (1.01) | 0.79 (1.03) | 0.88 (0.99) |
| 70.0 | 0.44 (1.01) | 0.64 (0.99) | 0.7 (1.01) | 0.82 (1.04) | 0.92 (1.0) |
| 75.0 | 0.43 (1.0) | 0.63 (0.97) | 0.7 (0.99) | 0.79 (1.02) | 0.92 (1.02) |
| 80.0 | 0.41 (1.0) | 0.61 (0.99) | 0.67 (1.01) | 0.75 (1.02) | 0.87 (1.0) |
| 85.0 | 0.35 (1.0) | 0.54 (1.0) | 0.63 (1.02) | 0.68 (1.02) | 0.79 (1.0) |
| 90.0 | 0.3 (1.0) | 0.47 (1.0) | 0.55 (1.03) | 0.57 (1.02) | 0.68 (1.02) |
| 95.0 | 0.18 (1.01) | 0.31 (1.01) | 0.39 (1.01) | 0.37 (1.03) | 0.45 (1.01) |
| 100.0 | -0.09 (1.01) | -0.01 (1.01) | 0.03 (1.04) | -0.06 (1.04) | -0.02 (1.01) |

TABLE V: z-scores of one-edge statistic for an SI process with $\zeta = 10.0$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|---------------------|--------------|-------------|--------------|-------------|--------------|
| percentage | | | | | |
| 5.0 | -0.02 (1.01) | 0.04 (1.04) | 0.06 (1.01) | 0.03 (1.03) | 0.02 (1.01) |
| 10.0 | -0.0 (1.01) | 0.07 (1.04) | 0.07 (1.0) | 0.03 (1.06) | 0.01 (1.04) |
| 15.0 | -0.02 (1.05) | 0.09 (1.04) | 0.07 (1.04) | 0.04 (1.02) | 0.05 (1.03) |
| 20.0 | -0.02 (1.06) | 0.1 (1.06) | 0.07 (1.08) | 0.05 (1.04) | 0.08 (1.02) |
| 25.0 | -0.03 (1.04) | 0.09 (1.05) | 0.06 (1.08) | 0.05 (1.04) | 0.11 (1.02) |
| 30.0 | -0.01 (1.02) | 0.11 (1.06) | 0.07 (1.05) | 0.07 (1.03) | 0.14 (1.01) |
| 35.0 | 0.0 (1.01) | 0.1 (1.04) | 0.08 (1.05) | 0.1 (1.0) | 0.15 (0.99) |
| 40.0 | 0.0 (1.0) | 0.09 (1.02) | 0.07 (1.06) | 0.11 (1.02) | 0.15 (1.0) |
| 45.0 | 0.0 (1.0) | 0.11 (1.03) | 0.05 (1.03) | 0.11 (1.05) | 0.13 (1.0) |
| 50.0 | 0.02 (1.02) | 0.1 (1.01) | 0.07 (1.04) | 0.11 (1.02) | 0.14 (1.02) |
| 55.0 | 0.04 (1.02) | 0.11 (0.99) | 0.08 (1.02) | 0.12 (1.0) | 0.14 (1.03) |
| 60.0 | 0.05 (1.03) | 0.1 (1.0) | 0.08 (1.0) | 0.1 (0.99) | 0.14 (1.02) |
| 65.0 | 0.04 (1.03) | 0.1 (1.02) | 0.08 (1.04) | 0.1 (1.01) | 0.13 (1.04) |
| 70.0 | 0.06 (1.05) | 0.1 (1.03) | 0.08 (1.02) | 0.11 (1.03) | 0.13 (1.04) |
| 75.0 | 0.05 (1.05) | 0.1 (1.04) | 0.07 (1.0) | 0.12 (1.02) | 0.12 (1.04) |
| 80.0 | 0.05 (1.07) | 0.11 (1.05) | 0.06 (1.0) | 0.12 (1.01) | 0.12 (1.02) |
| 85.0 | 0.04 (1.06) | 0.1 (1.03) | 0.05 (1.02) | 0.08 (1.02) | 0.11 (1.03) |
| 90.0 | 0.03 (1.05) | 0.07 (1.01) | 0.05 (1.0) | 0.08 (1.03) | 0.07 (1.02) |
| 95.0 | 0.03 (1.05) | 0.04 (1.01) | 0.01 (1.01) | 0.07 (1.02) | 0.04 (1.03) |
| 100.0 | 0.01 (1.04) | 0.03 (1.04) | -0.03 (1.01) | 0.02 (1.03) | -0.03 (1.01) |

TABLE VI: z-scores of one-edge statistic for an SI process with $\zeta = 100.0$ on a network of $N = 1000$ vertices.

IV. KS-TEST

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|--------------|--------------|--------------|--------------|--------------|
| 5.0 | 4.56 (2.12) | 5.14 (1.9) | 5.46 (1.86) | 5.53 (1.75) | 5.52 (1.62) |
| 10.0 | 7.09 (2.06) | 7.75 (1.81) | 7.77 (1.76) | 7.5 (1.64) | 7.43 (1.5) |
| 15.0 | 8.62 (1.98) | 9.33 (1.73) | 9.21 (1.64) | 8.47 (1.58) | 8.1 (1.51) |
| 20.0 | 9.64 (1.87) | 10.19 (1.72) | 9.9 (1.6) | 9.04 (1.53) | 8.53 (1.44) |
| 25.0 | 10.24 (1.88) | 10.7 (1.7) | 10.36 (1.56) | 9.2 (1.5) | 8.61 (1.42) |
| 30.0 | 10.58 (1.86) | 11.06 (1.67) | 10.47 (1.57) | 9.29 (1.46) | 8.68 (1.38) |
| 35.0 | 10.52 (1.83) | 11.25 (1.62) | 10.43 (1.49) | 9.25 (1.44) | 8.51 (1.35) |
| 40.0 | 10.25 (1.78) | 11.17 (1.64) | 10.33 (1.48) | 9.1 (1.41) | 8.3 (1.35) |
| 45.0 | 9.76 (1.77) | 10.95 (1.56) | 10.18 (1.47) | 8.97 (1.41) | 8.11 (1.29) |
| 50.0 | 9.08 (1.68) | 10.63 (1.57) | 9.87 (1.38) | 8.7 (1.32) | 7.74 (1.3) |
| 55.0 | 8.35 (1.57) | 10.3 (1.6) | 9.54 (1.41) | 8.28 (1.32) | 7.37 (1.28) |
| 60.0 | 7.49 (1.49) | 9.5 (1.45) | 9.06 (1.39) | 7.86 (1.25) | 7.04 (1.23) |
| 65.0 | 6.63 (1.42) | 8.66 (1.42) | 8.44 (1.33) | 7.45 (1.25) | 6.63 (1.23) |
| 70.0 | 5.71 (1.32) | 7.61 (1.46) | 7.71 (1.35) | 6.84 (1.17) | 6.14 (1.19) |
| 75.0 | 4.73 (1.28) | 6.34 (1.38) | 6.79 (1.26) | 6.14 (1.13) | 5.58 (1.19) |
| 80.0 | 3.75 (1.19) | 5.02 (1.28) | 5.73 (1.24) | 5.36 (1.13) | 4.93 (1.16) |
| 85.0 | 2.75 (1.14) | 3.76 (1.22) | 4.36 (1.28) | 4.38 (1.11) | 4.11 (1.14) |
| 90.0 | 1.77 (1.1) | 2.52 (1.17) | 2.85 (1.22) | 3.23 (1.1) | 3.08 (1.11) |
| 95.0 | 0.81 (1.05) | 1.27 (1.11) | 1.39 (1.16) | 1.71 (1.1) | 1.79 (1.09) |
| 100.0 | -0.11 (1.01) | 0.08 (1.07) | -0.02 (1.12) | -0.11 (1.05) | -0.06 (1.06) |

TABLE VII: z-scores of two-edge statistic for an SI process with $\zeta = 0.01$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|--------------|-------------|-------------|-------------|-------------|
| 5.0 | 1.54 (1.98) | 1.89 (1.95) | 2.25 (1.89) | 2.66 (1.86) | 2.85 (1.8) |
| 10.0 | 2.93 (1.86) | 3.47 (1.82) | 3.9 (1.78) | 4.34 (1.67) | 4.47 (1.62) |
| 15.0 | 4.03 (1.81) | 4.67 (1.76) | 5.01 (1.67) | 5.35 (1.61) | 5.47 (1.53) |
| 20.0 | 4.93 (1.75) | 5.63 (1.65) | 5.87 (1.57) | 6.09 (1.56) | 6.1 (1.46) |
| 25.0 | 5.54 (1.67) | 6.2 (1.61) | 6.43 (1.57) | 6.52 (1.49) | 6.43 (1.38) |
| 30.0 | 6.05 (1.66) | 6.67 (1.56) | 6.87 (1.53) | 6.87 (1.49) | 6.62 (1.37) |
| 35.0 | 6.48 (1.57) | 7.05 (1.52) | 7.29 (1.52) | 7.04 (1.46) | 6.68 (1.34) |
| 40.0 | 6.65 (1.54) | 7.24 (1.5) | 7.43 (1.48) | 7.16 (1.38) | 6.75 (1.3) |
| 45.0 | 6.77 (1.48) | 7.29 (1.47) | 7.43 (1.4) | 7.14 (1.37) | 6.69 (1.24) |
| 50.0 | 6.74 (1.43) | 7.28 (1.44) | 7.39 (1.33) | 7.09 (1.36) | 6.48 (1.25) |
| 55.0 | 6.55 (1.41) | 7.18 (1.42) | 7.21 (1.3) | 6.93 (1.25) | 6.3 (1.24) |
| 60.0 | 6.31 (1.36) | 6.92 (1.35) | 7.02 (1.34) | 6.7 (1.2) | 6.1 (1.2) |
| 65.0 | 5.89 (1.33) | 6.61 (1.35) | 6.67 (1.23) | 6.36 (1.18) | 5.79 (1.19) |
| 70.0 | 5.3 (1.28) | 6.11 (1.29) | 6.23 (1.21) | 5.98 (1.16) | 5.42 (1.15) |
| 75.0 | 4.61 (1.25) | 5.51 (1.24) | 5.72 (1.19) | 5.48 (1.16) | 4.97 (1.13) |
| 80.0 | 3.78 (1.2) | 4.72 (1.22) | 4.97 (1.14) | 4.87 (1.11) | 4.45 (1.12) |
| 85.0 | 2.83 (1.13) | 3.69 (1.19) | 4.06 (1.12) | 4.06 (1.07) | 3.74 (1.07) |
| 90.0 | 1.86 (1.07) | 2.49 (1.13) | 2.88 (1.09) | 3.05 (1.05) | 2.87 (1.03) |
| 95.0 | 0.92 (1.05) | 1.24 (1.08) | 1.47 (1.05) | 1.71 (1.03) | 1.7 (1.0) |
| 100.0 | -0.02 (1.01) | 0.05 (1.04) | 0.05 (1.01) | -0.02 (1.0) | -0.01 (1.0) |

TABLE VIII: z-scores of two-edge statistic for an SI process with $\zeta = 0.1$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|--------------|-------------|--------------|-------------|-------------|
| 5.0 | 0.17 (1.39) | 0.32 (1.46) | 0.39 (1.46) | 0.55 (1.41) | 0.64 (1.39) |
| 10.0 | 0.48 (1.34) | 0.69 (1.34) | 0.8 (1.35) | 1.07 (1.36) | 1.25 (1.33) |
| 15.0 | 0.74 (1.3) | 1.09 (1.34) | 1.22 (1.33) | 1.58 (1.3) | 1.76 (1.31) |
| 20.0 | 1.07 (1.32) | 1.37 (1.34) | 1.58 (1.29) | 1.98 (1.31) | 2.19 (1.27) |
| 25.0 | 1.31 (1.27) | 1.66 (1.29) | 1.93 (1.27) | 2.32 (1.34) | 2.57 (1.21) |
| 30.0 | 1.6 (1.29) | 1.94 (1.28) | 2.21 (1.28) | 2.56 (1.3) | 2.87 (1.24) |
| 35.0 | 1.77 (1.28) | 2.21 (1.29) | 2.53 (1.28) | 2.8 (1.24) | 3.12 (1.19) |
| 40.0 | 1.98 (1.25) | 2.4 (1.29) | 2.76 (1.29) | 2.98 (1.25) | 3.3 (1.2) |
| 45.0 | 2.15 (1.21) | 2.61 (1.25) | 2.9 (1.25) | 3.15 (1.23) | 3.4 (1.17) |
| 50.0 | 2.31 (1.15) | 2.72 (1.24) | 3.03 (1.23) | 3.25 (1.19) | 3.5 (1.16) |
| 55.0 | 2.39 (1.15) | 2.83 (1.21) | 3.09 (1.2) | 3.32 (1.17) | 3.54 (1.16) |
| 60.0 | 2.42 (1.16) | 2.85 (1.18) | 3.09 (1.18) | 3.35 (1.17) | 3.49 (1.14) |
| 65.0 | 2.43 (1.16) | 2.85 (1.16) | 3.05 (1.15) | 3.34 (1.16) | 3.51 (1.12) |
| 70.0 | 2.38 (1.14) | 2.79 (1.12) | 3.0 (1.14) | 3.23 (1.14) | 3.36 (1.12) |
| 75.0 | 2.26 (1.13) | 2.67 (1.11) | 2.88 (1.14) | 3.07 (1.13) | 3.22 (1.12) |
| 80.0 | 2.06 (1.1) | 2.47 (1.1) | 2.66 (1.11) | 2.84 (1.13) | 2.94 (1.1) |
| 85.0 | 1.78 (1.08) | 2.15 (1.09) | 2.31 (1.09) | 2.48 (1.09) | 2.59 (1.09) |
| 90.0 | 1.37 (1.07) | 1.68 (1.07) | 1.8 (1.05) | 2.0 (1.08) | 2.07 (1.06) |
| 95.0 | 0.78 (1.03) | 1.01 (1.05) | 1.08 (1.03) | 1.25 (1.05) | 1.33 (1.06) |
| 100.0 | -0.01 (1.01) | 0.03 (1.03) | -0.03 (0.99) | 0.04 (1.01) | 0.07 (1.03) |

TABLE IX: z-scores of two-edge statistic for an SI process with $\zeta = 1.0$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ percentage | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|-----------------------------------|--------------|--------------|-------------|--------------|--------------|
| 5.0 | nan (nan) | 0.08 (1.21) | 0.02 (1.07) | 0.09 (1.16) | 0.04 (0.99) |
| 10.0 | 0.04 (1.07) | 0.13 (1.14) | 0.12 (1.08) | 0.18 (1.12) | 0.14 (1.05) |
| 15.0 | 0.08 (1.09) | 0.16 (1.11) | 0.16 (1.05) | 0.26 (1.11) | 0.22 (1.09) |
| 20.0 | 0.1 (1.11) | 0.22 (1.09) | 0.23 (1.07) | 0.31 (1.1) | 0.32 (1.06) |
| 25.0 | 0.13 (1.07) | 0.24 (1.07) | 0.28 (1.08) | 0.4 (1.06) | 0.42 (1.09) |
| 30.0 | 0.18 (1.06) | 0.27 (1.05) | 0.32 (1.07) | 0.45 (1.03) | 0.48 (1.1) |
| 35.0 | 0.23 (1.04) | 0.35 (1.04) | 0.38 (1.06) | 0.53 (1.04) | 0.57 (1.09) |
| 40.0 | 0.28 (1.05) | 0.39 (1.06) | 0.43 (1.06) | 0.56 (1.06) | 0.61 (1.06) |
| 45.0 | 0.29 (1.09) | 0.41 (1.04) | 0.47 (1.06) | 0.62 (1.06) | 0.65 (1.05) |
| 50.0 | 0.31 (1.11) | 0.44 (1.04) | 0.53 (1.07) | 0.63 (1.07) | 0.73 (1.04) |
| 55.0 | 0.33 (1.06) | 0.47 (1.01) | 0.57 (1.06) | 0.69 (1.06) | 0.77 (1.04) |
| 60.0 | 0.37 (1.06) | 0.52 (1.01) | 0.59 (1.08) | 0.72 (1.03) | 0.82 (1.02) |
| 65.0 | 0.39 (1.06) | 0.54 (1.0) | 0.6 (1.06) | 0.73 (1.06) | 0.82 (1.0) |
| 70.0 | 0.38 (1.02) | 0.56 (0.99) | 0.62 (1.05) | 0.77 (1.07) | 0.86 (1.01) |
| 75.0 | 0.36 (1.03) | 0.55 (0.98) | 0.61 (1.03) | 0.73 (1.06) | 0.86 (1.04) |
| 80.0 | 0.35 (1.03) | 0.52 (0.99) | 0.58 (1.05) | 0.7 (1.04) | 0.81 (1.02) |
| 85.0 | 0.31 (1.02) | 0.46 (1.0) | 0.54 (1.04) | 0.64 (1.05) | 0.73 (1.0) |
| 90.0 | 0.25 (1.01) | 0.4 (1.0) | 0.46 (1.03) | 0.53 (1.04) | 0.61 (1.02) |
| 95.0 | 0.14 (1.01) | 0.25 (1.0) | 0.32 (1.01) | 0.35 (1.05) | 0.4 (1.01) |
| 100.0 | -0.08 (1.01) | -0.03 (0.99) | 0.0 (1.02) | -0.03 (1.05) | -0.03 (1.01) |

TABLE X: z-scores of two-edge statistic for an SI process with $\zeta = 10.0$ on a network of $N = 1000$ vertices.

| $\langle k \rangle$ | 2.0 | 3.0 | 4.0 | 6.0 | 8.0 |
|---------------------|--------------|-------------|--------------|-------------|--------------|
| percentage | | | | | |
| 5.0 | nan (nan) | 0.07 (1.17) | -0.04 (0.96) | 0.02 (1.08) | 0.05 (1.08) |
| 10.0 | 0.02 (1.03) | 0.07 (1.16) | 0.04 (1.02) | 0.05 (1.03) | 0.03 (1.08) |
| 15.0 | 0.01 (1.06) | 0.05 (1.06) | 0.06 (1.05) | 0.03 (1.02) | 0.07 (1.05) |
| 20.0 | -0.02 (1.07) | 0.08 (1.08) | 0.07 (1.1) | 0.05 (1.04) | 0.09 (1.04) |
| 25.0 | -0.02 (1.05) | 0.08 (1.06) | 0.06 (1.09) | 0.05 (1.02) | 0.09 (1.02) |
| 30.0 | -0.01 (1.04) | 0.1 (1.07) | 0.07 (1.07) | 0.06 (1.02) | 0.12 (1.02) |
| 35.0 | -0.01 (1.01) | 0.1 (1.04) | 0.06 (1.05) | 0.08 (0.98) | 0.13 (1.0) |
| 40.0 | -0.0 (1.01) | 0.09 (1.03) | 0.04 (1.08) | 0.1 (1.01) | 0.13 (1.0) |
| 45.0 | -0.01 (1.03) | 0.11 (1.02) | 0.01 (1.05) | 0.09 (1.04) | 0.11 (1.01) |
| 50.0 | 0.01 (1.06) | 0.1 (1.0) | 0.04 (1.05) | 0.09 (1.01) | 0.12 (1.04) |
| 55.0 | -0.0 (1.03) | 0.11 (0.99) | 0.04 (1.03) | 0.1 (1.01) | 0.12 (1.05) |
| 60.0 | 0.01 (1.04) | 0.1 (1.0) | 0.06 (1.02) | 0.08 (1.0) | 0.13 (1.03) |
| 65.0 | 0.0 (1.04) | 0.08 (1.02) | 0.06 (1.07) | 0.09 (1.02) | 0.12 (1.03) |
| 70.0 | 0.01 (1.05) | 0.09 (1.05) | 0.05 (1.04) | 0.1 (1.05) | 0.12 (1.04) |
| 75.0 | 0.02 (1.04) | 0.08 (1.05) | 0.04 (1.01) | 0.12 (1.05) | 0.12 (1.03) |
| 80.0 | 0.02 (1.07) | 0.09 (1.05) | 0.04 (1.0) | 0.11 (1.02) | 0.11 (1.02) |
| 85.0 | 0.01 (1.04) | 0.08 (1.02) | 0.02 (1.02) | 0.07 (1.01) | 0.1 (1.03) |
| 90.0 | 0.01 (1.03) | 0.04 (1.01) | 0.03 (1.01) | 0.07 (1.03) | 0.06 (1.01) |
| 95.0 | 0.01 (1.04) | 0.02 (1.01) | -0.01 (1.01) | 0.07 (1.03) | 0.02 (1.02) |
| 100.0 | -0.02 (1.04) | 0.0 (1.02) | -0.06 (1.0) | 0.02 (1.03) | -0.04 (1.01) |

TABLE XI: z-scores of two-edge statistic for an SI process with $\zeta = 100.0$ on a network of $N = 1000$ vertices.




FIG. 9: KS-test of the results for an voter-model process with $\zeta = 0.01$ on a network of $N = 1000$ vertices.




FIG. 10: KS-test of the results for a voter-model process with $\zeta = 0.1$ on a network of $N = 1000$ vertices.




FIG. 11: KS-test of the results for a voter-model process with $\zeta = 1$ on a network of $N = 1000$ vertices.




FIG. 12: KS-test of the results for a voter-model process with $\zeta = 10$ on a network of $N = 1000$ vertices.




FIG. 13: KS-test of the results for a voter-model process with $\zeta = 100$ on a network of $N = 1000$ vertices.




FIG. 14: KS-test of the results for a SI process with $\zeta = 0.01$ on a network of $N = 1000$ vertices.





FIG. 15: KS-test of the results for a SI process with $\zeta = 0.1$ on a network of $N = 1000$ vertices.

FIG. 16: KS-test of the results for a SI process with $\zeta = 1$ on a network of $N = 1000$ vertices.




FIG. 17: KS-test of the results for a SI process with $\zeta = 10$ on a network of $N = 1000$ vertices.




FIG. 18: KS-test of the results for a SI process with $\zeta = 100$ on a network of $N = 1000$ vertices.




FIG. 19: Comparison of Mahalanobis distance of the results for an SI process on a network of $N = 1000$ vertices. The best statistically significant statistic for a given percentage, ζ and $\langle k \rangle$ is stated as "lc" for largest component, "one" for one-edge, "two" for two-edge, etc. A z-score greater than 1 is considered to be statistically significant, otherwise an "Na" value is put in the table.




FIG. 20: Comparison of Mahalanobis distance of the results for a voter model process on a network of $N = 1000$ vertices. The best statistically significant statistic for a given percentage, ζ and $\langle k \rangle$ is stated as "lc" for largest component, "one" for one-edge, "two" for two-edge, etc. A z-score greater than 1 is considered to be statistically significant, otherwise an "Na" value is put in the table.