

**PHILOSOPHICAL TRANSACTIONS
OF THE ROYAL SOCIETY B**
BIOLOGICAL SCIENCES

**Historic collections as a tool for assessing the global
pollinator crisis**

Journal:	<i>Philosophical Transactions B</i>
Manuscript ID	Draft
Article Type:	Research
Date Submitted by the Author:	n/a
Complete List of Authors:	Bartomeus, Ignasi; Estacion Biologica de Donana CSIC Stavert, Jamie; The University of Auckland, Ward, Darren; Landcare Research New Zealand Aguado, Oscar; Andrena Iniciativas y Estudios Medio Ambientales
Issue Code (this should have already been entered but please contact the Editorial Office if it is not present):	MUSEUM
Subject:	Ecology < BIOLOGY
Keywords:	Museums, biodiversity, global change, bees, hoverflies, butterflies

SCHOLARONE™
Manuscripts

Historic collections as a tool for assessing the global pollinator crisis

Bartomeus, I.^{1*}, Stavert, J.R.², Ward, D.^{2,3}, and Aguado, O.⁴

¹ Estación Biológica de Doñana (EBD-CSIC), Avda. Américo Vespucio 26, Isla de la Cartuja, E-41092 Sevilla, Spain

² Centre for Biodiversity and Biosecurity, School of Biological Sciences, The University of Auckland, Auckland, New Zealand

³ Landcare Research, Auckland, New Zealand

⁴ Andrena Iniciativas y Estudios Medio Ambientales, Valladolid, Spain

*Correspondence: nacho.bartomeus@gmail.com

Author contributions: IB wrote the initial draft. DW and OA provided data. IB and JS analysed the data. All authors contributed to writing the manuscript.

Running head: Historic collections and pollinators

Abstract

There is increasing concern about the decline of pollinators worldwide. However, despite reports that pollinator declines are widespread, data are scarce and often geographically and taxonomically biased. These biases limit robust inference about any potential pollinator crisis. Non-structured and opportunistic historical specimen collection data provide the only source of historical information which can serve as a baseline for identifying pollinator declines. Specimens historically collected and preserved in museums not only provide information on where and when species were collected, but also contain other ecological information such as species interactions and morphological traits. Here, we provide a synthesis of how researchers have used historical data to identify long-term changes in biodiversity, species abundances, morphology and pollination services. Despite recent advances, we show that information on the status and trends of most pollinators is absent, but we highlight opportunities and limitations to progress the assessment of pollinator declines globally. Finally, we demonstrate different approaches to analysing museum collection data using two contrasting case studies from distinct geographical regions (New Zealand and Spain) for which long-term pollinator declines have never been assessed. There is immense potential for museum specimens to play a central role in assessing the extent of the global pollination crisis.

Keywords: Museums, biodiversity, global change, bees, hoverflies, butterflies.

1
2
3
4 36 **Introduction**

5
6
7 37 Animal pollinators are a critical component of both natural and agricultural ecosystems
8
9 38 worldwide, given their role in plant reproduction [1] and food security [2]. As with many
10
11 39 other taxa, pollinators are vulnerable to a range of anthropogenic disturbances, which can
12
13
14 40 cause local and regional population declines or even extinctions. The vulnerability of
15
16 41 pollinators was identified several decades ago, and was popularized in 1996 by the
17
18 42 influential book “The forgotten pollinators” [3]. However, early accounts of pollinator
19
20 43 declines were somewhat anecdotal, given the lack of pollinator population data at that
21
22 44 time. These initial claims triggered the first efforts to assess this potential issue and
23
24
25 45 included the formation of a US National Academy of Science (NAS) panel in 2006, which
26
27 46 was commissioned to assess the extent of pollinator declines. The NAS report concluded
28
29 47 that “For most pollinator species [...] the paucity of long-term population data and the
30
31 48 incomplete knowledge of even basic taxonomy and ecology make definitive assessment of
32
33 49 status exceedingly difficult” [4]. Since then, studies on pollinator responses to various
34
35 50 global change drivers have multiplied rapidly. Researchers have now developed strong
36
37 51 consensus that disturbances such as habitat destruction, land-use intensification, chemical
38
39 52 exposure, exotic species and climate change are causing pollinator declines, and often act
40
41 53 synergistically [5,6]. Yet, the current status and population trends of most pollinator
42
43 54 species worldwide remain unknown. For example, a recent IUCN report concluded that
44
45 55 even for Europe’s comparatively well-studied bee fauna, greater than 55% of bee species
46
47 56 fell into the “Data Deficient” category [7]. For countries outside of Europe and the US, data
48
49
50 57 on pollinator populations is almost non-existent.
51
52
53
54
55
56
57
58
59
60

One of the main barriers to identifying long-term pollinator population trends is that pollinators are incredibly taxonomically diverse and include bees, flies, butterflies, beetles, birds, bats and lizards [8]. Additionally, many pollinators are highly mobile, short-lived and small, which makes monitoring their populations difficult. Bees are generally regarded as the most important pollinator group due to their abundance, pollination efficiency and widespread distribution [9]. However, bees are diverse, with more than 20,000 species currently described worldwide, and often require expert taxonomists for identification. Furthermore, the uneven distribution of researchers has resulted in geographical biases in bee decline research [10], as well as taxonomic biases toward species that are easier to identify, such as bumblebees [11,12].

One solution to overcoming these barriers is the use of space-for-time substitutions, where researchers compare pollinator populations across environmental gradients. Despite critiques on the robustness of this approach [13,14], these studies currently provide the most extensive source of pollinator population data. For example, researchers have recently estimated bee richness declines for every country in Europe using predictions from models of pollinator associations with different land-use types [15]. A second important method is the use of data collected from pollinator monitoring programs, which are often driven by citizen scientists. This approach was inspired by successful butterfly monitoring programs [16] and is currently being extended to other pollinator taxa. However, these programs require significant time to generate long-term datasets and cannot be used to assess historic pollinator populations. Finally, the most practical approach for assessing long-term historical pollinator population trends is to use historical

information on species occurrences, which is often archived in museum collections [e.g. 17].

In this review, we first assess current evidence for pollinator richness declines and present a roadmap outlining a strategy for using historical collection data to fill current knowledge gaps. We highlight the major technical difficulties involved in using historical collection data and demonstrate several approaches for analysing different types of collection data to assess long-term pollinator population trends. Finally, we highlight the need to move beyond simple biological diversity descriptors and unleash the power of historical data to assess changes in species interactions, ecosystem functioning and evolutionary changes through time.

Current evidence on pollinator declines

At a global scale, current evidence of pollinator declines is highly limited with most data restricted to the US and Europe. It is unsurprising that studies on pollinator declines are biased towards developed western countries, which have also been subject to extensive anthropogenic disturbance. For example, in the UK and the Netherlands, a citizen science based study using both observations and museum collection data detected strong richness declines for bees, hoverflies and flowering plants [18]. In the Netherlands, museum data have also revealed simultaneous plant and pollinator declines [19]. Specifically, bee species with the strongest host plant preferences (i.e., specialists) displayed the strongest declines and thus, were most threatened with extinction. However, it is important to note that even for these two countries, local estimates of pollinator richness are biased toward large cities

1
2
3 101 and regions dominated by agriculture, and thus lack data for well-preserved natural areas.

4
5 102 Further exploration of this dataset revealed that for declining pollinator taxa, the trend has

6
7
8 103 attenuated in recent decades [20].

9
10
11 104 Although studies of local pollinator communities often detect richness declines, regional

12
13 105 richness may remain relatively stable. For example, regional estimates for bee species

14
15 106 richness changes in the eastern US show moderate declines [17] and very few regional

16
17 107 extinctions [21]. This is a pattern also detected in the UK, where relatively few regional bee

18
19 108 extinctions have been reported [22]. These regional findings are in stark contrast with the

20
21 109 widespread local extinctions reported in local studies. For example, Burkle et al. [23]

22
23 110 compared historical observations of bee species' occurrences in a large forested ecosystem

24
25 111 with remaining forest remnants and reports several local extinctions. However, it is

26
27 112 important to note that there is strong concordance between local extinctions and regional

28
29 113 declines [24], suggesting that local extinctions are indicators of regional population

30
31 114 declines.

32
33
34
35 115 Reported declines for bumblebees are the most severe of all pollinator taxa. For example,

36
37 116 declines of up to 18% in local bumblebee richness have been reported for Belgium and the

38
39 117 Netherlands [20]. In other parts of Europe, local richness declines range from 5% in Great

40
41 118 Britain [20] to 42% in Denmark [25]. In the USA, reported bumble decline are also severe

42
43 119 with estimates ranging between 25% [26] and 30% [17]. However, studies on species

44
45 120 richness changes for other pollinator taxa are both scarce and geographically restricted.

46
47 121 For butterflies, the only evidence of richness declines comes from Europe. Butterfly species

48
49 122 richness has declined substantially in the Netherlands and Belgium since the 1950's,

1
2
3 123 although declines in Great Britain have been less severe [20]. In Belgium, another study
4
5 124 [27] found that richness declines have been severe (approximately 30%), although this
6
7
8 125 study assessed richness changes over a longer time period (early 1900's to 2000)
9
10 126 compared to [20] (1950-69 vs. 1970-80 and 1970-89 vs. 1990-2009). In parts of Germany,
11
12 127 up to 70% declines in local butterfly richness have been reported [28]. Compared with
13
14
15 128 other insect pollinator taxa, there are very few studies on hoverfly species richness
16
17 129 changes, which are all restricted to Europe. In Belgium, Great Britain and the Netherlands,
18
19 130 hoverfly richness changes have been modest [20]. In the Netherlands, moderate increases
20
21 131 in hoverfly species richness have been shown, whereas in Great Britain no significant
22
23 132 directional changes were detected [18]. Furthermore, directionality (richness increase or
24
25 133 decrease) varies depending on the time period assessed. For example, hoverfly richness
26
27 134 decreased in Belgium by approximately 6% from 1950-69 to 1970-80, but increased by
28
29 135 approximately 10% between 1970-89 and 1990-2009 [20].
30
31
32
33
34 136 For illustrative purposes, we mapped the findings of this studies in Figure 1 to show the
35
36 137 strong contrast between bee species richness worldwide, with bee diversity hotspots in
37
38 138 Mediterranean countries, against the paucity of countries for which we have any local or
39
40 139 regional data on bee, hoverfly or butterfly declines (see raw data in Sup Mat 1). Despite
41
42 140 outside of Europe and the US and for non-insect taxa, there are very few or no studies on
43
44 141 pollinator declines using historical records, there are species-specific examples of historical
45
46 142 losses from different parts of the world (e.g., *Bombus dalbhomii*; [29]).
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 1: Global map showing a) bee species richness per area (Data from www.discoverlife.org) calculated as the residuals of the log-log regression between bee species richness per country and country size. This correction accounts for the species-area relationship. Warmer colours indicate higher bee diversity. Note that some African countries may have incomplete listed faunas and that Alaska is included with USA values. Countries with available historical changes in (b) bee, (c) hoverfly and (d) butterfly richness within the last 100 years. Warmer colours indicate steeper average declines. Countries without data are coloured in white.

Using historical collection specimen records to fill knowledge gaps

Estimates of pollinator declines are lacking for most countries worldwide (Figure 1). The use of historic collection data may be the most effective tool for filling these gaps. The core aim of museums is to conserve and curate historic collections. Thus, they serve as a precious repository for specimens, and at the same time, often ensure higher quality taxonomic identification. Yet, the major bottleneck for researchers wanting to use these data is the lack of digitization. Digitizing old collection specimens is not a trivial task and requires expertise to (i) ensure proper taxonomic identification [30–32], (ii) geo-locate the coordinates of collection events (e.g. <http://www.geonames.org>) and (iii) store the data in a properly curated database [33]. Undertaking this process for tens or hundreds of thousands of museum collection specimens can be a daunting task and requires specialized personnel. While some tasks can only be undertaken by people with specialist skills (e.g., taxonomists), new technologies and citizen science can speed up the collection digitization process. High resolution photos of specimens and associated labels can be uploaded to the internet, where the task of image transcription can be distributed across hundreds or thousands of volunteers (e.g., <https://www.zooniverse.org/>). In addition, new algorithms have been created that allow location geo-referencing based on vernacular names (e.g. <https://geoparser.io>). However, achieving this requires adequate funding [34].

Where digitization has been completed, the data provide a rich source of information, allowing assessment of the current status and long-term trends of pollinator populations [17,19,35]. This is despite the fact that museum collections often have a number of biases, including unknown sampling effort, personal interests of collectors and the curatorial

1
2
3 174 techniques used. For example, collectors tend to target rare or unusual over common taxa,
4
5 175 discard damaged individuals or only accession a certain number of individuals. In addition,
6
7 176 collections are often made opportunistically, leading to a spatial biases where difficult to
8
9 177 access areas are under-sampled or conversely, where samples are biased towards easily
10
11 178 accessed locations (e.g., towns/cities and/or roadsides). Further, museum collection data
12
13 179 can only be used to determine where species are present and not where they are absent.
14
15 180 However, given adequate sample sizes and appropriate statistical techniques, most biases
16
17 181 can be accounted for [e.g. 17,36,37].
18
19
20
21
22
23

24 182 **The way forward: Prioritizing the low hanging fruit.**

25
26
27 183 As we have shown, there is a paucity of countries for which historical data is available
28
29 184 (Figure 1), and hence can be used as baseline for assessing pollinator population declines.
30
31 185 While ideally one would aim to digitize all museum collection records, this is unlikely in the
32
33 186 near future, predominantly due to funding constraints. Here we show how researchers can
34
35 187 optimize the use of historical collection data to assess long-term pollinator population
36
37 188 changes.
38
39
40
41
42

43 189 GBIF (<https://www.gbif.org/>) is a central repository for global species occurrence data.
44
45 190 Much of these data come from museums, private collections and government research
46
47 191 institutes, but several other sources are also integrated. In combination with the popular
48
49 192 statistical language R [38], GBIF can be directly queried into your computer [39] and data
50
51 193 availability can be checked for the region of interest. Focusing on bee taxa, we show here
52
53 194 the number of modern and historic bee records currently available for different countries
54
55
56
57
58
59
60

1
2
3 195 (Figure 2a, Sup mat 2). Thirty-seven countries have more than 1800 records in each time
4
5 196 period, making these data potentially analyzable without further data collection effort (see
6
7
8 197 Figure 2b and c for an initial exploration). However, a proper analysis of this dataset would
9
10 198 require a careful inspection of the data, as we detail below for two specific countries (Spain
11
12 199 and New Zealand). In contrast, most countries fall short in one or both axes of Figure 2a.
13
14 200 For example, a variety of countries located in different continents such as Switzerland, Sri
15
16 201 Lanka, Nicaragua or Zimbabwe have a decent number of recent records, but lack historical
17
18 202 collections. In this cases, researchers should prioritize the digitalization of old material
19
20 203 before embarking on data analyses. For this end, it is also important to note that historical
21
22 204 records are not always vouchered in local museums (i.e., many European and USA
23
24 205 museums contain large collections of pollinators from other countries). Finally, it's
25
26 206 remarkable that more than 192 countries have less than 1000 records for each of both time
27
28 207 periods, making them poor candidates for analysing long-term pollinator population
29
30 208 trends. Aside from bees, similar exploratory analyses can easily be conducted for other
31
32 209 taxa.
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60




Figure 2. Exploration of available data for bee records showing: (a) The number of bee occurrences before 1980 and after 1980 in GBIF for each country. The upper right quadrant (records in black) contains well covered countries with New Zealand (NZ) and Spain (ES) marked in red (see below). For well covered countries, we show preliminary comparisons of the rarefied number of species in both time periods and show that for most countries (21 out of 28) the number of species recorded is slightly lower (average of 10% richness decline; red lines) for recent time periods (b). Data is log transformed for visualization purposes. A more careful analysis of this data would help complete the map of global declines (c). In this map we plot the % change in species recorded in GFIF for the available

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

countries to show the potential geographic coverage. Note that this data is likely to contain strong undetected biases, as we explore below.

As stated above, once historical collection datasets are made available, researchers must identify any potential biases. We explore this process with two contrasting dataset examples (Spain and New Zealand). In the Spanish dataset, most of the data comes from a few specific locations and was collected by a few specific teams. Hence, the geographical coverage is not representative. Even worst, historical and modern collections do not overlap spatially, making any inference impossible to interpret. In this case, we contacted the original collectors of the historical data to define their sampling protocols. We then resurveyed the same sites (35 years after the original surveys) using the same sampling protocols. In contrast, the New Zealand dataset includes a wide suite of collectors and collection locations but shows no obvious biases in geographical and taxonomic coverage through time. We complemented GBIF data with further museum collections for bees and flies and analyze the regional richness changes through time. For these two case studies, we provide annotated R scripts as examples of analysis for different dataset types (Sup Mat 3). These different analytical approaches allow us to reveal long-term trends in pollinator populations for regions with contrasting sampling histories. We hope this resource will encourage researchers to analyse data for regions where current information on pollinator declines is lacking.

241 Case study one: Spain

242 Spain provides an interesting study system because its natural habitats have been
243 transformed extensively by humans over a long time period, but land-use is not as
244 intensive compared with many other European countries. In addition, Spain is a bee
245 diversity hotspot (Figure 1a) and maintains a relatively heterogeneous landscape. Spain
246 has already digitalized a large amount of pollinator occurrence data for both historic and
247 recent periods (Figure 2a). However, visual inspection of the data revealed clustering
248 around a few localities. Further, historic records did not spatially match recent records,
249 making comparisons difficult. For this dataset, most of the historic records were located
250 around Valladolid and were collected by Enrique Asensio and collaborators. There has
251 been no recent sampling of bees in this area. However, we found that Enrique
252 systematically sampled six independent locations and that additional historical data were
253 available at the “Museo Nacional de Ciencias Naturales” and other minor collections.
254 Digitization of these records, along with a re-survey of the original sampling locations
255 provided an excellent dataset for a before and after comparison of bee communities.

256 In brief, after cleaning taxonomic names for possible typos and synonyms using the taxize
257 package [40], we checked for sampling completeness for both time periods and compared
258 rarefied species richness for each site before and after 1980 with a paired t-test
259 (rarefaction at 1000 specimens). We found that there were a reduced number of species at
260 sites after 1980 (mean difference 20.27 species; 95 percent confidence interval: -1.03,
261 41.58; $t = 2.44$, $df = 5$, $P = 0.06$). However, this trend was highly dependent on site identity,
262 as two out of six sites showed no richness declines. Interestingly, these two localities were

the two that has experienced less land use changes (both are natural areas embedded into agro-ecosystems). In contrast the other 4 localities suffered large urban or agricultural intensification. In addition, species lost in the re-surveys are not a random selection of species, but are clustered in a few genera. For example, Andrenidae and their parasites (e.g. *Nomada*) showed the strongest declines whereas Halictidae tend to be more stable (Sup mat 4). This pattern of winners and losers of land use intensification is in accordance with findings elsewhere [17], indicating that some clades are more sensitive to disturbance than others.




Figure 3. Comparison of historic collections (1980's) and modern re-surveys (2016) of the rarefied richness of bees at six Spanish localities.

Case study two: New Zealand

In contrast to Spain, New Zealand is an isolated oceanic archipelago, with a distinctive pollinator biota and a unique history of human occupation. Much of New Zealand's

pollinator fauna is also relatively depauperate. For example, New Zealand has only 27 native bee species [41], which is a fraction of nearby Australia's c. 1600 species [42]. However, New Zealand has a surprisingly high diversity of flies (Diptera), which are important pollinators in many ecosystems [43]. Thus, New Zealand provides a unique system to study long-term changes in pollinator communities, and is unlike continental Europe and the US, which have been the focus of an overwhelming majority of pollinator decline studies.

In global terms, human colonisation of New Zealand was relatively recent (c. 740 y) [44]. Before human arrival, New Zealand was predominately forested, but has since been dramatically altered by people. Early Māori settlers cleared forests by burning and more recently, European colonists cleared large tracts of remaining forests and drained low-lying wetlands for agriculture, mostly before 1900 [45]. Therefore, human activity likely affected pollinator communities in New Zealand long before surveys and specimen collections began. Nevertheless, we can use museum records to identify trends in pollinator communities during New Zealand's more recent history.

We used New Zealand bee collection records gathered from multiple sources, including university, research institute, museum and private collections. Collection records from the New Zealand Arthropod Collection (NZAC) are freely available online (<https://scd.landcareresearch.co.nz/>). Fly pollinator data was obtained from three participating New Zealand museums and covers two families (Calliphoridae and Syrphidae) that contain important fly pollinators. Collections for the bee and fly datasets span over 100 years (early 1900s to late 2000s).

1
2
3 299 We followed protocols outlined in [17] to analyse the New Zealand data at the regional
4
5 300 level. First, we filtered our original datasets so that data used for analyses only included
6
7 301 independent collection events. To do this, we removed specimens collected at the same
8
9 302 location, on the same date, and by the same collector. We found our data had reasonable
10
11 303 coverage across time periods, although there was a peak in collection occurrences from
12
13 304 1960-1980. Further exploration of the New Zealand native bee data raised doubts on
14
15 305 collection completeness in records prior to 1970, so we removed these records from
16
17 306 further analyses. We accounted for differences in collection effort through binning
18
19 307 collection records by time so that each bin had a similar number of records but a different
20
21 308 number of years. We then estimated richness for each time period bin by rarefying all bins
22
23 309 to an equal number of specimens and calculated the mean species richness \pm SE for each
24
25 310 bin. Finally, we estimated the significance of change in richness using a permutation test
26
27 311 that randomly reordered time periods and calculated the correlation between time period
28
29 312 and species richness. Thus, reported P-values were the proportion of permutations that
30
31 313 had higher or lower correlations compared to the correlation between richness and the
32
33 314 actual chronological time period sequence.
34
35
36
37
38
39
40
41 315 Second, to determine if the probability of finding a species in the collection changed over
42
43 316 time, we used a general linear model with a binomial distribution and a logit link. For
44
45 317 species that showed overdispersion, we used a quasi-binomial distribution. Further, we
46
47 318 only included species in this analysis for which we had 30 or more records. To account for
48
49 319 differences in sampling effort between years, we weighted each year by the total number of
50
51 320 samples collected that year.
52
53
54
55
56
57
58
59
60

We found that rarefied richness for native bees was stable through time. Exotic bees showed an increase in rarefied richness, but this trend was non-significant (P-value for both natives and exotic bees > 0.05). In contrast, native fly richness declined, whereas exotic fly richness increased, although results for these groups were also non-significant (P-values for both groups > 0.05). Note that rarefied richness is sensitive to species evenness, so increases in rarefied richness over time may actually indicate increased species evenness and vice-versa for decreased richness.




Figure 4. Changes in rarefied species richness for different pollinator groups in New Zealand over time. All trends were non-significant ($\alpha = 0.05$).

However, at the species level, we found that 11 out of 27 bee species increased in relative occurrence over time (10 native and one exotic) and three bee species declined in relative occurrence (one native and two exotic) (Figure 3). Interestingly, the two exotic bee species that declined in relative occurrence were both in the genus *Bombus*, which were intentionally introduced into New Zealand for the pollination of crops. Native bees that increased in relative occurrence were mostly from the genus *Leioproctus*, which are medium sized, ground nesting solitary bees. Only one out of 14 fly species increased in relative occurrence, which was exotic, whereas four species decreased in occurrence (three native and one exotic). Native flies that decreased in relative occurrence were all Syrphidae in the genus *Helophilus*.




Figure 5. Model estimated changes (± 1 SE) in the relative occurrence frequency of different New Zealand bee and fly species in museum collections over time.

Beyond species occurrences

A recent study found that more than 90% of the papers investigating pollinator responses to land-use change focused solely on richness and abundance descriptors [9]. But in addition to local (*alpha*) diversity and regional (*gamma*) diversity, researchers need to assess changes in turnover between sites (*beta* diversity). Environmental changes often result in a few “winner” species and many “losers” species [17]. Identifying winners and losers is critical as the few winners are often exotic and represent a subset of traits that facilitate survival in highly modified environments [46]. These changes can have important effects for pollination of native plant species and crops [47].

In addition, museum specimen collections can provide much more information besides species occurrence records, given that such information is recorded when digitizing collections. This is particularly important for identifying mechanisms of decline and adaptation. For example, recording the date of collection is particularly important for tracking of phenological advances congruent with contemporary climate change [48]. In addition, pollinator specimen labels often include information about the host plant on which the specimen was collected. This information critical for understanding past and present species interactions [49]. Aside from this information, bee specimens often contain pollen loads trapped on hairs, from which past visitation events can be identified [50]. Finally, museum specimens can be measured to track evolutionary changes by measuring

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

the traits of specimen. This approach has been already used to investigate tongue length [51] and body size [52] changes in response to climate and land-use change. Finally, plant herbariums can also contain indirect evidence of pollinator and pollination declines [53], a basic information for linking pollinator declines with its consequences for ecosystem functioning.

Conclusions

Unleashing the power of museum collection data to answer pressing ecological and evolutionary questions is at our hands, but requires the coordinated effort of many actors. Using two case studies, we show that strong collaboration between museum curators and ecologists is key to understanding data and treating it appropriately. To progress our understanding of the global pollination crisis, researchers and curators must aim to digitize museum collection data and make it readily available in a format that is widely accessible. Centralization of regional and national museum collection data in existing global platforms, such as GBIF, would facilitate free and widespread access. However, datasets could also be stored in alternative webpages or database repositories (e.g., university and museum webpages or Dryad) providing they are thoroughly documented and easily retrieved and combined with other datasets using open science tools [54].

We must revolutionize the way that researchers collaborate with museums, in order to foster healthy bidirectional relationships. For example, ecological researchers collect massive amounts of specimens, but these are often inappropriately vouchered [55,56], rendering them less useful for future research. To improve this process, strong

384 communication between museums and researchers is required. However, this can only be
385 achieved with adequate funding and recognition that accurate data recording and long-
386 term preservation are critical for research [57].

387 To identify global trends in pollinator declines we require robust data, collected from
388 diverse geographic regions. It is also crucial that these data are analysed appropriately.
389 This requires researches to identify biases and to any fill taxonomic and geographic gaps
390 where possible. We need to place increased emphasis on quantifying pollinator declines in
391 regions outside of the US and Europe, and for pollinator groups other than bees. For the US
392 and Europe, there have been few regional bee extinctions [17,22] but in disturbed
393 ecosystems, declines are widespread [15,18]. For most other pollinator taxa and regions
394 throughout the world we know almost nothing. Moving forward, the first step for many
395 taxa will be to identify and describe species. Only then can we begin to document pollinator
396 declines.

397 **Acknowledgements**

398 We thank Curro Molina, Carola Warner, Patrick McQuinn, and Crona McMonagle for data
399 entry and Gregorio Aguado for carrying out the Spanish re-sampling. We thank Barry
400 Donovan for providing New Zealand bee collection records and E. Asensio for sharing his
401 historical data and knowledge. We thank the "Museo Nacional de Ciencias Naturales",
402 specially Mercedes Paris, ITACyL (Instituto Tecnológico Agrario de Castilla y León),
403 Canterbury Museum, the New Zealand Arthropod Collection and the Museum of New
404 Zealand Te Papa Tongarewa for access to historical collections. IB was funded by a

1
2
3 405 “Fundación Banco Bilbao Vizcaya Argentaria” (FBBVA) project. DW was funded through
4
5 406 Landcare Research within the Characterising New Zealand’s Land Biota Portfolio.
6
7
8 407
9
10
11 408 *The datasets supporting this article have been uploaded as part of the supplementary*
12
13 409 *material and will be deposited at dryad or Figshare upon acceptance.*
14
15
16
17 410 *We have no competing interests*
18
19
20
21

22 411 **References**
23
24
25
26 412

27
28 413 1. Ollerton J, Winfree R, Tarrant S. 2011 How many flowering plants are pollinated by
29
30 414 animals? *Oikos* **120**, 321–326.
31
32
33
34 415 2. Klein AM, Vaissiere BE, Cane JH, Steffan-Dewenter I, Cunningham SA, Kremen C,
35
36 416 Tscharntke T. 2007 Importance of pollinators in changing landscapes for world crops.
37
38 417 *Proceedings of the Royal Society B: Biological Sciences* **274**, 303–313.
39
40
41
42 418 3. Buchmann SL, Nabhan GP. 1996 *The forgotten pollinators*.
43
44
45 419 4. National Research Council. 2007 *Status of Pollinators in North America*. Washington, DC:
46
47 420 The National Academies Press. (doi:[doi:10.17226/11761](https://doi.org/10.17226/11761))
48
49
50
51 421 5. Goulson D, Nicholls E, Botías C, Rotheray EL. 2015 Bee declines driven by combined
52
53 422 stress from parasites, pesticides, and lack of flowers. *Science* **347**, 1255957.
54
55
56
57
58
59
60

- 423 6. Potts SG, Biesmeijer JC, Kremen C, Neumann P, Schweiger O, Kunin WE. 2010 Global
424 pollinator declines: Trends, impacts and drivers. *Trends in Ecology and Evolution* **25**, 345–
425 353.
- 426 7. Nieto A *et al.* 2014 European red list of bees. *Luxembourg: Publication Office of the*
427 *European Union* **98**.
- 428 8. Ollerton J. 2017 Pollinator Diversity: Distribution, Ecological Function, and Conservation.
429 *Annual Review of Ecology, Evolution, and Systematics* **48**, 353–376. (doi:[10.1146/annurev-ecolsys-110316-022919](https://doi.org/10.1146/annurev-ecolsys-110316-022919))
- 431 9. Winfree R, Bartomeus I, Cariveau DP. 2011 Native pollinators in anthropogenic systems.
432 *Annual Review of Ecology and Systematics* **42**, 1–21.
- 433 10. Martin LJ, Blossey B, Ellis E. 2012 Mapping where ecologists work: biases in the global
434 distribution of terrestrial ecological observations. *Frontiers in Ecology and the Environment*
435 **10**, 195–201. (doi:[doi:10.1890/110154](https://doi.org/10.1890/110154))
- 436 11. Archer CR, Pirk CWW, Carnevalheiro LG, Nicolson SW. 2014 Economic and ecological
437 implications of geographic bias in pollinator ecology in the light of pollinator declines. *Oikos*
438 **123**, 401–407. (doi:[doi:10.1111/j.1600-0706.2013.00949.x](https://doi.org/10.1111/j.1600-0706.2013.00949.x))
- 439 12. De Palma A *et al.* 2016 Predicting bee community responses to land-use changes:
440 Effects of geographic and taxonomic biases. *Scientific Reports* **6**, 31153.
- 441 13. Franca F, Louzada J, Korasaki V, Griffiths H, Silveira JM, Barlow J. 2016 Do space-for-
442 time assessments underestimate the impacts of logging on tropical biodiversity? An
443 Amazonian case study using dung beetles. *Journal of Applied Ecology* **53**, 1098–1105.

1
2
3 444 14. De Palma A *et al.* 2018 Challenges With Inferring How Land-Use Affects Terrestrial
4
5 445 Biodiversity: Study Design, Time, Space and Synthesis. *Next Generation Biomonitoring* **58**,
6
7 446 163.
8
9
10
11 447 15. De Palma A, Kuhlmann M, Bugter R, Ferrier S, Hoskins AJ, Potts SG, Roberts SP,
12
13 448 Schweiger O, Purvis A. 2017 Dimensions of biodiversity loss: Spatial mismatch in land-use
14
15 449 impacts on species, functional and phylogenetic diversity of European bees. *Diversity and*
16
17 450 *Distributions* **23**, 1435–1446.
18
19
20
21 451 16. Swaay CA van, Nowicki P, Settele J, Strien AJ van. 2008 Butterfly monitoring in Europe:
22
23 452 methods, applications and perspectives. *Biodiversity and Conservation* **17**, 3455–3469.
24
25
26
27 453 17. Bartomeus I, Ascher JS, Gibbs J, Danforth BN, Wagner DL, Hedtke SM, Winfree R. 2013
28
29 454 Historical changes in northeastern US bee pollinators related to shared ecological traits.
30
31 455 *Proceedings of the National Academy of Sciences of the United States of America* **110**, 4656–
32
33 456 4660. (doi:[10.1073/pnas.1218503110](https://doi.org/10.1073/pnas.1218503110))
34
35
36
37 457 18. Biesmeijer JC *et al.* 2006 Parallel Declines in Pollinators and Insect-Pollinated Plants in
38
39 458 Britain and the Netherlands. *Science* **313**, 351–354. (doi:[10.1126/science.1127863](https://doi.org/10.1126/science.1127863))
40
41
42
43 459 19. Scheper J, Reemer M, Kats R van, Ozinga WA, Linden GT van der, Schaminée JH, Siepel H,
44
45 460 Kleijn D. 2014 Museum specimens reveal loss of pollen host plants as key factor driving
46
47 461 wild bee decline in The Netherlands. *Proceedings of the National Academy of Sciences* **111**,
48
49 462 17552–17557.
50
51
52
53
54
55
56
57
58
59
60

- 463 20. Carvalheiro LG *et al.* 2013 Species richness declines and biotic homogenisation have
464 slowed down for NW-European pollinators and plants. *Ecology Letters* **16**, 870–878.
465 (doi:[10.1111/ele.12121](https://doi.org/10.1111/ele.12121))
- 466 21. Colla SR *et al.* 2012 Documenting Persistence of Most Eastern North American Bee
467 Species (Hymenoptera: Apoidea: Anthophila) to 19902009. *Journal of the Kansas*
468 *Entomological Society* **85**, 14–22. (doi:[10.2317/JKES110726.1](https://doi.org/10.2317/JKES110726.1))
- 469 22. Ollerton J, Erenler H, Edwards M, Crockett R. 2014 Extinctions of aculeate pollinators in
470 Britain and the role of large-scale agricultural changes. *Science* **346**, 1360–1362.
- 471 23. Burkle LA, Marlin JC, Knight TM. 2013 Plant-Pollinator Interactions over 120 Years:
472 Loss of Species, Co-Occurrence, and Function. *Science* **339**, 1611–1615.
473 (doi:[10.1126/science.1232728](https://doi.org/10.1126/science.1232728))
- 474 24. Bartomeus I, Winfree R. 2013 Pollinator declines: reconciling scales and implications
475 for ecosystem services. *F1000Research* **2**.
- 476 25. Dupont YL, Damgaard C, Simonsen V. 2011 Quantitative historical change in bumblebee
477 (*Bombus* spp.) assemblages of red clover fields. *PLoS ONE* **6**.
478 (doi:[10.1371/journal.pone.0025172](https://doi.org/10.1371/journal.pone.0025172))
- 479 26. Grixti JC, Wong LT, Cameron SA, Favret C. 2009 Decline of bumble bees (*Bombus*) in the
480 North American Midwest. *Biological Conservation* **142**, 75–84.
481 (doi:[10.1016/j.biocon.2008.09.027](https://doi.org/10.1016/j.biocon.2008.09.027))
- 482 27. Maes D, Van Dyck H. 2001 Butterfly diversity loss in Flanders (north Belgium): Europe's
483 worst case scenario? *Biological conservation* **99**, 263–276.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

28. Wenzel M, Schmitt T, Weitzel M, Seitz A. 2006 The severe decline of butterflies on western German calcareous grasslands during the last 30 years: A conservation problem. *Biological Conservation* **128**, 542–552. (doi:[10.1016/j.biocon.2005.10.022](https://doi.org/10.1016/j.biocon.2005.10.022))

29. Morales CL, Arbetman MP, Cameron SA, Aizen MA. 2013 Rapid ecological replacement of a native bumble bee by invasive species. *Frontiers in Ecology and the Environment* **11**, 529–534.

30. Meier R, Dikow T. 2004 Significance of Specimen Databases from Taxonomic Revisions for Estimating and Mapping the Global Species Diversity of Invertebrates and Repatriating Reliable Specimen Data. *Conservation Biology* **18**, 478–488. (doi:[doi:10.1111/j.1523-1739.2004.00233.x](https://doi.org/10.1111/j.1523-1739.2004.00233.x))

31. Huber JT. 1998 The importance of voucher specimens, with practical guidelines for preserving specimens of the major invertebrate phyla for identification. *Journal of Natural History* **32**, 367–385.

32. Thomson SA *et al.* 2018 Taxonomy based on science is necessary for global conservation. *PLoS biology* **16**, e2005075.

33. Hart EM *et al.* 2016 Ten simple rules for digital data storage. *PLoS computational biology* **12**, e1005097.

34. Wiggins GB, Marshall SA, Downes JA. 1991 Importance of research collections of terrestrial arthropods. A brief prepared by the Biological Survey of Canada (Terrestrial Arthropods). *Bulletin of the Entomological society of Canada* **23**, 16.

- 504 35. Young BE, Auer S, Ormes M, Rapacchiolo G, Schweitzer D, Sears N. 2017 Are pollinating
505 hawk moths declining in the Northeastern United States? An analysis of collection records.
506 *PloS one* **12**, e0185683.
- 507 36. Keil P, Biesmeijer JC, Barendregt A, Reemer M, Kunin WE. 2011 Biodiversity change is
508 scale-dependent: an example from Dutch and UK hoverflies (Diptera, Syrphidae).
509 *Ecography* **34**, 392–401.
- 510 37. Pearce JL, Boyce MS. 2006 Modelling distribution and abundance with presence-only
511 data. *Journal of applied ecology* **43**, 405–412.
- 512 38. R Core Team. 2017 *R: A language and environment for statistical computing*. R
513 Foundation for Statistical Computing, Vienna, Austria.
- 514 39. Chamberlain S, Ram K, Barve V, McGlinn D. 2014 rgbif: Interface to the Global
515 Biodiversity Information Facility API. *R package version 0.7 7*.
- 516 40. Chamberlain S, Szoecs E, Foster Z, Arendsee Z. 2018 Package 'taxize'.
- 517 41. Donovan BJ. 2007 Apoidea (Insecta: Hymenoptera). *Fauna of New Zealand* **57**, 295.
- 518 42. Ascher JS, Pickering J. 2015 Discover Life bee species guide and world checklist
519 (Hymenoptera: Apoidea: Anthophila).
- 520 43. Newstrom L, Robertson AW. 2005 Progress in understanding pollination systems in
521 New Zealand. *New Zealand Journal of Botany* **43**, 1–59.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

44. Wilmshurst JM, Anderson AJ, Higham TF, Worthy TH. 2008 Dating the late prehistoric dispersal of Polynesians to New Zealand using the commensal Pacific rat. *Proceedings of the National Academy of Sciences* **105**, 7676–7680.

45. Perry GL, Wilmshurst JM, McGlone MS. 2014 Ecology and long-term history of fire in New Zealand. *New Zealand Journal of Ecology*, 157–176.

46. Stavert JR, Pattemore DE, Gaskett AC, Beggs JR, Bartomeus I. 2017 Exotic species enhance response diversity to land-use change but modify functional composition. *Proceedings of the Royal Society B: Biological Sciences* **284**. (doi:[20170788](https://doi.org/10.1098/rspb.2017.0788))

47. Stavert JR, Pattemore DE, Bartomeus I, Gaskett AC, Beggs JR. 2018 Exotic flies maintain pollination services as native pollinators decline with agricultural expansion. *Journal of Applied Ecology*

48. Bartomeus I, Ascher JS, Wagner D, Danforth BN, Colla S, Kornbluth S, Winfree R. 2011 Climate-associated phenological advances in bee pollinators and bee-pollinated plants. *Proceedings of the National Academy of Sciences of the United States of America* **108**, 20645–20649.

49. Tylianakis JM, Didham RK, Bascompte J, Wardle DA. 2008 Global change and species interactions in terrestrial ecosystems. *Ecology Letters* **11**, 1351–1363. (doi:[10.1111/j.1461-0248.2008.01250.x](https://doi.org/10.1111/j.1461-0248.2008.01250.x))

50. Kleijn D, Raemakers I. 2008 A retrospective analysis of pollen host plant use by stable and declining bumble bee species. *Ecology* **89**, 1811–1823.

51. Miller-Struttmann NE *et al.* 2015 Functional mismatch in a bumble bee pollination mutualism under climate change. *Science* **349**, 1541–1544.
52. Oliveira MO, Freitas BM, Scheper J, Kleijn D. 2016 Size and sex-dependent shrinkage of Dutch bees during one-and-a-half centuries of land-use change. *PloS one* **11**, e0148983.
53. Pauw A, Hawkins JA. 2011 Reconstruction of historical pollination rates reveals linked declines of pollinators and plants. *Oikos* **120**, 344–349.
54. Dallas T. 2016 helminthR: an R interface to the London Natural History Museum's host-parasite database. *Ecography* **39**, 391–393.
55. Packer L, Monckton SK, Onuferko TM, Ferrari RR. 2018 Validating taxonomic identifications in entomological research. *Insect Conservation and Diversity* **11**, 1–12.
56. Wheeler TA. 2003 *The Role of Voucher Specimens in Validating Faunistic and Ecological Research: A Brief*. Biological Survey of Canada (Terrestrial Arthropods).
57. Ward DF, Leschen RA, Buckley TR. 2015 More from ecologists to support natural history museums. *Trends in ecology & evolution* **30**, 373–374.