

– an invitation –

On the scope of the dynamical method in commutative algebra

Arbeitstagung Verona–München 2023

Munich
January 18th, 2022

Ingo Blechschmidt

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

A primer to the dynamical method


Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. Assume not.

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. Assume not. Then there is a maximal ideal \mathfrak{m} .



maximal among the proper ideals


$$\neg(1 \in \mathfrak{m})$$

$$x \in \mathfrak{m} \quad \vee \quad 1 \in \mathfrak{m} + (x)$$

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. **Assume not.** Then there is a **maximal ideal** \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} .




$$\neg(1 \in \mathfrak{m})$$
$$x \in \mathfrak{m} \quad \vee \quad 1 \in \mathfrak{m} + (x)$$

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. **Assume not.** Then there is a **maximal ideal** \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} . Since A/\mathfrak{m} is a field, this is a contradiction to basic linear algebra. \square



$$\neg(1 \in \mathfrak{m})$$
$$x \in \mathfrak{m} \quad \vee \quad 1 \in \mathfrak{m} + (x)$$

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. Assume not. Then there is a maximal ideal \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} . Since A/\mathfrak{m} is a field, this is a contradiction to basic linear algebra. \square

Three questions

- 1 *Is there also a constructive proof?*

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. **Assume not.** Then there is a **maximal ideal** \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} . Since A/\mathfrak{m} is a field, this is a contradiction to basic linear algebra. \square

Three questions

1 *Is there also a constructive proof?*

Yes, by completeness for coherent logic.

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. Assume not. Then there is a maximal ideal \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} . Since A/\mathfrak{m} is a field, this is a contradiction to basic linear algebra. \square

Three questions

1 *Is there also a constructive proof?*

Yes, by completeness for coherent logic.

2 *Do we know a constructive proof?*

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. **Assume not.** Then there is a **maximal ideal** \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} . Since A/\mathfrak{m} is a field, this is a contradiction to basic linear algebra. \square

Three questions

1 *Is there also a constructive proof?*

Yes, by completeness for coherent logic.

2 *Do we know a constructive proof?*

Write $M = \begin{pmatrix} x \\ y \end{pmatrix}$. By surjectivity, have u, v with $u \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $v \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Hence $1 = (vy)(ux) = (uy)(vx) = 0$.

A primer to the dynamical method

Thm. Let M be a surjective matrix with more rows than columns over a ring A . Then $1 = 0$ in A .

Proof. Assume not. Then there is a maximal ideal \mathfrak{m} . The matrix is surjective over A/\mathfrak{m} . Since A/\mathfrak{m} is a field, this is a contradiction to basic linear algebra. \square

Three questions

1 *Is there also a constructive proof?*

Yes, by completeness for coherent logic.

2 *Do we know a constructive proof?*

Write $M = \begin{pmatrix} x \\ y \end{pmatrix}$. By surjectivity, have u, v with $u \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix}$ and $v \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$. Hence $1 = (vy)(ux) = (uy)(vx) = 0$.

3 *Can the constructive proof be **extracted** from the classical one?*

Yes, by the dynamical method (and others).

The status of maximal ideals

- ▶ In **classical mathematics**, every ring has a maximal ideal.
 - *Zorn's lemma*

The status of maximal ideals

- ▶ In **classical mathematics**, every ring has a maximal ideal.
 - *Zorn's lemma*
- ▶ Without **Zorn**, every **countable** ring has a maximal ideal.
 - *Iterative construction given enumeration x_0, x_1, \dots [Krull 1929]:*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \begin{cases} \mathfrak{m}_n + (x_n), & \text{if } 1 \notin \mathfrak{m}_n + (x_n), \\ \mathfrak{m}_n, & \text{else.} \end{cases}$$

The status of maximal ideals

- ▶ In **classical mathematics**, every ring has a maximal ideal.

- *Zorn's lemma*

- ▶ Without **Zorn**, every **countable** ring has a maximal ideal.

- *Iterative construction given enumeration x_0, x_1, \dots [Krull 1929]:*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \begin{cases} \mathfrak{m}_n + (x_n), & \text{if } 1 \notin \mathfrak{m}_n + (x_n), \\ \mathfrak{m}_n, & \text{else.} \end{cases}$$

- *Also constructively! [Krivine 1996], [Berardi–Valentini 2004]*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \mathfrak{m}_n + \underbrace{(\{x_n \mid 1 \notin \mathfrak{m}_n + (x_n)\})}_{\text{a certain subsingleton set}}$$

The status of maximal ideals

- ▶ In **classical mathematics**, every ring has a maximal ideal.
 - *Zorn's lemma*
- ▶ Without **Zorn**, every **countable** ring has a maximal ideal.
 - *Iterative construction given enumeration x_0, x_1, \dots [Krull 1929]:*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \begin{cases} \mathfrak{m}_n + (x_n), & \text{if } 1 \notin \mathfrak{m}_n + (x_n), \\ \mathfrak{m}_n, & \text{else.} \end{cases}$$

- *Also constructively! [Krivine 1996], [Berardi-Valentini 2004]*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \mathfrak{m}_n + \underbrace{(\{x_n \mid 1 \notin \mathfrak{m}_n + (x_n)\})}_{\text{a certain subsingleton set}}$$



The status of maximal ideals

- ▶ In **classical mathematics**, every ring has a maximal ideal.

- *Zorn's lemma*

- ▶ Without **Zorn**, every **countable** ring has a maximal ideal.

- *Iterative construction given enumeration x_0, x_1, \dots [Krull 1929]:*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \begin{cases} \mathfrak{m}_n + (x_n), & \text{if } 1 \notin \mathfrak{m}_n + (x_n), \\ \mathfrak{m}_n, & \text{else.} \end{cases}$$

- *Also constructively! [Krivine 1996], [Berardi–Valentini 2004]*

$$\mathfrak{m}_0 = \{0\}, \quad \mathfrak{m}_{n+1} = \mathfrak{m}_n + \underbrace{(\{x_n \mid 1 \notin \mathfrak{m}_n + (x_n)\})}_{\text{a certain subsingleton set}}$$

The quotient A/\mathfrak{m} is a residue field: noninvertible implies zero.

$$\neg(1 \in \mathfrak{m})$$

$$\neg(1 \in \mathfrak{m} + (x)) \implies x \in \mathfrak{m}$$

Forcing

► Forcing in commutative algebra

- $A \rightsquigarrow A[X]$ adjoining an indeterminate
- $A \rightsquigarrow A[x^{-1}]$ forcing an element to become invertible
- $A \rightsquigarrow A/(x)$ forcing an element to become zero


► Forcing in classical set theory

- $V \rightsquigarrow V[G]$ adjoining a generic filter of a forcing poset \mathbb{P}
e.g. adding a cardinal between \aleph_0 and \mathfrak{c} ,
adding a random real,
collapsing two cardinals, ...

► Forcing in constructive mathematics

- $V \rightsquigarrow V^{\neg\neg}$ forcing LEM
- $V \rightsquigarrow \text{Sh}(X)$ adjoining a generic point of X
- $V \rightsquigarrow V[\mathbb{T}]$ adjoining a generic \mathbb{T} -model


Parametrized mathematics



local “Every real symmetric matrix does have an eigenvector.” ✓

global “For every continuous family of symmetric matrices,
eigenvectors can locally be picked continuously.” ?

Parametrized mathematics




local “Every real symmetric matrix does have an eigenvector.” ✓

global “For every continuous family of symmetric matrices, eigenvectors can locally be picked continuously.” ?

“Let X be a topological space and let $A : X \rightarrow M_n^{\text{sym}}(\mathbb{R})$ be a continuous map to the space of symmetric $(n \times n)$ -matrices. Then there is an open covering $\bigcup_{i \in I} U_i$ of X such that for all indices $i \in I$, there is a continuous map $v : U_i \rightarrow \mathbb{R}^n$ such that for all $x \in U_i$, the vector $v(x)$ is an eigenvector of $A(x)$.”

Parametrized mathematics




local “Every real symmetric matrix does have an eigenvector.” ✓

global “For every continuous family of symmetric matrices, eigenvectors can locally be picked continuously.” ✗

“Let X be a topological space and let $A : X \rightarrow M_n^{\text{sym}}(\mathbb{R})$ be a continuous map to the space of symmetric $(n \times n)$ -matrices. Then there is an open covering $\bigcup_{i \in I} U_i$ of X such that for all indices $i \in I$, there is a continuous map $v : U_i \rightarrow \mathbb{R}^n$ such that for all $x \in U_i$, the vector $v(x)$ is an eigenvector of $A(x)$.”

Parametrized mathematics




local “Every real symmetric matrix does **not not** have an eigenvector.” ✓

global “For every continuous family of symmetric matrices,
on a dense open eigenvectors can locally be picked continuously.” ✓

“Let X be a topological space and let $A : X \rightarrow M_n^{\text{sym}}(\mathbb{R})$ be a continuous map to the space of symmetric $(n \times n)$ -matrices. Then there is an open covering $\bigcup_{i \in I} U_i$ **of a dense open subset** $U \subseteq X$ such that for all indices $i \in I$, there is a continuous map $v : U_i \rightarrow \mathbb{R}^n$ such that for all $x \in U_i$, the vector $v(x)$ is an eigenvector of $A(x)$.”


Parametrized mathematics



local “Let M be a finitely generated module over a field k .
Then M is finite free.” ✓

global “Let M be a finitely generated module over a ring A .
Then M^\sim is finite locally free.” ?

Parametrized mathematics




local “Let M be a finitely generated module over a field k .
Then M is finite free.” ✓

global “Let M be a finitely generated module over a ring A .
Then M^\sim is finite locally free.” ?

“Let M be a finitely generated module over an arbitrary commutative ring A . Then there is a partition $1 = f_1 + \cdots + f_n \in A$ of unity such that, for each index i , the localized module $M[f_i^{-1}]$ is finite free over $A[f_i^{-1}]$.”

Parametrized mathematics




local “Let M be a finitely generated module over a field k .
Then M is finite free.” ✓

global “Let M be a finitely generated module over a ring A .
Then M^\sim is finite locally free.” ✗

“Let M be a finitely generated module over an arbitrary commutative ring A . Then there is a partition $1 = f_1 + \cdots + f_n \in A$ of unity such that, for each index i , the localized module $M[f_i^{-1}]$ is finite free over $A[f_i^{-1}]$.”

Parametrized mathematics



local “Let M be a finitely generated module over a field k .
Then M is **not not** finite free.” ✓

global “Let M be a finitely generated module over a ring A .
Then M^\sim is finite locally free **on a dense open**.” ✓

“Let M be a finitely generated module over an arbitrary commutative ring A . **If $f = 0$ is the only element of A such that $M[f^{-1}]$ is finite free over $A[f^{-1}]$, then $1 = 0$ in A .**”

Finite approximations to ideal objects

- Approximate **maps** $\mathbb{N} \rightarrow X$ by their finite prefixes.

Given a finite list σ , be prepared to ...

- 1 make it more defined: $\{\sigma ::^r x \mid x \in X\}$

- Approximate **enumerations** $\mathbb{N} \rightarrow X$ by their finite prefixes.

Given a finite list σ , be prepared to ...

- 1 make it more defined: $\{\sigma ::^r x \mid x \in X\}$
- 2 ensure that a value x occurs: $\{\sigma \uparrow\uparrow \tau \mid \tau \in X^*, x \in \sigma \uparrow\uparrow \tau\}$

- Approximate **prime ideals** by finitely generated ideals.

Given a f.g. ideal \mathfrak{a} , be prepared to ...

- 1 add the individual factors in case $xy \in \mathfrak{a}$: $\{\mathfrak{a} + (x), \mathfrak{a} + (y)\}$
- 2 collapse in case $1 \in \mathfrak{a}$: \emptyset

- Approximate **local algebras** by finitely presented rings.

Given a f.p. ring A , be prepared to ...

- 1 invert the individual summands in case $x + y$ is invertible in A : $\{A[x^{-1}], A[y^{-1}]\}$
- 2 collapse in case $1 = 0$ in A : \emptyset

The generic enumeration

For any monotone predicate P on finite lists, we inductively define what it means that

*no matter how a **given list** σ evolves to a **better approximation** σ' ,
eventually $P(\sigma')$ will hold*

by the following clauses.

- 0 If $P(\sigma)$, then $P \mid \sigma$.
- 1 If $P \mid \sigma ::^r x$ for all $x \in X$, then $P \mid \sigma$.
- 2 If $P \mid \sigma \uparrow\uparrow \tau$ for all $\tau \in X^*$ such that $x \in \sigma \uparrow\uparrow \tau$, then $P \mid \sigma$.

Notation. Write “ $\nabla\sigma. P(\sigma)$ ” for $P \mid \sigma$.

Examples. (in case $X = \mathbb{R}$)

- ✓ $\nabla\sigma. \text{length}(\sigma) \geq 5$
- ✗ $\nabla\sigma. \text{length}(\sigma) \geq 2 \wedge \sigma[0] = \sigma[1]$
- ✓ $\forall x \in \mathbb{R}. \nabla\sigma. \exists n \in \mathbb{N}. \sigma[n] = x \wedge (\nabla\sigma. \exists m \in \mathbb{N}. \sigma[m] = \sin(n))$

Soundness of the ∇ -translation. If $\Gamma \vdash \varphi$, then $\Gamma^\nabla \vdash \varphi^\nabla$.

```

module _ (A : Set) where

open import Data.List
open import Data.List.Membership.Propositional
open import Data.Product

data Eventually (P : List A → Set) : List A → Set where
  now
    : {σ : List A}
      → P σ
      → Eventually P σ
  later
    : {σ : List A} {a : A}
      → ((τ : List A) → a ∈ (σ ++ τ) → Eventually P (σ ++ τ))
      → Eventually P σ

State : (List A → Set) → (List A → Set)
State P σ = ((τ : List A) → Σ[ v ∈ List A ] P (σ ++ τ ++ v))

```

U:** Countable.agda All L1 <N> (Agda:Checked +5 Undo-Tree)

U:%*- *All Done* All L1 <M> (AgdaInfo Undo-Tree)

Agda formalization available.