

Tensor categories

Ingo Blechschmidt

8th February 2014

1 Basics

Example 1.1. The category Mod_R of modules over a commutative ring R is the archetypical example of a tensor category.

Definition 1.2. A *tensor category* (\mathcal{C}, \otimes) (i.e. a monoidal category with symmetric braiding) consists of

- a category \mathcal{C} ,
- a functor $\otimes : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ (the tensor operation),
- an object $1 \in \mathcal{C}$ (the unit object),
- natural isomorphisms $X \otimes (Y \otimes Z) \xrightarrow{\phi_{XYZ}} (X \otimes Y) \otimes Z$ (the associator),
- natural isomorphisms $X \otimes Y \xrightarrow{\psi_{XY}} Y \otimes X$ (the braiding),
- natural isomorphisms $1 \otimes X \xrightarrow{\lambda_X} X$ and $X \otimes 1 \xrightarrow{\rho_X} X$ (the unitors),

such that

- the braiding is symmetric: $\psi_{YX} \circ \psi_{XY} = \text{id}_{X \otimes Y}$ and
- the following coherence conditions are satisfied:

$$XXX$$

The definitions mimics the definition of a monoid, only “one level up”. Demanding that the tensor operation is associative and commutative on the nose (e.g. $X \otimes (Y \otimes Z) = (X \otimes Y) \otimes Z$) would be *evil* in the technical sense, i.e. not invariant under equivalence of categories. The isomorphism classes of a tensor category form a commutative monoid.

Example 1.3. In Mod_R , ϕ_{XYZ} is given by

$$x \otimes (y \otimes z) \mapsto (x \otimes y) \otimes z.$$

If $-1 \neq 1 \in R$, introducing a sign here will cause the pentagon diagram to fail to commute by a sign.

Example 1.4. Let \mathcal{C} be a category with finite products. Then (\mathcal{C}, \times) is a tensor category, with $\phi, \psi, \lambda, \rho$ given by the universal property of the product. In particular, (Set, \times) is a tensor category.

Example 1.5. Let \mathcal{C} be a tensor category. Then \mathcal{C}^{op} is in a natural way a tensor category, by using the inverses of the given natural isomorphisms.

Remark 1.6. In a general tensor category, there is no natural morphism $X \rightarrow X \otimes X$.

The coherence conditions in the definition are needed for the following reason: In the category of modules, we are used to dropping all parentheses when dealing with iterated tensor products. This is justified because between any two given groupings, e. g.

$$X \otimes ((Y \otimes Z) \otimes (T \otimes U)) \quad \text{and} \quad ((X \otimes Y) \otimes (Z \otimes T)) \otimes U,$$

we have a *canonical* isomorphism. A classical theorem of Mac Lane guarantees that the stated coherence conditions suffice to render any “reasonable” diagram commutative:

Theorem 1.7 (Mac Lane). *Any (formal) diagram in a tensor category built by*

$$\otimes, \text{id}, \phi, \psi, \lambda, \rho, \phi^{-1}, \psi^{-1}, \lambda^{-1}, \rho^{-1}$$

(in which both sides have the same permutation) commutes.

The two caveats are the following: Firstly, in a given tensor category, there may hold certain *identities* between objects for no general abstract reason. For example, for some totally unrelated objects X, Y, A, B , it might hold that $X \otimes Y = A \otimes B$. Using those identities we can form diagrams which we do *not* expect to commute. Mac Lane’s coherence theorem does not make any statement about those diagrams.

To understand the restriction about permutations, consider the diagram

$$\psi_{XX} \stackrel{?}{=} \text{id}_{X \otimes X}.$$

We do not expect this diagram to be commutative; the permutations associated to both sides are not equal: $(1, 2) \neq \text{id}$.

Theorem 1.8 (Joyal, Street). *If the graphical depictions of given morphisms of a tensor category are “the same” (in 4D space), the morphisms are equal.*

Example 1.9. XXX

Theorem 1.10 (Mac Lane). *Any tensor category may be strictified, i. e. is equivalent as a tensor category to a strict tensor category: a category in which ϕ, λ, ρ (but not ψ) are identities.*

2 Structure in tensor categories

Definition 2.1. An *internal Hom* between objects X, Y of a tensor category \mathcal{C} consists of

- an object $\underline{\text{Hom}}(X, Y) \in \mathcal{C}$ and
- a morphism $\underline{\text{Hom}}(X, Y) \otimes X \xrightarrow{\text{ev}} Y$ (evaluation morphism)

such that this pair is terminal among such pairs, i. e. such that for any object $T \in \mathcal{C}$ and a [fake] evaluation morphism $T \otimes X \xrightarrow{\widetilde{\text{ev}}} Y$ there exists an unique morphism $f : T \rightarrow \underline{\text{Hom}}(X, Y)$ such that the following diagram commutes:

$$\text{ev} \circ (f \otimes \text{id}_X) = \widetilde{\text{ev}}.$$

Example 2.2. In (Set, \times) , the internal Homs are given by the usual Hom sets. The evaluation morphism is given by $(f, x) \mapsto f(x)$.

Example 2.3. In Mod_R , the internal Homs are given by the Hom sets equipped with the usual module structure.

Remark 2.4. If an internal Hom $\underline{\text{Hom}}(X, Y)$ exists for all objects $Y \in \mathcal{C}$ (and if appropriate choice principles are available), the internal Hom can be made into a functor $\underline{\text{Hom}}(X, _) : \mathcal{C} \rightarrow \mathcal{C}$ which is right adjoint to taking tensor product with X :

$$_ \otimes X \dashv \underline{\text{Hom}}(X, _)$$

Remark 2.5. The relation with the usual Hom (which is only a set) is the following:

$$\text{Hom}(1, \underline{\text{Hom}}(X, Y)) \cong \text{Hom}(1 \otimes X, Y) \cong \text{Hom}(X, Y).$$

Remark 2.6. In (Set, \amalg) , internal Homs do not exist in general: This is because in general, $_ \amalg X$ does not preserve colimits and so cannot be a left adjoint.

Definition 2.7. 1. A *dual* of an object X is an internal Hom $X^\vee := \underline{\text{Hom}}(X, 1)$.
2. The *dual* of a morphism $f : X \rightarrow Y$ is the unique morphism $f^t : Y^\vee \rightarrow X^\vee$ rendering the diagram commutative (if X^\vee and Y^\vee exist).

Example 2.8. In Mod_R , $f^t : \theta \mapsto \theta \circ f$.

Proposition 2.9. *In any tensor category, the set $\text{End}(1) = \text{Hom}(1, 1)$ is a commutative monoid with respect to composition of morphisms.*

Proof. The results holds even if there would be no braiding: On $\text{End}(1)$, the tensor product induces a second binary operation. By the coherence conditions, this operation commutes with the operation given by composition, so by the famous Eckmann–Hilton theorem, both operations coincide and are commutative. \square

Example 2.10. In Mod_R , $\text{End}(1) \cong R$.

Example 2.11. In (Set, \amalg) , $\text{End}(1) = \{\text{id}\}$.

3 Tensor functors

Definition 3.1. A *tensor functor* $F : (\mathcal{C}, \otimes) \rightarrow (\mathcal{C}', \otimes')$ consists of

1. a functor $F : \mathcal{C} \rightarrow \mathcal{C}'$,
2. natural isomorphisms $FX \otimes' FY \xrightarrow{c_{XY}} F(X \otimes Y)$ and
3. an isomorphism $1' \xrightarrow{e} F1$

such that the following coherence conditions are satisfied:

$$XXX$$

Example 3.2. The forgetful functor $\text{Rep}_k(G) \rightarrow \text{Vect}_k$ of the category of finite-dimensional k -linear representations of a group (or group scheme) G is a tensor functor.

Example 3.3. Extension of scalars defines a tensor functor $\text{Mod}_R \rightarrow \text{Mod}_S$.

Example 3.4. A quantum field theory determines a tensor functor $\text{Cob}_d \rightarrow \text{Vect}_k$.

Definition 3.5. A *morphism of tensor functors* $\eta : (F, c, e) \rightarrow (\tilde{F}, \tilde{c}, \tilde{e})$ consists of a natural transformation $\eta : F \rightarrow \tilde{F}$ which is compatible with the coherence isomorphisms:

$$XXX$$

4 Rigid tensor categories

Definition 4.1. A tensor category \mathcal{C} is *rigid* iff

- all internal Homs exist,
- the natural morphisms

$$\underline{\text{Hom}}(X_1, Y_1) \otimes \underline{\text{Hom}}(X_2, Y_2) \longrightarrow \underline{\text{Hom}}(X_1 \otimes X_2, Y_1 \otimes Y_2)$$

are isomorphisms and

- all objects $X \in \mathcal{C}$ are *reflexive* (i. e. the natural map $X \rightarrow X^{\vee\vee}$ is an isomorphism).

Example 4.2. The category $\text{Vect}_k^{\text{fd}}$ of finite-dimensional is rigid.

Example 4.3. More generally, the category $\text{Mod}_R^{\text{fin. free}}$ of finitely free R -modules is rigid.

Example 4.4. Changing toposes, the category of locally free $\mathcal{O}_{\text{Spec } R}$ modules is rigid. (This category is equivalent to category of finitely generated projective R -modules, by the tilde construction.)

Example 4.5. The category $\text{Rep}_k(G)$ is rigid.

Remark 4.6. Let \mathcal{C} be a rigid category. Then the functor $\mathcal{C} \rightarrow \mathcal{C}^{\text{op}}, X \mapsto X^\vee$ is an equivalence of tensor categories.

In a rigid tensor category, we can define the notion of traces:

Definition 4.7. 1. The *trace* of an endomorphism $f : X \rightarrow X$ in a rigid tensor category is the following element of $\text{End}(1)$:

$$f \in \text{Hom}(X, X) \cong \text{Hom}(1, \underline{\text{Hom}}(X, X)) \cong \text{Hom}(1, X^\vee \otimes X) \rightarrow \text{Hom}(1, 1) \ni: \text{tr} f.$$

2. The *rank* (or *Euler characteristic*) of an object X is the trace of the identity morphism on X .

Trace and rank satisfy the relations you expect:

Lemma 4.8. 1. $\text{tr}(f \otimes f') = \text{tr}(f) \circ \text{tr}(f')$, $\text{tr}(f \circ g) = \text{tr}(g \circ f)$.

2. $\text{rk}(X \otimes X') = \text{rk}(X) \circ \text{rk}(X')$, $\text{rk}(1) = \text{id}_1$.

Lemma 4.9. In a rigid tensor category, a pair $(Y, Y \otimes X \xrightarrow{\text{ev}} 1)$ is a dual of X iff there exists a morphism $1 \xrightarrow{\varepsilon} X \otimes Y$ such that the following triangle identities hold:

$$XXX$$

Proof. Given a pair $(T, T \otimes X \xrightarrow{\widetilde{\text{ev}}} 1)$, construct $f : T \rightarrow Y$ as follows:

$$XXX$$

Then the relevant diagram commutes, since

$$XXX.$$

Use the second triangle identity to show uniqueness.

For the converse direction, construct ε by dualizing ev . □

Remark 4.10. Rigidity is only necessary for the converse direction: A pair $(Y, Y \otimes X \xrightarrow{\text{ev}} 1)$ which satisfies the condition stated in lemma is always a dual—and in fact, because the condition is symmetric, the coevaluation morphism exhibits X as a dual of Y . In particular, X is reflexive.

Proposition 4.11. Let $F : \mathcal{C} \rightarrow \mathcal{C}'$ be a tensor functor, with \mathcal{C} rigid. Let $X, Y \in \mathcal{C}$. Then an internal Hom $\underline{\text{Hom}}(FX, FY)$ exists in \mathcal{C}' and the natural morphism

$$F(\underline{\text{Hom}}(X, Y)) \longrightarrow \underline{\text{Hom}}(FX, FY)$$

is an isomorphism.

Proof. It is enough to show that in \mathcal{C}' a dual of FX exists and that the natural morphism

$$F(X^\vee) \longrightarrow (FX)^\vee$$

is an isomorphism. This is obvious by the lemma, as the given condition is preserved by F . \square

Corollary 4.12. $\mathrm{tr}F(f) = F(\mathrm{tr}f)$, $\mathrm{rk}FX = F(\mathrm{rk}X)$.

Proposition 4.13. *In a rigid category, $_ \otimes X$ is not only a left, but also a right adjoint:*

$$\underline{\mathrm{Hom}}(X, _) \dashv _ \otimes X \dashv \underline{\mathrm{Hom}}(X, _).$$

Corollary 4.14. *In a rigid category, the tensor operation is continuous and cocontinuous.*

5 Abelian tensor categories

Definition 5.1. An *abelian tensor category* is a tensor category (\mathcal{C}, \otimes) such that \mathcal{C} is abelian and \otimes is bi-additive.

Remark 5.2. In an abelian category, $\mathrm{End}(1)$ is a commutative ring.

Lemma 5.3. *Let \mathcal{C} be an abelian tensor category. Let $X \in \mathcal{C}$.*

1. *Assume that the unit 1 is a simple object and that \mathcal{C} is rigid. Then:*

$$X \not\cong 0 \implies X^\vee \otimes X \rightarrow 1 \text{ is epic.}$$

2. *Assume that $1 \not\cong 0 \in \mathcal{C}$. Then:*

$$X \not\cong 0 \iff X^\vee \otimes X \rightarrow 1 \text{ is epic.}$$

Proof. 1. Because the unit is simple, the natural morphism $X^\vee \otimes X \rightarrow 1$ is either zero or epic. (This follows by considering the image of the morphism.) Under the correspondence

$$\mathrm{Hom}(X^\vee \otimes X, 1) \cong \mathrm{Hom}(X, X^{\vee\vee}),$$

the morphism corresponds to the natural morphism $X \rightarrow X^{\vee\vee}$. If this was zero, X were zero as well.

2. Assume $X \cong 0$. Then we have an epic morphism $0 \rightarrow 1$. By terminality of the zero object, this is an isomorphism. \square

Remark 5.4. The rigidity is necessary: In $\mathrm{Mod}_{\mathbb{Z}}$, the object $\mathbb{Z}/(2)$ is not isomorphic to the zero object, but the morphism $(\mathbb{Z}/(2))^\vee \otimes \mathbb{Z}/(2) \rightarrow \mathbb{Z}$ has zero codomain.

4. Applying this observation to $X = V$, it follows that $U^\perp \otimes V \cong V$.

Applying it to $X = U$, it follows that $U^\perp \otimes U \cong 0$: Let $T \hookrightarrow U$ with $T \otimes U \cong 0$. By exactness of tensoring with T , the sequence

$$0 \longrightarrow T \otimes U \longrightarrow T \longrightarrow T \otimes V \longrightarrow 0$$

is exact. Since $T \otimes U \cong 0$, we have

$$T \cong T \otimes V \hookrightarrow U \otimes V \cong 0.$$

The “ \hookrightarrow ” is—again—by exactness of tensoring (with V).

5. By exactness of tensoring with U^\perp , the sequence

$$0 \longrightarrow U^\perp \otimes U \longrightarrow U^\perp \longrightarrow U^\perp \otimes V \longrightarrow 0$$

is exact. By the previous observation, this shows $U^\perp \cong V$.

6. By applying the five lemma to the diagram

$$XXX,$$

it follows that $1 \cong U \oplus U^\perp$. □

Remark 5.9. Rigidity is necessary: In $\text{Mod}_{\mathbb{Z}}$, the unit object is not simple but admits to non-trivial decompositions.

Corollary 5.10. *In a rigid abelian category, the unit object is simple iff $\text{End}(1)$ is a field.*

Proof. For the “only if” direction, let $f \in \text{End}(1)$. As the unit object is simple, f is either zero or an isomorphism (i. e. invertible in the ring $\text{End}(1)$). This follows by considering kernel and image of f .

For the “if” direction, let $U \hookrightarrow 1$. By the lemma, there exists a projection operator $P : 1 \rightarrow 1$ with $\text{im}(P) = U$. If P is zero in $\text{End}(1)$, $U = 0$; if P is invertible, $U = 1$ (as subobjects of 1). □

6 The Tannaka reconstruction theorem

Theorem 6.1. *Let \mathcal{C} be a rigid abelian tensor category. Let $k := \text{End}(1)$ be a field. Let $\omega : \mathcal{C} \rightarrow \text{Vect}_k^{\text{fd}}$ be an exact faithful k -linear tensor functor. Then:*

1. *The functor $\underline{\text{Aut}}^\otimes(\omega) : \text{Alg}_k \rightarrow \text{Set}$, given by*

$$R \longmapsto \text{set of tensor automorphisms of } (_ \otimes_k R) \cdot \omega,$$

is the functor of points of an affine group scheme G .

2. *A certain functor $\mathcal{C} \rightarrow \text{Rep}_k(G)$ induced by ω is an equivalence of tensor categories.*