

Sacituzumab Govitecan in Hormone Receptor–Positive/Human Epidermal Growth Factor Receptor 2–Negative Metastatic Breast Cancer

Hope S. Rugo, MD¹; Aditya Bardia, MD, MPH²; Frederik Marmé, MD, PhD³; Javier Cortes, MD, PhD^{4,5}; Peter Schmid, MD, PhD⁶; Delphine Loirat, MD, PhD⁷; Olivier Trédan, MD, PhD⁸; Eva Ciruelos, MD, PhD⁹; Florence Dalenc, MD, PhD¹⁰; Patricia Gómez Pardo, MD¹¹; Komal L. Jhaveri, MD¹²; Rosemary Delaney, MPH¹³; Olivia Fu, MD¹⁴; Lanjia Lin, PhD¹⁵; Wendy Verret, PhD¹³; and Sara M. Tolaney, MD, MPH¹⁶; on behalf of the TROPICS-02 Study Investigators

abstract

PURPOSE Hormone receptor–positive (HR+) human epidermal growth factor receptor 2–negative (HER2–) endocrine-resistant metastatic breast cancer is treated with sequential single-agent chemotherapy with poor outcomes. Sacituzumab govitecan (SG) is a first-in-class antibody-drug conjugate with an SN-38 payload targeting trophoblast cell-surface antigen 2, an epithelial antigen expressed in breast cancer.

METHODS In this global, randomized, phase III study, SG was compared with physician's choice chemotherapy (eribulin, vinorelbine, capecitabine, or gemcitabine) in endocrine-resistant, chemotherapy-treated HR+/HER2– locally recurrent inoperable or metastatic breast cancer. The primary end point was progression-free survival (PFS) by blinded independent central review.

RESULTS Patients were randomly assigned to receive SG (n = 272) or chemotherapy (n = 271). The median age was 56 years, 95% had visceral metastases, and 99% had a prior cyclin-dependent kinase 4/6 inhibitor, with three median lines of chemotherapy for advanced disease. Primary end point was met with a 34% reduction in risk of progression or death (hazard ratio, 0.66 [95% CI, 0.53 to 0.83; *P* = .0003]). The median PFS was 5.5 months (95% CI, 4.2 to 7.0) with SG and 4.0 months (95% CI, 3.1 to 4.4) with chemotherapy; the PFS at 6 and 12 months was 46% (95% CI, 39 to 53) v 30% (95% CI, 24 to 37) and 21% (95% CI, 15 to 28) v 7% (95% CI, 3 to 14), respectively. Median overall survival (first planned interim analysis) was not yet mature (hazard ratio, 0.84; *P* = .14). Key grade ≥ 3 treatment-related adverse events (SG v chemotherapy) were neutropenia (51% v 38%) and diarrhea (9% v 1%).

CONCLUSION SG demonstrated statistically significant PFS benefit over chemotherapy, with a manageable safety profile in patients with heavily pretreated, endocrine-resistant HR+/HER2– advanced breast cancer and limited treatment options.

J Clin Oncol 40:3365-3376. © 2022 by American Society of Clinical Oncology

Creative Commons Attribution Non-Commercial No Derivatives 4.0 License

ASSOCIATED CONTENT

Appendix

Data Supplement
Protocol

Author affiliations and support information (if applicable) appear at the end of this article.

Accepted on July 28, 2022 and published at ascopubs.org/journal/jco on August 26, 2022; DOI <https://doi.org/10.1200/JCO.22.01002>

INTRODUCTION

Although endocrine therapy combined with cyclin-dependent kinase 4/6 inhibitors (CDK4/6i) has extended overall survival (OS) for metastatic hormone receptor–positive (HR+) human epidermal growth factor receptor 2–negative (HER2–) breast cancer to over 5 years in the first-line setting¹⁻⁵ and combinations with phosphoinositide 3-kinase or mammalian target of rapamycin inhibitors offer benefit in subsequent treatment lines,⁶ endocrine resistance eventually develops. Sequential single-agent chemotherapy is the next therapeutic option but is associated with declining response rates and disease control and increased toxicity.⁶⁻¹⁰

Sacituzumab govitecan (SG) is a first-in-class trophoblast cell-surface antigen 2 (Trop-2)–directed antibody-drug conjugate (ADC), consisting of a humanized anti-Trop-2 monoclonal antibody conjugated to the active metabolite of irinotecan, SN-38,¹¹ via a hydrolyzable CL2A linker.¹² Trop-2 is a transmembrane calcium signal transducer highly expressed in solid tumors, especially HR+/HER2– and triple-negative breast cancers (with a prevalence of > 90%), and linked to tumor progression and poor prognosis.¹³⁻¹⁵ Internalization of Trop-2–bound SG delivers SN-38 into the tumor cell through hydrolysis of the linker.¹⁶ Because SN-38 is a membrane-permeable free molecule released in the tumor microenvironment, it

CONTEXT

Key Objective

Hormone receptor–positive (HR+) human epidermal growth factor receptor 2–negative (HER2–) breast cancer is treated with sequential endocrine therapy combined with targeted agents, followed by sequential single-agent chemotherapy, with declining response rates and increased toxicity. This randomized, global phase III study evaluates sacituzumab govitecan (SG), a trophoblast cell-surface antigen 2–directed antibody-drug conjugate, versus single-agent chemotherapy in HR+/HER2– advanced breast cancer.

Knowledge Generated

SG demonstrated a significant improvement in progression-free survival over chemotherapy (median, 5.5 months v 4.0 months; hazard ratio, 0.66; $P = .0003$) in patients who have received prior endocrine-based therapy, including cyclin-dependent kinase 4/6 inhibitors and ≥ 2 prior chemotherapy regimens in the metastatic setting. The SG safety profile was manageable and consistent with previous studies, with neutropenia and diarrhea as the most common treatment-related adverse events.

Relevance (K.D. Miller)

SG is a treatment option for patients with heavily pretreated, endocrine-resistant HR+/HER2– advanced breast cancer.*

*Relevance section written by JCO Senior Deputy Editor Kathy D. Miller, MD.

may elicit antitumor effects in adjacent non–Trop-2-expressing tumor cells (bystander effect).¹⁶

SG received full authorization from the Food and Drug Administration and European Medicines Agency for patients with metastatic triple-negative breast cancer who had received at least two prior chemotherapies (at least one for metastatic disease) and has accelerated approval for locally advanced or metastatic urothelial cancer.¹⁷⁻¹⁹ In the phase I/II IMMU-132-01 basket study, SG showed encouraging activity and safety in 54 patients with HR+/HER2– metastatic breast cancer (MBC) who progressed on at least one line of endocrine therapy and at least one prior chemotherapy in the metastatic setting.²⁰ The objective response rate was 31.5%, and the median progression-free survival (PFS) and OS were 5.5 and 12 months, respectively (CDK4/6i-pretreated group, 25%, 3.8 and 11 months).

Here, we provide the primary results of TROPiCS-02, a global, randomized, open-label, multicenter phase III study of SG versus single-agent chemotherapy in patients with locally recurrent inoperable or metastatic HR+/HER2– breast cancer (Data Supplement, online only).

METHODS

Patients

Eligible patients had histologically locally confirmed measurable HR+/HER2– MBC and 2-4 prior systemic chemotherapy regimens for metastatic disease. (Neo)adjuvant therapy for early-stage disease qualified as one of the required prior chemotherapy regimens if the development of unresectable, locally advanced, or metastatic disease occurred within 12 months of therapy (early relapse). Patients must have previously received at least one taxane, at least

one anticancer hormonal treatment, and at least one CDK4/6i, reflecting standard clinical practice. Additional details are provided in the Data Supplement.

Trial Design and Treatment

Patients were randomly assigned 1:1 to receive 10 mg/kg of SG (Trodely; Gilead Sciences Inc, Foster City, CA) intravenously once weekly on day 1 and day 8 every 21 days or chemotherapy of physician's choice determined before random assignment (eribulin, capecitabine, gemcitabine, or vinorelbine). The chemotherapy agents included in this trial, and their recommended doses, were in accordance with locally approved prescribing information or according to National Comprehensive Cancer Network Clinical Practice Guidelines in Oncology (National Comprehensive Cancer Network Guidelines) for Breast Cancer.²¹ Recommended doses are summarized below: eribulin, 1.4 mg/m² (North America) or 1.23 mg/m² (Europe) intravenously once weekly on days 1 and 8 every 21 days; vinorelbine, 25 mg/m² intravenously once weekly; gemcitabine, 800-1,200 mg/m² intravenously once weekly on days 1, 8, and 15 every 28 days; and capecitabine, 1,000-1,250 mg/m² orally twice daily for 2 weeks followed by a 1 week rest period, every 21 days.

Random assignment was stratified by number of prior chemotherapy regimens for metastatic disease, visceral metastases, and prior endocrine treatment in the metastatic setting for at least 6 months. Patients were treated until disease progression, unacceptable toxicity, withdrawal of consent, or per investigator's decision that it was in the patient's best interest to discontinue. Treatment beyond progression was permitted if deemed clinically beneficial by the investigator. Additional details are provided in the Data Supplement.

Trial Oversight

The study was approved by national regulatory authorities and each investigational site's institutional review/ethics committee before implementation and was compliant with the Declaration of Helsinki and International Council for Harmonisation Good Clinical Practice Guidelines. All patients provided written informed consent.

End Points

The primary end point was PFS as determined by blinded independent central review (BICR) per the RECIST v1.1.²² Secondary end points included OS, objective response, clinical benefit rate, duration of response, patient-reported outcomes, and safety (Data Supplement).

Assessments

The primary and secondary end points were measured by computed tomography or magnetic resonance imaging, conducted every 6 weeks for the first 54 weeks and every 12 weeks thereafter. Additional details are provided in the Data Supplement.

Safety and tolerability were assessed in all treated patients throughout the study, with severity of adverse events (AEs) graded using National Cancer Institute Common Terminology Criteria for Adverse Events v5.0. Patients were allowed premedications (eg, antipyretics and H1 blockers) for prevention of infusion reactions and supportive medications for the prevention and treatment of chemotherapy-induced nausea, vomiting, and diarrhea.

Statistical Analysis

Anticipated recruitment was 520 patients; sample size calculation was based on treatment group comparisons, with PFS as the primary end point and OS as the key secondary end point. For PFS, assuming a hazard ratio (HR) of 0.70, 350 events of progression or death were needed to detect a statistically significant PFS difference between treatment groups at a two-sided α of .05 with 92% power. For OS, assuming a HR of 0.73, 438 events were needed to detect a statistically significant difference with 87% power at a two-sided α of .05. OS would be formally tested sequentially if PFS is statistically significant, and then objective response and quality of life (QoL) if the prior end point in the hierarchy is significant. The study was planned to have two interim analyses and a final analysis of OS, with 272 events targeted for the first interim and .036 α to be spent on the basis of Lan DeMets α spending function that approximated a Pocock approach. Analysis of OS was combined with the primary PFS analysis since the required OS events for the first interim analysis were reached earlier than PFS events (329 events occurred).

AEs summarized were treatment-emergent unless otherwise specified (defined as any AEs that began or worsened on or after study drug administration through 30 days after the last dose of study drug).

The intent-to-treat population includes all randomly assigned patients (efficacy population). Stratification factors used in random assignment were applied to all stratified analyses. All patients who received at least one dose of study drug were included in safety analyses. Additional details are provided in the Data Supplement.

RESULTS

Patient Characteristics

From May 2019 to April 2021, 543 patients with HR+/HER2- locally inoperable or MBC were enrolled in 91 centers across North America (United States and Canada) and Europe (the United Kingdom, France, Spain, Italy, Germany, Belgium, and the Netherlands). Patients were randomly assigned to the SG group (272 patients) or chemotherapy group (271 patients; 48% eribulin, 23% vinorelbine, 21% gemcitabine, and 8% capecitabine; Data Supplement). In total, 268 patients (99%) in the SG group and 249 patients (92%) in the chemotherapy group received study treatment. Of the 26 patients randomly assigned but not treated, 16 withdrew consent (one for SG and 15 for chemotherapy).

The median age was 56 years (range, 27-86; Table 1). Most had visceral metastases (95%) and received endocrine therapy in the metastatic setting for at least 6 months (86%); 40% of patients received CDK4/6i therapy for more than 12 months. Patients received a median of 3 (range, 0-8) prior lines of chemotherapy in the metastatic setting (57% at least three lines). At the data cutoff date (January 3, 2022), 18 patients (7%) in the SG group and four (1.5%) in the chemotherapy group remained on study treatment. Patients discontinued study treatment primarily because of progressive disease (SG group, 77%; chemotherapy group, 73%; Fig 1).

Efficacy

As of the data cutoff date, the median duration of follow-up was 10.2 months (11.3 months with SG and 9.8 months with chemotherapy). The primary end point of PFS was met with a 34% reduction in risk of progression or death (HR, 0.66; 95% CI, 0.53 to 0.83; $P = .0003$; 329 events); the median PFS determined by BICR was 5.5 months (95% CI, 4.2 to 7.0) for SG and 4.0 months (95% CI, 3.1 to 4.4) for chemotherapy (Fig 2A and Table 2). The reduction in risk of progression or death was consistent with local investigator assessment (HR, 0.73; 95% CI, 0.60 to 0.88; $P = .001$; Data Supplement).

In landmark analyses, the PFS rates at 6 and 12 months (SG v chemotherapy) were 46% versus 30% and 21% versus 7%, respectively (Fig 2A and Table 2). The PFS benefit for SG over chemotherapy was maintained in most predefined subgroups, including patients with three or more prior chemotherapy regimens in the metastatic setting, visceral metastases, and age 65 years or older (Fig 3).

Because PFS results were statistically significant, OS was sequentially tested. The median OS (first planned interim

TABLE 1. Baseline Characteristics and Treatment History of Patients

Characteristic	SG (n = 272)	Chemotherapy (n = 271)	All (N = 543)
Female, No. (%)	270 (99)	268 (99)	538 (99)
Median age, years (range)	57 (29-86)	55 (27-78)	56 (27-86)
Race or ethnic group, No. (%)			
White	184 (68)	178 (66)	362 (67)
Black	8 (3)	13 (5)	21 (4)
Asian	11 (4)	5 (2)	16 (3)
Others ^a	0	5 (2)	5 (1)
Not specified ^b	69 (25)	70 (26)	139 (26)
ECOG PS, No. (%)			
0	116 (43)	126 (46)	242 (45)
1	156 (57)	145 (54)	301 (55)
Visceral metastases at baseline, No. (%)	259 (95)	258 (95)	517 (95)
Liver metastases, ^c No. (%)	229 (84)	237 (87)	466 (86)
De novo MBC, No. (%)	78 (29)	60 (22)	138 (25)
Median time from initial metastatic diagnosis to random assignment, months (range)	48.5 (1.2-243.8)	46.6 (3.0-248.8)	47.8 (1.2-248.8)
Prior chemotherapy in the (neo)adjuvant setting, No. (%)	173 (64)	184 (68)	357 (66)
Prior endocrine therapy in the metastatic setting > 6 months, No. (%)			
Yes	235 (86)	234 (86)	469 (86)
No	37 (14)	37 (14)	74 (14)
Prior CDK4/6i use, months, No. (%)			
≤ 12	161 (59)	166 (61)	327 (60)
> 12	106 (39)	102 (38)	208 (38)
Unknown	5 (2)	3 (1)	8 (1)
Median prior chemotherapy regimens in the metastatic setting, No. (%) ^d	3 (0-8) ^d	3 (1-5) ^d	3 (0-8) ^d
0	1 (< 1)	0	1 (< 1)
1	8 (3)	2 (1)	10 (2)
2	104 (38)	118 (43)	222 (41)
≥ 3	159 (58)	151 (56)	310 (57)
Median prior chemotherapy regimens, No. (range)	4 (1-9)	4 (2-7)	4 (1-9)
Median prior anticancer regimens, ^e No. (range)	7 (3-17)	7 (3-16)	7 (3-17)
Setting of prior anticancer regimens, ^e No. (%)			
Neoadjuvant	67 (25)	62 (23)	129 (24)
Adjuvant	186 (68)	206 (76)	392 (72)
Advanced/metastatic	272 (100)	271 (100)	543 (100)
Others/unknown	12 (4)	9 (3)	21 (4)
Most common prior anticancer therapy, ^e No. (%)			
Palbociclib	238 (88)	228 (84)	466 (86)
Capecitabine	226 (83)	234 (86)	460 (85)
Fulvestrant	235 (86)	223 (82)	458 (84)
Cyclophosphamide	204 (75)	209 (77)	413 (76)
Paclitaxel	210 (77)	196 (72)	406 (75)
Letrozole	185 (68)	210 (77)	395 (73)
Tamoxifen	160 (59)	165 (61)	325 (60)

(continued on following page)

TABLE 1. Baseline Characteristics and Treatment History of Patients (continued)

Characteristic	SG (n = 272)	Chemotherapy (n = 271)	All (N = 543)
Doxorubicin ^e	149 (55)	134 (49)	283 (52)
Exemestane	142 (52)	134 (49)	276 (51)
Everolimus	117 (43)	115 (42)	232 (43)
Docetaxel	106 (39)	120 (44)	226 (42)
Eribulin ^e	97 (36)	97 (36)	194 (36)
Anastrozole	87 (32)	69 (25)	156 (29)
Epirubicin	78 (29)	94 (35)	172 (32)
Fluorouracil	66 (24)	77 (28)	143 (26)
Most common prior anticancer therapy class in the metastatic setting, ^e No. (%)			
Endocrine therapy	268 (99)	269 (99)	537 (99)
CDK4/6i	267 (98)	270 (> 99)	537 (99)
Targeted agent	181 (67)	172 (63)	353 (65)
Immunotherapy	21 (8)	15 (6)	36 (7)
Chemotherapy	271 (> 99)	271(100)	542 (> 99)
Most common prior chemotherapy agent in the metastatic setting, ^e No. (%)			
Capecitabine	221 (81)	232 (86)	453 (83)
Paclitaxel	174 (64)	147 (54)	321 (59)
Eribulin ^e	95 (35)	88 (33)	183 (34)
ER expression, No. (%) ^f			
< 1%	2 (1)	5 (2)	7 (1)
1%-10%	12 (4)	15 (6)	27 (5)
> 10%	258 (95)	246 (91)	504 (93)
Unknown	0	5 (2)	5 (1)
PR expression, No. (%) ^f			
< 1%	103 (38)	101 (37)	204 (38)
1%-10%	45 (17)	44 (16)	89 (16)
> 10%	124 (46)	120 (44)	244 (45)
Unknown	0	6 (2)	6 (1)

Abbreviations: CDK4/6i, cyclin-dependent kinase 4/6 inhibitor; ECOG PS, Eastern Cooperative Oncology Group performance status; ER, estrogen receptor; MBC, metastatic breast cancer; PR, progesterone receptor; SG, sacituzumab govitecan.

^aIncludes American Indian or Alaska Native, Native Hawaiian or other Pacific Islander.

^bNot reported indicates local regulators who did not allow collection of race or ethnicity information.

^cPresence of baseline target/nontarget liver lesion per RECIST1.1 by local investigator review.

^dThe reported number of prior therapies was miscounted at screening for some patients. Nine patients had fewer or more prior chemotherapy regimens in the metastatic setting than the specified inclusion criteria and were included in the intention-to-treat population.

^eAnticancer regimens refer to any treatment regimen that was used to treat breast cancer in any setting and includes endocrine therapy and everolimus. Eribulin includes the preferred drug name eribulin and eribulin mesylate. Doxorubicin includes the preferred drug name doxorubicin, pegylated liposomal doxorubicin hydrochloride, liposomal doxorubicin, doxorubicin hydrochloride, liposomal doxorubicin hydrochloride, and pegylated liposomal doxorubicin.

^fPer protocol, hormone receptor status was to be documented from locally recurrent or metastatic sites and the few cases that did not have protocol deviations.

analysis) was 13.9 months (95% CI, 12.7 to 15.4) for SG and 12.3 months (95% CI, 10.8 to 14.2) for chemotherapy (HR for death, 0.84; 95% CI, 0.67 to 1.06; $P = .14$; Fig 2B). These results are not yet mature; further follow-up is ongoing. Because OS was not significant, objective response and QoL end points—which fell after OS in the hierarchy—were not formally tested.

The percentage of patients with objective response by BICR was 21% with SG and 14% with chemotherapy (Table 2), of which two patients (1%) and no patients achieved a complete response, respectively. Clinical benefit rate was higher with SG than with chemotherapy (34% v 22%). The median time to response was 2.9 (range, 1.2-11.3) months with SG and 2.7 (range, 1.2-10.5) months with




FIG 1. CONSORT diagram. ^aPatients in the chemotherapy group were randomly assigned to eribulin (n = 130), vinorelbine (n = 63), gemcitabine (n = 56), or capecitabine (n = 22). AE, adverse event; SG, sacituzumab govitecan.

chemotherapy. The median duration of response was 7.4 months (95% CI, 6.5 to 8.6) with SG and 5.6 months (95% CI, 3.8 to 7.9) with chemotherapy.

For patient-reported outcomes, the European Organization for Research and Treatment of Cancer QoL questionnaire completion rate was at least 85% through cycle 13 day 1 for both SG and chemotherapy and was generally comparable across assessments between treatment groups. Median time to deterioration was longer for SG versus chemotherapy for global health status/QoL (4.0 v 2.9 months; HR, 0.74; 95% CI, 0.59 to 0.91) and fatigue (2.1 v 1.4 months; HR, 0.76; 95% CI, 0.62 to 0.93). Median time to deterioration for pain was similar (3.7 v 3.4 months; HR, 0.92; 95% CI, 0.74 to 1.14).

Safety

In the safety population, the median duration of treatment was 4.1 (range, 0.03-24.2) months and 2.3 (range, 0.03-22.3) months with SG and chemotherapy, respectively. Patients in the SG group received a mean of 8.2 treatment cycles (range, 1.0-35.0), with a median relative dose intensity of 99%.

An AE summary is found in the Data Supplement. The most common treatment-related AEs of any grade (> 25% incidence) with SG versus chemotherapy were neutropenia (70% v 54%), diarrhea (57% v 16%), nausea (55% v 31%), alopecia (46% v 16%), fatigue (37% v 29%), and anemia (34% v 25%). The most common grade 3 or higher treatment-related AEs (> 5% incidence) were neutropenia (51% v 38%), leukopenia (9% v 5%), diarrhea (9% v 1%), anemia (6% v 3%), and fatigue (6% v 2%; Table 3). There

was a low incidence of treatment-related febrile neutropenia (5% v 4%), interstitial lung disease (0% v 1%), and neuropathy (9% v 15%). For additional results about growth factor use, see the Data Supplement.

Given the difference in treatment durations in the SG and chemotherapy groups, a summary of exposure-adjusted incidence rates (EAIRs) for AEs is provided in the Data Supplement. Although the EAIRs for common AEs of any grade, such as diarrhea, alopecia, and nausea, were higher with SG versus chemotherapy, the EAIRs for other common AEs of any grade, including neutropenia, anemia, and fatigue, were similar between treatment groups (EAIR difference [95% CI]: diarrhea: 2.29 per patient-years of exposure [PYE; 1.72 to 2.87], alopecia: 1.23 per PYE [0.80 to 1.68]; nausea: 0.85 per PYE [0.30 to 1.39]; neutropenia: 0.75 per PYE [-0.16 to 1.66]; anemia: 0 per PYE [-0.37 to 0.35]; fatigue: -0.36 per PYE [-0.82 to 0.07]).

Serious treatment-related AEs were reported in 37 patients (14%) in the SG group and 25 patients (10%) in the chemotherapy group. The most common ($\geq 2\%$ incidence) serious treatment-related AEs for SG were diarrhea (5%), febrile neutropenia (4%), neutropenia (3%), and neutropenic colitis (2%); for chemotherapy, they were febrile neutropenia (4%), pneumonia (2%), nausea (2%), and dyspnea (2%). AEs leading to study treatment discontinuations occurred in 17 patients (6%) in the SG group and 11 patients (4%) in the chemotherapy group. Dose delays and reductions can be found in the Data Supplement. Although six patients experienced AEs leading to death in the SG group, only one had a treatment-related AE leading




FIG 2. Efficacy outcomes in the intent-to-treat population. (A and B) PFS (final analysis) and OS (first planned interim analysis), respectively, in the intent-to-treat population (all randomly assigned patients). PFS was determined by blinded independent central review according to RECIST, version 1.1. HR, hazard ratio; OS, overall survival; PFS, progression-free survival; SG, sacituzumab govitecan.

to death (septic shock because of neutropenic colitis). The AEs leading to death in the remaining five patients included (*n* = 1 each) arrhythmia, COVID-19 pneumonia, pulmonary embolism, pneumonia, and nervous system disorder. The patients with fatal infections of COVID-19 pneumonia and pneumonia were not neutropenic at the time of event onset. No mechanistic or etiologic pattern was identified for these AEs (Data Supplement). No AEs leading to death were reported in the chemotherapy group.

DISCUSSION

Patients with metastatic HR+/HER2- breast cancer ultimately develop endocrine resistance, and treatment options are limited to sequential single-agent chemotherapy. In this phase III trial of patients with heavily pretreated, locally recurrent inoperable, or metastatic HR+/HER2- breast cancer, SG, a Trop-2-directed ADC, demonstrated significant improvement in PFS versus chemotherapy with a 34% reduction in risk of disease

TABLE 2. Summary of Treatment Efficacy (per blinded independent central review)

Efficacy Outcome	SG (n = 272)	Chemotherapy (n = 271)
Median PFS, months (95% CI),	5.5 (4.2 to 7.0)	4.0 (3.1 to 4.4)
HR; <i>P</i> (95% CI)	0.66 (0.53 to 0.83); <i>P</i> = .0003	
PFS rate, %, months (95% CI)		
6	46 (39 to 53)	30 (24 to 37)
12	21 (15 to 28)	7 (3 to 14)
Median OS, months (95% CI)	13.9 (12.7 to 15.4)	12.3 (10.8 to 14.2)
HR; <i>P</i> (95% CI)	0.84 (0.67 to 1.06); <i>P</i> = .14	
Objective response rate, No. (%)	57 (21)	38 (14)
Best overall response, No. (%)		
Complete response	2 (1)	0
Partial response	55 (20)	38 (14)
Stable disease	142 (52)	106 (39)
Stable disease ≥ 6 months	35 (13)	21 (8)
Progressive disease	58 (21)	76 (28)
Not evaluable	15 (6)	51 (19)
CBR, ^a No. (%)	92 (34)	59 (22)
Median DOR, months (95% CI)	7.4 (6.5 to 8.6)	5.6 (3.8 to 7.9)

Abbreviations: CBR, clinical benefit rate; DOR, duration of response; HR, hazard ratio; OS, overall survival; PFS, progression-free survival; SG, sacituzumab govitecan.

^aCBR is defined as the percentage of patients with a confirmed best overall response of complete response, partial response, and stable disease ≥ 6 months.

progression or death (HR, 0.66; *P* = .0003) and a higher proportion of patients who are alive and progression-free at all landmark time points with nonoverlapping confidence intervals compared with chemotherapy.

The population in this study had progressive disease and extensive prior chemotherapy treatment in the advanced setting (median prior lines of chemotherapy = 3), and uniquely, all patients had received prior CDK4/6i, reflecting standard of care and allowing assessment of efficacy post-CDK4/6i treatment.

Benefit with SG was seen across most of the prespecified subgroups, including patients who received at least three prior chemotherapies for metastatic disease, had visceral metastases, and were age 65 years or older. The OS results (16% reduction in risk of death; *P* = .14) are not yet mature in the first planned interim analysis. Overall, results are consistent with those from the HR+/HER2-cohort of the phase I/II IMMUNO-132-01 basket study,²⁰ and performance of the control group is as previously reported.⁶⁻¹⁰ Additional preplanned follow-up will provide more clarity into the survival benefit of SG in this patient population.

With limited advancements for treatments in this later-line setting, the most recent phase III trials in a similar patient population were EMBRACE,⁸ which led to the approval of eribulin in 2010,²³ and Study 301.²⁴ A pooled analysis

demonstrated a significant PFS improvement (4.1 v 3.4 months; HR, 0.84; *P* = .03) with eribulin, but no significant OS benefit (15.7 v 13.5 months; HR, 0.87; *P* = .06) versus other chemotherapies in patients with HR+ MBC.²⁵ Comparison with this pooled analysis is complicated by the fact that these two studies involved different patient populations with regard to the extent of prior chemotherapy for advanced disease (eligibility, EMBRACE: 2-5; Study 301: up to 2), HER2 status, and lack of prior use of CDK 4/6i.

Recently, the phase III DESTINY-Breast04 trial compared the ADC trastuzumab deruxtecan with chemotherapy of physician's choice in a patient population that partially overlaps with those enrolled in TROPiCS-02. Trastuzumab deruxtecan significantly improved both PFS and OS in patients with HR+/HER2-low MBC;²⁶ however, there are important differences in the study populations that limit comparisons with this trial. DESTINY-Breast04 only included patients with HER2-low (immunohistochemistry 1+ or immunohistochemistry 2+/in situ hybridization-negative) less heavily pretreated disease (median number of prior chemotherapies in the metastatic setting = 1), along with other differences. It is clear that the standard of care now includes treatment with ADCs for a number of breast cancer subtypes; sequential efficacy is yet to be evaluated.

SG demonstrated a manageable safety profile, with a low incidence of treatment discontinuation because of AEs




FIG 3. Subgroup analysis of PFS. Early relapse is defined as relapse to metastatic disease within 1 year of the end of (neo)adjuvant chemotherapy. Patients without chemotherapy in the (neo)adjuvant setting are not considered as early relapse. CDK, cyclin-dependent kinase; ECOG PS, Eastern Cooperative Oncology Group performance status; HR, hazard ratio; NE, not evaluable; PFS, progression-free survival; SG, sacituzumab govitecan.

(6%). The most clinically relevant grade 3 or 4 AEs with SG were neutropenia and diarrhea, adequately managed with established supportive care measures, as previously reported.²⁷ The incidence of treatment-related febrile neutropenia and neuropathy was low with SG; no interstitial lung disease was reported in the SG arm of this trial.

This study has some potential limitations. A number of patients randomly assigned to the chemotherapy group were not treated (n = 22; 8%), likely because of patient preference not to receive standard chemotherapy. Most patients had visceral metastases (95%), consistent with aggressive disease, and had received multiple lines of chemotherapy, which are the factors associated with shorter PFS and higher neutropenia risk. The heterogeneity

of prior treatments and chemotherapy of physician's choice options might have affected efficacy findings. Hormone receptor status was determined locally at any stage of disease, which has historically presented challenges for accurate assessment. The study did not require real-time BICR assessment of progressive disease, potentially increasing censoring.

SG is a Trop-2-directed ADC that demonstrated significant clinical benefit and manageable safety compared with standard chemotherapy in a phase III trial of patients with unresectable locally advanced or metastatic heavily pretreated, endocrine-resistant HR+/HER2- breast cancer, a population with limited treatment options. The magnitude of PFS benefit should be considered in the

TABLE 3. Summary of Treatment-Related AEs of Any Grade ($\geq 10\%$) and Worst Grade 2 or Grade ≥ 3 ($\geq 5\%$) in the Safety Population (all patients who received ≥ 1 dose of study treatment)

Treatment-Related AE ^a	SG (n = 268)			Chemotherapy (n = 249)		
	All Grade	Grade 2	Grade ≥ 3	All Grade	Grade 2	Grade ≥ 3
Hematologic, No. (%)						
Neutropenia ^b	188 (70)	45 (17)	136 (51)	134 (54)	29 (12)	94 (38)
Anemia ^c	91 (34)	44 (16)	17 (6)	62 (25)	31 (12)	8 (3)
Leukopenia ^d	37 (14)	7 (3)	23 (9)	23 (9)	8 (3)	13 (5)
Lymphopenia ^e	31 (12)	11 (4)	10 (4)	25 (10)	7 (3)	8 (3)
Febrile neutropenia	14 (5)	0	14 (5)	11 (4)	0	11 (4)
GI, No. (%)						
Diarrhea	152 (57)	56 (21)	25 (9)	41 (16)	12 (5)	3 (1)
Nausea	148 (55)	56 (21)	3 (1)	77 (31)	23 (9)	7 (3)
Vomiting	50 (19)	12 (4)	1 (< 1)	30 (12)	8 (3)	4 (2)
Constipation	49 (18)	8 (3)	0	36 (14)	8 (3)	0
Abdominal pain	34 (13)	12 (4)	2 (1)	17 (7)	4 (2)	0
Others, No. (%)						
Alopecia	123 (46)	105 (39)	0	41 (16)	18 (7)	0
Fatigue	100 (37)	37 (14)	15 (6)	73 (29)	31 (12)	6 (2)
Asthenia	53 (20)	26 (10)	5 (2)	37 (15)	19 (8)	2 (1)
Decreased appetite	41 (15)	9 (3)	1 (< 1)	34 (14)	13 (5)	1 (< 1)
Neuropathy ^f	23 (9)	8 (3)	3 (1)	38 (15)	16 (6)	6 (2)

NOTE. Assessed in the safety population.

Abbreviations: AE, adverse event; SG, sacituzumab govitecan.

^aPatients may report more than one event per preferred term. AEs were coded using Medical Dictionary for Regulatory Activities v24.0, and AE severity was graded per National Cancer Institute Common Terminology Criteria for Adverse Events v5.0.

^bCombined preferred terms of neutropenia and neutrophil count decreased.

^cCombined preferred terms of anemia, hemoglobin decreased, and red blood cell count decreased.

^dCombined preferred terms of leukopenia and WBC count decreased.

^eCombined preferred terms of lymphopenia and lymphocyte count decreased.

^fCombined preferred terms of gait disturbance, hypoesthesia, muscular weakness, neuropathy peripheral, paresthesia, and peripheral sensory neuropathy.

context of the totality of efficacy data from this trial in the late-line setting, which all favored SG over chemotherapy, with a substantially higher proportion of patients alive and progression-free at all landmark time points. This novel agent directed to a highly expressed target may represent an important treatment option for these patients, addressing a critical unmet medical need. In accordance with the NCCN Guidelines for Breast Cancer, SG is a preferred therapy option for patients with HR+/HER2– cancers after prior

treatment including endocrine therapy, a CDK4/6i and at least two lines of chemotherapy (including a taxane) for advanced breast cancer.²¹ Neutropenia and diarrhea are known AEs with SG and should be prevented and managed according to established guidelines.²⁸⁻³² Additional phase III studies evaluating SG in HR+ breast cancer are underway, including GBG102-SASCIA (ClinicalTrials.gov identifier: [NCT04595565](#))³³ and EVER-132-002 (ClinicalTrials.gov identifier: [NCT04639986](#)).

AFFILIATIONS

¹Department of Medicine, University of California San Francisco Helen Diller Family Comprehensive Cancer Center, San Francisco, CA

²Medical Oncology, Massachusetts General Hospital Cancer Center, Harvard Medical School, Boston, MA

³Medical Faculty Mannheim, Heidelberg University, University Hospital Mannheim, Heidelberg, Germany

⁴Medical Oncology Department, International Breast Cancer Center, Pangaea Oncology, Quironsalud Group, Madrid and Barcelona, Spain

⁵Faculty of Biomedical and Health Sciences, Department of Medicine, Universidad Europea de Madrid, Madrid, Spain

⁶Barts Cancer Institute, Queen Mary University of London, London, United Kingdom

⁷Medical Oncology Department and D3i, Institut Curie, Paris, France

⁸Medical Oncology Department, Centre Léon Bérard, Lyon, France

⁹Medical Oncology, Hospital Universitario 12 de Octubre, Madrid, Spain

¹⁰Institut Claudius Regaud, IUCT-Oncopole, Toulouse, France

¹¹Hospital Universitari Vall D'Hebron, Barcelona, Spain

¹²Memorial Sloan-Kettering Cancer Center, New York, NY

¹³Department of Clinical Development, Gilead Sciences Inc, Foster City, CA

¹⁴Department of Global Patient Safety, Gilead Sciences Inc, Foster City, CA

¹⁵Department of Biostatistics, Gilead Sciences Inc, Foster City, CA

¹⁶Department of Medical Oncology, Dana-Farber Cancer Institute, Boston, MA

CORRESPONDING AUTHOR

Hope S. Rugo, MD, University of California San Francisco Helen Diller Family Comprehensive Cancer Center, 1825 4th St, San Francisco, CA 94158; e-mail: Hope.Rugo@ucsf.edu.

EQUAL CONTRIBUTION

H.S.R. and A.B. contributed equally to this work.

SUPPORT

Supported by Gilead Sciences Inc.

CLINICAL TRIAL INFORMATION

NCT03901339 and EudraCT 2018-004201-33.

AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Disclosures provided by the authors are available with this article at DOI <https://doi.org/10.1200/JCO.22.01002>.

REFERENCES

1. Finn RS, Boer K, Bondarenko I, et al: Overall survival results from the randomized phase 2 study of palbociclib in combination with letrozole versus letrozole alone for first-line treatment of ER+/HER2- advanced breast cancer (PALOMA-1, TRIO-18). *Breast Cancer Res Treat* 183:419-428, 2020
2. Slamon DJ, Neven P, Chia S, et al: Overall survival with ribociclib plus fulvestrant in advanced breast cancer. *N Engl J Med* 382:514-524, 2020
3. Sledge GW Jr, Toi M, Neven P, et al: The effect of abemaciclib plus fulvestrant on overall survival in hormone receptor-positive, ERBB2-negative breast cancer that progressed on endocrine therapy-MONARCH 2: A randomized clinical trial. *JAMA Oncol* 6:116-124, 2020
4. Gao JJ, Cheng J, Bloomquist E, et al: CDK4/6 inhibitor treatment for patients with hormone receptor-positive, HER2-negative, advanced or metastatic breast cancer: A US Food and Drug Administration pooled analysis. *Lancet Oncol* 21:250-260, 2020
5. Gennari A, Andre F, Barrios CH, et al: ESMO Clinical Practice Guideline for the diagnosis, staging and treatment of patients with metastatic breast cancer. *Ann Oncol* 32:1475-1495, 2021
6. Rugo HS, Rumble RB, Macrae E, et al: Endocrine therapy for hormone receptor-positive metastatic breast cancer: American Society of Clinical Oncology guideline. *J Clin Oncol* 34:3069-3103, 2016
7. Cazzaniga ME, Pinotti G, Montagna E, et al: Metronomic chemotherapy for advanced breast cancer patients in the real world practice: Final results of the VICTOR-6 study. *Breast* 48:7-16, 2019
8. Cortes J, O'Shaughnessy J, Loesch D, et al: Eribulin monotherapy versus treatment of physician's choice in patients with metastatic breast cancer (EMBRACE): A phase 3 open-label randomised study. *Lancet* 377:914-923, 2011
9. Twelves C, Awada A, Cortes J, et al: Subgroup analyses from a phase 3, open-label, randomized study of eribulin mesylate versus capecitabine in pretreated patients with advanced or metastatic breast cancer. *Breast Cancer (Auckl)* 10:77-84, 2016
10. Yuan P, Hu X, Sun T, et al: Eribulin mesilate versus vinorelbine in women with locally recurrent or metastatic breast cancer: A randomised clinical trial. *J Cancer* 112:57-65, 2019
11. Mathijssen RH, van Alphen RJ, Verweij J, et al: Clinical pharmacokinetics and metabolism of irinotecan (CPT-11). *Clin Cancer Res* 7:2182-2194, 2001
12. Bardia A: A closer look at sacituzumab govitecan-hziy. *Clin Adv Hematol Oncol* 18:715-717, 2020
13. Vidula N, Yau C, Rugo HS: Trop2 gene expression (Trop2e) in primary breast cancer (BC): Correlations with clinical and tumor characteristics. *J Clin Oncol* 35: 1075-1075, 2017
14. Bardia A, Tolane SM, Punie K, et al: Biomarker analyses in the phase III ASCENT study of sacituzumab govitecan versus chemotherapy in patients with metastatic triple-negative breast cancer. *Ann Oncol* 32:1148-1156, 2021
15. Ambrogio F, Fornili M, Boracchi P, et al: Trop-2 is a determinant of breast cancer survival. *PLoS One* 9:e96993, 2014
16. Goldenberg DM, Stein R, Sharkey RM: The emergence of trophoblast cell-surface antigen 2 (TROP-2) as a novel cancer target. *Oncotarget* 9:28989-29006, 2018
17. TRODELVY (Sacituzumab Govitecan-Hziy). Prescribing Information. Gilead Sciences, Inc, 2021. https://www.gilead.com/-/media/files/pdfs/medicines/oncology/trodelvy/trodelvy_pi.pdf
18. Gilead Sciences Inc: Sacituzumab govitecan receives positive CHMP opinion as 2L treatment for adult patients with metastatic triple-negative breast cancer, 2021. <https://www.gilead.com/news-and-press/press-room/press-releases/2021/10/sacituzumab-govitecan-receives-positive-chmp-opinion-as-2l-treatment-for-adult-patients-with-metastatic-triple-negative-breast-cancer>
19. Gilead Sciences Inc: Trodelvy® (sacituzumab govitecan) granted European Commission marketing authorization for treatment of metastatic triple-negative breast cancer in second line, 2021. <https://www.gilead.com/news-and-press/press-room/press-releases/2021/11/trodelvy-sacituzumab-govitecan-granted>

AUTHOR CONTRIBUTIONS

Conception and design: Aditya Bardia, Javier Cortes, Peter Schmid, Sara M. Tolane

Provision of study materials or patients: Aditya Bardia, Frederik Marmé, Javier Cortes, Delphine Loirat, Eva Ciruelos, Florence Dalenc

Collection and assembly of data: Hope S. Rugo, Aditya Bardia, Frederik Marmé, Delphine Loirat, Olivier Trédan, Eva Ciruelos, Patricia Gómez Pardo, Rosemary Delaney, Sara M. Tolane

Data analysis and interpretation: Hope S. Rugo, Aditya Bardia, Peter Schmid, Delphine Loirat, Eva Ciruelos, Florence Dalenc, Komal L.

Jhaveri, Olivia Fu, Lanjia Lin, Wendy Verret, Sara M. Tolane

Manuscript writing: All authors

Final approval of manuscript: All authors

Accountable for all aspects of the work: All authors

ACKNOWLEDGMENT

We thank the patients and their caregivers for helping us realize the possibilities of this research. Thanks to the dedicated clinical trial investigators and their devoted team members who participated in this trial. The study was sponsored by Gilead Sciences Inc and was designed through a collaboration of the sponsor and the lead investigators. The authors would also like to thank Jin You, PharmD, and Dustin Chernick, PhD, of Gilead Sciences for their editorial assistance in preparation of this manuscript. Medical writing and editorial assistance were provided by Shala Thomas, PhD, CMPP, at Team 9 Science. The TROPICS-02 Study Principal Investigators are listed in [Appendix 1](#) (online only).

european-commission-marketing-authorization-for-treatment-of-metastatic-triplenegative-breast-cancer-in-sec

20. Kalinsky K, Diamond JR, Vahdat LT, et al: Sacituzumab govitecan in previously treated hormone receptor-positive/HER2-negative metastatic breast cancer: Final results from a phase I/II, single-arm, basket trial. *Ann Oncol* 31:1709-1718, 2020
21. National Comprehensive Cancer Network Clinical Practice Guidelines: Breast Cancer, V.4.2022. https://www.nccn.org/professionals/physician_gls/pdf/breast.pdf
22. Eisenhauer EA, Therasse P, Bogaerts J, et al: New response evaluation criteria in solid tumours: Revised RECIST guideline (version 1.1). *Eur J Cancer* 45: 228-247, 2009
23. HALAVEN (Eribulin Mesylate). Prescribing Information. Eisai, 2021. <http://www.halaven.com/-/media/Files/Halaven/HALAVEN-Full-Prescribing-Information.pdf?v=20211013>
24. Kaufman PA, Awada A, Twelves C, et al: Phase III open-label randomized study of eribulin mesylate versus capecitabine in patients with locally advanced or metastatic breast cancer previously treated with an anthracycline and a taxane. *J Clin Oncol* 33:594-601, 2015
25. Pivot X, Marme F, Koenigsberg R, et al: Pooled analyses of eribulin in metastatic breast cancer patients with at least one prior chemotherapy. *Ann Oncol* 27: 1525-1531, 2016
26. Modi S, Jacot W, Yamashita T, et al: Trastuzumab deruxtecan in previously treated HER2-low advanced breast cancer. *N Engl J Med* 387:9-20, 2022
27. Bardia A, Hurvitz SA, Tolaney SM, et al: Sacituzumab govitecan in metastatic triple-negative breast cancer. *N Engl J Med* 384:1529-1541, 2021
28. American Society of Clinical Oncology, Kris MG, Hesketh PJ, et al: American Society of Clinical Oncology guideline for antiemetics in oncology: Update 2006. *J Clin Oncol* 24:2932-2947, 2006
29. Benson AB III, Ajani JA, Catalano RB, et al: Recommended guidelines for the treatment of cancer treatment-induced diarrhea. *J Clin Oncol* 22:2918-2926, 2004
30. Jordan K, Feyer P, Holler U, et al: Supportive treatments for patients with cancer. *Dtsch Arztebl Int* 114:481-487, 2017
31. National Comprehensive Cancer Network Clinical Practice Guidelines: Hematopoietic Growth Factors, V.1.2022. https://www.nccn.org/professionals/physician_gls/pdf/growthfactors.pdf
32. National Comprehensive Cancer Network Clinical Practice Guidelines: Palliative Care, V.1.2022. https://www.nccn.org/professionals/physician_gls/pdf/palliative.pdf
33. Marme F, Stickeler E, Furlanetto J, et al: Phase III postneoadjuvant study evaluating sacituzumab govitecan, an antibody drug conjugate in primary HER2-negative breast cancer patients with high relapse risk after standard neoadjuvant treatment: SASCIA. *J Clin Oncol* 39:1075-1075, 2021



AUTHORS' DISCLOSURES OF POTENTIAL CONFLICTS OF INTEREST

Sacituzumab Govitecan in Hormone Receptor–Positive/Human Epidermal Growth Factor Receptor 2–Negative Metastatic Breast Cancer

The following represents disclosure information provided by authors of this manuscript. All relationships are considered compensated unless otherwise noted. Relationships are self-held unless noted. I = Immediate Family Member, Inst = My Institution. Relationships may not relate to the subject matter of this manuscript. For more information about ASCO's conflict of interest policy, please refer to www.asco.org/rwc or ascopubs.org/jco/authors/author-center.

Open Payments is a public database containing information reported by companies about payments made to US-licensed physicians ([Open Payments](https://openpaymentsdata.cms.gov)).

Hope S. Rugo

Honoraria: Puma Biotechnology, Mylan, Samsung Bioepis, Chugai Pharma, Blueprint Medicines, Napo Pharmaceuticals

Research Funding: OBI Pharma (Inst), Pfizer (Inst), Novartis (Inst), Lilly (Inst), Genentech (Inst), Merck (Inst), Odonate Therapeutics (Inst), Daiichi Sankyo (Inst), Sermonix Pharmaceuticals (Inst), AstraZeneca (Inst), Gilead Sciences (Inst), Ayala Pharmaceuticals (Inst), Astellas Pharma (Inst), Seattle Genetics (Inst), MacroGenics (Inst), Boehringer Ingelheim (Inst), Polyphor (Inst)

Open Payments Link: <https://openpaymentsdata.cms.gov/summary>

Aditya Bardia

Consulting or Advisory Role: Novartis (Inst), Genentech, Pfizer, Spectrum Pharmaceuticals, BioTheragnostics, Merck, Radius Health (Inst), Immunomedics (Inst), Genentech/Roche (Inst), Pfizer (Inst), Innocrin Pharma (Inst), Sanofi, Puma Biotechnology, Daiichi Sankyo/AstraZeneca, Foundation Medicine, Philips Healthcare

Research Funding: Genentech (Inst), Novartis (Inst), Pfizer (Inst), Merck (Inst), Sanofi (Inst), Radius Health (Inst), Immunomedics (Inst), AstraZeneca/Daiichi Sankyo (Inst)

Open Payments Link: <https://openpaymentsdata.cms.gov/physician/523675>

Frederik Marmé

Honoraria: Roche/Genentech, Novartis, Pfizer, AstraZeneca, Tesaro, Clovis Oncology, Eisai, Celgene, Genomic Health, PharmaMar, Amgen, MSD Oncology, Immunomedics (Inst), Settle Genetics, Myriad Genetics, Pierre Fabre, GlaxoSmithKline, AGENDIA, Lilly, Gilead Sciences

Consulting or Advisory Role: AstraZeneca (Inst), Tesaro, Pfizer, Roche (Inst), Genomic Health, CureVac, Amgen, Vaccibody (Inst), Immunomedics (Inst), Eisai, GlaxoSmithKline, Gilead Sciences

Research Funding: Roche/Genentech (Inst), Novartis (Inst), AstraZeneca (Inst), Tesaro (Inst), Clovis Oncology (Inst), MSD Oncology (Inst), Vaccibody (Inst), Gilead Sciences (Inst), GlaxoSmithKline (Inst)

Travel, Accommodations, Expenses: Roche, Pfizer, AstraZeneca

Javier Cortes

Stock and Other Ownership Interests: MedSIR, Nektar, Leuko (I)

Honoraria: Novartis, Eisai, Celgene, Pfizer, Roche, Samsung, Lilly, Merck Sharp & Dohme, Daiichi Sankyo

Consulting or Advisory Role: Celgene, Cellectis Biotech, AstraZeneca, Roche, Settle Genetics, Daiichi Sankyo, ERYTECH Pharma, Polyphor, Athenex, Lilly, Servier, Merck Sharp & Dohme, GlaxoSmithKline, Leuko, Clovis Oncology, Bioasis, Boehringer Ingelheim, Ellipses Pharma, HiberCell, BiolInvent, GEMoaB, Gilead Sciences, Menarini, Zymeworks, Reveal Genomics

Research Funding: ARIAD (Inst), AstraZeneca (Inst), Baxalta (Inst), Bayer (Inst), Eisai (Inst), Guardant Health (Inst), Merck Sharp & Dohme (Inst), Pfizer (Inst), Puma Biotechnology (Inst), Queen Mary University of London (Inst), Roche (Inst), Piquor (Inst)

Patents, Royalties, Other Intellectual Property: Pharmaceuticals Combinations of A PI3k Inhibitor and a Microtubule Destabilizing Agent. Javier Cortes Castan, Alejandro Piris Gimenez, Violeta Srra Elizalde. WO 2014/199294 A, Her2 as a predictor of response to dual HER2 blockade in the absence of cytotoxic therapy, Aleix Prat, Antonio Llombart, Javier Cortes, US 2019/0338368 A1

Travel, Accommodations, Expenses: Roche, Pfizer, Eisai, Novartis, Daiichi Sankyo, Gilead Sciences

Peter Schmid

Employment: Roche (I)

Honoraria: AstraZeneca, Bayer, Boehringer Ingelheim, Merck, Novartis, Pfizer, Puma Biotechnology, Roche, Eisai, Celgene

Consulting or Advisory Role: Genentech/Roche (I), AstraZeneca, Merck, Boehringer Ingelheim, Bayer, Pfizer, Novartis, Eisai, Celgene, Puma Biotechnology

Research Funding: AstraZeneca (Inst), Astellas Pharma (Inst), OncoGenex (Inst), Genetech (Inst), Novartis (Inst), Roche (Inst), Medivation

Delphine Loirat

Honoraria: MSD, Gilead Sciences, AstraZeneca, Lilly

Consulting or Advisory Role: Roche, MSD Oncology, AstraZeneca, Novartis, Pfizer, Immunomedics, Gilead Sciences, 4D Pharma, Lilly

Travel, Accommodations, Expenses: Roche, AstraZeneca, MSD, Pfizer, Gilead Sciences

Olivier Trédan

Consulting or Advisory Role: Roche, Pfizer, Lilly, AstraZeneca, MSD Oncology, Daiichi Sankyo Europe GmbH, Eisai Europe, Sandoz-Novartis, Seattle Genetics, Pierre Fabre, Gilead Sciences

Research Funding: Roche (Inst), Bristol Myers Squibb (Inst), MSD Oncology (Inst), AstraZeneca (Inst), Novartis (Inst), Bayer (Inst),

Travel, Accommodations, Expenses: Roche, Novartis, Pfizer, Lilly, AstraZeneca, MSD Oncology

Eva Ciruelos

Consulting or Advisory Role: Roche, Pfizer, AstraZeneca, Novartis, Lilly, MSD Oncology, Daiichi Sankyo/AstraZeneca

Speakers' Bureau: Lilly, Roche, Pfizer, Daiichi Sankyo/AstraZeneca, Novartis

Travel, Accommodations, Expenses: Roche, Pfizer

Komal L. Jhaveri

Consulting or Advisory Role: Novartis, Pfizer, AstraZeneca, Jounce Therapeutics, Synthron, IntelliSphere, Bristol Myers Squibb, Genentech, AbbVie, Lilly, Blueprint Medicines, Seattle Genetics, Daiichi Sankyo, BioTheragnostics, Sun Pharma Advanced Research Company, Taiho Oncology, Sanofi, Gilead Sciences

Research Funding: Novartis (Inst), Genentech (Inst), Debiopharm Group (Inst), ADC Therapeutics (Inst), Pfizer (Inst), Novita Pharmaceuticals (Inst), Clovis Oncology (Inst), Lilly (Inst), Zymeworks (Inst), Immunomedics (Inst), Puma Biotechnology (Inst), VelosBio/Merck (Inst), AstraZeneca (Inst)

Travel, Accommodations, Expenses: Taiho Pharmaceutical, Jounce Therapeutics, Pfizer, AstraZeneca, IntelliSphere, Lilly, Gilead Sciences

Rosemary Delaney

Employment: Gilead Sciences

Stock and Other Ownership Interests: Immunomedics (I), Gilead Sciences

Olivia Fu

Employment: Gilead Sciences

Stock and Other Ownership Interests: Gilead Sciences

Travel, Accommodations, Expenses: Gilead Sciences

Lanjia Lin

Employment: Gilead Sciences

Stock and Other Ownership Interests: Gilead Sciences

Travel, Accommodations, Expenses: Gilead Sciences

Wendy Verret

Employment: Roche/Genentech, Gilead Sciences

Sara M. Tolaney

This author is a member of the *Journal of Clinical Oncology* Editorial Board. Journal policy recused the author from having any role in the peer review of this manuscript.

Consulting or Advisory Role: Novartis, Pfizer, Merck, Lilly, Nektar, NanoString Technologies, AstraZeneca, Puma Biotechnology, Genentech, Eisai, Sanofi, Bristol Myers Squibb, Paxman, Seattle Genetics, Odonate Therapeutics, G1 Therapeutics, OncoPep, Kyowa Hakko Kirin, Samsung Bioepis, CytomX Therapeutics, Daiichi Sankyo, Athenex, Immunomedics/Gilead, Mersana, Certara Inc, 4D Pharma, Ellipses Pharma, OncoSec, Chugai Pharma, BeyondSpring Pharmaceuticals, OncXerna Therapeutics, Infinity Pharmaceuticals, Zymeworks, Zentalis, BluePrint Medicines, Reveal Genomics

Research Funding: Genentech/Roche (Inst), Merck (Inst), Exelixis (Inst), Pfizer (Inst), Lilly (Inst), Novartis (Inst), Bristol Myers Squibb (Inst), Eisai (Inst), AstraZeneca (Inst), NanoString Technologies (Inst), Cyclacel (Inst), Nektar (Inst), Immunomedics (Inst), Odonate Therapeutics (Inst), Sanofi (Inst), Settle Genetics (Inst)

No other potential conflicts of interest were reported.

APPENDIX 1. LIST OF TROPICS-02 STUDY PRINCIPAL INVESTIGATORS

Study Investigator	Study Institution
Maysa Abu-Khalaf	Thomas Jefferson University
Foluso Ademuyiwa	Washington University School of Medicine
Clarence Adoo	HonorHealth Research Institute
Philippe Aftimos	Institut Jules Bordet
Samreen Ahmed	University Hospitals of Leicester NHS Trust
Cristina Alencar	Virginia Oncology Associates
Anne Armstrong	The Christie NHS Foundation Trust
Cristina Arqueros	Hospital de la Santa Creu i de Sant Pau
Jean-Pierre Ayoub	Centre Hospitalier De l'Universite De Montreal CHUM
Aditya Bardia ^a	Massachusetts General Hospital
J. Thaddeus Beck	Highlands Oncology Group
Giampaolo Bianchini	Ospedale San Raffaele
Sibel Blau	Northwest Medical Specialties, PLLC
Nathalie Bonnin	Hospices Civils de Lyon
Adam Brufsky	UPMC Hillman Cancer Center
Christina Brzezniak	Virginia Cancer Specialists, PC
Giuseppe Cairo	Ospedale Vito Fazzi di Lecce
Lourdes Calvo Martinez	Complejo Hospitalario Universitario A Coruña
Luigi Cavanna	Azienda Unità Sanitaria Locale di Piacenza-Ospedale Guglielmo da Saliceto
Marina Elena Cazzaniga	Azienda Ospedaliera San Gerardo di Monza
Eva Ciruelos	Hospital Universitario 12 de Octubre
Patrick Cobb	Saint Vincent Frontier Cancer Center
Javier Cortes Castan ^a	Hospital Ruber Internacional
Ricardo Costa	Moffitt Cancer Center
Florence Dalenc	Institut Claudius Regaud
Maaïke de Boer	Maastricht University Medical Center
Neelam Desai	Beth Israel Deaconess Medical Center
Jennifer Diamond	University of Colorado
Erion Dobi	Hôpital Jean Minjoz
Lara Durna	Kaiser Permanente Medical Center
Peter Fasching	Universitätsklinik Erlangen
Gianluigi Ferretti	Istituto Regina Elena
Nelly Firmin	Institut Regional du Cancer de Montpellier
Keerthi Gogineni	Emory University
Patricia Gomez Pardo	Hospital Universitari Vall D'Hebron
Julien Grenier	Institut Sainte Catherine
Erika Hamilton	Tennessee Oncology
Susanna Hegewisch-Becker	Onkologische Schwerpunktpraxis Eppendorf
(continued on following page)	

(continued)

Study Investigator	Study Institution
Stephanie Henry	CHU UCL NAMUR—Sainte Elisabeth
Dawn Hershman	Columbia University Medical Center
Claudine Isaacs	Georgetown University
Jean-Philippe Jacquin	Institut de Cancerologie Lucien Neuwirth
Komal Jhaveri	Memorial Sloan-Kettering Cancer Center
Virginia Kaklamani	University of Texas Health Science Center at San Antonio
Hans-Christian Kolberg	Marienhospital Bottrop
Amy Krie	Virginia Piper Cancer Center (Alliant Health)
Christian Kurbacher	Gynakologisches Zentrum Bonn
Charles Kurkul	Texas Oncology PA
Sylvain Ladoire	Centre Georges François Leclerc
Sabine Linn	Antoni van Leeuwenhoekziekenhuis
Delphine Loirat	Institut Curie
Rafael Lopez	Hospital Clínico Universitario de Santiago de Compostela—CHUS
Kristina Lübke	DIAKOVERE Krankenhaus gGmbH Henriettenstift—Standort Kirchrode
Hans-Joachim Luck	Gynakologisch-Onkologische Praxis Hannover
Elisabeth Luporsi	Hôpital de Mercy
Ling Ma	Rocky Mountain Cancer Center
Rolf Mahlberg	Klinikum Mutterhaus der Borromäerinnen
Frederik Marne ^a	Universitätsmedizin Mannheim
Miguel Martin Jimenez	Hospital General Universitario Gregorio Marañón
Eduardo Martinez	Hospital Provincial de Castellón
Ingrid Mayer	Vanderbilt University
Kelly McCann	University of California Los Angeles (UCLA)
Serafin Morales	Hospital Universitari Arnau De Vilanova (Huav)
Sarah Mougalian	Yale University
Rita Nanda	University of Chicago
Polly Niravath	Houston Methodist Hospital
Yelena Novik	New York University School of Medicine
Shannon O'Connor	Maryland Oncology Hematology
Alicia Okines	The Royal Marsden NHS Foundation Trust
Coral Omene	Rutgers Cancer Institute of New Jersey
Rianne Oosterkamp	Medisch Centrum Haaglanden Antoniushove
Joyce O'Shaughnessy	Texas Oncology
Steven Papish	Summit Medical Group
Ritesh Parajuli	University of California Irvine (UCIMC)
Felicity Paterson	Royal Surrey County Hospital NHS Foundation Trust
Michel Pavic	Centre Hospitalier De l'Université De Sherbrooke Hôpital Fleurimont
Rebecca Pedersini	Azienda Ospedaliera Spedali Civili di Brescia
Alejandra Perez	Sylvester Comprehensive Cancer Center
Timothy Pluard	St Luke's Cancer Institute
(continued on following page)	

(continued)

Study Investigator	Study Institution
David Potter	Masonic Cancer Center
Paola Pozzi	Ospedale di Desio
Kevin Punie	Universitaire Ziekenhuis Leuven
Fabricio Racca	Hospital Quironsalud Barcelona
Daniel Rayson	Queen Elizabeth II Health Sciences Center
Mattea Reinisch	Kliniken Essen-Mitte
Paul Richards	Blue Ridge Cancer Care
Elizabeth Riley	James Graham Brown Cancer Center
David Riseberg	Mercy Medical Center
Regan Rostorfer	University of Florida
Hope Rugo ^a	University of California San Francisco
Manuel Ruiz Borrego	Hospital Universitario Virgen del Rocío
Sagar Sardesai	Stefanie Spielman Comprehensive Breast Center
Peter Schmid ^a	Barts Health NHS Trust
Andreas Schneeweiss	Nationales Centrum für Tumorerkrankungen Heidelberg
Holger Schultz	HELIOS Klinikum Berlin-Buch
Lee Schwartzberg	West Clinic
Lasika Seneviratne	Los Angeles Hematology Oncology Medical Group
Priyanka Sharma	University of Kansas
Sasha Strain	Texas Oncology PA
Rachel Swart	Arizona Oncology Associates
Karen Tedesco	New York Oncology Hematology—Amsterdam Cancer Center
Hans Tesch	Centrum für Hämatologie und Onkologie Bethanien
Amy Tiersten	Mount Sinai
Sara Tolaney ^a	Dana Farber Cancer Institute
Olivier Trédan	Centre Leon Berard
Nisha Unni	University of Texas Southwestern Medical Center
Thierry Velu	Chirec Cancer Institute
Hermann Voss	Städtisches Klinikum Dessau
Grace Wang	Baptist Health—Miami Cancer Institute
Rudolf Weide	Praxisklinik für Hämatologie und Onkologie Koblenz
Duncan Wheatley	Royal Cornwall Hospital Trust
Kay Yeung	University of California San Diego
Amelia Zelnak	Northside Hospital Atlanta

^aDenotes members of the TROPiCS-02 scientific steering committee.