

Cellular-Level Effects of Swimming in Chlorinated Pools: Evidence-Based Research Report

Executive Summary

Swimming in chlorinated pools exposes individuals to disinfection by-products (DBPs) that form when chlorine reacts with organic matter from swimmers' bodies. Recent peer-reviewed research demonstrates measurable cellular-level effects, including DNA damage and respiratory epithelial changes, after exposure to chlorinated pool water. Studies show increases in genotoxicity biomarkers (micronuclei and urinary mutagenicity) following 40-minute swimming sessions. However, these effects appear to be transient and may be mitigated through natural DNA repair mechanisms. While current evidence suggests short-term exposure risks are manageable for healthy individuals, long-term effects require further investigation, particularly for vulnerable populations.

Definition & Context

Chlorinated Swimming Pool Exposure refers to human contact with water treated with chlorine-based disinfectants, primarily through swimming, but also via inhalation of airborne chemicals and dermal absorption. Swimming pools typically maintain 1-4 ppm free chlorine levels as recommended by health agencies. When chlorine interacts with organic matter (sweat, skin cells, urine, cosmetics), it forms over 100 different disinfection by-products including trihalomethanes (THMs), haloacetic acids, and chloramines. These DBPs are absorbed through the skin, lungs, and inadvertently ingested during swimming activities.

Common Settings:

- Indoor and outdoor public swimming pools
- Private residential pools
- Competitive swimming facilities
- Aquatic therapy centers
- Hotel and recreational facility pools

Benefits / Positive Effects

Benefit	Why It Matters	Supporting Evidence
Pathogen Control	Prevents waterborne diseases and infections	CDC guidelines confirm chlorination effectively kills bacteria, viruses, and parasites
Safe Recreation	Enables safe aquatic activities for millions	Public health data shows dramatic reduction in pool-related disease outbreaks with proper chlorination
Exercise Facilitation	Supports cardiovascular and muscle health through swimming	Well-established exercise physiology research on swimming benefits
Accessibility	Makes pool use safe for vulnerable populations	Properly maintained pools reduce infection risks for immunocompromised individuals

Risks / Negative Effects

Risk	Potential Impact	Evidence / Source
DNA Damage (Genotoxicity)	Moderate	Research shows increased micronuclei and urinary mutagenicity after 40-min exposure (Environmental Health Perspectives, 2010)
Respiratory Epithelial Changes	Mild-Moderate	Increased CC16 protein indicating lung epithelium permeability changes (UAB/CREAL study)
Cellular Cytotoxicity	Moderate	Pool water demonstrates higher cell-killing capacity than potable water at lower concentrations
Gene Expression Changes	Mild-Moderate	Blood RNA alterations detected after chlorinated pool swimming (ScienceDirect, 2017)
Immune System Perturbation	Mild	Acute alterations in cytokine and chemokine secretion patterns observed
Mutagenicity Exposure	Moderate	Urine mutagenicity increases significantly after swimming, correlating with bromoform levels
Asthma Exacerbation	Moderate	Chloramines in indoor pools linked to respiratory symptoms, particularly in children
Potential Cancer Risk	Uncertain/Long-term	Epidemiological associations with bladder cancer, though causation not established

Toxicity & Safety Profile

Short-term Exposure (Single Session):

- DNA damage markers increase measurably but appear reversible
- Respiratory symptoms mild and transient
- No acute systemic toxicity at normal pool concentrations

Long-term Exposure (Regular Swimming):

- Effects poorly understood and require further research
- Occupational lifeguards show higher DNA damage levels
- Chronic respiratory symptoms possible with poor indoor air quality

Exposure Thresholds:

- **Free Chlorine:** 1-4 ppm (CDC recommended range)
- **Combined Chlorine (Chloramines):** ≤ 0.4 ppm (health department limits)
- **pH:** 7.0-7.8 (optimal range for disinfection efficiency)

Health Agencies' Positions:

- **CDC:** Maintains that benefits of swimming outweigh risks when pools are properly maintained
- **EPA:** Regulates DBPs in drinking water but limited pool water standards
- **WHO:** Acknowledges DBP concerns but emphasizes proper pool management
- No specific cellular-level exposure limits established by major agencies

Vulnerable Groups:

- **Children:** Higher surface area-to-body weight ratio increases exposure; developing respiratory systems more susceptible
- **Pregnant Women:** Potential concerns for fetal development, though evidence limited
- **Asthmatics:** Increased risk of respiratory symptoms from chloramines
- **Immunocompromised:** May be more susceptible to both infections and chemical effects
- **Professional Swimmers/Lifeguards:** Chronic high-level exposure concerns

Environmental Impact

Air Quality: Indoor pools generate chloramines that degrade air quality and can cause respiratory irritation. Poor ventilation concentrates these compounds.

Water Systems: DBPs from pool backwash may enter wastewater systems, though treatment typically removes most compounds.

Wildlife: Pool chemicals can be toxic to aquatic life if improperly discharged; chlorinated water should be dechlorinated before environmental release.

Who Should / Should Not Be Exposed

Generally Safe For:

- Healthy adults and children with proper pool maintenance
- Occasional recreational swimmers
- Competitive swimmers with adequate recovery periods

Increased Caution Recommended:

- **Pregnant Women:** Consult healthcare providers, especially for frequent exposure
- **Severe Asthmatics:** Consider alternative exercise options or well-ventilated pools
- **Children Under 3:** Limited exposure time and ensure proper pool maintenance
- **Professional Athletes:** Monitor for cumulative effects with regular health assessments

Should Avoid or Minimize:

- Individuals with severe respiratory conditions
- Those with documented chemical sensitivities
- People with compromised DNA repair mechanisms (rare genetic conditions)

Safer Alternatives or Mitigation Strategies

Pool Selection:

- Choose well-ventilated outdoor pools when possible
- Select facilities with proper maintenance and water quality testing
- Avoid overcrowded pools with visible chloramine odors

Personal Protection:

- Shower before and immediately after swimming
- Use swim caps and goggles to reduce chemical contact
- Minimize time in poorly ventilated indoor pools
- Stay hydrated to support natural detoxification

Alternative Disinfection:

- UV-treated pools (reduced chemical load)
- Saltwater pools (generate chlorine differently)
- Ozone-treated facilities
- Natural swimming pools (where available)

Facility Management:

- Proper ventilation systems in indoor facilities
- Regular water testing and chemical balancing
- Enforcement of hygiene rules (mandatory showers)
- Use of secondary disinfection methods

Comparison

Option	Pros	Cons	Best For
Chlorinated Pools	Effective pathogen control, widely available, cost-effective	DNA damage risk, respiratory effects, chemical odors	General public recreation, competitive swimming
UV + Low Chlorine	Reduced chemical exposure, effective disinfection	Higher initial costs, requires backup disinfection	Health-conscious swimmers, sensitive individuals
Saltwater Pools	Gentler on skin/eyes, reduced chloramine formation	Still produces chlorine, maintenance complexity	Frequent swimmers, those with skin sensitivities
Natural Pools	No chemical exposure, ecological benefits	Limited pathogen control, seasonal availability	Environmentally conscious swimmers, low-intensity use
Open Water	No chemical exposure, natural environment	Weather dependent, potential biological hazards	Experienced swimmers, seasonal use

Sources

1. Kogevinas, M. et al. (2010). Genotoxic Effects in Swimmers Exposed to Disinfection By-products in Indoor Swimming Pools. Environmental Health Perspectives.
<https://pmc.ncbi.nlm.nih.gov/articles/PMC2974689/>
2. University Autonomous Barcelona (UAB). Swimming in chlorinated pools can damage DNA. Research dissemination. <https://www.uab.cat/web/news-detail/swimming-in-chlorinated-pools-can-damage-dna-1345680342044.html?articleId=1284964586805>
3. Richardson, S.D. et al. (2010). What's in the Pool? A Comprehensive Identification of Disinfection By-products and Assessment of Mutagenicity. Environmental Health Perspectives.
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2974688/>
4. Centers for Disease Control and Prevention (2025). Guidelines for Keeping Your Pool Safe and Healthy. <https://www.cdc.gov/healthy-swimming/safety/what-you-can-do-to-stay-healthy-in-swimming-pools.html>
5. Villanueva, C. (2019). DNA-damaging disinfection by-products found in pool water. Science News.
<https://www.sciencenews.org/article/dna-damaging-disinfection-products-found-pool-water>

6. CDC (2025). Chloramines and Pool Operation. <https://www.cdc.gov/healthy-swimming/toolkit/chloramines-and-pool-operation.html>
7. EPA (2024). Recreational Water Quality Criteria and Methods. <https://www.epa.gov/wqc/recreational-water-quality-criteria-and-methods>
8. Font-Ribera, L. et al. (2017). Gene expression changes in blood RNA after swimming in a chlorinated pool. Environment International. <https://www.sciencedirect.com/science/article/abs/pii/S100107421631261X>
9. Parenting Science (2021). Swimming pools and health risks: A parent's evidence-based guide. <https://parentingscience.com/swimming-pools/>
10. Ma, S. et al. (2015). Health Effects from Swimming Training in Chlorinated Pools and the Corresponding Metabolic Stress Pathways. PLOS One. <https://pmc.ncbi.nlm.nih.gov/articles/PMC4351252/>