

Low dose ketamine IV infusion

Following Complex Spinal Surgery



Use Lock-Box on ward
(controlled drug)

Do usual post-op Vital Signs

To LOCK SCREEN on BBraun pump

Scroll down to OPTIONS, ◀ select

DATA LOCK, ◀ select

Choose level 1 or 2

Enter Code: **2222**

Do **NOT** display poster in patient area with **code** written.

IV Ketamine low dose may continue on wards

100mg in 50mL (2mg/mL) running at **1.5mL/hr** (3mg/hr)

Pain Team/Anaesthetist will review daily and advise whether to continue.

Contact Acute Pain Team / Anaesthetist for:

- Instructions to continue before syringe is finished
 - may run for up to 72hrs as indicated
- Any concerns with pain, sedation or any adverse effects.



Key for
lock-box is
on CD key
bunch

Ward staff may administer pre-filled syringe.

Will require concurrent simple analgesics plus appropriate opioid analgesics for surgical pain.

Indication:

Following complex spinal surgery

Following major spinal surgery and
patient is opioid tolerant or dependent

Difficult to manage neuropathic pain
post-operatively

Acute exacerbation of chronic pain
who present for major or complex
spinal surgery

Contraindication: NEVER give to
these patients.

Patient has 'brain' pathology or has
had Brain Surgery

Raised intracranial / intraocular
pressure

Poorly controlled cardio-vascular
disease including Uncontrolled
Hypertension

Severe liver disease

Active psychosis

Porphyria (metabolic disorder)

Stroke

Pregnancy