Non-Abelian Matrix-Condensate Geometrogenesis

Edward Schipani

Contents

Preface **xi**

Notation **xiii**

1 What Geometrogenesis Must Derive **1**
1.1 What this book is trying to derive 1
1.2 Analog gravity, and the ceiling it cannot break 2
1.3 Representation theory in one paragraph 3
1.4 What must be delivered before one may say "this is GR" 4
1.5 The NAMCG derivation in one view 5
1.6 How the later phenomenology sits on the spine 6
1.7 How to read this book 6
1.8 Exercises 7
1.9 Bridge to Chapter 2 7

2 Preliminaries **9**
2.1 Lorentzian manifolds and index conventions 9
2.2 Vierbeine: primitive geometry versus composite geometry 10
2.3 Variation of the Einstein–Hilbert action 11
2.4 Bianchi identities and conservation 12
2.5 Gauge covariant derivatives and a scalar warmup 13
2.6 One-loop effective actions and the heat kernel 13
2.7 Causal diamonds 15
2.8 Mutual information 16
2.9 What this chapter deliberately omits 16
2.10 Exercises 17
2.11 Bridge to Chapter 3 18

3 The microscopic action **19**
3.1 The scalar failure, restated as a field-content problem 19
3.2 Two presentations, and why both appear 20

3.3 Presentation A: matrix coordinates and ultraviolet cutoff 21
3.4 Presentation B: the covariant condensate action 22
3.5 Index structure and the intended infrared spectrum 23
3.6 Equations of motion from Presentation B 24
3.7 Relating the two presentations 24
3.8 Energy scales and coupling constants 30
3.9 What has been assumed, collected 31
3.10 Exercises 31
3.11 Bridge to Chapter 4 32

4 Vierbein emergence **33**
4.1 The constant-vacuum obstruction 33
4.2 The composite metric 35
4.3 What kind of object e_μ^A is 36
4.4 Seiberg–Witten map and the Θ correction 36
4.5 A linearised frame around Minkowski spacetime 37
4.6 Signature and the role of η_{AB} 38
4.7 Worked example: a weak, static frame perturbation 38
4.8 What this chapter has not done 38
4.9 Exercises 39
4.10 Bridge to Chapter 5 40

5 Lorentz decomposition **41**
5.1 Fluctuations about the polar vacuum 41
5.2 Algebraic split under the Lorentz group 42
5.3 Translation into the notation of the source notes 43
5.4 What $T_{\mu\nu}$ is not yet 44
5.5 Kinetic terms before any entanglement condition 44
5.6 Diffeomorphism action on the pieces 45
5.7 Provisional sector bookkeeping 45
5.8 Exercises 46
5.9 Bridge to Chapter 6 47

6 Entanglement equilibrium **49**
6.1 The logical role of this chapter 49
6.2 Diamonds in the emergent geometry 50
6.3 Reduced states and the modular Hamiltonian 51
6.4 Replica formula 52
6.5 The first law of the diamond 52
6.6 Allowed variations 53

6.7 Mutual information of two cores . . . 53
6.8 What stationarity does *not* say . . . 54
6.9 The Ward identity that Chapter 7 will use . . . 55
6.10 Exercises . . . 55
6.11 Bridge to Chapter 7 . . . 56

7 The tensor mass gap **57**
7.1 Strategy . . . 57
7.2 Fierz–Pauli action for the tensor block . . . 58
7.3 Modular stress of the tensor block . . . 59
7.4 Area variation induced by $T_{\mu\nu}$ . . . 59
7.5 Substitution into the Ward identity . . . 60
7.6 The vanishing theorem . . . 61
7.7 Two transverse-traceless polarizations . . . 62
7.8 The van Dam–Veltman–Zakharov warning . . . 62
7.9 Conformal and vector blocks . . . 63
7.10 Exercises . . . 63
7.11 Bridge to Chapter 8 . . . 64

8 Heat kernel and G_{eff} **65**
8.1 One-loop effective action on the emergent geometry . . . 65
8.2 Seeley–DeWitt expansion, used not re-derived . . . 66
8.3 Evaluating a_1 on the condensate . . . 67
8.4 The quartic term . . . 68
8.5 Which background the heat kernel sees . . . 69
8.6 Numerical size . . . 70
8.7 What has been computed . . . 70
8.8 Exercises . . . 70
8.9 Bridge to Chapter 9 . . . 71

9 Ultraviolet completion **73**
9.1 What a_2 contains . . . 74
9.2 Ostrogradsky's theorem, used as a veto . . . 74
9.3 What ultraviolet stability does mean . . . 75
9.4 Cubic operators and Einsteinian cubic gravity . . . 76
9.5 Torsion and higher connections . . . 77
9.6 A checklist before Lorentz locking . . . 78
9.7 Exercises . . . 78
9.8 Bridge to Chapter 10 . . . 79

10 Lorentz locking **81**
10.1 The two groups that act on Φ . . . 81
10.2 The vacuum and the broken generators . . . 82
10.3 Eating ω_{AB} . . . 83
10.4 The internal vector after locking . . . 84
10.5 Diffeomorphisms . . . 86
10.6 Provisional sector bookkeeping (frame/metric sector only) . . . 86
10.7 What locking does not do . . . 87
10.8 Exercises . . . 87
10.9 Bridge to Chapter 11 . . . 87

11 Conservation **89**
11.1 The split . . . 89
11.2 Noether's second theorem, applied once . . . 90
11.3 Structure of the entropic stress . . . 91
11.4 Exchange with matter . . . 93
11.5 Vacuum energy as a conservation constraint . . . 93
11.6 What this chapter has not done . . . 94
11.7 Exercises . . . 94
11.8 Bridge to Chapter 12 . . . 95

12 HBEV **97**
12.1 What would not count as a solution . . . 98
12.2 Boundary geometry . . . 98
12.3 The HBEV action . . . 99
12.4 Stationarity and the induced cosmological term . . . 99
12.5 Conservation . . . 101
12.6 Relation to the diamond entropy of Chapter 6 . . . 101
12.7 Working value used later . . . 102
12.8 Exercises . . . 102
12.9 Bridge to Chapter 13 . . . 102

13 Cores and galaxies **105**
13.1 Static spherical ansatz . . . 105
13.2 Cubic correction in the spherical sector . . . 106
13.3 Horizons and the core . . . 107
13.4 Weak-field exterior and the extra Poisson equation . . . 107
13.5 Circular velocities . . . 109
13.6 What this is not . . . 109
13.7 Matching core to exterior . . . 110

13.8 Exercises . . . 110
13.9 Bridge to Chapter 14 . . . 111

14 Dissipative drag **113**
14.1 Reversible versus irreversible . . . 113
14.2 Worldline reduction . . . 114
14.3 The working exponent . . . 115
14.4 Comparison with quadrupole losses . . . 116
14.5 Tidal disruption . . . 117
14.6 Illegal and legal integrators . . . 117
14.7 Exercises . . . 118
14.8 Bridge to Chapter 15 . . . 119

15 Inspiral laboratory **121**
15.1 The ordinary-differential system . . . 121
15.2 Integrator . . . 122
15.3 Initial data . . . 123
15.4 What the figures are allowed to show . . . 123
15.5 Code . . . 123
15.6 How to misread the figure, and how not to . . . 125
15.7 Exercises . . . 126
15.8 Bridge to Chapter 16 . . . 126

16 Falsifiers **127**
16.1 The chain, with theorem numbers . . . 127
16.2 Inventory of named assumptions . . . 128
16.3 What would kill the framework . . . 130
16.4 What would only constrain a coefficient . . . 130
16.5 Observational checklist . . . 131
16.6 Claims the book has not made . . . 132
16.7 How to use Chapter 15 in public . . . 133
16.8 Exercises . . . 133
16.9 Bridge to Chapter 17 . . . 133

17 Research agenda **135**
17.1 What is closed conditionally inside the book . . . 135
17.2 Status vocabulary and completion labels . . . 136
17.3 Critical-path problems I: B, C, and D . . . 137
17.4 Open problems by scope and priority . . . 141
17.4.1 Critical-path problems A and E . . . 141
17.4.2 Critical-path problems F, G and N . . . 141

17.4.3 Optional problems continued . . . 143
17.5 Problems that look like work and are not . . . 145
17.6 A recommended order of work . . . 145
17.7 What "finished" would mean . . . 146
17.8 Exercises . . . 148
17.9 Bridge to the appendices . . . 148

A Seeley–DeWitt Coefficients at Working Thickness 149
A.1 Laplace-type operators . . . 149
A.2 Small-time expansion . . . 150
A.3 Proper-time integrals . . . 150
A.4 Worked example: a non-minimal scalar . . . 151
A.5 Fibre traces on the locked block . . . 152
A.6 What this appendix will not do . . . 152
A.7 Exercises . . . 152

B Modular Hamiltonians and the Replica Formula 155
B.1 States, reductions, and von Neumann entropy . . . 155
B.2 Relative entropy and mutual information . . . 156
B.3 The replica formula . . . 156
B.4 Bisognano–Wichmann and small diamonds . . . 157
B.5 Fixed volume . . . 157
B.6 Boundary entropy versus diamond entropy . . . 157
B.7 What this appendix will not do . . . 158
B.8 Exercises . . . 158

C Verlet Laboratory and Energy Identity 159
C.1 Reduced equation . . . 159
C.2 One step . . . 159
C.3 Mechanical energy, work, and balance . . . 160
C.4 Angular momentum . . . 160
C.5 Initial data and units . . . 161
C.6 How to run the script . . . 161
C.7 What this appendix will not do . . . 161
C.8 Exercises . . . 161

D SPARC Rotation Curves: A Coefficient-Level Consistency Check 163
D.1 What is being tested . . . 163
D.2 SPARC comparison . . . 163
D.3 Numerical implementation . . . 164
D.4 What the comparison is and is not . . . 165

E A Provisional Bridge from Mutual Information to the RAR 167
E.1 What the two sides already share . . . 167
E.2 Why the first NAMCG-shaped interpolator failed . . . 168
E.3 An algebraic identity, not a derivation . . . 169
E.4 A modular conjecture . . . 169
E.5 What this bridge excludes . . . 170
E.6 How the bridge would be confirmed or killed . . . 170
E.7 One paragraph for Chapter 13 . . . 171
E.8 Exercises . . . 171

F Einstein Gravity Plus a Conserved Extra Tensor 173
F.1 Einstein . . . 173
F.2 NAMCG infrared equation . . . 173
F.3 Weak field . . . 174
F.4 What this does and does not change . . . 174
F.5 One-line summary . . . 174

G Heat-Kernel Area Term on a Causal Diamond 175
G.1 Replica formula and cone expansion . . . 175
G.2 What is not derived . . . 176
G.3 Status . . . 176

H Status of the Open Problems 177
H.1 Problem A: matrix-to-modular map . . . 177
H.2 Problem B: replica Ward identity . . . 178
H.3 Problem C: conformal mode and Cassini . . . 199
H.4 Problem D: local Lorentz . . . 201
H.5 Problem E: signature . . . 214
H.6 Problem F: coefficients A, B, C . . . 215
H.7 Problem G: HBEV . . . 259
H.8 Problem H: modular origin of the interpolating function . . . 259
H.9 Problem I: dynamical origin of the drag . . . 260
H.10 Problem J: spherical ECG solutions . . . 262
H.11 Problem N: control of the effective expansion . . . 264
H.12 What this appendix does not do . . . 266

I Gaia DR3 Wide Binaries: A Coefficient-Level Consistency Check 267
I.1 What is being compared . . . 267
I.2 What the data show . . . 268
I.3 What this is not . . . 268
I.4 Status . . . 268

J Infrared Limits That Follow from the Proved Chain **269**
J.1 Limits that are obtained . . . 269
J.2 Limits that are not obtained . . . 270
J.3 Status . . . 271

K Heat-Kernel Response to an External Field Strength **273**
K.1 Vanishing of the linear term in a_1 . . . 273
K.2 Quadratic appearance in a_2 . . . 274
K.3 Precision constraints on conformal mixing . . . 274
K.4 What is not shown . . . 274
K.5 Status . . . 274

L A Phenomenological Redshift-Dependent Multiplier (Outside the Proved Chain) **277**
L.1 The inserted multiplier . . . 277
L.2 Illustrative high-redshift comparison . . . 277
L.3 What the illustration is not . . . 278
L.4 Status . . . 278

Conclusion **279**
What has been obtained . . . 279
What has not been obtained . . . 279
Next steps . . . 280

Preface

This book develops Non-Abelian Matrix-Condensate Geometrogenesis (NAMCG) as a single derivation chain: from a microscopic matrix order parameter to an emergent Lorentzian metric, massless spin-2 dynamics, covariantly conserved entanglement backreaction, and a set of astrophysical corollaries.

The text is written for any reader who will follow an argument in order. The working language is the Einstein equation, geodesic deviation, and the idea of a Lagrangian. Quantum field theory is reviewed where it is used. Entanglement entropy is developed from scratch.

Two reading rules are in force throughout.

First, a composite metric is not general relativity until the infrared spectrum contains two massless transverse-traceless polarizations, the Einstein–Hilbert prefactor is computed from microscopic data, and every extra stress tensor satisfies $\nabla^\mu T^{\text{total}}_{\mu\nu} = 0$. Analog-gravity metrics and phenomenological drag terms are treated as insufficient.

Second, numerical laboratories in later chapters are reduced models. They illustrate conservation and dissipation after the covariant theory has been stated. They do not replace that theory.

Chapter 1 states the derivation problem. Chapter 2 installs only the geometry and field theory consumed by later proofs. Chapters 3–11 are the core. Chapters 12–15 are applications. Chapter 16 restates the chain. Chapter 17 lists limitations and falsifiers.

Lake Ontario, 2026-09-19
Edward Schipani, Independent Researcher

Notation

Throughout, spacetime indices $\mu, \nu, \ldots$ are coordinate indices, $A, B, \ldots \in \{0, 1, 2, 3\}$ are tangent-space Lorentz indices, and $I, J, \ldots$ are internal gauge indices of the matrix condensate. The metric signature is $(-, +, +, +)$. Repeated indices are summed.

Symbol	Meaning	Introduced
$\Phi^I{}_A$	matrix order parameter	Ch. 3
v	condensate vacuum expectation value	Ch. 3
$\hat{X}^\mu$	matrix coordinates	Ch. 3
$\Theta^{\mu\nu}$	non-commutativity tensor	Ch. 3
Λ_{UV}	ultraviolet cutoff $\sim \Theta^{-1/2}$	Ch. 3
e^A_μ	emergent vierbein $\langle \partial_\mu \Phi^{IA} \rangle$	Ch. 4
$g_{\mu\nu}$	composite metric $\eta_{AB} e^A_\mu e^B_\nu$	Ch. 4
$T_{\mu\nu}$	symmetric trace-free shear mode	Ch. 5
S_D	entanglement entropy of causal diamond D	Ch. 6
H_D	modular Hamiltonian $-\ln \rho_D$	Ch. 6
U_{ent}	mutual-information potential	Ch. 6
G_{eff}	emergent Newton coupling	Ch. 8
$T^{\mathrm{ent}}_{\mu\nu}$	entropic stress tensor	Ch. 11
$\mathbf{a}_{\mathrm{irrev}}$	irreversible drag acceleration	Ch. 11
S_{HBEV}	holographic boundary-entropy action	Ch. 12
Λ_{eff}	regulated cosmological constant	Ch. 12
r_0	core-regularization scale in $f(r)$	Ch. 13

Where a later chapter needs a second meaning for a symbol, the local definition is restated at the start of that section. The table above is the default.

Chapter 1

What Geometrogenesis Must Derive

Learning outcomes. After this chapter the reader should be able to:

1. state the geometrogenesis problem as a derivation task, not a metaphor;
2. explain why analog-gravity metrics are not general relativity;
3. identify the representation content that a microscopic order parameter must carry if massless spin-2 modes are to exist;
4. list the consistency conditions treated as non-negotiable in this book: diffeomorphism invariance, $\nabla^\mu T^{\rm total}_{\mu\nu} = 0$, a bounded infrared Hamiltonian, and an explicit map from microscopic couplings to $G_{\rm eff}$;
5. see why Chapter 2 is required before any later proof can be stated.

1.1 What this book is trying to derive

General relativity is an infrared theory of a dynamical Lorentzian metric $g_{\mu\nu}$ whose vacuum excitations are two massless, transverse-traceless spin-2 polarizations, and whose coupling to matter is fixed by local energy-momentum conservation and the equivalence principle. In standard presentations the metric is primitive. One writes an Einstein–Hilbert action

$$S_{\rm EH}[g] = \frac{1}{16\pi G}\int \mathrm{d}^4x\,\sqrt{-g}\,R \tag{1.1}$$

and proceeds.

Geometrogenesis asks a more primitive question:

> Can $g_{\mu\nu}$, the Einstein–Hilbert dynamics, and the two graviton polarizations be obtained as the long-wavelength collective description of a microscopic quantum system that does not contain a metric as a fundamental field?

That question is the subject of this book. The framework developed here is Non-Abelian Matrix-Condensate Geometrogenesis (NAMCG). The microscopic object is not a single complex scalar. It is a matrix-valued order parameter $\Phi^I{}_A(x)$ carrying an internal gauge index I and an emergent tangent-space index $A \in \{0, 1, 2, 3\}$. The metric is a composite,

$$e^A_\mu(x) \equiv \langle \partial_\mu \Phi^{IA}(x) \rangle, \qquad g_{\mu\nu}(x) = \eta_{AB}\, e^A_\mu(x)\, e^B_\nu(x). \tag{1.2}$$

The later chapters argue that entanglement equilibrium on causal diamonds forces the infrared mass of the symmetric trace-free shear mode to vanish, that a heat-kernel expansion supplies an explicit formula for $G_{\rm eff}$, and that the entanglement backreaction stress tensor remains covariantly conserved.

Those are theorems *inside the framework*. This first chapter does not prove them. It makes precise what would count as a proof, and why the obvious scalar shortcuts fail.

1.2 Analog gravity, and the ceiling it cannot break

A large and useful literature constructs effective metrics for perturbations on condensed-matter backgrounds. The standard example is a weakly interacting Bose gas with order parameter $\Psi = \sqrt{\rho}\, e^{i\theta}$. Phase fluctuations θ obey, after linearization, a wave equation

$$\frac{1}{\sqrt{-g_{\rm ac}}} \partial_\mu \Big(\sqrt{-g_{\rm ac}}\, g^{\mu\nu}_{\rm ac} \partial_\nu \theta \Big) = 0 \tag{1.3}$$

in an acoustic metric $g^{\mu\nu}_{\rm ac}$ built from the background density and flow. Horizons, ergoregions, and Hawking-like spectra can be engineered in the laboratory.

This is analog gravity. It is not geometrogenesis of general relativity, for four structural reasons.

Only one scalar mode. The Goldstone field of a single complex scalar is a scalar. The acoustic metric is a rewriting of that scalar wave equation. It does not create two independent tensor polarizations.

Screening. Gross–Pitaevskii interactions introduce a healing length ξ. The static potential felt by a localized source is of Yukawa or modified-Bessel type, not a $1/r$ Newtonian potential with radiative $1/r$ tails.

Wrong infrared gravity. A relativistic scalar condensate coupled to an auxiliary metric generically yields Nordström gravity (conformal, with no bending of light in the pure-scalar sector) or scalar–tensor gravity. The Einstein tensor is not selected as the unique linearized kinetic operator for a massless spin-2 field.

Dissipation inserted by hand. Binary inspiral in general relativity follows from quadrupole radiation in a theory with massless spin-2 waves. Replacing that derivation by a Peters formula or a drag term in Newton's equation may fit a curve. It does not derive the curve from the microscopic action.

NAMCG is written to climb over that ceiling, not to decorate it.




Figure 1.1: A scalar order parameter supplies an acoustic metric. It does not supply the two massless spin-2 polarizations of general relativity.

1.3 Representation theory in one paragraph

Local Lorentz invariance acts on tangent-space indices. A graviton is a symmetric, traceless, divergenceless tensor $h_{\mu\nu}$ with two on-shell degrees of freedom in four dimensions. A single scalar ϕ can produce a conformal mode proportional to $g_{\mu\nu}\delta\phi$, and perhaps a vector current if a global symmetry is gauged, but it cannot produce an irreducible shear tensor as an independent propagating excitation of the order parameter itself.

That is the representation-theoretic reason the microscopic field is promoted to $\Phi^I{}_A$. The tangent index A is not cosmetic. It is the handle on which local $\mathrm{SO}(3,1)$ will later act. Chapter 5 decomposes

$$\delta\Phi^I{}_A = \frac{1}{4}\eta_{\mu\nu}\delta g^{\mu\nu}\, I + A_\mu \gamma^\mu + T_{\mu\nu} \tag{1.4}$$

into conformal, vector, and symmetric trace-free pieces. Chapter 7 is the argument that $T_{\mu\nu}$ becomes massless.

If that argument fails, NAMCG collapses back to analog gravity. The book is organized so the reader can see exactly where it would fail.

1.4 What must be delivered before one may say "this is GR"

A derivation of infrared general relativity from a microscopic model is not complete unless all of the following are obtained, not assumed.

1. **Composite metric.** $g_{\mu\nu}$ is defined from microscopic expectation values. It is not put into the fundamental action as an independent Einstein metric.

2. **Local Lorentz frames.** The vacuum must lock internal indices to tangent indices so that e_μ^A transforms as a vierbein and no physical Goldstone scalars remain. That is Chapter 10.

3. **Two massless TT modes.** The infrared kinetic operator of $T_{\mu\nu}$ must be the Lichnerowicz / Einstein operator, with $m_T^2 = 0$. That is Chapters 6–7.

4. **Coefficient map.** Matching the physical G_{eff} requires specified gauge-fixed operators, determinant weights, sector thresholds, and a scheme. Chapter 8 gives a conditional truncated identity; denote the continued curvature coefficient for those inputs by C_R^L; its physical value remains open in Problem F:

$$C_R^L[\{\mathcal{O}_s, w_s, \Lambda_{\mathrm{UV}\,s}, \mu_s\}, \mathcal{S}] \quad \text{is not yet evaluated.} \tag{1.5}$$

5. **Bianchi consistency.** Whatever extra stress is attributed to entanglement or condensate backreaction must satisfy

$$\nabla^\mu \big(T_{\mu\nu}^{\text{matter}} + T_{\mu\nu}^{\text{tensor}} + T_{\mu\nu}^{\text{ent}}\big) = 0. \tag{1.6}$$

   That is Chapter 11. Without it, one has a non-conservative force, not geometry.

6. **Ultraviolet control.** Higher-derivative terms generated by integrating out the condensate cannot leave propagating Ostrogradsky ghosts in the infrared. That is Chapter 9.

7. **Vacuum energy.** The same heat-kernel expansion produces a term $c_0\Lambda_{\mathrm{UV}}^4$. A framework that claims to recover Einstein's equation must say why this is not a Planck-scale cosmological constant. That is Chapter 12.

8. **Phenomenology as corollary.** Rotation curves, regular cores, inspiral drag, and tidal disruption are allowed only after items 1–7. They are not evidence that items 1–7 have been proved.

This checklist is the spine of Parts II–IV. Parts V–VI apply it.

1.5 The NAMCG derivation in one view

The logical order used from Chapter 3 onward is

$$\begin{aligned}\Phi^I{}_A &\longrightarrow \langle\Phi\rangle = v\,\delta^I{}_A \longrightarrow e^A_\mu = \langle\partial_\mu \Phi^{IA}\rangle \\ &\longrightarrow g_{\mu\nu} = \eta_{AB} e^A_\mu e^B_\nu \longrightarrow \delta\Phi = \text{conformal} + \text{vector} + T_{\mu\nu} \\ &\longrightarrow \delta S_D = 0 \longrightarrow m_T \to 0 \longrightarrow S_{\rm EH}[g] + S^{(\text{backreaction})}_{\rm NAMCG}.\end{aligned} \tag{1.7}$$




Figure 1.2: The NAMCG derivation spine used in this book. A single scalar phase cannot lock internal indices to local Lorentz frames.

Two comments are essential already.

First, the entanglement step is doing real work. In a generic spontaneously broken matrix model, $T_{\mu\nu}$ may be massive. The central physical claim of the book is that Jacobson's stationarity condition on dyadic causal diamonds,

$$\delta S_D = \delta\langle H_D\rangle = 0, \tag{1.8}$$

is not an extra thermodynamic slogan. It is a Ward identity of the microscopic theory, and that identity forbids $m_T^2 \neq 0$ in the infrared.

Second, the entropic drag must not be mistaken for a derived result. The inspiral integrator of Chapter 15 is a reduced Newtonian laboratory for the phenomenological law

$$\mathbf{F}_{\rm diss} = -\eta\,\frac{\mu}{d^4}\,\mathbf{v}. \tag{1.9}$$

Its bookkeeping does not establish that this force is the reduction of a covariantly conserved NAMCG stress. That microscopic connection remains open in Problem H.9. If the reader treats the Python drag as the theory, the book has failed.

1.6 How the later phenomenology sits on the spine

Once the infrared Einstein sector exists, NAMCG adds three controlled deformations, each with a named chapter.

Holographic Boundary-Entropy Vacuum (HBEV). A boundary functional involving $K/8\pi G$ and an entropy density $S_{\rm ent}(h_{ij}, \phi)$ is used to regulate $\Lambda_{\rm eff}$. This is how the book addresses the $c_0\Lambda_{\rm UV}^4$ term rather than ignoring it.

Einsteinian cubic gravity. A cubic curvature invariant whose linearization is Einstein-like on maximally symmetric backgrounds [1] motivates a regular-core model for compact objects (a phenomenological profile, not an established exact solution),

$$f(r) = 1 - \frac{2Mr^2}{r^3 + r_0^3} - \frac{\Lambda_{\rm eff}}{3} r^2, \tag{1.10}$$

so that $f(r) \to 1 - \Lambda_{\rm eff} r^2/3$ as $r \to 0$. This does not imply nonlinear or sourced stability, nor derive the profile from the ECG action.

Vacuum backreaction in galaxies and compact binaries. The same conserved extra tensor sources a modification of Poisson's equation; a phenomenological drag in compact binaries is tested as a separate assumption, not derived from that tensor.

None of these is a substitute for Chapters 3–11. They are tests of whether the extra tensors have the right long-distance falloff and the right conservation law.

1.7 How to read this book

Chapters 2–5 are definitional. Read them with a pen. If a symbol is used later with two meanings, the fault is in those chapters and should be caught there.

Chapters 6–11 are the proof core. Each theorem is written with hypotheses first. If a hypothesis is an assumption of the framework rather than a lemma, it is labeled as such.

Chapters 12–15 are applications. Numerical sections flag reduced models in the first paragraph.

Chapter 16 restates the whole derivation as a single chain. Chapter 17 lists what would falsify the framework. Read Chapter 17 before deciding that the theory is closed.

Prerequisites.

What you need. Special relativity, the Euler–Lagrange equation, and the Einstein equation as a field equation for a metric. Linear algebra at the level of commutators and traces. Quantum field theory is helpful but is introduced at the point of use. Entanglement entropy is built in Chapter 6 and Appendix B. If a later proof needs more than that, the extra tool is installed in Chapter 2 or in an appendix, not presupposed as coursework.

1.8 Exercises

1. **Acoustic metric.** Starting from the continuity and Bernoulli equations of an irrotational barotropic fluid, derive $g^{\mu\nu}_{\text{ac}}$ and show that linearized phonons see that metric. Where does a healing length appear if the fluid is a Gross–Pitaevskii condensate?

2. **Mode count.** In $3+1$ dimensions, how many on-shell degrees of freedom does a massive symmetric tensor have? A massless one with gauge invariance $\delta h_{\mu\nu} = \partial_\mu \xi_\nu + \partial_\nu \xi_\mu$? Why does that counting matter for Section 1.3?

3. **Scalar-tensor ceiling.** Consider $S = \int \sqrt{-g}(\phi R - (\partial\phi)^2 - V(\phi))$. Linearize about Minkowski spacetime and a constant ϕ_0. Show that the propagating excitations are a massless spin-2 field mixed with a scalar. What extra condition would be needed to freeze the scalar?

4. **Conservation as a constraint.** Suppose a two-body equation $\mu\ddot{\mathbf{r}} = -\nabla U + \mathbf{a}_{\text{irrev}}$ is written in Minkowski space. What property must $\mathbf{a}_{\text{irrev}}$ have if there is to exist a stress tensor $T^{\text{ent}}_{\mu\nu}$ with $\partial^\mu(T^{\text{matter}}_{\mu\nu} + T^{\text{ent}}_{\mu\nu}) = 0$? A full curved-space answer is Chapter 11; a Newtonian answer is enough here.

5. **Checklist audit.** Take any analog-gravity paper you know and score it against the eight items of Section 1.4. This exercise is the standard the book applies to itself.

1.9 Bridge to Chapter 2

Every later proof is a sentence in a language that has not yet been fixed: manifolds and vierbeine, Einstein–Hilbert variation and Bianchi identities, heat kernels, causal diamonds, and modular Hamiltonians. Chapter 1 stated the derivation problem. Chapter 2 installs the language at the exact thickness later chapters consume—no more, because a reader who must first finish a complete treatise on general relativity will never reach the matrix condensate.

The first object Chapter 2 defines is the one this chapter kept using without ceremony: a Lorentzian manifold with a non-degenerate metric of signature $(-,+,+,+)$, and the difference between a primitive metric and a composite vierbein bilinear.

Chapter 2

Mathematical Preliminaries

Learning outcomes. After this chapter the reader should be able to:

1. distinguish a primitive metric from a composite vierbein bilinear, and write the conversion between $g_{\mu\nu}$ and e^A_μ;
2. vary the Einstein–Hilbert action to obtain $G_{\mu\nu} = 8\pi G\, T_{\mu\nu}$ and state the contracted Bianchi identity that forces $\nabla^\mu T_{\mu\nu} = 0$;
3. expand a functional determinant to second Seeley–DeWitt order and recognize which coefficient multiplies R;
4. define a causal diamond, its modular Hamiltonian, and the first variation δS_D;
5. enter Chapter 3 already possessing every geometric sentence used in the matrix action and the later Ward identities.

This chapter is a toolkit, not a treatise. Every section exists because a later proof calls it by name. Results that are used as black boxes are labeled as such. Results that NAMCG must derive, rather than import, are flagged and postponed.

2.1 Lorentzian manifolds and index conventions

A spacetime in this book is a smooth four-manifold M equipped with a non-degenerate symmetric tensor $g_{\mu\nu}$ of signature $(-,+,+,+)$. The inverse is $g^{\mu\nu}$, and $g = \det(g_{\mu\nu})$. The volume form is $\sqrt{-g}\,\mathrm{d}^4x$.

Coordinate indices μ, ν, ρ, σ transform under diffeomorphisms. Tangent-space (Lorentz) indices $A, B, \ldots$ transform under local $\mathrm{SO}(3,1)$. Internal gauge indices $I, J, \ldots$ transform under the microscopic gauge group of the condensate. These three alphabets are never mixed silently.

The Levi-Civita connection compatible with $g_{\mu\nu}$ is

$$\Gamma^{\rho}_{\mu\nu} = \frac{1}{2} g^{\rho\sigma} (\partial_\mu g_{\nu\sigma} + \partial_\nu g_{\mu\sigma} - \partial_\sigma g_{\mu\nu}). \tag{2.1}$$

The Riemann tensor, Ricci tensor, and Ricci scalar follow the conventions

$$\begin{aligned} R^{\rho}{}_{\sigma\mu\nu} &= \partial_\mu \Gamma^{\rho}_{\nu\sigma} - \partial_\nu \Gamma^{\rho}_{\mu\sigma} + \Gamma^{\rho}_{\mu\lambda}\Gamma^{\lambda}_{\nu\sigma} - \Gamma^{\rho}_{\nu\lambda}\Gamma^{\lambda}_{\mu\sigma}, \\ R_{\mu\nu} &= R^{\rho}{}_{\mu\rho\nu}, \qquad R = g^{\mu\nu} R_{\mu\nu}. \end{aligned} \tag{2.2}$$

The Einstein tensor is

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2} g_{\mu\nu} R. \tag{2.3}$$

Remark 2.1 (Mostly-minus versus mostly-plus). Particle-physics texts often use $(+,-,-,-)$. Every energy condition and every sign in a heat-kernel coefficient is convention-dependent. If a formula in a cited paper looks off by a sign, check the signature before inventing new physics.

2.2 Vierbeine: primitive geometry versus composite geometry

A vierbein (tetrad) is a local frame field e^{A}_{μ} satisfying

$$g_{\mu\nu} = \eta_{AB}\, e^{A}_{\mu} e^{B}_{\nu}, \qquad \eta_{AB} = \mathrm{diag}(-1,+1,+1,+1). \tag{2.4}$$

The inverse frame e^{μ}_{A} obeys $e^{A}_{\mu} e^{\mu}_{B} = \delta^{A}_{B}$ and $e^{A}_{\mu} e^{\nu}_{A} = \delta^{\mu}_{\nu}$. Local Lorentz transformations act as

$$e^{A}_{\mu}(x) \;\longmapsto\; \Lambda^{A}{}_{B}(x)\, e^{B}_{\mu}(x), \qquad \Lambda^{T}\eta\Lambda = \eta. \tag{2.5}$$

For a given invertible frame, the torsion-free metric-compatible spin connection is unique. Equation (2.6) is the standard definition of the Levi-Civita spin connection:

$$\nabla_\mu e^{A}_{\nu} = \partial_\mu e^{A}_{\nu} - \Gamma^{\rho}_{\mu\nu} e^{A}_{\rho} + \omega_{\mu}{}^{A}{}_{B} e^{B}_{\nu} = 0. \tag{2.6}$$

In NAMCG an independent Lorentz connection is introduced in Chapter 10, (10.3); whether it equals this torsion-free connection is Problem D, not a result of the NAMCG dynamics.

Primitive metric. In textbook general relativity one takes $g_{\mu\nu}$ as the fundamental field. A vierbein is an optional rewriting, useful for spinors.

Composite metric. In NAMCG the logical order is reversed. A microscopic matrix field $\Phi^I{}_A$ is fundamental. After condensation one defines

$$e^A_\mu(x) \equiv \langle \partial_\mu \Phi^{IA}(x) \rangle \tag{2.7}$$

and only then reconstructs $g_{\mu\nu}$ by (2.4). Equation (2.7) is not justified in this chapter. It is the definition that Chapter 4 must earn.

The distinction matters for counting degrees of freedom. A primitive symmetric tensor in four dimensions has ten components, of which diffeomorphisms remove four and local Lorentz transformations are absent because there is no independent frame. A primitive vierbein has sixteen components; local Lorentz invariance removes six, diffeomorphisms remove four, and six remain, matching the metric. When the vierbein is composite, those sixteen numbers are not independent fields. They are expectation values of derivatives of Φ. Gauge redundancies of Φ and of the emergent geometry must be matched, not double counted. That matching is Chapter 10.

2.3 Variation of the Einstein–Hilbert action

The Einstein–Hilbert action with a cosmological term and a matter action is

$$S[g] = \frac{1}{16\pi G} \int_M \mathrm{d}^4x \, \sqrt{-g} \, (R - 2\Lambda) + S_{\mathrm{m}}[g, \Psi]. \tag{2.8}$$

Under a metric variation $g_{\mu\nu} \to g_{\mu\nu} + \delta g_{\mu\nu}$ one uses

$$\delta g^{\mu\nu} = -g^{\mu\alpha} g^{\nu\beta} \delta g_{\alpha\beta}, \qquad \delta\sqrt{-g} = \frac{1}{2}\sqrt{-g}\, g^{\mu\nu} \delta g_{\mu\nu}, \tag{2.9}$$

and the Palatini identity

$$\delta R_{\mu\nu} = \nabla_\rho(\delta\Gamma^\rho_{\mu\nu}) - \nabla_\nu(\delta\Gamma^\rho_{\mu\rho}). \tag{2.10}$$

The covariant-derivative terms become a boundary integral. On a manifold without boundary, or after adding the Gibbons–Hawking–York term

$$S_{\mathrm{GHY}} = \frac{1}{8\pi G} \int_{\partial M} \mathrm{d}^3x \, \sqrt{|h|} \, K, \tag{2.11}$$

the bulk stationarity condition is

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G \, T_{\mu\nu}, \tag{2.12}$$

where the matter stress tensor is defined by

$$T_{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta S_{\rm m}}{\delta g^{\mu\nu}}. \tag{2.13}$$

The same definition will be used in Chapter 11 for the entropic contribution $T^{\rm ent}_{\mu\nu}$, with $S_{\rm m}$ replaced by the entanglement functional.

Remark 2.2 (Boundary terms are not optional in this book). Chapter 12 promotes a boundary entropy functional to a dynamical object. The combination $K/8\pi G$ appearing in (2.11) is therefore not an afterthought. It is the gravitational piece of the HBEV boundary action.

2.4 Bianchi identities and conservation

The Riemann tensor obeys

$$\nabla_{[\lambda}R_{\rho\sigma]\mu\nu} = 0. \tag{2.14}$$

Contracting twice yields the contracted Bianchi identity

$$\nabla^{\mu}G_{\mu\nu} = 0. \tag{2.15}$$

If the field equation is $G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G\, T_{\mu\nu}$, then necessarily

$$\nabla^{\mu}T_{\mu\nu} = 0. \tag{2.16}$$

This is not a courtesy to matter. It is the geometric content of diffeomorphism invariance.

In an emergent theory the same identity becomes a constraint on the microscopic construction. Suppose the total effective stress tensor is split as

$$T^{\rm total}_{\mu\nu} = T^{\rm matter}_{\mu\nu} + T^{\rm tensor}_{\mu\nu} + T^{\rm ent}_{\mu\nu}. \tag{2.17}$$

Geometry still requires $\nabla^{\mu}T^{\rm total}_{\mu\nu} = 0$. It is not enough that ordinary matter be conserved by itself if the extra pieces dump momentum into the world. Chapter 11 exists to prove that the NAMCG split satisfies (2.16). Until that proof, any inspiral "drag" is a phenomenological force and not a geometric backreaction.

Noether's second theorem makes the same point in field language. If an action is invariant under local diffeomorphisms generated by an arbitrary vector field ξ^{μ}, the Euler–Lagrange tensor $\mathcal{E}^{\mu\nu}$ of the metric obeys $\nabla_{\mu}\mathcal{E}^{\mu\nu} = 0$ identically, off shell as an identity of the tensor densities involved, and on shell as conservation of $T^{\mu\nu}$. Mass terms that break $\delta_{\xi}h_{\mu\nu} = \nabla_{\mu}\xi_{\nu} + \nabla_{\nu}\xi_{\mu}$ destroy that identity. That observation is the skeleton of Chapter 7.

2.5 Gauge covariant derivatives and a scalar warmup

Let $A_\mu = A_\mu^a T^a$ be a connection on a principal bundle with compact gauge group, generators T^a anti-Hermitian in the conventions of this book. The covariant derivative on a field Φ in a representation ρ is

$$D_\mu \Phi = \partial_\mu \Phi - i A_\mu \Phi \tag{2.18}$$

(up to the representation matrices, restored whenever the index structure requires them). The field strength is

$$F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]. \tag{2.19}$$

A complex scalar with Mexican-hat potential,

$$S[\phi] = \int \mathrm{d}^4x \, \sqrt{-g}\Big(- g^{\mu\nu} (D_\mu \phi)^* D_\nu \phi - \frac{\lambda}{4} \big(|\phi|^2 - v^2\big)^2 \Big), \tag{2.20}$$

condenses at $|\langle \phi \rangle| = v$. The phase is eaten by A_μ (Abelian Higgs / Stueckelberg mechanism), and the physical spectrum contains a massive vector and a massive radial scalar. There is no graviton. Here $g_{\mu\nu}$ is a fixed Lorentzian background of signature $(-,+,+,+)$, so time derivatives enter with positive energy. The earlier form $+|D_\mu\phi|^2 - V$ belongs to the $(+,-,-,-)$ convention and is not used in this book. The Euclidean action has $+g^{\mu\nu}(D_\mu\phi)^* D_\nu \phi + V$.

This warmup is here for two reasons. First, the NAMCG potential in Chapter 3 is the non-Abelian matrix analogue of (2.20). Second, the failure mode is already visible: a single complex scalar has one Goldstone phase. After it is eaten, nothing remains that can be identified with a symmetric trace-free tensor $T_{\mu\nu}$.

2.6 One-loop effective actions and the heat kernel

Let $\mathcal{O}$ be a second-order elliptic (or hyperbolic, after Wick rotation) differential operator on a bundle over M, typically of Laplace type,

$$\mathcal{O} = -g^{\mu\nu} D_\mu D_\nu + E, \tag{2.21}$$

with bundle endomorphism E. The one-loop effective action obtained by integrating out a fluctuation η is

$$W[g] = \frac{1}{2} \mathrm{Tr} \, \ln \mathcal{O} = -\frac{1}{2} \int_0^\infty \frac{\mathrm{d}s}{s} \, \mathrm{Tr} \, e^{-s\mathcal{O}}, \tag{2.22}$$

up to local counterterms and the usual $i\varepsilon$ / Wick-rotation caveats. The operator $e^{-s\mathcal{O}}$ is the heat kernel. Its diagonal admits the Seeley–DeWitt expansion

$$\langle x|e^{-s\mathcal{O}}|x\rangle \sim \frac{1}{(4\pi s)^2}(a_0(x) + a_1(x)s + a_2(x)s^2 + \cdots) \qquad (s \to 0^+), \tag{2.23}$$

in four dimensions. For a Laplace-type operator the first coefficients are

$$\begin{aligned} a_0 &= 1, \\ a_1 &= \frac{1}{6}R - E, \\ a_2 &= \frac{1}{180}(R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - R_{\mu\nu}R^{\mu\nu} + \tfrac{5}{2}R^2) + \frac{1}{2}E^2 - \frac{1}{6}RE + \cdots, \end{aligned} \tag{2.24}$$

where the omitted terms involve derivatives of E and the bundle curvature. The precise tensor combination in a_2 will be written in full in Appendix A when it is needed for ghost analysis. What Chapter 8 requires is only this fact:

Proposition 2.3 (Which coefficient contributes to the curvature term)**.** *In the derivative expansion for specified Laplace-type sectors, the term linear in curvature and quadratic in the ultraviolet cutoff receives a contribution from a_1. After a cutoff Λ_{UV} is imposed on the proper-time integral* (2.22)*, the truncated expansion contains a contribution of the schematic form*

$$W[g] \supset \int \mathrm{d}^4x\,\sqrt{-g}(c_0\Lambda_{\mathrm{UV}}^4 + c_2\Lambda_{\mathrm{UV}}^2 R + \cdots). \tag{2.25}$$

The coefficient c_2 contributes to the conditional matching target for $1/(16\pi G_{\mathrm{eff}})$; this proposition does not determine it or establish a physical Newton coupling. The coefficient c_0 contributes to the vacuum-energy problem under the conventions stated in Chapter 8.

Proof sketch. Insert (2.23) into (2.22) and cut the proper-time integral at $s \sim \Lambda_{\mathrm{UV}}^{-2}$. The a_0/s^3 and a_1/s^2 terms give the quartic and quadratic cutoff dependence term by term in the truncated expansion. The a_1 curvature contribution depends on the endomorphism E, sector weights, and thresholds; it need not remain nonzero in the completed sum. Chapter 8 gives the conditional identity, while Problem F requires the missing sector calculation. □

Two warnings, both of which the later text will spend chapters respecting.

1. The quartic term $c_0\Lambda_{\mathrm{UV}}^4$ is not small. Declaring $\Lambda_{\mathrm{eff}} \approx 10^{-120}\rho_{\mathrm{Pl}}$ without a mechanism is not a calculation. Chapter 12 is that mechanism, or it is a failure.

2. If the fluctuation operator is fourth order, as in the quadratic action of Chapter 9, the heat-kernel calculus changes and Ostrogradsky poles appear. One cannot

quote a_1 from a second-order Laplacian and then use a $\Box^2$ kinetic term without further work.

2.7 Causal diamonds

Let p and q be events in M with q in the chronological future of p. The causal diamond (Alexandrov set)

$$D(p,q) = J^+(p) \cap J^-(q) \tag{2.26}$$

is the intersection of the causal future of p with the causal past of q. Its boundary ∂D consists of two null cones joined at a spacelike 2-surface $\mathcal{B}$ (the edge, or bifurcation surface in the small-diamond limit).

A *small* causal diamond is one whose proper size is short compared with the local curvature radius, so that the metric inside D may be written in Riemann normal coordinates

$$g_{\mu\nu}(x) = \eta_{\mu\nu} - \frac{1}{3} R_{\mu\rho\nu\sigma}(0) x^\rho x^\sigma + O(x^3). \tag{2.27}$$

Jacobson's thermodynamic derivation of Einstein's equation uses precisely such diamonds: the matter flux across ∂D is compared with the variation of entanglement entropy of the diamond interior.

Dyadic diamonds $D_{j,n}$ are a nested family obtained by repeated bisection of a reference diamond. They are bookkeeping. The physics is in the statement that first variations of entropy vanish for arbitrary small deformations of ∂D, not in the dyadic label.

Definition 2.4 (Diamond entanglement functional)**.** Let $\rho_D = \mathrm{Tr}_{M\setminus D}\rho$ be the reduced state of the microscopic theory on the diamond interior. The von Neumann entropy

$$S_D = -\mathrm{Tr}(\rho_D \ln \rho_D) \tag{2.28}$$

is the diamond entanglement functional. The modular Hamiltonian is

$$H_D = -\ln \rho_D, \tag{2.29}$$

so that $S_D = \langle H_D \rangle$ up to the additive constant $\mathrm{Tr}(\rho_D \ln \rho_D) + \langle \ln \rho_D \rangle = 0$ written in that form.

For a free field in Rindler space, H_D is proportional to the boost generator (Bisognano–Wichmann). For a small diamond in a curved background the same statement holds approximately, with Unruh temperature

$$T_{\mathrm{Unruh}} = \frac{\kappa}{2\pi}, \tag{2.30}$$

where κ is the surface gravity of ∂D as seen by the conformal Killing vector that fixes $\mathcal{B}$.

This book does *not* take Jacobson's equation of state as an axiom imported from thermodynamics of horizons. Chapter 6 derives $\delta S_D = 0$ from the microscopic matrix theory, or states clearly where an assumption remains. The definitions above are only the language in which that derivation is written.

2.8 Mutual information

For two regions A and B,

$$I(A:B) = S(A) + S(B) - S(A \cup B). \tag{2.31}$$

Mutual information is non-negative, vanishes if and only if $\rho_{AB} = \rho_A \otimes \rho_B$ (for finite-dimensional algebras), and is invariant under local unitaries on A or B. In NAMCG the macroscopic potential

$$U_{\text{ent}} = -\lambda\, I(A:B) = -\lambda\big[S(A) + S(B) - S(A \cup B)\big] \tag{2.32}$$

is the interaction energy assigned to two condensed cores A and B. The sign is chosen so that positive mutual information lowers the potential: entanglement is attractive in the effective description.

Varying the static U_{ent} gives the conservative interaction. It does not by itself produce a dissipative force; deriving drag requires the dynamical response calculation specified in Problem H.9. Because $I(A:B)$ is built from diffeomorphism scalars $S(\,\cdot\,)$, the conservative variation is not free to point in an arbitrary non-conservative direction. Making this remark precise is the content of Theorem 2 in the ultraviolet note, rewritten as a textbook proof in Chapter 11.

2.9 What this chapter deliberately omits

The following topics are real, and they are not treated here.

- A complete development of spinor bundles and the Dirac operator. NAMCG as formulated in the source notes is bosonic at the microscopic level. Fermionic matter can be coupled to the emergent vierbein later; it is not needed to obtain $G_{\mu\nu}$.
- Full renormalization of quantum gravity as a fundamental metric theory. The metric is not fundamental here. The ultraviolet object is the matrix condensate, with cutoff Λ_{UV}.

- A self-contained treatise on Tomita–Takesaki modular theory. Definition 2.4 and the replica formula $S = (1 - n\partial_n) \ln \operatorname{Tr} \rho^n|_{n=1}$ are sufficient for Chapter 6. Appendix B expands them.

- Numerical relativity. The integrators of Chapter 15 are ordinary-differential toy models, announced as such.

If a later proof appears to need a tool that is not in this chapter, the defect is in the later proof or in this chapter, and should be repaired rather than papered over with a citation.

2.10 Exercises

1. **Volume variation.** Derive both identities in (2.9) from $\delta g = g\, g^{\mu\nu} \delta g_{\mu\nu}$ and $g^{\mu\alpha} g_{\alpha\nu} = \delta^\mu_\nu$.

2. **Einstein tensor from variation.** Starting from (2.8) with $\Lambda = 0$ and no boundary, obtain $G_{\mu\nu} = 8\pi G\, T_{\mu\nu}$. Keep the Palatini terms long enough to see them become a total derivative.

3. **Contracted Bianchi.** Contract (2.14) twice and obtain $\nabla^\mu G_{\mu\nu} = 0$. Do not use the field equations.

4. **Frame versus metric degrees of freedom.** Count local components of e^A_μ and of $g_{\mu\nu}$. Subtract local Lorentz transformations and diffeomorphisms. Confirm that both descriptions have six physical components off shell in the usual sense of geometric gravity, before any equations of motion.

5. **Composite-frame warning.** Suppose $e^A_\mu = \langle \partial_\mu \Phi^{IA} \rangle$ and Φ is subject to a global internal rotation $\Phi \mapsto U\Phi$. What happens to e^A_μ? What additional structure is required before this rotation can be identified with a local Lorentz transformation of the emergent frame? (The answer is Chapter 10; write the question cleanly now.)

6. **Heat-kernel dimensions.** In four dimensions, assign canonical mass dimensions to a_0, a_1, a_2 and to the proper-time variable s. Show that $a_0 \Lambda^4_{\mathrm{UV}}$ and $a_1 \Lambda^2_{\mathrm{UV}}$ are the only two local terms generated by (2.23) that can appear as relevant or marginal couplings of mass dimension four or less, aside from logarithms.

7. **Small diamond.** In Riemann normal coordinates (2.27), compute the spatial 3-volume of a small diamond of radius ℓ to order $\ell^5 R$. This is the geometric ingredient in Jacobson's original argument.

8. **Mutual information positivity.** Prove $I(A:B) \geq 0$ from strong subadditivity, or from relative entropy $S(\rho_{AB}\|\rho_A \otimes \rho_B)$. Why does the sign in (2.32) then correspond to an attractive effective potential?

2.11 Bridge to Chapter 3

The language is now in place. A metric may be primitive or composite. If it is composite, it is built from a vierbein, and the vierbein must come from a field that carries a tangent index. The Einstein tensor is the Euler–Lagrange tensor of (2.8), and Bianchi forces conservation. Newton's constant, if it is emergent, lives in the a_1 heat-kernel coefficient. Entanglement lives on causal diamonds.

What is missing is the field. Chapter 3 introduces the non-Abelian matrix order parameter $\Phi^I{}_A$, the matrix-coordinate algebra $[\hat{X}^\mu, \hat{X}^\nu] = i\Theta^{\mu\nu}$, and the microscopic action whose condensation will produce the vierbein of (2.7). After that chapter, geometry is no longer an independent ingredient. It is an output.

Chapter 3

Matrix Coordinates and the Microscopic Action

Learning outcomes. After this chapter the reader should be able to:

1. write the non-commutative coordinate algebra $[\hat{X}^\mu, \hat{X}^\nu] = i\Theta^{\mu\nu}$ and state how $\Theta^{\mu\nu}$ supplies an ultraviolet cutoff;
2. write both microscopic actions used in the NAMCG notes—the pregeometric matrix action and the covariant condensate-field action—and say what each is for;
3. explain why the order parameter must carry an internal index I and a tangent index A;
4. expand the Mexican-hat matrix potential about its vacuum and identify the radial and Goldstone directions;
5. enter Chapter 4 knowing which object will condense into a vierbein.

Chapter 2 supplied geometry as language. This chapter supplies the field. After this point the metric is not an independent ingredient. It is something the field must produce.

3.1 The scalar failure, restated as a field-content problem

A single complex scalar ϕ transforming as a singlet under every tangent-space representation can condense. It can eat one Goldstone phase into a gauge field. It cannot furnish a local Lorentz frame. The infrared excitations of such a condensate are, at best,

- a radial Higgs scalar,

- a massive or massless vector, if a gauge field is present,
- phonons that admit an acoustic metric rewrite.

There is no symmetric trace-free tensor $T_{\mu\nu}$ hiding in ϕ. Seeking that tensor in a more elaborate potential for the same ϕ is a representation-theoretic error, not a calculational one.

NAMCG therefore promotes the order parameter to a rectangular (or square, after a choice of basis) matrix field

$$\Phi^I{}_A(x), \tag{3.1}$$

where $I = 1, \ldots, N$ labels an internal gauge representation and $A = 0, 1, 2, 3$ labels an emergent tangent space. The field is a linear map from the tangent space of an auxiliary frame into the internal space, or the reverse, according to index placement. The placement used in this book is: I up, A down on Φ, unless a formula from the source notes is being quoted verbatim.

The rest of the chapter writes an action for Φ and for a closely related pregeometric object, the matrix coordinates $\hat{X}^\mu$.

3.2 Two presentations, and why both appear

The original notes use two microscopic languages. They are not identical, and treating them as interchangeable by slogan would produce a confused book.

Presentation A: pregeometric matrix coordinates. There is, at the outset, no manifold. The dynamical objects are Hermitian operators $\hat{X}^\mu$ acting on a Hilbert space $\mathcal{H}$. Spacetime points, if they appear at all, appear as eigenvalue data of a complete commuting subset of these operators, or as a derived continuum after condensation.

Presentation B: condensate field on an auxiliary background. There is an auxiliary manifold with a non-dynamical metric $g^{(0)}_{\mu\nu}$. The dynamical object is the field $\Phi^I{}_A(x)$. The physical metric $g_{\mu\nu}$ is not $g^{(0)}_{\mu\nu}$. The auxiliary metric is a scaffold for writing derivatives and heat kernels. Chapter 8 writes its heat kernel on the emergent metric under Assumption 8.4.

This book treats Presentation A as the ontological starting point and Presentation B as the background-field language appropriate after a condensate has selected a notion of local coordinates. Chapter 4 is the bridge. Until that bridge is built, the two actions below are written separately and not added together.

Assumption 3.1 (Auxiliary background is non-dynamical)**.** Whenever Presentation B is in use, $g^{(0)}_{\mu\nu}$ is a fixed scaffold, usually taken flat, $g^{(0)}_{\mu\nu} = \eta_{\mu\nu}$. It is not varied in the

Einstein–Hilbert sense. Varying the *emergent* metric $g_{\mu\nu}$ defined in Chapter 4 is a different operation.

3.3 Presentation A: matrix coordinates and ultraviolet cutoff

Definition 3.2 (Matrix coordinates)**.** Let $\mathcal{H}$ be a separable Hilbert space and let $\hat{X}^\mu$, $\mu = 0, 1, 2, 3$, be Hermitian operators on $\mathcal{H}$. The fundamental kinematic relation of Presentation A is

$$[\hat{X}^\mu, \hat{X}^\nu] = i\Theta^{\mu\nu}, \tag{3.2}$$

where $\Theta^{\mu\nu} = -\Theta^{\nu\mu}$ is an antisymmetric tensor with dimensions of length squared. If $\Theta^{\mu\nu}$ is a multiple of the identity on $\mathcal{H}$, it is a c-number background. In the fully dynamical theory it may itself be an operator; the notes treat the leading infrared analysis with a c-number $\Theta^{\mu\nu}$.

The scale

$$\Lambda_{\mathrm{UV}} \sim \Theta^{-1/2}, \qquad \Theta \equiv \sqrt{\tfrac{1}{2}\Theta_{\mu\nu}\Theta^{\mu\nu}}, \tag{3.3}$$

is the ultraviolet cutoff of the theory. Two operators that fail to commute cannot be simultaneously diagonalized, so the notion of a sharply localized event (x^0, x^1, x^2, x^3) is meaningless below $\Lambda_{\mathrm{UV}}^{-1}$. This is the NAMCG replacement for a hard lattice.

Remark 3.3 (What $\Theta^{\mu\nu}$ is not). Equation (3.2) is not yet a solution of the cosmological-constant problem, not yet a black-hole core, and not yet a graviton. It is a kinematic regulator. Claims that begin "because spacetime is non-commutative…" and end at galactic rotation curves are skipping the work of Chapters 4–14.

A Yang–Mills-type matrix action for these coordinates, including a cubic structure-constant coupling to a background condensate tensor $C_{\mu\nu\lambda}$, is

$$\begin{aligned} S_{\mathrm{matrix}} = &-\frac{1}{4g_{\mathrm{YM}}^2}\mathrm{Tr}\big([\hat{X}^\mu, \hat{X}^\nu][\hat{X}_\mu, \hat{X}_\nu]\big) + \frac{1}{3!}\mathrm{Tr}\big(C_{\mu\nu\lambda}[\hat{X}^\mu, \hat{X}^\nu]\hat{X}^\lambda\big) \\ &+ S_{\mathrm{matter}}(\Psi, \hat{X}^\mu). \end{aligned} \tag{3.4}$$

The first term is the IKKT / reduced-Yang–Mills kinetic term. On a background satisfying (3.2) it is proportional to $\mathrm{Tr}(\Theta_{\mu\nu}\Theta^{\mu\nu})$ plus fluctuation pieces. The cubic term selects a preferred vacuum condensation structure; its detailed index symmetries are constrained by the requirement that the vacuum preserve a Lorentzian interpretation after Chapter 4.

Matter fields Ψ in S_{matter} are functions of the matrices $\hat{X}^\mu$ in the sense of the Weyl map associated with $\Theta^{\mu\nu}$. This book does not need the full star-product calculus until

phenomenology with explicit localized sources appears in Chapter 14. The existence of such a map is enough for now.

Definition 3.4 (Extended matrix coordinates (conditional extension))**.** Let $D \geq 4$ and let $\hat{X}^I$, $I = 0, \ldots, D-1$, be Hermitian operators on $\mathcal{H}$ with $[\hat{X}^I, \hat{X}^J] = i\Theta^{IJ}$, $\Theta^{IJ} = -\Theta^{JI}$. Matrix indices are contracted with the fixed constant form $\eta_{IJ} = \mathrm{diag}(-1, 1, \ldots, 1)$, exactly as $\eta_{\mu\nu}$ is used in the first (quartic) term and in the matter term of (3.4), read with I, J in place of μ, ν; the cubic term is not extended here. The definition reduces, for $D = 4$, to Definition 3.2. In the semiclassical regime a configuration is a map $y \mapsto x^I(y)$ from a four-dimensional manifold carrying a nondegenerate Poisson bivector $\theta^{\alpha\beta}$, with $\Theta^{IJ} = \theta^{\alpha\beta}\partial_\alpha x^I \partial_\beta x^J$ and $\mathrm{Tr} \to \int \mathrm{d}^4 y\, \rho$, where $\rho = 1/|\operatorname{Pf}\theta|$. Two tensors are then defined from the configuration, not from the action: the induced tensor $g_{\alpha\beta} = \eta_{IJ}\partial_\alpha x^I \partial_\beta x^J$ and the operator tensor $K^{\mu\nu} = \theta^{\alpha\mu} g_{\alpha\beta} \theta^{\beta\nu}$, which is the kinetic tensor of a fluctuation $[\hat{X}^I, \cdot\,]$ in the Poisson limit. This definition adds no fundamental metric: the form η_{IJ} contracts matrix indices only, and its Lorentzian signature is retained as Assumption 4.6. The index I of this definition is not the internal index of $\Phi^I{}_A$ in Presentation B, and the $D-4$ transverse coordinates are not covered by Assumption 5.1. For a real four-dimensional leaf and a real Lorentzian g, K^{-1} is nowhere proportional to g, for every $D \geq 4$ (Appendix H); neither g nor K^{-1} is asserted here to be the emergent metric of Equation (4.9) or the metric of Assumption 8.4.

3.4 Presentation B: the covariant condensate action

Once local coordinates x^μ have been introduced as labels on an auxiliary manifold, the dynamical variable is the matrix field $\Phi^I{}_A(x)$. Let A_μ be a connection on the internal bundle, with covariant derivative

$$D_\mu \Phi = \partial_\mu \Phi - i A_\mu \Phi \tag{3.5}$$

acting on the I index. The microscopic action used for the heat-kernel analysis is

$$S_{\mathrm{micro}}[\Phi, A] = \int \mathrm{d}^4 x \, \sqrt{-g^{(0)}}\, \mathrm{Tr}\Big(- g^{(0)\mu\nu} (D_\mu \Phi)^\dagger D_\nu \Phi - V(\Phi^\dagger \Phi) \Big). \tag{3.6}$$

The trace is over internal indices. Tangent indices on Φ are contracted through the potential and through the identification of A with a frame index after condensation; they are not traced in the same sum. In the signature $(-,+,+,+)$ of Chapter 2 the kinetic term is written with the minus sign so that time derivatives have positive energy. This is a correction to the Presentation-B action as stated; that the matrix action of Presentation A reproduces this sign is part of the unproved dictionary (Assumption 3.6) and remains to be derived.

The potential that triggers condensation is the matrix Mexican hat

$$V(\Phi^\dagger\Phi) = \frac{\lambda}{4}\mathrm{Tr}\big(\Phi^\dagger\Phi - v^2\mathbb{I}\big)^2. \tag{3.7}$$

Its minima lie on the manifold

$$\Phi^\dagger\Phi = v^2\mathbb{I} \tag{3.8}$$

in the internal space, up to unitary rotations. A representative vacuum used throughout the book is

$$\langle \Phi^I{}_A \rangle = v\,\delta^I{}_A, \tag{3.9}$$

which is available when the internal dimension is at least four and a basis has been chosen that identifies four internal directions with the tangent directions. If $N > 4$, the remaining internal directions must either decouple or be locked by additional Yukawa-like couplings. The source notes assume that a vacuum of the form (3.9) exists. This book records that as an assumption.

Assumption 3.5 (Existence of a frame-locking vacuum)**.** There exists a stable vacuum (3.9) (or a locally equivalent vacuum) in which four internal directions are identified with the four tangent directions. Goldstone modes associated with the broken generators are available to be eaten by a connection, as in Chapter 10.

3.5 Index structure and the intended infrared spectrum

Under an internal gauge transformation $\Omega(x)$ and a local Lorentz transformation $\Lambda(x)$, the field is required to transform as

$$\Phi^I{}_A(x) \;\longmapsto\; \Omega^I{}_J(x)\,\Phi^J{}_B(x)\,\Lambda^B{}_A(x). \tag{3.10}$$

Equation (3.10) is the transformation law that makes Φ capable of locking the two groups to a diagonal subgroup. It is a choice of field content. It is not derived from Presentation A alone; it is the additional structure that Presentation B puts in by hand so that a vierbein can exist.

Fluctuations about (3.9) will be decomposed in Chapter 5 into

$$\delta\Phi^I{}_A = \frac{1}{4}\eta_{\mu\nu}\delta g^{\mu\nu}\, I + A_\mu\gamma^\mu + T_{\mu\nu}. \tag{3.11}$$

The only fact needed here is that the last piece exists because Φ has two indices that can be symmetrized and made traceless after the vacuum has identified I with A. A scalar has no such piece.

3.6 Equations of motion from Presentation B

Varying (3.6) with respect to $\Phi^\dagger$ gives

$$D_\mu(\sqrt{-g^{(0)}}\, g^{(0)\mu\nu} D_\nu \Phi) - \sqrt{-g^{(0)}}\, \frac{\lambda}{2} \Phi(\Phi^\dagger \Phi - v^2 \mathbb{I}) = 0. \tag{3.12}$$

Varying with respect to A_μ produces a matrix current conservation law

$$D_\mu(\Phi(D^\mu \Phi)^\dagger - (D^\mu \Phi)\Phi^\dagger) = 0, \tag{3.13}$$

up to the conventional factor of i in the current definition. These equations determine the condensate profile on the auxiliary background. They do not yet determine the Einstein tensor of the emergent metric. That requires integrating out fluctuations, which is Chapter 8, after the emergent metric has been defined in Chapter 4 and the tensor mode identified in Chapter 5.

3.7 Relating the two presentations

A complete derivation from (3.4) to (3.6) would construct, from the spectra of $\hat{X}^\mu$, a commutative manifold, a Weyl map, and a slowly varying matrix-valued field on that manifold whose covariant derivative reproduces the commutator $[\hat{X}^\mu, \hat{\Phi}]$. That derivation is not supplied in full in the source notes. The working relation used in this book is the following.

Assumption 3.6 (Infrared dictionary)**.** In a condensate phase in which the matrices $\hat{X}^\mu$ are, to leading order, commutative coordinates $x^\mu \mathbb{I}$ plus fluctuations, there exists a slowly varying field $\Phi^I{}_A(x)$ such that

$$[\hat{X}_\mu, \Phi] \longleftrightarrow i\Theta_{\mu\nu} D^\nu \Phi \qquad \text{plus higher-derivative corrections in } \Theta. \tag{3.14}$$

Under this dictionary the quadratic commutator term in (3.4) is required to generate the kinetic term in (3.6), including its sign in the signature $(-,+,+,+)$; this is a matching condition, not an established result. Λ_{UV} remains the cutoff of the derivative expansion.

Assumption 3.6 is the most important unproved bridge in the microscopic setup. Chapter 17 lists a sharper derivation of this dictionary as an open problem. Every later heat-kernel formula is a calculation *within* Presentation B, contingent on the dictionary.

Remark 3.7 (Constant-Θ Moyal check (conditional; not closure))*.* Assume a Moyal

realization with constant, real, antisymmetric $\Theta^{\mu\nu}$, and define

$$(f \star g)(x) = f(x) \exp\left(\frac{i}{2}\Theta^{\mu\nu}\overleftarrow{\partial_\mu}\overrightarrow{\partial_\nu}\right) g(x).$$

For general symbols,

$$[f,g]_\star = i\Theta^{\mu\nu}(\partial_\mu f)(\partial_\nu g) - \frac{i}{24}\Theta^{\mu\nu}\Theta^{\rho\sigma}\Theta^{\alpha\beta}(\partial_\mu\partial_\rho\partial_\alpha f)(\partial_\nu\partial_\sigma\partial_\beta g) + O(\Theta^5).$$

For the coordinate symbol x_μ, derivatives of order greater than one vanish, so

$$[x_\mu, \phi]_\star = i\Theta_{\mu\nu}\partial^\nu\phi$$

exactly in this realization. This verifies the coordinate-background kinematic identity, but not the full dictionary (3.14) for fluctuating matrix coordinates, a gauge connection, or nonconstant Θ.

There is a further restriction if the two-commutator contraction is to reproduce a Lorentzian kinetic metric. With signature $(-,+,+,+)$, define

$$K_\Theta^{\nu\rho} := \eta_{\alpha\beta}\Theta^{\alpha\nu}\Theta^{\beta\rho}, \qquad N^\mu{}_\nu := \Theta^{\mu\alpha}\eta_{\alpha\nu}.$$

Then N is a real Lorentz generator and $K_\Theta^{\nu\rho} = -(N^2)^\nu{}_\sigma\eta^{\sigma\rho}$. Thus a nonzero metric-proportional contraction $K_\Theta^{\nu\rho} = c\,\eta^{\nu\rho}$, $c \neq 0$, would require $N^2 = -c\,\mathbb{I}$. In a Lorentz frame, writing N in terms of boost and rotation vectors $\mathbf{E}, \mathbf{B}$, its characteristic polynomial is

$$p(z) = z^4 + (|\mathbf{B}|^2 - |\mathbf{E}|^2)z^2 - (\mathbf{E}\cdot\mathbf{B})^2.$$

But $N^2 = -c\,\mathbb{I}$ would give $p(z) = (z^2 + c)^2$, whose constant term $c^2 > 0$, whereas the displayed Lorentz-generator polynomial has constant term $-(\mathbf{E}\cdot\mathbf{B})^2 \leq 0$. Hence no nonzero c is possible. If $c = 0$, $K_\Theta = 0$ says that the image of the antisymmetric map Θ is totally null. In Lorentzian four-space a totally null subspace has dimension at most one, while the rank of an antisymmetric matrix is even; therefore $\Theta = 0$. So no nonzero constant real Θ contracted with η yields a tensor proportional to $\eta^{\nu\rho}$. For example, if only $\Theta^{12} = -\Theta^{21} = \theta$ is nonzero, then $K_\Theta^{\nu\rho} = \theta^2\,\mathrm{diag}(0,1,1,0)$.

This obstruction concerns only the stated constant-Θ contraction. Matching a Lorentzian kinetic term would need a different specified mechanism, such as a dynamical or nonconstant background, a fluctuation-generated metric, or an additional tangent-index contraction from a derived field structure. It does not rule out those possibilities or an enlarged matrix construction, and it does not derive the kinetic term, its sign, a coframe, or local Lorentz symmetry. The matrix-derived criterion of Problem D remains open. Problems B, C, D, F and G remain open.

Remark 3.8 (Constant-Θ trace and quadratic fluctuations (conditional; not closure)).

Assume the irreducible Weyl representation of the constant, nondegenerate Heisenberg algebra

$$[\bar{X}^\mu, \bar{X}^\nu] = i\Theta^{\mu\nu}\mathbb{I},$$

and use the Weyl convention

$$\hat{f} := \int \frac{\mathrm{d}^4k}{(2\pi)^4}\, \widetilde{f}(k) e^{ik_\mu \bar{X}^\mu}.$$

For Schwartz symbols, the Weyl trace formula and Moyal trace identity are

$$\mathrm{Tr}_{\mathcal{H}}\, \hat{f} = \frac{1}{(2\pi)^2 |\operatorname{Pf}\Theta|} \int \mathrm{d}^4x\, f(x), \qquad \int \mathrm{d}^4x\, f \star g = \int \mathrm{d}^4x\, fg.$$

Thus the constant-Θ trace density in this representation is $(2\pi)^{-2}|\operatorname{Pf}\Theta|^{-1}$.

For the fluctuation calculation, restrict to the $C_{\mu\nu\lambda} = 0$ branch of (3.4) and omit matter fluctuations. The central background is a classical solution of the pure quartic matrix equation because its commutators are central. Write

$$\hat{X}^\mu = \bar{X}^\mu + \hat{A}^\mu, \qquad D^\mu = [\bar{X}^\mu, \cdot], \qquad f^{\mu\nu} = D^\mu \hat{A}^\nu - D^\nu \hat{A}^\mu.$$

Assume the fluctuation symbols $a^\mu(x)$ decay rapidly enough for the trace formula and integrations by parts to apply, so the linear term vanishes. The quadratic matrix action is

$$S^{(2)}_{\mathrm{YM}} = -\frac{1}{4g^2_{\mathrm{YM}}} \mathrm{Tr}(f_{\mu\nu} f^{\mu\nu}).$$

The exact constant-Θ coordinate identity gives

$$f^{\mu\nu} \longleftrightarrow i\mathcal{F}^{\mu\nu}_\Theta, \qquad \mathcal{F}^{\mu\nu}_\Theta := \Theta^{\mu\rho}\partial_\rho a^\nu - \Theta^{\nu\rho}\partial_\rho a^\mu.$$

Consequently,

$$\boxed{S^{(2)}_{\mathrm{YM}} = \frac{1}{4g^2_{\mathrm{YM}}(2\pi)^2|\operatorname{Pf}\Theta|} \int \mathrm{d}^4x\, \mathcal{F}_{\Theta,\mu\nu}\mathcal{F}^{\mu\nu}_\Theta}$$

with indices contracted using $\eta_{\mu\nu}$. In particular,

$$\mathcal{F}_{\Theta,\mu\nu}\mathcal{F}^{\mu\nu}_\Theta = 2K^{\rho\sigma}_\Theta \partial_\rho a_\nu\, \partial_\sigma a^\nu - 2\Theta^{\mu\rho}\Theta_\nu{}^\sigma \partial_\rho a^\nu\, \partial_\sigma a_\mu, \qquad K^{\rho\sigma}_\Theta := \eta_{\mu\nu}\Theta^{\mu\rho}\Theta^{\nu\sigma}.$$

The second, mixed term is part of the vector kinetic operator; this is not the scalar operator of Remark 3.9.

Under the linearized matrix-gauge transformation

$$\delta a^\mu = \Theta^{\mu\rho}\partial_\rho \lambda,$$

the antisymmetry of Θ and commutativity of partial derivatives give $\delta\mathcal{F}_\Theta^{\mu\nu} = 0$. In Fourier space, writing $q^\mu = \Theta^{\mu\rho}k_\rho$, the quadratic kernel is proportional to

$$\mathcal{M}_{\alpha\beta}(k) = q^2\eta_{\alpha\beta} - q_\alpha q_\beta, \qquad \mathcal{M}_{\alpha\beta}q^\beta = 0, \qquad q^2 = K_\Theta^{\rho\sigma}k_\rho k_\sigma.$$

For the usual flat scaffold $g^{(0)}_{\mu\nu} = \eta_{\mu\nu}$, the constant-$\Theta$ obstruction of Remark 3.7 gives $K_\Theta \neq c\, g^{(0)}$ for nonzero real Θ. Thus this quadratic action does not have the $g^{(0)}$-contracted form used in Presentation B. Its sign does not establish a healthy Lorentzian energy.

For $g^{(0)}_{\mu\nu} = \eta_{\mu\nu}$, let

$$P(k) := q^2 = K_\Theta^{\rho\sigma}k_\rho k_\sigma.$$

This is a characteristic-level statement for the displayed quadratic kernel, not a claim about the full gauge-fixed PDE. When $q^2 \neq 0$, $\mathcal{M} = q^2\eta - qq$ has rank three with kernel spanned by q; its determinant on the quotient is proportional to $(q^2)^3$. When $q \neq 0$ is null, $\mathcal{M} = -qq$ has rank one, so the rank of $\mathcal{M}$ drops to one and its kernel grows to dimension three.

A degenerate example is the rank-two electric block

$$\Theta^{01} = E \neq 0, \qquad \Theta^{\mu\nu} = 0 \quad \text{for all other independent components.}$$

It gives

$$K_\Theta = \mathrm{diag}(E^2, -E^2, 0, 0), \qquad P(k) = E^2(k_0^2 - k_1^2).$$

The characteristic roots in scaffold time x^0 are real, $k_0 = \pm k_1$, but the polynomial is degenerate: it contains no k_2 or k_3. When $k_0 = k_1 = 0$, $q^\mu = 0$ and the entire quadratic vector kernel vanishes, even for nonzero transverse momenta. This is a characteristic-level observation, not a claim of well-posedness for a gauge-fixed initial-value problem.

If Θ is nondegenerate, K_Θ is a nondegenerate Lorentzian form by congruence. For any η-timelike covector n_ρ, antisymmetry gives $n_\mu\Theta^{\mu\rho}n_\rho = 0$, so $q^\mu = \Theta^{\mu\rho}n_\rho$ is orthogonal to n. Since Θ is invertible and $n \neq 0$, $q \neq 0$, and hence $P(n) = q^2 > 0$. Therefore no η-timelike covector is a hyperbolic direction for this characteristic polynomial. The polynomial does have hyperbolic directions with $P(n) < 0$, but they are not η-timelike. Neither the degenerate nor nondegenerate case establishes a healthy Lorentzian energy or a well-posed gauge-fixed problem.

This is an action for the matrix-coordinate symbol a^μ, not for $\Phi^I{}_A$. It does not construct $\Phi^I{}_A$, a coframe, or a tangent-space connection, and makes no claim for nonconstant Θ. Problems B, C, D, F and G remain open.

Remark 3.9 (A varying-Θ Lorentzian toy geometry (conditional; not a matrix derivation)). The exact coordinate-commutator identity in Remark 3.7 is for constant Θ

and is not asserted for the varying bivector used here; a nonconstant Poisson bivector requires a corresponding deformation-quantization product, whose coordinate commutators can include derivatives of Θ.

As a geometric example, choose by hand

$$\Theta = a\,\partial_0 \wedge \partial_1 + b(y)\,\partial_2 \wedge \partial_3, \qquad y = x^2, \qquad b(y) = b_0 e^{\kappa y^2},$$

with real $a \neq 0$ and $b_0 > 0$. This is a Poisson bivector: the two blocks have disjoint coordinates, and b depends only on a coordinate within its own two-dimensional block. Its Pfaffian is $ab(y) \neq 0$. The algebraic tensor

$$K_\Theta^{\nu\rho} := \eta_{\alpha\beta}\Theta^{\alpha\nu}\Theta^{\beta\rho} = \mathrm{diag}(a^2, -a^2, b(y)^2, b(y)^2)$$

is built from this chosen Θ. If, as an additional normalization and metric-identification assumption, one sets

$$K_\Theta^{\nu\rho} = \ell_*^4\,\tilde{g}^{\nu\rho},$$

then

$$\mathrm{d}\tilde{s}^2 = \ell_*^4\left[a^{-2}(\mathrm{d}x^0)^2 - a^{-2}(\mathrm{d}x^1)^2 + b(y)^{-2}\big((\mathrm{d}y)^2 + (\mathrm{d}x^3)^2\big)\right].$$

In the signature $(+,-,+,+)$ of $\tilde{g}$, x^1 is the odd-signed coordinate. The two-dimensional conformal factor is $e^\sigma = \ell_*^2/b(y)$, so this chosen metric has scalar curvature

$$\tilde{R}(y) = -2e^{-2\sigma}\frac{\mathrm{d}^2\sigma}{\mathrm{d}y^2} = \frac{4\kappa b_0^2}{\ell_*^4}e^{2\kappa y^2}.$$

It is nonflat for $\kappa \neq 0$.

For completeness, one orthonormal coframe for this chosen metric is obtained by setting

$$L = \frac{\ell_*^2}{|a|}, \qquad f(y) = \frac{\ell_*^2}{b(y)} > 0, \qquad \eta_{\hat{a}\hat{b}} = \mathrm{diag}(-,+,+,+),$$

and choosing

$$\tilde{e}^{\hat{0}} = L\,\mathrm{d}x^1, \qquad \tilde{e}^{\hat{1}} = L\,\mathrm{d}x^0, \qquad \tilde{e}^{\hat{2}} = f(y)\,\mathrm{d}y, \qquad \tilde{e}^{\hat{3}} = f(y)\,\mathrm{d}x^3.$$

The swap of the first two coordinate labels assigns the negative frame direction to x^1. This coframe obeys

$$\tilde{g}_{\mu\nu} = \eta_{\hat{a}\hat{b}}\,\tilde{e}^{\hat{a}}{}_\mu\tilde{e}^{\hat{b}}{}_\nu$$

and reproduces the displayed metric. It is one choice; other orthonormal coframes are related locally by Lorentz transformations preserving $\eta_{\hat{a}\hat{b}}$. This freedom follows from the metric description and is not a gauge symmetry derived from the matrix action.

Define the torsion-free, metric-compatible spin connection by

$$\mathrm{d}\tilde{e}^{\hat{a}} + \tilde{\omega}^{\hat{a}}{}_{\hat{b}} \wedge \tilde{e}^{\hat{b}} = 0, \qquad \tilde{\omega}_{\hat{a}\hat{b}} = -\tilde{\omega}_{\hat{b}\hat{a}},$$

with curvature convention

$$\tilde{\mathcal{R}}^{\hat{a}}{}_{\hat{b}} = \mathrm{d}\tilde{\omega}^{\hat{a}}{}_{\hat{b}} + \tilde{\omega}^{\hat{a}}{}_{\hat{c}} \wedge \tilde{\omega}^{\hat{c}}{}_{\hat{b}}.$$

Writing $q = f'/f = -2\kappa y$, the only nonzero connection one-forms are

$$\tilde{\omega}^{\hat{3}}{}_{\hat{2}} = q\,\mathrm{d}x^3, \qquad \tilde{\omega}^{\hat{2}}{}_{\hat{3}} = -q\,\mathrm{d}x^3,$$

or, with both frame indices raised, $\tilde{\omega}^{\hat{2}\hat{3}} = 2\kappa y\,\mathrm{d}x^3$. The only independent curvature two-form is

$$\tilde{\mathcal{R}}^{\hat{2}}{}_{\hat{3}} = 2\kappa\,\mathrm{d}y \wedge \mathrm{d}x^3 = \frac{2\kappa}{f(y)^2}\,\tilde{e}^{\hat{2}} \wedge \tilde{e}^{\hat{3}}.$$

It gives

$$\tilde{R} = \frac{4\kappa}{f(y)^2} = \frac{4\kappa b_0^2 e^{2\kappa y^2}}{\ell_*^4},$$

in agreement with the scalar curvature above. This is the Levi–Civita connection determined by the chosen toy metric, not a derivation of the independent gauge connection or its dynamics from the matrix action.

For comparison, assume in addition that the continuum trace uses the symplectic density

$$\rho(y) = \frac{1}{|\operatorname{Pf}\Theta|} = \frac{1}{|a|b(y)}$$

and that the scalar kinetic term is built from the two-commutator tensor $K_\Theta^{\mu\nu}$. These are extra assumptions for nonconstant Θ; neither the trace map nor this kinetic term is derived here. The corresponding divergence operator is

$$\mathcal{L}_\Theta\phi := \rho^{-1}\partial_\mu(\rho K_\Theta^{\mu\nu}\partial_\nu\phi) = a^2(\partial_0^2 - \partial_1^2)\phi + b(y)^2(\partial_y^2 + \partial_3^2)\phi + 2\kappa y\, b(y)^2\partial_y\phi.$$

Define instead a conformally related metric g_ρ by

$$g_\rho^{\mu\nu} := \rho K_\Theta^{\mu\nu}, \qquad (g_\rho)_{\mu\nu} = \operatorname{diag}\left(\frac{b}{|a|}, -\frac{b}{|a|}, \frac{|a|}{b}, \frac{|a|}{b}\right).$$

It is Lorentzian and $\sqrt{|\det g_\rho|} = 1$, whence

$$\Box_{g_\rho} = \partial_\mu(\rho K_\Theta^{\mu\nu}\partial_\nu), \qquad \mathcal{L}_\Theta = \rho^{-1}\Box_{g_\rho}.$$

Thus the Poisson-trace operator is a position-dependent scalar multiple of this metric's

Laplace–Beltrami operator, not exactly the Laplace–Beltrami operator itself. Since $\rho \neq 0$, the two operators have the same homogeneous zero-mode equation, but are different differential operators and need not have the same spectrum or heat kernel.

The metric g_ρ is conformal to the hand-chosen $\tilde{g}$:

$$g_\rho^{\mu\nu} = \ell_*^4 \rho\, \tilde{g}^{\mu\nu}.$$

Its scalar curvature, in the same curvature convention as above, is

$$R[g_\rho] = -\frac{b(y)}{|a|}(2\kappa + 6\kappa^2 y^2),$$

which differs in general from $\tilde{R}$. Accordingly, $\tilde{R}$ is not the curvature associated with this assumed Poisson-trace operator through g_ρ. Neither curvature is derived from the matrix action by this comparison. This is only an algebraic construction: $b(y)$, κ, and the metric identification are chosen by hand. Accordingly, $\tilde{R}$ is a property of those choices, not a prediction of the matrix action or the emergent curvature $R[g]$ of Chapter 4, and not the heat-kernel curvature contribution of Chapter 8. A metric identification does not by itself supply a matrix-derived coframe, a dynamical connection, or a local Lorentz gauge symmetry. The coframe above is chosen rather than matrix-derived, and its Lorentz-related freedom is not derived from the matrix action. The matrix-derived criterion of Problem D remains open. Problems B, C, D, F and G remain open.

3.8 Energy scales and coupling constants

The dimensionful quantities introduced so far are

$$\Lambda_{\mathrm{UV}}, \qquad v, \qquad \lambda, \qquad g_{\mathrm{YM}}, \tag{3.15}$$

together with the auxiliary-background curvature scale, which is set to zero when $g^{(0)} = \eta$. Their intended roles are:

Quantity	Role
Λ_{UV}	kinematic cutoff from $\Theta^{\mu\nu}$
v	condensate scale; appears in the working conditional G_{eff} parameterization
λ	controls the radial Higgs mass $m_\sigma \sim \sqrt{\lambda}\, v$
g_{YM}	internal gauge coupling; enters the connection that eats Goldstones

Chapter 8 gives a conditional truncated heat-kernel identity. Define C_R^L as the

Lorentzian curvature coefficient only after the specified sector sum and continuation:

$$C_R^L[\{\mathcal{O}_s, w_s, \Lambda_{\mathrm{UV}\,s}, \mu_s\}, \mathcal{S}] \quad \text{is not evaluated here.} \tag{3.16}$$

The scale hierarchy needed to integrate out the radial mode is separate from this coefficient matching:

$$m_\sigma \ll \Lambda_{\mathrm{UV}}, \qquad v \lesssim \Lambda_{\mathrm{UV}}. \tag{3.17}$$

These are working hierarchy assumptions, not consequences of a computed Newton coupling. No Planck-cutoff inference, cancellation, or numerical value of G_{eff} follows before Problem F is completed.

3.9 What has been assumed, collected

The microscopic theory, as used from here on, rests on:

1. the coordinate algebra (3.2) with cutoff (3.3);
2. the matrix action (3.4) at the pregeometric level;
3. the condensate action (3.6)–(3.7) after coordinates exist;
4. Assumption 3.1, that $g^{(0)}$ is a scaffold;
5. Assumption 3.5, that a vacuum $\langle \Phi^I{}_A \rangle = v\delta^I{}_A$ exists;
6. Assumption 3.6, the infrared commutator dictionary.

Everything in Chapters 4–11 is a calculation from this list, or an additional assumption announced as such. The list is long enough that the theory is not "derived from nothing." It is derived from a definite microscopic ansatz. That is still a stronger starting point than inserting $g_{\mu\nu}$ by hand, provided the later chapters do the work advertised in Chapter 1.

3.10 Exercises

1. **Dimensions.** Assign mass dimensions to $\hat{X}^\mu$, $\Theta^{\mu\nu}$, Φ, v, λ, and g_{YM} in four dimensions, with $\hbar = c = 1$. Confirm (3.3).
2. **Vacuum of the potential.** Show that (3.7) is bounded below and that its minima satisfy (3.8). Linearize $\Phi = v\,\mathbb{I} + \sigma + i\pi$ in the internal space (ignoring tangent structure) and read off m_σ^2 and m_π^2.
3. **Why $N \geq 4$.** If the internal dimension is $N = 2$, can a vacuum of the form (3.9) exist? What infrared representation content is then missing?

4. **Commutator to derivative.** In the Moyal plane with constant $\Theta^{12} = -\Theta^{21} = \theta$ and all other components zero, verify the leading term of (3.14) on a plane-wave matrix $\Phi = e^{ik\cdot\hat{X}}$. Where do higher powers of θ appear?

5. **IKKT vacuum energy.** Evaluate the first term of (3.4) on the c-number background (3.2). Show that it is a constant proportional to $\mathrm{Tr}(\Theta_{\mu\nu}\Theta^{\mu\nu})$. This constant is a preview of the $c_0\Lambda_{\mathrm{UV}}^4$ problem.

6. **Transformation law.** Starting from (3.10) and the representative vacuum (3.9), compute the infinitesimal variation of Φ under a simultaneous internal rotation and local Lorentz transformation that preserve the vacuum. Which generators are broken? (This is the algebra of Chapter 10, prepared here.)

7. **No graviton in the scalar truncation.** Set $N = 1$ and drop the tangent index, reducing Φ to a complex scalar. Show that the fluctuation spectrum of (3.6) contains no spin-2 state. This is the Chapter 1 ceiling written as a homework problem.

3.11 Bridge to Chapter 4

The field exists, the potential prefers a vacuum $\langle \Phi^I{}_A \rangle = v\delta^I{}_A$, and the coordinate algebra supplies a cutoff. Nothing yet is a metric. Chapter 4 defines the emergent vierbein as the condensate gradient

$$e_\mu^A(x) \equiv \langle \partial_\mu \Phi^{IA}(x) \rangle, \tag{3.18}$$

reconstructs $g_{\mu\nu} = \eta_{AB} e_\mu^A e_\nu^B$, and writes the Seiberg–Witten map that relates the non-commutative background $\Theta^{\mu\nu}$ to the leading correction $\mathcal{G}_{\mu\nu}(\langle \hat{X} \rangle)$. After that chapter, geometry is an output of the action written here.

Chapter 4

Condensation, Vierbein Emergence, and the Seiberg–Witten Map

Learning outcomes. After this chapter the reader should be able to:

1. explain why a strictly constant vacuum $\langle \Phi^I{}_A \rangle = v\delta^I{}_A$ cannot, by itself, produce a non-degenerate vierbein from $\langle \partial_\mu \Phi \rangle$;
2. write the polar decomposition $\Phi = (v + \sigma)\Lambda$, state the polar-derivative prescription of the source notes as an unverified proposal that does not supply a frame (Problem C), and state the frame of Repair II;
3. reconstruct $g_{\mu\nu} = \eta_{AB} e^A_\mu e^B_\nu$ and state the non-degeneracy condition;
4. use the leading Seiberg–Witten map to relate $\Theta^{\mu\nu}$ to a correction of the emergent metric;
5. enter Chapter 5 with a definite background about which to expand $\delta\Phi$.

Chapter 3 introduced the field and a preferred vacuum. This chapter turns that vacuum into geometry conditionally: it posits the Repair II spacetime-dependent condensate as the frame, since the Repair I prescription does not supply one. No Einstein equation is derived here. The only output is a composite metric, plus the dictionary that connects it to the non-commutative parameter $\Theta^{\mu\nu}$.

4.1 The constant-vacuum obstruction

The representative vacuum of Assumption 3.5 is

$$\langle \Phi^I{}_A \rangle = v\, \delta^I{}_A. \tag{4.1}$$

If this expectation value is independent of x, then

$$\langle \partial_\mu \Phi^{IA} \rangle = 0. \tag{4.2}$$

The definition advertised in the source notes,

$$\mathcal{E}^A{}_{\mu I}(x) \equiv \langle \partial_\mu \Phi^{IA}(x) \rangle, \tag{4.3}$$

then yields the zero tensor. A zero tensor is not a frame.

This is not a notational quibble. A textbook that passes from (4.1) to a non-degenerate $g_{\mu\nu}$ without an intermediate step is asserting an identity that is false. There are two candidate repairs. Only Repair II is carried through in this book; they must be kept distinct.

Repair I: the proposed polar-derivative prescription. Write the field in polar form, as in the ultraviolet note,

$$\Phi^{IA}(x) = \big(v + \sigma(x)\big)\, \Lambda^A{}_I(x), \tag{4.4}$$

where σ is a radial singlet and $\Lambda(x)$ takes values in the broken coset that rotates internal indices into tangent indices. Even if v is constant, $\partial_\mu \Phi$ need not vanish:

$$\partial_\mu \Phi^{IA} = (\partial_\mu \sigma)\, \Lambda^A{}_I + (v + \sigma)\, \partial_\mu \Lambda^A{}_I. \tag{4.5}$$

A vacuum with $\langle \sigma \rangle = 0$ and a slowly varying $\langle \Lambda \rangle$ has $\langle \partial_\mu \Lambda \rangle \neq 0$, but this is a candidate frame only in the sense of the source notes; see the discussion after Repair II.

Repair II: identify the vacuum itself with a frame. Promote the representative vacuum to a spacetime-dependent but still "pure Higgs" configuration

$$\langle \Phi^I{}_A(x) \rangle = v\, e^I{}_A(x), \tag{4.6}$$

and treat $e^I{}_A$ as the fundamental condensate profile. In this repair, equation (4.3) is abandoned as a definition. The vierbein *is* the normalized vacuum,

$$e^A_\mu(x) \equiv \frac{1}{v} \langle \Phi^A{}_\mu(x) \rangle, \tag{4.7}$$

after the internal and tangent bases have been identified.

Repair II is simpler and is the standard Higgs-gravity pattern. Repair I is what the notes write when they insist on (4.3). The two are not equivalent. Repair II defines a frame directly from the condensate and, wherever Assumption 4.4 holds, supplies a non-degenerate $e^A{}_\mu$. Repair I, as written in Definition 4.1, leaves the index I

uncontracted, and Appendix H (Section H.3) shows that no Lorentz-equivariant linear contraction of $\partial_\mu\Lambda$ produces a vierbein fluctuation. Definition 4.1 is therefore kept only as a record of the prescription in the source notes; it is not a constructed frame. In the rest of the book the frame is the Repair II frame (4.7), or a non-degenerate coframe assumed separately (Assumption 4.4); statements that use the explicit form (4.8) are conditional on a repair of Repair I that this book does not supply.

Definition 4.1 (Repair I prescription of the source notes)**.** In a polar configuration (4.4) with $\langle\sigma\rangle = 0$, define the three-index object

$$\mathcal{E}^A{}_{\mu I}(x) \equiv \langle\partial_\mu \Lambda^A{}_I(x)\rangle. \tag{4.8}$$

By (4.5) this is $v^{-1}\langle\partial_\mu \Phi^{IA}\rangle$ at $\sigma = 0$ (with $\langle\partial_\mu\sigma\rangle = 0$), which is the prescription the source notes write as a vierbein, with the internal index I left uncontracted and no identification of I with a tangent index used. It is not a vierbein: it carries the extra index I, and no Lorentz-equivariant linear contraction of it yields a vierbein fluctuation (Appendix H, Section H.3). The factor $1/v$ is the normalisation the notes use when Φ has the engineering dimension of v; if a formula in the notes omits $1/v$, it is using units $v = 1$.

Remark 4.2 (The two repairs do not agree)*.* The expansion $\Lambda^A{}_I = \delta^A{}_I + v^{-1}\omega^A{}_{I\mu}x^\mu + \cdots$ makes $\partial_\mu\Lambda^A{}_I$ a constant three-index tensor, antisymmetric in (A, I). It lies in $V \otimes \mathfrak{so}(3,1)$, while a vierbein perturbation lies in $V \otimes V$; the two share no irreducible summand (Appendix H, Section H.3). Repair I therefore does not reproduce Repair II at first order. A corrected Repair I would need additional background tensors or an independent coframe, which is part of Problem C.

4.2 The composite metric

Definition 4.3 (Composite metric)**.** Given a non-degenerate frame e^A_μ, taken from Repair II, (4.7), or supplied separately as an assumption (Assumption 4.4),

$$g_{\mu\nu}(x) \equiv \eta_{AB}\, e^A_\mu(x)\, e^B_\nu(x). \tag{4.9}$$

Non-degeneracy is not automatic.

Assumption 4.4 (Non-degenerate condensate frame)**.** On the region of spacetime under consideration, e^A_μ has non-vanishing determinant. Equivalently, the four covectors $e^A{}_\mu \mathrm{d}x^\mu$ are linearly independent.

If Assumption 4.4 fails, $g_{\mu\nu}$ is degenerate and there is no Lorentzian geometry to discuss. Vanishing of $\det e$ is a physical possibility—a condensate defect, a region

where the frame-locking vacuum has not formed—not a gauge artifact. This book restricts attention to the condensed phase in which Assumption 4.4 holds.

The inverse metric and inverse frame are defined in the usual way once non-degeneracy is granted,

$$g^{\mu\nu} e^A_\mu e^B_\nu = \eta^{AB}, \qquad e^\mu_A e^A_\nu = \delta^\mu_\nu. \tag{4.10}$$

Local Lorentz transformations act on the frame index,

$$e^A_\mu \;\longmapsto\; \Lambda^A{}_B e^B_\mu, \tag{4.11}$$

and leave $g_{\mu\nu}$ invariant. Diffeomorphisms act on the coordinate index. These two redundancies are the reason a composite metric can have the same geometric content as a primitive metric, provided the Goldstone counting of Chapter 10 works.

4.3 What kind of object e^A_μ is

At this stage e^A_μ is a condensate profile, not a solution of Einstein's equation. It is closer to a magnetization than to a gravitational field. Three consequences follow.

1. There is no a priori reason for e^A_μ to satisfy the torsion-free condition (2.6). If torsion is to vanish, that is a dynamical statement about the independent connection (whose dynamics remain open in Problem D), not a definition.

2. There is no a priori reason for small disturbances of $g_{\mu\nu}$ to propagate as massless spin-2 waves. That is Chapter 7.

3. The auxiliary scaffold $g^{(0)}_{\mu\nu}$ of Presentation B is still present in the microscopic action. It must not be confused with (4.9). Heat-kernel calculations in Chapter 8 are conditional on the emergent- background identification; its precise role in matching the physical G_{eff} remains part of Problem F.

4.4 Seiberg–Witten map and the Θ correction

Presentation A still has a non-commutative parameter $\Theta^{\mu\nu}$. After a commutative manifold has been introduced, the open-string / non-commutative-gauge-theory dictionary of Seiberg and Witten relates an ordinary metric $g_{\mu\nu}$ and a two-form background to an open-string effective metric. The source notes write the leading relation

$$G_{\mu\nu}(x) = \big((g+\Theta)^{-1}\big)_{\mu\nu} + \mathcal{G}_{\mu\nu}(\langle\hat{X}\rangle). \tag{4.12}$$

Two clarifications are required before (4.12) is used.

Matrix versus form. As written, $g+\Theta$ adds a symmetric tensor to an antisymmetric tensor. The honest object is the mixed tensor

$$(g+\Theta)_\mu{}^\nu = g_{\mu\rho}\eta^{\rho\nu} + \Theta_\mu{}^\nu, \tag{4.13}$$

or, equivalently, the linear map $g+\Theta : T^*M \to TM$ after one index of Θ is raised with the auxiliary metric. The inverse in (4.12) is the inverse of that map. In components, for constant Θ on a flat auxiliary background,

$$\left((g+\Theta)^{-1}\right)_{\mu\nu} = g_{\mu\nu} - g_{\mu\alpha}\Theta^{\alpha\beta}g_{\beta\nu} + O(\Theta^2). \tag{4.14}$$

The first correction is symmetric if one takes the symmetric part, which is the part that can enter a metric. The antisymmetric remainder is a two-form background, not a correction to $g_{\mu\nu}$.

The extra term $\mathcal{G}_{\mu\nu}$. The piece $\mathcal{G}_{\mu\nu}(\langle\hat{X}\rangle)$ in (4.12) is the backreaction of the matrix condensate on the emergent geometry, beyond the kinematic Θ correction. At this stage of the book it is a name, not a formula. Chapters 8 and 11 compute pieces of it: the Einstein–Hilbert term from the heat kernel, and the entropic stress tensor from mutual information. Until those calculations exist, $\mathcal{G}_{\mu\nu}$ must not be assigned a phenomenology.

Definition 4.5 (Working infrared metric)**.** Unless a higher-order Θ correction is being tracked explicitly, "the emergent metric" means (4.9). Equation (4.12) is the next-order kinematic correction plus a placeholder for dynamical backreaction.

4.5 A linearised frame around Minkowski spacetime

The fluctuation chapters need a background. Choose coordinates and a local Lorentz gauge so that the condensed frame is the trivial frame plus a small disturbance,

$$e_\mu^A = \delta_\mu^A + \tfrac{1}{2}h^A{}_\mu + \omega^A{}_\mu, \tag{4.15}$$

where $h_{\mu\nu} \equiv \eta_{AB}\delta_\mu^A h^B{}_\nu$ is symmetric and $\omega_{\mu\nu}$ is antisymmetric. Then

$$g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu} + O(h^2, \omega^2, h\omega). \tag{4.16}$$

The antisymmetric piece ω is a local Lorentz Goldstone. Under the assumptions of Chapter 10, its six orbit variables are removable in unitary gauge; this does not determine the dynamics of the independent diagonal connection. The symmetric piece $h_{\mu\nu}$ is the candidate graviton, still un-decomposed and possibly massive. Chapter 5 splits it; Chapter 7 asks whether its mass vanishes.

This expansion assumes that a Minkowski condensate exists: a vacuum in which $e^A_\mu = \delta^A_\mu$ globally, or in a region large compared with the scale of later experiments. Existence of that vacuum is part of Assumption 4.4, not a new principle.

4.6 Signature and the role of η_{AB}

Equation (4.9) borrows its signature from η_{AB}. Nothing in the potential (3.7) selected $(-,+,+,+)$ rather than $(+,+,+,+)$. A Euclidean condensate is a priori as natural as a Lorentzian one.

Assumption 4.6 (Lorentzian condensate)**.** The locked vacuum is Lorentzian: $\eta_{AB} = \mathrm{diag}(-1,+1,+1,+1)$. Dynamical emergence of this signature from a Euclidean matrix model is not proved in the source notes and is not pretended here. Chapter 10 shows how local $\mathrm{SO}(3,1)$ can arise *given* this η_{AB}.

A reader looking for a derivation of Lorentzian signature from nothing will not find it in this book. What the book does derive, contingent on Assumption 4.6, is the dynamics of the resulting Lorentzian metric.

4.7 Worked example: a weak, static frame perturbation

Take the Repair II frame, or any assumed non-degenerate frame, with the constant background $e^A{}_\mu = \delta^A{}_\mu$, and a small coordinate shift $x^\mu \to x^\mu + \zeta^\mu(x)$. The frame changes by its Lie derivative,

$$\delta e^A{}_\mu = \zeta^\nu \partial_\nu \delta^A{}_\mu + \partial_\mu \zeta^\nu \, \delta^A{}_\nu = \partial_\mu \zeta^A. \tag{4.17}$$

The associated metric perturbation is

$$h_{\mu\nu} = \partial_\mu \zeta_\nu + \partial_\nu \zeta_\mu \tag{4.18}$$

to first order. This configuration is geometry in name only: it is a coordinate artifact. A physical graviton must be a disturbance that cannot be written as (4.18). Isolating that disturbance is the point of the irreducible decomposition in Chapter 5.

The example demonstrates diffeomorphism redundancy of a Repair II (or assumed) frame. It does not derive a frame from Φ by Definition 4.1.

4.8 What this chapter has not done

The following statements are *not* theorems of Chapter 4.

- $g_{\mu\nu}$ obeys Einstein's equation.

- $h_{\mu\nu}$ contains two massless TT polarizations.
- A physical value of G_{eff} has not been obtained; it requires the specified sector calculation of Problem F.
- Torsion vanishes.
- The signature is dynamical.
- $\mathcal{G}_{\mu\nu}$ has been computed.

What has been done is narrower and indispensable: a composite metric has been defined from a non-degenerate condensate frame (Assumption 4.4); the constant-vacuum obstruction is addressed only by positing, in Repair II, a spacetime-dependent condensate that is identified with the frame (4.7) (no such profile is derived), while the Repair I formula of Definition 4.1 is not a constructed frame (Problem C, Appendix H); and the linearized Minkowski background used in every later proof has been fixed.

4.9 Exercises

1. **Zero frame.** Prove that a strictly constant $\langle\Phi\rangle$ yields $\langle\partial_\mu\Phi\rangle = 0$. This is the obstruction of Section 4.1 as a two-line exercise.
2. **Polar derivative.** Derive (4.5) from (4.4). Under what condition on Λ is e^A_μ proportional to a Maurer–Cartan form $\Lambda^{-1}\partial_\mu\Lambda$?
3. **Non-degeneracy.** Let $e^A_\mu = \mathrm{diag}(1, 1, 1, \varepsilon)$. Compute $g_{\mu\nu}$ and $\det g$. What fails as $\varepsilon \to 0$? Interpret the limit as a condensate defect.
4. **Symmetric part of the Θ correction.** From (4.14), extract the symmetric $O(\Theta)$ correction to $g_{\mu\nu}$ and the antisymmetric remainder. Which object can enter geodesic motion at this order?
5. **Linearized metric.** Starting from (4.15), derive (4.16) and show that ω_{AB} drops out of $g_{\mu\nu}$ at first order. Where does ω re-enter at second order?
6. **Pure gauge.** Show that (4.18) is the linearized diffeomorphism action on $\eta_{\mu\nu}$. Write one explicit $h_{\mu\nu}$ that cannot be written in this form. (Any TT wave will do.)
7. **Auxiliary versus emergent.** In Presentation B the microscopic action uses $g^{(0)}$. In this chapter the geometry is g. Write one sentence stating what must be assumed in order to replace $g^{(0)}$ by g inside a heat kernel, and why the displayed action does not supply it. The full answer is Chapter 8, Assumption 8.4.

4.10 Bridge to Chapter 5

The background is a condensed frame, taken Minkowski for the linearized theory,

$$\bar{e}_\mu^A = \delta_\mu^A, \qquad \bar{g}_{\mu\nu} = \eta_{\mu\nu}. \tag{4.19}$$

The next object is the fluctuation $\delta\Phi^I{}_A$ about the polar vacuum (4.4). Chapter 5 decomposes that fluctuation under local Lorentz transformations into a conformal piece, a vector piece, and a symmetric trace-free tensor $T_{\mu\nu}$. Only after that decomposition is it meaningful to ask which piece might be a graviton.

Chapter 5

Irreducible Lorentz Decomposition of Fluctuations

Learning outcomes. After this chapter the reader should be able to:

1. convert a fluctuation $\delta\Phi^I{}_A$ to a spacetime tensor using the background frame;
2. split that tensor into trace, antisymmetric, and symmetric trace-free pieces, and count components;
3. match the split to the schematic formula used in the source notes, $\delta\Phi^I{}_A = \frac{1}{4}\eta_{\mu\nu}\delta g^{\mu\nu} I + A_\mu \gamma^\mu + T_{\mu\nu}$;
4. distinguish algebraic irreducibility from the later dynamical conditions $\nabla^\mu T_{\mu\nu} = 0$ and $m_T^2 = 0$;
5. enter Chapter 6 knowing which piece is the candidate graviton.

Chapter 4 fixed a condensed background frame $\bar{e}^A_\mu = \delta^A_\mu$ (the Repair II frame, or an assumed non-degenerate frame) and a composite Minkowski metric. This chapter is linear algebra on the tangent space. No entanglement condition is used. No mass gap is shown to vanish. The only task is to name the pieces of $\delta\Phi$.

5.1 Fluctuations about the polar vacuum

Start from the polar field (4.4),

$$\Phi^{IA}(x) = \big(v + \sigma(x)\big)\, \Lambda^A{}_I(x), \tag{5.1}$$

and expand about the Minkowski condensate

$$\langle\sigma\rangle = 0, \qquad \langle\Lambda^A{}_I\rangle = \delta^A{}_I, \qquad \bar{e}^A_\mu = \delta^A_\mu. \tag{5.2}$$

Write

$$\Phi^{IA} = v\,\delta^{IA} + \delta\Phi^{IA}, \tag{5.3}$$

with $\delta\Phi$ first order in σ and in the deviation of Λ from the identity. For the rest of the linearized theory it is convenient to lower the internal index with the background identification $I \leftrightarrow A$ and convert the remaining tangent index to a coordinate index with $\bar{e}$:

$$\delta\Phi_{\mu\nu} \equiv \bar{e}^{A}{}_{\mu}\,\delta\Phi_{IA}\,\bar{e}^{I}{}_{\nu}. \tag{5.4}$$

When $N = 4$ and the vacuum of Assumption 3.5 locks every internal direction to a tangent direction, $\delta\Phi_{\mu\nu}$ is a genuine 4×4 spacetime tensor. If $N > 4$, (5.4) captures only the locked 4×4 block. The orthogonal internal block is a collection of extra Higgs fields. They are assumed heavy and integrated out; they are not candidate gravitons.

Assumption 5.1 (Locked block dominates the infrared)**.** Fluctuations in internal directions orthogonal to the four locked axes have masses of order v or Λ_{UV} and may be discarded from the infrared Lorentz decomposition.

5.2 Algebraic split under the Lorentz group

At a fixed event, $\delta\Phi_{\mu\nu}$ is an element of $\mathbb{R}^4 \otimes \mathbb{R}^4$. The Lorentz group $\mathrm{SO}(3,1)$ acts on each factor. The tensor product decomposes into irreducible pieces

$$\mathbf{4} \otimes \mathbf{4} = \mathbf{1} \oplus \mathbf{6} \oplus \mathbf{9}, \tag{5.5}$$

namely a scalar, an antisymmetric tensor, and a symmetric traceless tensor. Explicitly,

$$\delta\Phi_{\mu\nu} = \frac{1}{4}\eta_{\mu\nu}\psi + F_{\mu\nu} + T_{\mu\nu}, \tag{5.6}$$

where

$$\begin{aligned} \psi &= \eta^{\alpha\beta}\delta\Phi_{\alpha\beta}, \\ F_{\mu\nu} &= \frac{1}{2}(\delta\Phi_{\mu\nu} - \delta\Phi_{\nu\mu}), \\ T_{\mu\nu} &= \frac{1}{2}(\delta\Phi_{\mu\nu} + \delta\Phi_{\nu\mu}) - \frac{1}{4}\eta_{\mu\nu}\psi. \end{aligned} \tag{5.7}$$

By construction

$$T_{\mu\nu} = T_{\nu\mu}, \qquad \eta^{\mu\nu}T_{\mu\nu} = 0, \qquad F_{\mu\nu} = -F_{\nu\mu}. \tag{5.8}$$

Component count in four dimensions:

Piece	Symbol	Components
trace scalar	ψ	1
antisymmetric tensor	$F_{\mu\nu}$	6
symmetric traceless tensor	$T_{\mu\nu}$	9
total	$\delta\Phi_{\mu\nu}$	16

This is algebra. It uses no field equation and no gauge condition.

5.3 Translation into the notation of the source notes

The microscopic notes write

$$\delta\Phi^{I}{}_{A} = \frac{1}{4}\eta_{\mu\nu}\delta g^{\mu\nu}\, I + A_\mu\gamma^\mu + T_{\mu\nu}. \tag{5.9}$$

Equation (5.9) is (5.6) after three identifications.

Trace piece. On the linearized background of Section 4.5,

$$\psi = \eta^{\mu\nu}h_{\mu\nu} + O(\omega) = \delta g^{\mu}{}_{\mu}. \tag{5.10}$$

The first term of (5.9) is therefore the conformal (trace) mode of the composite metric. The identity matrix I in the notes is the locked vacuum $\delta^{I}{}_{A}$, converted to $\eta_{\mu\nu}$ by (5.4).

Antisymmetric / vector piece. The notes' identification of part of $F_{\mu\nu}$ with a vector current $A_\mu\gamma^\mu$ is schematic, not a disjoint decomposition of the six components of $F_{\mu\nu}$. In the present bookkeeping, those six components are the Goldstone orbit variables of Proposition 10.2; the internal gauge field A_μ is a separate field and is counted separately. The notes use Clifford generators γ^μ satisfying

$$\{\gamma^\mu, \gamma^\nu\} = 2\eta^{\mu\nu} \tag{5.11}$$

. The notes' field-strength notation is

$$F_{\mu\nu} = \nabla_\mu A_\nu - \nabla_\nu A_\mu + \omega_{\mu\nu}, \tag{5.12}$$

. Here this notation is schematic, not a decomposition into independent fields. The Lorentz-connection dynamics and the dynamics of A_μ are not fixed by this algebraic split.

Symmetric traceless piece. The tensor $T_{\mu\nu}$ in (5.9) is identical to the $T_{\mu\nu}$ of (5.7). It is the only algebraic piece with the index symmetries of a graviton.

5.4 What $T_{\mu\nu}$ is not yet

The source notes also write

$$T^{\mu}{}_{\mu} = 0, \qquad \nabla^{\mu} T_{\mu\nu} = 0. \tag{5.13}$$

The first of these is (5.8) and has already been imposed. The second is not algebraic. It is four differential conditions.

Those four conditions are the linearized statement that $T_{\mu\nu}$ is taken in de Donder / Lorentz-like gauge, or, on shell, that a massive spin-2 field has five rather than nine components. They are not free to impose until one knows the kinetic operator of $T_{\mu\nu}$.

Statement	Type	Status in this chapter
$T_{\mu\nu} = T_{\nu\mu}$	algebraic	imposed
$T^{\mu}{}_{\mu} = 0$	algebraic	imposed
$\nabla^{\mu} T_{\mu\nu} = 0$	gauge or on-shell	*not* imposed
$m_T^2 = 0$	dynamical	Chapter 7
$\Box T_{\mu\nu} = 0$ with two TT modes	dynamical	Chapter 7

A massive symmetric traceless tensor in four dimensions has five on-shell degrees of freedom (the Fierz–Pauli count). A massless one with the gauge invariance

$$\delta_\xi T_{\mu\nu} = \nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu - \tfrac{1}{2}\eta_{\mu\nu} \nabla \cdot \xi \tag{5.14}$$

has two. The difference is the entire physical question of Chapters 6 and 7. It cannot be settled by rearranging indices.

5.5 Kinetic terms before any entanglement condition

Inserting (5.6) into the quadratic expansion of S_{micro} produces, schematically,

$$S^{(2)} = S^{(2)}_{\psi}[\psi] + S^{(2)}_{A}[A] + S^{(2)}_{\omega}[\omega] + S^{(2)}_{T}[T] + S^{(2)}_{\text{mix}}. \tag{5.15}$$

Mixing terms of the form $\psi\, \partial\partial T$ and $A\, \partial T$ would be the usual linearized scalar–tensor and vector–tensor mixings. The quadratic expansion of S_{micro} on the flat scaffold contains no ψ–T mixing: its kinetic term contracts derivatives only through $g^{(0)\mu\nu}$, and the trace and η-traceless parts are orthogonal in its kinetic and potential forms. The internal connection A_μ mixes only with the anti-Hermitian part of $\delta\Phi$, with the derivative index contracted with that of A_μ; which components of $T_{\mu\nu}$ lie in that part depends on the identification of $T_{\mu\nu}$ in $\delta\Phi$ and on the internal gauge group. A $\psi\, \partial\partial T$ structure requires an additional ingredient (Appendix H, Section H.6). Two structural facts will be used later.

1. The tensor block has the generic form

$$S_T^{(2)} = \int \mathrm{d}^4x \Big(-\tfrac{1}{2}\nabla_\rho T_{\mu\nu}\nabla^\rho T^{\mu\nu} - \tfrac{1}{2}m_T^2 T_{\mu\nu}T^{\mu\nu} + \cdots \Big), \tag{5.16}$$

   in the signature $(-,+,+,+)$, where the ellipsis contains the additional derivative structures required by Fierz–Pauli if a mass term is present (the $T^\mu{}_\mu$ piece has already been projected out). The mass m_T^2 is a function of v, λ, and the condensate alignment. Nothing in Chapter 3 forces it to vanish.

2. The conformal block $S_\psi^{(2)}$ is the scalar-tensor sector. If ψ remains light, the infrared theory is not pure general relativity even if $m_T = 0$. Killing or gapping ψ is a separate requirement, recorded as such in Chapter 7 and again in Chapter 17.

Equation (5.16) is the action whose mass term Chapter 7 must kill. Writing it here prevents that later argument from looking as if $T_{\mu\nu}$ arrived already massless.

5.6 Diffeomorphism action on the pieces

An infinitesimal diffeomorphism generated by ξ^μ acts on the composite metric as

$$\delta_\xi g_{\mu\nu} = \nabla_\mu \xi_\nu + \nabla_\nu \xi_\mu. \tag{5.17}$$

On the linearized Minkowski background this is $\delta_\xi h_{\mu\nu} = \partial_\mu \xi_\nu + \partial_\nu \xi_\mu$. Passing through (5.6),

$$\begin{aligned} \delta_\xi \psi &= 2\partial\cdot\xi, \\ \delta_\xi F_{\mu\nu} &= 0 \quad \text{(to first order, after the symmetric projection)}, \\ \delta_\xi T_{\mu\nu} &= \partial_\mu \xi_\nu + \partial_\nu \xi_\mu - \tfrac{1}{2}\eta_{\mu\nu}\partial\cdot\xi. \end{aligned} \tag{5.18}$$

The last line is the gauge transformation that a massless spin-2 field must possess. If $m_T^2 \neq 0$, (5.18) is not a symmetry of (5.16), and the Ward identity that Chapter 6 will write on a causal diamond is obstructed. That obstruction is the lever used to force $m_T \to 0$.

5.7 Provisional sector bookkeeping

Component counts in the algebraic split are distinct from on-shell polarization counts. The independent diagonal connection B, torsion modes, and internal vector A_μ are not part of $\delta\Phi_{\mu\nu}$ and are counted separately.

Sector	Count and status
algebraic split	$\psi : 1, \quad F_{\mu\nu} : 6, \quad T_{\mu\nu} : 9; \quad$ total : 16
$F_{\mu\nu}$	Six Goldstone orbit variables (ω_{AB}); removable in unitary gauge (Ch. 10). No full-spectrum claim follows.
ψ	One component; gapped or constrained only under Assumption 7.5.
$T_{\mu\nu}$	Nine algebraic components. The full massive Fierz–Pauli theory has four on-shell divergence constraints and five polarizations; this is not established here for the NAMCG tensor block. Two TT helicities follow only under the separately established massless FP/diffeomorphism-gauge limit (Theorem 7.3); this is not a smooth $m_T \to 0$ limit of all massive polarizations.
Outside $\delta\Phi_{\mu\nu}$	A_μ: four components, omitted from the low-energy frame/metric-sector bookkeeping only if its specified quadratic operator gaps them; otherwise its spectrum is open. B and torsion remain unresolved (Problem D).

No full-theory mode count follows from this bookkeeping. The final "2" is only a metric-sector target.

5.8 Exercises

1. **Projectors.** Write the three projectors on $\mathbb{R}^4 \otimes \mathbb{R}^4$ that extract ψ, $F_{\mu\nu}$, and $T_{\mu\nu}$ from $\delta\Phi_{\mu\nu}$. Verify they are complementary and idempotent.

2. **Component count.** Confirm the $1 + 6 + 9 = 16$ split by writing every independent component of a general real 4×4 matrix in an explicit basis.

3. **Fierz–Pauli count.** A massive spin-2 field in $d = 4$ has how many on-shell degrees of freedom? Derive the number from SO(3) little-group representations. Where does the number five sit in the table of Section 5.7?

4. **Massless count.** Repeat for a massless spin-2 field with gauge invariance (5.14). Recover two helicities.

5. **Trace versus conformal mode.** From (4.16) and (5.10), show that ψ is the linearized conformal factor of $g_{\mu\nu}$ plus a term involving ω. In what gauge does the extra term drop?

6. **Diffeomorphism on $T_{\mu\nu}$.** Derive the third line of (5.18) from (5.17) and (5.7).

7. $N > 4$**.** Suppose $N = 8$. How many extra internal scalars appear outside the locked 4×4 block? Why are they not gravitons?

8. **Danger of imposing** $\nabla^\mu T_{\mu\nu} = 0$ **early.** Take (5.16) with $m_T \neq 0$ and impose $\partial^\mu T_{\mu\nu} = 0$ off shell as a constraint. Show that the constraint is consistent with the massive equation of motion, but that the massless limit $m_T \to 0$ must be taken *before* residual gauge transformations are used to reach two polarizations. This is why Chapter 7 cannot be skipped.

5.9 Bridge to Chapter 6

The candidate graviton has a name, $T_{\mu\nu}$, nine algebraic components, and a generic mass m_T. The conformal mode has a separate conditional gap assumption, and the internal vector is a distinct field whose spectrum remains conditional. Neither is identified with the gravitational-wave polarizations.

Nothing in the microscopic potential (3.7) sets $m_T = 0$. The mechanism proposed by NAMCG is not a further tuning of λ. It is the entanglement-stationarity condition $\delta S_D = 0$ on causal diamonds, imposed on the same fluctuations just decomposed. Chapter 6 defines that condition from the modular Hamiltonian of the matrix condensate and shows how it acts as a Ward identity. Chapter 7 applies the identity to $T_{\mu\nu}$ and obtains $m_T \to 0$.

Chapter 6

Entanglement Equilibrium and the Modular Hamiltonian

Learning outcomes. After this chapter the reader should be able to:

1. write the reduced state ρ_D, the modular Hamiltonian H_D, and the von Neumann entropy S_D of a causal diamond in the matrix condensate;
2. state Jacobson's entanglement-equilibrium condition $\delta S_D = 0$ and distinguish it from the Einstein equation it will later imply;
3. obtain the first variation $\delta S_D = \delta\langle H_D\rangle$ at fixed diamond volume, including the meaning of that constraint;
4. define the mutual-information potential U_{ent} used for two condensed cores;
5. enter Chapter 7 with a Ward identity ready to be applied to $T_{\mu\nu}$.

Chapter 5 named the candidate graviton $T_{\mu\nu}$ and left it with a generic mass m_T. The mass is not removed by a further term in $V(\Phi^\dagger\Phi)$. It is removed, if at all, by a stationarity condition on entanglement. This chapter writes that condition in a form that can be varied. The variation itself is Chapter 7.

6.1 The logical role of this chapter

There are two different claims that are easy to conflate.

1. **Jacobson's theorem** (in the form used here). For a quantum field on a prescribed geometry, the first-order stationarity of vacuum entanglement in small geodesic balls, at fixed volume, is equivalent to the semiclassical Einstein equation. This direction assumes a metric and a quantum field, and obtains $G_{\mu\nu} \propto \langle T_{\mu\nu}\rangle$.

2. **The NAMCG claim.** The microscopic condensate state is itself an entanglement-stationary state on the diamonds of its *emergent* metric. Stationarity is therefore a property of ρ, not an extra thermodynamic axiom pasted onto a metric theory. Applied to the tensor block of Chapter 5, it forbids $m_T^2 \neq 0$.

Claim 1 is a theorem in the literature, with stated hypotheses (conformal matter at first order; a conjecture for non-conformal matter). Claim 2 is the load-bearing physical hypothesis of NAMCG. This chapter formulates Claim 2 so that it can be used. It does not pretend that Claim 2 has been derived from the path integral of Chapter 3 with the same status as Claim 1.

A recent operator-algebraic treatment [2] gives a rigorous form of the matter side of Claim 1 for a free real scalar field: on a bifurcate Killing horizon, the relative entropy between the vacuum and a coherent excitation equals the null energy flux across the horizon. Its passage to the Einstein equation takes the metric, a local Minkowski approximation, the Raychaudhuri equation and the relation $S = A/4$ as inputs. It therefore belongs to Claim 1, and it is restricted to free fields and coherent states. It does not support Claim 2 and does not derive an area term for the emergent metric; that is the content of Problems A and B in Appendix H.1–H.2.

Assumption 6.1 (Microscopic entanglement equilibrium)**.** Let ρ be the vacuum state of the matrix condensate in the condensed phase of Chapters 3–4. For every sufficiently small causal diamond D constructed from the emergent metric $g_{\mu\nu}$,

$$\delta S_D = 0 \tag{6.1}$$

at fixed spatial volume of the maximal slice of D, to first order in variations of the condensate that preserve the local vacuum character of ρ.

The remainder of the chapter unpacks every symbol in that statement.

6.2 Diamonds in the emergent geometry

The causal diamond $D(p,q) = J^+(p) \cap J^-(q)$ was defined in (2.26). From here on the causal structure is that of the *emergent* metric (4.9), not of the auxiliary scaffold $g^{(0)}$. For a linearized analysis about Minkowski spacetime the distinction is invisible at zeroth order and appears at first order through $h_{\mu\nu}$.

A small diamond of radius ℓ is one for which $\ell^2 \|R_{\mu\nu\rho\sigma}\| \ll 1$. In Riemann normal coordinates (2.27) its maximal-slice three-volume admits the expansion

$$V(D) = \frac{4}{3}\pi\ell^3 - \frac{4\pi\ell^5}{15}\left(\tfrac{1}{3}R + \text{anisotropic Ricci terms}\right) + O(\ell^7). \tag{6.2}$$

"Fixed volume" in Assumption 6.1 means that the first variation is taken subject to $\delta V(D) = 0$. The geometric meaning of that constraint is that a conformal breathing of the diamond is not counted as an independent entanglement variation; it has been used to fix the comparison background. This is the same gauge-like restriction Jacobson imposes, and it is why the conformal mode ψ of Chapter 5 and the tensor mode $T_{\mu\nu}$ enter the first law differently.

Dyadic labels $D_{j,n}$ are a nested bookkeeping device. Nothing physical depends on bisection. Every identity written for a generic small D applies to each dyadic diamond.

6.3 Reduced states and the modular Hamiltonian

Let $\mathcal{A}_D$ be the algebra of condensate observables supported in D. In Presentation B these are functions of $\Phi(x)$ and $A_\mu(x)$ for $x \in D$. In Presentation A they are the operators generated by the spectral projections of $\hat{X}^\mu$ whose eigenvalues lie in D, after the dictionary of Assumption 3.6.

Definition 6.2 (Diamond state and modular Hamiltonian)**.** Given a global state ρ of the condensate,

$$\rho_D = \mathrm{Tr}_{\mathcal{A}'_D} \rho, \tag{6.3}$$

where $\mathcal{A}'_D$ is the commutant (the complement algebra). The modular Hamiltonian and von Neumann entropy are

$$H_D = -\ln \rho_D, \qquad S_D = -\mathrm{Tr}(\rho_D \ln \rho_D) = \langle H_D \rangle_{\rho_D}. \tag{6.4}$$

For a free field in Rindler space, the Bisognano–Wichmann theorem identifies H_D with $2\pi/\kappa$ times the boost generator that fixes the edge $\mathcal{B} = \partial D \cap \Sigma$. For a small diamond in a slowly varying background the same identification holds to leading order in ℓ:

$$H_D = \frac{2\pi}{\kappa} \int_{\Sigma \cap D} \mathrm{d}\Sigma^\mu \, \xi^\nu T^{\mathrm{op}}_{\mu\nu} + H_{\mathrm{area}} + O(\ell^3). \tag{6.5}$$

Here ξ^μ is the approximate conformal Killing vector of D, κ is its surface gravity at $\mathcal{B}$, $T^{\mathrm{op}}_{\mu\nu}$ is the microscopic stress tensor of the condensate fluctuations, and H_{area} is the area term that will later match $A/(4G_{\mathrm{eff}})$.

Equation (6.5) is the working formula of this book. It is exact only for free fields in Rindler space. For interacting matrix fluctuations it is the first term in an operator product expansion on the diamond. Higher terms are absorbed into the $O(\ell^3)$ remainder and into the definition of H_{area}.

6.4 Replica formula

When a direct construction of ρ_D is unavailable, the replica trick supplies S_D. For integer $n \geq 1$,

$$S_D = \lim_{n\to 1}(1 - n\partial_n) \ln \mathrm{Tr}\,(\rho_D)^n, \tag{6.6}$$

and $\mathrm{Tr}\,(\rho_D)^n$ is computed as a path integral of the microscopic action (3.6) on an n-sheeted cover of M branched over the edge $\mathcal{B}$. The first variation δS_D then becomes a variation of that branched partition function with respect to the background frame. In the Euclidean replica formulation, after Wick rotation, the path integral uses $S_E = \int \mathrm{d}^4x\,\sqrt{g}\,\mathrm{Tr}\big(g^{\mu\nu}(D_\mu\Phi)^\dagger D_\nu\Phi + V\big)$ as the continuation of (3.6). Matching the corrected Lorentzian sign of (3.6) to Presentation A remains unproved under Assumption 3.6.

Two remarks are enough for Chapter 7.

- The replica construction is well defined in Presentation B, where a path integral exists. In Presentation A it requires the dictionary of Assumption 3.6.

- Differentiating (6.6) at $n = 1$ produces an insertion of the modular Hamiltonian. The first variation $\delta S_D = \delta\langle H_D\rangle$ used below is therefore not a second formula; it is the replica formula at first order.

6.5 The first law of the diamond

Vary Definition 6.2 at fixed diamond, then allow the diamond itself to vary with the metric. The result, to first order and at fixed volume, is the entanglement first law

$$\delta S_D = \delta\langle H_D\rangle = \frac{2\pi}{\kappa}\int_{\Sigma\cap D} \mathrm{d}\Sigma^\mu\,\xi^\nu\,\delta\langle T^{\mathrm{op}}_{\mu\nu}\rangle + \frac{\delta A}{4G_{\mathrm{eff}}} + O(\ell^3). \tag{6.7}$$

The area variation δA is computed from the emergent metric. The identification of its prefactor with $1/(4G_{\mathrm{eff}})$ is not justified in this chapter; it is the Bekenstein–Hawking relation that Chapter 8 will match to the heat kernel. Until then, $1/(4G_{\mathrm{eff}})$ is a name for the coefficient of δA.

Assumption 6.1 now reads

$$\frac{2\pi}{\kappa}\int_{\Sigma\cap D} \mathrm{d}\Sigma^\mu\,\xi^\nu\,\delta\langle T^{\mathrm{op}}_{\mu\nu}\rangle + \frac{\delta A}{4G_{\mathrm{eff}}} = 0, \tag{6.8}$$

up to $O(\ell^3)$, for every small diamond and every allowed variation of the condensate.

If $\delta\langle T^{\mathrm{op}}_{\mu\nu}\rangle$ were the stress tensor of ordinary matter on a prescribed geometry, and if δA were computed from a prescribed Einstein metric, (6.8) would be Jacobson's

derivation of $G_{\mu\nu} = 8\pi G_{\rm eff}\,\langle T_{\mu\nu}\rangle$. In NAMCG both $\delta\langle T^{\rm op}\rangle$ and δA are functionals of the *same* fluctuation $\delta\Phi$. That is the leverage used in Chapter 7.

6.6 Allowed variations

Not every deformation of Φ is an allowed variation in Assumption 6.1. The hypothesis is a statement about perturbations that remain locally vacuum-like: they do not inject an arbitrary excited-state entropy into D.

Definition 6.3 (Allowed variation)**.** A first-order deformation $\delta\Phi$ is *allowed* if

1. it is smooth on a neighbourhood of D;
2. it preserves the polar vacuum to zeroth order outside a compact region, so that ρ remains a condensed-phase vacuum at infinity;
3. its restriction to each algebraic block of Chapter 5 may be chosen independently, after gauge fixing of diffeomorphisms and local Lorentz transformations.

Condition 3 is the reason the tensor, vector, and conformal blocks can be tested separately. If a variation supported only on $T_{\mu\nu}$ violates (6.8) unless $m_T = 0$, that is a constraint on the tensor block alone.

6.7 Mutual information of two cores

The same entropy data define an interaction between two disjoint condensed regions A and B (two compact-object cores, two stellar interiors, two laboratory samples).

Definition 6.4 (Entropic potential)**.**

$$U_{\rm ent}(A,B) = -\lambda I(A:B) = -\lambda\big[S(A) + S(B) - S(A\cup B)\big], \tag{6.9}$$

with $\lambda > 0$ a microscopic coupling of dimension energy.

Because $I(A:B) \geq 0$, one has $U_{\rm ent} \leq 0$. Increasing the mutual information lowers the potential: the effective force is attractive. For two small balls whose proper separation d is large compared with their radii but small compared with the curvature radius,

$$I(A:B) \;\sim\; c_{p_*}\left(\frac{\ell_A\ell_B}{d^2}\right)^{p_*} \qquad (d \gg \ell_A, \ell_B). \tag{6.10}$$

Here p_* is the smallest sum of scaling dimensions of operators whose product has vacuum quantum numbers, and c_{p_*} is a dimensionless, theory- and shape-dependent coefficient. For the exchange of a pair of operators of dimension Δ_*, $p_* = 2\Delta_*$; other

channels can set a different leading power. Cardy [3] derives this scaling for the Rényi mutual information of compact regions in the ground state of a flat-space conformal field theory, and gives the von Neumann limit explicitly for spherical regions in the free-scalar case in $2+1$ and $3+1$ dimensions, where for $3+1$ dimensions $p_* = 2$, i.e. a tail proportional to $\ell_A^2 \ell_B^2 / d^4$. Applying (6.10) to the von Neumann mutual information of other theories assumes that the $n \to 1$ continuation holds, and applying it to NAMCG cores, which need not be vacuum balls of a conformal theory, is a further assumption. Neither p_* nor c_{p_*} is computed here for NAMCG. The static $U_{\rm ent}$ defines a conservative interaction. Its separation dependence alone does not determine a velocity-dependent dissipative force or a retarded response. The Newtonian laboratory of Chapter 15 adopts

$$\mathbf{a}_{\rm irrev} = -\eta \, \frac{1}{d^4} \, \mathbf{v} \tag{6.11}$$

as a phenomenological working law. Equation (6.11) is not derived from (6.10); its microscopic dynamical origin is Problem H.9. It is flagged here so that Chapter 11 knows which entropy functional it must vary covariantly.

The conservation question is already visible. Mutual information is built from scalars $S(\,\cdot\,)$. A covariant variation of $U_{\rm ent}$ with respect to the metric defines a tensor $T^{\rm ent}_{\mu\nu}$ by the same rule as (2.13). Whether $\nabla^\mu T^{\rm ent}_{\mu\nu} = 0$ holds on shell is not automatic from the sign of λ. It is a calculation, reserved for Chapter 11.

6.8 What stationarity does *not* say

Assumption 6.1 does not assert any of the following.

- That S_D is a maximum rather than a saddle. Jacobson's original language is maximization at fixed volume; the identities used below need only first-order stationarity.
- That the area term in (6.7) has already been matched to $G_{\rm eff}$. That match is Chapter 8.
- That $m_T = 0$. Stationarity is a condition on variations. Whether the tensor kinetic operator is compatible with that condition is Chapter 7.
- That $\Lambda_{\rm eff}$ is small. Vacuum entropy of the diamond includes a piece that will source a cosmological term. Chapter 12 treats it.
- That the replica path integral has been computed at two loops. The book uses the first law (6.7) as the working identity, not a finished replica evaluation of S_D for the matrix potential (3.7).

6.9 The Ward identity that Chapter 7 will use

Collecting (6.8), Definition 6.3, and the split (5.6), the usable statement is:

Proposition 6.5 (Entanglement Ward identity)**.** *Under Assumptions 6.1 and 3.6, every allowed variation supported on a single algebraic block ψ, A_μ, or $T_{\mu\nu}$ satisfies*

$$\int_{\Sigma\cap D} \mathrm{d}\Sigma^\mu\, \xi^\nu \delta\langle T^{\mathrm{op}}_{\mu\nu}\rangle + \frac{\kappa}{8\pi G_{\mathrm{eff}}}\,\delta A = 0 \tag{6.12}$$

to first order in ℓ, at fixed $V(D)$. The operator $T^{\mathrm{op}}_{\mu\nu}$ is the metric variation of the quadratic action of that block.

The proof of Proposition 6.5 is the chain of this chapter: Definition 6.2 to the first law (6.7) to Assumption 6.1, restricted to one block by Definition 6.3. No property of $T_{\mu\nu}$ beyond its being one algebraic block has been used. That is intentional. Chapter 7 substitutes the quadratic action (5.16) into (6.12) and reads off $m_T = 0$.

6.10 Exercises

1. **Volume constraint.** From (6.2), show that a pure conformal variation $\delta g_{\mu\nu} = \chi\eta_{\mu\nu}$ changes $V(D)$ at order ℓ^3. Why must this variation be excluded, or compensated, if stationarity is imposed at fixed volume?

2. **Rindler check.** In Rindler space with Unruh temperature $T = \kappa/2\pi$, verify that $\delta S = \delta E/T$ for a boost-energy perturbation is the restriction of (6.7) to a diamond that has become a Rindler wedge.

3. **Replica derivative.** Starting from (6.6), show that $\partial_n \ln \mathrm{Tr}\,\rho^n|_{n=1} = \langle \ln\rho\rangle$, and hence that $S = \langle H\rangle$ at $n = 1$.

4. **Allowed versus thermal variations.** Give one example of a $\delta\Phi$ that changes the particle content of D (a one-particle excitation) and explain why Assumption 6.1 is silent about it.

5. **Mutual information positivity.** Prove $I(A:B) \geq 0$ from relative entropy. Deduce $U_{\mathrm{ent}} \leq 0$ for $\lambda > 0$.

6. **Static information versus drag.** Explain why a static power law for $I(A:B)$ does not determine the low-frequency retarded force response or its drag exponent. What dynamical information is required to calculate that response?

7. **Two metrics.** Write one paragraph explaining why the diamond of this chapter is built from $g_{\mu\nu}$ rather than from $g^{(0)}_{\mu\nu}$. What would go wrong, conceptually, if stationarity were imposed on auxiliary-scaffold diamonds instead?

6.11 Bridge to Chapter 7

The candidate graviton $T_{\mu\nu}$ has a quadratic action (5.16) with an unknown mass m_T. The condensate vacuum is assumed to obey the entanglement Ward identity (6.12) on every small diamond of the emergent geometry. Chapter 7 substitutes a compactly supported tensor variation into that identity. The area term tracks the Einstein operator. The stress-tensor term tracks $m_T^2 T_{\mu\nu}$. Compatibility of the two forces $m_T = 0$, after which the surviving kinetic operator is the linearized Einstein operator and the infrared spectrum contains two massless TT polarizations—provided the conformal and vector blocks are separately gapped or eaten, as already flagged.

Chapter 7

Vanishing of the Tensor Mass Gap

Learning outcomes. After this chapter the reader should be able to:

1. write the Fierz–Pauli quadratic action for $T_{\mu\nu}$ and the linearized Einstein operator that remains when $m_T = 0$;
2. substitute a compactly supported tensor variation into the entanglement Ward identity and see where m_T^2 appears;
3. state Theorem 7.3 with its hypotheses, and say which hypotheses are still assumptions of the framework;
4. count the two remaining TT polarizations;
5. list what must still be done for the conformal and vector blocks before the infrared theory is pure GR.

The candidate graviton now has a name, a quadratic action, and a Ward identity. This chapter puts the last of those into the second and reads off $m_T = 0$.

7.1 Strategy

On a Minkowski condensate the tensor block is a linear field $T_{\mu\nu}$ with the algebraic symmetries (5.8). Its quadratic action has the generic structure (5.16). The entanglement Ward identity, Proposition 6.5, must hold for every allowed variation supported on that block.

The geometric side of the identity, δA, depends on $T_{\mu\nu}$ only through the linearized metric

$$h_{\mu\nu} = T_{\mu\nu} + \tfrac{1}{4}\eta_{\mu\nu}\psi, \tag{7.1}$$

and, in a gauge where the conformal piece is held fixed by the fixed-volume constraint of Assumption 6.1, through $T_{\mu\nu}$ alone. The modular side, $\delta\langle T^{\rm op}_{\mu\nu}\rangle$, is the metric stress of the quadratic tensor action. If that action contains a mass term, the modular side contains a piece proportional to $m_T^2 T_{\mu\nu}$. The geometric side does not. Compatibility for arbitrary compactly supported $T_{\mu\nu}$ forces $m_T^2 = 0$.

That is the argument. The rest of the chapter writes it so that each step can be rejected independently.

7.2 Fierz–Pauli action for the tensor block

The unique two-derivative quadratic action for a massive symmetric tensor that is ghost-free about Minkowski spacetime is the Fierz–Pauli action. For a general symmetric $h_{\mu\nu}$, in the signature $(-,+,+,+)$, the full action contains the following additional trace-dependent terms, where $h = h^{\mu}{}_{\mu}$:

$$-\partial_\mu h^{\mu\nu}\partial_\nu h + \tfrac{1}{2}\partial_\rho h\,\partial^\rho h + \tfrac{1}{2}m_T^2 h^2.$$

On restriction to a traceless field $T_{\mu\nu}$, it reduces to

$$\begin{aligned} S_{\rm FP}[T] = \int \mathrm{d}^4x\Big(&-\tfrac{1}{2}\partial_\rho T_{\mu\nu}\partial^\rho T^{\mu\nu} + \partial^\rho T_{\rho\mu}\partial_\sigma T^{\sigma\mu} \\ &- \tfrac{1}{2}m_T^2 T_{\mu\nu}T^{\mu\nu}\Big). \end{aligned} \tag{7.2}$$

In this signature the transverse-traceless spatial modes have kinetic term $+\frac{1}{2}\dot{T}_{ij}^2$. The derivative structure is required. A generic mass term $m^2(T_{\mu\nu}T^{\mu\nu} - \alpha T^2)$ with $\alpha \neq 1$ (before the trace projection) propagates a ghost. With the trace already zero the remaining danger is a wrong-sign kinetic term, which (7.2) does not have.

The unrestricted Euler–Lagrange equation of the full Fierz–Pauli action, evaluated at $h_{\mu\nu} = T_{\mu\nu}$ with vanishing trace, is

$$\mathcal{E}_{\mu\nu}[T] - m_T^2 T_{\mu\nu} = 0, \tag{7.3}$$

where

$$\begin{aligned} \mathcal{E}_{\mu\nu}[T] \equiv\ &\Box T_{\mu\nu} - \partial_\mu\partial^\rho T_{\rho\nu} - \partial_\nu\partial^\rho T_{\rho\mu} \\ &+ \eta_{\mu\nu}\partial^\rho\partial^\sigma T_{\rho\sigma} \end{aligned} \tag{7.4}$$

is the linearized Einstein operator on a traceless tensor about Minkowski spacetime. (The linearized Einstein tensor on a general $h_{\mu\nu}$ contains additional trace pieces; they are absent here because $T^{\mu}{}_{\mu} = 0$.)

For $m_T \neq 0$, the divergence of the full equation gives $\partial^\mu h_{\mu\nu} = \partial_\nu h$. Combining this with its trace gives $h = 0$; only then does the trace of the equation imply $\partial^\rho\partial^\sigma T_{\rho\sigma} = 0$,

yielding five polarizations for the full Fierz–Pauli theory. Varying (7.2) with $T^{\mu}{}_{\mu} = 0$ imposed in the action instead gives the traceless projection, with $\frac{1}{2}\eta_{\mu\nu}\partial^{\rho}\partial^{\sigma}T_{\rho\sigma}$. Its divergences give $(\Box + 2m_T^2)\partial^{\rho}\partial^{\sigma}T_{\rho\sigma} = 0$, not the constraint. Thus the displayed trace-constrained action alone does not establish the Fierz–Pauli constraints or five-polarization count, and the NAMCG tensor block is not shown to have the full Fierz–Pauli trace sector and constraint structure. Whether and how the ψ sector supplies them is unresolved and is a dependency of Problem C.

When $m_T = 0$, (7.2) is invariant under

$$\delta_{\xi}T_{\mu\nu} = \partial_{\mu}\xi_{\nu} + \partial_{\nu}\xi_{\mu} - \tfrac{1}{2}\eta_{\mu\nu}\partial\cdot\xi \tag{7.5}$$

with $\partial\cdot\xi$ arranged so that the trace remains zero. For the full Fierz–Pauli theory, when $m_T \neq 0$ that invariance is absent and the on-shell count is five degrees of freedom; this count is not established here for the NAMCG tensor block.

7.3 Modular stress of the tensor block

The operator $T^{\mathrm{op}}_{\mu\nu}$ in Proposition 6.5 is defined by the same rule as (2.13), applied to the full Fierz–Pauli action treated as a matter action on the background metric that defines the diamond. The following relation holds for that full theory; it is not established here for the NAMCG tensor block. To first order about Minkowski spacetime, evaluated on the full-theory constraint surface,

$$\delta\langle T^{\mathrm{op}}_{\mu\nu}\rangle = \mathcal{E}_{\mu\nu}[T] - m_T^2 T_{\mu\nu} + \partial^{\rho}\Sigma_{\rho\mu\nu}, \tag{7.6}$$

where $\Sigma_{\rho\mu\nu}$ is a total-derivative superpotential coming from the fact that the canonical and metric stress tensors differ by an improvement. On a diamond the superpotential contributes only at the edge $\mathcal{B}$. That edge contribution is absorbed into the area term of the first law, which is already present in (6.12). What remains in the bulk integral is the improved stress

$$\delta\langle T^{\mathrm{op}}_{\mu\nu}\rangle_{\mathrm{bulk}} = \mathcal{E}_{\mu\nu}[T] - m_T^2 T_{\mu\nu}. \tag{7.7}$$

7.4 Area variation induced by $T_{\mu\nu}$

The edge $\mathcal{B}$ of a small diamond of radius ℓ is a 2-sphere of area $4\pi\ell^2$ in the Minkowski background. A metric perturbation $h_{\mu\nu} = T_{\mu\nu}$ changes that area by

$$\delta A = \int_{\mathcal{B}} \mathrm{d}A\, \tfrac{1}{2}\sigma^{\mu\nu}T_{\mu\nu} + O(\ell^3\|T\|, \ell\|T\|^2), \tag{7.8}$$

where $\sigma^{\mu\nu}$ is the induced metric on $\mathcal{B}$. Jacobson's geometric identity converts the same variation, after integration against the conformal Killing vector and use of the fixed-volume constraint, into

$$\frac{\kappa}{8\pi G_{\text{eff}}}\,\delta A = -\int_{\Sigma\cap D} \mathrm{d}\Sigma^{\mu}\,\xi^{\nu}\,\mathcal{E}_{\mu\nu}[T] + O(\ell^3). \tag{7.9}$$

The right-hand side contains the Einstein operator and does *not* contain m_T. The area of an emergent 2-surface knows about the metric perturbation. It does not know about a potential term in the fluctuation action.

Equation (7.9) is imported from the geometry of small diamonds, not re-derived here. Its hypotheses are: Riemann normal coordinates to the order kept in (2.27), fixed volume, and the identification of the area prefactor with G_{eff}. The latter remains conditional on the specified sector calculation in Problem F; Chapter 8 gives only the truncated heat-kernel identity.

7.5 Substitution into the Ward identity

Insert (7.7) and (7.9) into (6.12). The Einstein-operator terms cancel, and what remains is

$$\int_{\Sigma\cap D} \mathrm{d}\Sigma^{\mu}\,\xi^{\nu}(-m_T^2 T_{\mu\nu}) + O(\ell^3) = 0. \tag{7.10}$$

The conformal Killing vector ξ^{ν} of a small diamond is timelike in the interior and vanishes on the edge. Its leading expression in Riemann normal coordinates is

$$\xi^{\mu} = \frac{\kappa}{2\ell}\big((\ell^2 - t^2 - r^2)\partial_t^{\mu} + 2t\,x^{\mu}\big) + O(R_{\alpha\beta\gamma\delta}\ell^3). \tag{7.11}$$

Lemma 7.1 (Arbitrary tensor sources)**.** *Compactly supported, smooth, traceless, symmetric variations $T_{\mu\nu}$ are allowed in the sense of Definition 6.3. In particular they may be chosen independently in each small diamond, up to the fixed-volume constraint already used.*

Proof. The tensor block was split off algebraically in Chapter 5. Gauge fixing of diffeomorphisms and local Lorentz transformations removes ψ-mixing and ω-mixing at this order but does not constrain the nine components of $T_{\mu\nu}$ beyond smoothness and compact support. Fixed volume constrains the trace sector, which has already been projected out. □

Lemma 7.2 (Integral identity implies point identity)**.** *If* (7.10) *holds for every small diamond and every allowed $T_{\mu\nu}$, then*

$$m_T^2 T_{\mu\nu}(x) = 0 \tag{7.12}$$

at every event x in the condensed phase.

Proof. Suppose $m_T^2 T_{\mu\nu}$ is not identically zero. Then there is an event x_0 and a pair of indices at which the tensor is nonzero. Choose a diamond centered at x_0 with ℓ small enough that $T_{\mu\nu}$ does not change sign in the relevant component on $\Sigma \cap D$, and choose the time orientation so that ξ^ν has a nonvanishing contraction with that component. The integrand of (7.10) is then of definite sign on a set of positive measure, and the $O(\ell^3)$ remainder cannot cancel it as $\ell \to 0$. □

7.6 The vanishing theorem

Theorem 7.3 (Vanishing of the tensor mass gap)**.** *Assume:*

1. *the microscopic setup of Chapter 3, including Assumptions 3.1–3.6;*
2. *the condensed frame of Chapter 4, including Assumptions 4.4 and 4.6;*
3. *the algebraic split of Chapter 5 and Assumption 5.1;*
4. *entanglement equilibrium, Assumption 6.1, in the form of Proposition 6.5;*
5. *the Fierz–Pauli quadratic action* (7.2) *as the two-derivative tensor block;*
6. *the geometric identity* (7.9) *for small diamonds.*

Then $m_T^2 = 0$ throughout the condensed phase. The quadratic tensor action reduces to the linearized Einstein–Hilbert action on traceless perturbations,

$$S_{\mathrm{EH}}^{(2)}[T] = \int \mathrm{d}^4x \, \tfrac{1}{2} T^{\mu\nu} \mathcal{E}_{\mu\nu}[T], \tag{7.13}$$

and is invariant under (7.5)*.*

Proof. Lemma 7.2 gives $m_T^2 T_{\mu\nu} = 0$ at every event. The tensor block is not the zero field: linearized gravitational waves, if they exist, are particular solutions for $T_{\mu\nu}$. Hence $m_T^2 = 0$. Substituting into (7.2) produces (7.13) after an integration by parts, boundary terms being discarded for compactly supported variations. Gauge invariance (7.5) is the standard invariance of the linearized Einstein operator on a traceless tensor. □

Remark 7.4 (What the theorem does not hide). Hypothesis 4 is the physical input of NAMCG. Hypothesis 5 excludes non-Fierz–Pauli two-derivative structures and higher-derivative quadratic terms; the latter return in Chapter 9. Hypothesis 6 borrows a geometric identity used in Jacobson's argument. The cancellation of the Einstein-operator terms in (7.10) also uses that the coefficient of $\mathcal{E}_{\mu\nu}$ in the stress

response (7.7) equals the coefficient of $\mathcal{E}_{\mu\nu}$ in the area term (7.9). This equality is part of the content of Hypotheses 5 and 6 as written; it is not supplied by Hypothesis 4 or by the identification of the area prefactor with G_{eff} (Section H.2). The theorem is therefore a compatibility statement: a Fierz–Pauli tensor block cannot sit in an entanglement-stationary condensate unless it is massless. It is not a derivation of Assumption 6.1 from the path integral.

7.7 Two transverse-traceless polarizations

With $m_T = 0$, the equation $\mathcal{E}_{\mu\nu}[T] = 0$ plus the gauge invariance (7.5) is linearized vacuum GR in the traceless sector. Choose the de Donder gauge

$$\partial^\mu T_{\mu\nu} = 0. \tag{7.14}$$

Residual gauge transformations with $\Box\xi^\mu = 0$ remain. They remove four more components. Together with the four conditions (7.14) and the four conditions implicit in taking the trace to vanish and staying in the symmetric sector after gauge fixing, one is left with two independent components. In a plane- wave basis they are the helicities ± 2,

$$T_{\mu\nu}(x) = \mathrm{Re}\big(\epsilon^{\pm 2}_{\mu\nu}\, e^{ik\cdot x}\big), \qquad k^2 = 0, \quad k^\mu \epsilon_{\mu\nu} = 0, \quad \epsilon^\mu{}_\mu = 0. \tag{7.15}$$

This is the infrared graviton advertised in Chapter 1. It has been obtained as a fluctuation of the condensate, not as a fundamental metric field, contingent on Theorem 7.3.

7.8 The van Dam–Veltman–Zakharov warning

A massive spin-2 field with m_T small but nonzero does not approach the massless theory smoothly in four dimensions. The discontinuity is visible in the linearized deflection of light and in the tensor-to-scalar ratio of a static source. Therefore it is not enough to argue that m_T is "small compared with Λ_{UV}." Entanglement equilibrium is being asked to set m_T identically to zero in the infrared, not to suppress it by a loop factor.

If a later calculation produced $m_T \sim v\, e^{-\Lambda_{\text{UV}}/v}$ or any other parametrically small but nonzero value, the theory would fail Solar-System tests unless an additional Vainshtein mechanism were demonstrated. No such mechanism is claimed here. Theorem 7.3 is an identity-zero statement.

7.9 Conformal and vector blocks

Theorem 7.3 is silent about ψ and A_μ.

Vector block. The internal gauge connection A_μ is distinct from the independent Lorentz connection ω_μ and from their diagonal stabilizer combination. Its mass, if any, depends on kinetic normalizations and the residual internal symmetry, neither of which is specified here. If the internal gauge group has an unbroken subgroup, the corresponding field is massless. Neither case is claimed to produce a second graviton. Chapter 10 identifies the broken-direction combination and the unitary-gauge removal of the Goldstone orbit variables; it does not settle the diagonal connection's dynamics (Problem D).

Conformal block. The scalar ψ is more dangerous. If it remains massless it is a Brans–Dicke field, and the infrared theory is scalar–tensor gravity. Entanglement equilibrium at fixed volume was designed to remove a conformal breathing from the comparison diamond; it does not, by itself, gap ψ as a propagating excitation.

Assumption 7.5 (Gapped conformal mode)**.** The radial Higgs σ of the polar decomposition, and the conformal metric mode ψ mixed with it, acquire a mass $m_\psi \sim \sqrt{\lambda}\, v$ from the potential (3.7) and decouple from the infrared tensor sector.

Assumption 7.5 is the scalar-tensor ceiling of Chapter 1, reappeared as a named hypothesis. If it fails, NAMCG predicts a light scalar and is constrained by Cassini-type bounds on ω_{BD}. Chapter 17 records this as a falsifier, not as a detail.

7.10 Exercises

1. **Fierz–Pauli ghost.** Restore a trace piece $T^\mu{}_\mu = \psi$ and replace the mass term by $m^2(T_{\mu\nu}T^{\mu\nu} - \alpha\psi^2)$. Show that $\alpha \neq 1$ produces a sixth, ghostlike excitation. This is why (7.2) is not an arbitrary massive action.

2. **Gauge invariance at $m_T = 0$.** Verify that (7.5) leaves (7.13) invariant. Show the mass term in (7.2) is not invariant.

3. **Einstein operator on a TT wave.** Insert (7.15) into (7.4) and recover $k^2 = 0$.

4. **Sign in the Ward identity.** Track the sign from (7.7) through (7.9) to (7.10). A sign error here would force $m_T^2 = 0$ by the same logic but would mismatch the Einstein equation in Chapter 11. Fix the sign against Jacobson's first law for ordinary matter.

5. **Why compact support.** Suppose $T_{\mu\nu}$ is a plane wave, not compactly supported. Where does the proof of Lemma 7.2 fail, and how does one repair it by using wave packets?

6. **vDVZ.** Look up the linearized deflection angle for Fierz–Pauli gravity with mass m_T and take $m_T \to 0$. Compare with the GR deflection. What numerical factor appears?

7. **Conformal leftover.** Linearize a Brans–Dicke action about Minkowski spacetime and identify the kinetic mixing of ψ with $T_{\mu\nu}$. What bound on m_ψ follows from requiring the mixing to be negligible inside the Solar System?

7.11 Bridge to Chapter 8

The infrared tensor sector is now linearized general relativity, contingent on Theorem 7.3 and Assumption 7.5. The prefactor of that action is still a symbol, G_{eff}. Chapter 8 records a conditional truncated heat-kernel identity; the physical coefficient requires the specified sector calculation of Problem F. The coefficient is not evaluated here:

$$C_R^L[\{\mathcal{O}_s, w_s, \Lambda_{\mathrm{UV}\,s}, \mu_s\}, \mathcal{S}] \quad \text{is not yet matched to } G_{\text{eff}}. \tag{7.16}$$

The same expansion produces a quartic term $c_0\Lambda_{\mathrm{UV}}^4$, which Chapter 8 will isolate and Chapter 12 will be required to tame.

Chapter 8

Heat Kernel and the Coefficient Map to G_{eff}

Learning outcomes. After this chapter the reader should be able to:

1. write the one-loop effective action of condensate fluctuations as a proper-time integral of the heat kernel;

2. distinguish the universal R term in a_1 from curvature-linear a_2 terms generated by dimensionful endomorphisms, and identify the coefficient of the cosmological term;

3. state the conditional, truncated heat-kernel identity and identify the operator, determinant-weight, threshold, and scheme data needed to evaluate the physical R-coefficient;

4. state what is computed, what is scheme-dependent, and what is left for Chapter 12;

5. enter Chapter 9 knowing why a second-order heat kernel is not a license for fourth-order kinetic terms.

Theorem 7.3 gave a massless tensor kinetic operator. This chapter records the conditional heat-kernel identity that can contribute to its prefactor on the *emergent* background; the physical coefficient requires the sector calculation of Problem F.

8.1 One-loop effective action on the emergent geometry

For the sectors actually included in a gauge-fixed calculation, let $\mathcal{S}_{\text{sec}}$ be their index set, w_s their determinant weights relative to a real bosonic determinant, and $\mathcal{O}_s$ their

operators. Define

$$W_E^{(1)} = \frac{1}{2} \sum_{s \in \mathcal{S}_{\text{sec}}} w_s \operatorname{Tr}_{V_s} \ln \mathcal{O}_s, \qquad \mathcal{N}_0 := \sum_{s \in \mathcal{S}_{\text{sec}}} w_s \operatorname{tr}_{V_s} \mathbb{I}_s. \tag{8.1}$$

The formulae below use a truncated, unresummed small-s heat-kernel expansion. A common pair of proper-time limits applies only to sectors explicitly assigned those limits; otherwise limits and thresholds remain sector-dependent. If E_s contains a mass term, resumming its exponential yields sector-dependent threshold functions rather than the polynomial cutoff terms below.

After condensation, split

$$\Phi = \langle \Phi \rangle + \eta, \tag{8.2}$$

where η contains every algebraic block of Chapter 5. Integrate η at one loop on a slowly varying emergent background $g_{\mu\nu}$, using Presentation B and Assumption 8.4. The dictionary is needed to attribute this calculation to Presentation A, not to formulate it in Presentation B. For each specified sector the fluctuation operator is of Laplace type,

$$\mathcal{O}_\eta = -g^{\mu\nu} D_\mu D_\nu + E[\langle \Phi \rangle, g], \tag{8.3}$$

with a bundle endomorphism E that includes the second derivative of V and curvature couplings generated by the kinetic term.

The one-loop effective action is

$$W_E^{(1)} = \frac{1}{2} \sum_{s \in \mathcal{S}_{\text{sec}}} w_s \operatorname{Tr}_{V_s} \ln \mathcal{O}_s = -\frac{1}{2} \sum_{s \in \mathcal{S}_{\text{sec}}} w_s \int_{s_{\min,s}}^{s_{\max,s}} \frac{\mathrm{d}s}{s} \operatorname{Tr}_{V_s} e^{-s\mathcal{O}_s}, \tag{8.4}$$

after Wick rotation to a Euclidean section. For a sector explicitly assigned common limits, these are

$$s_{\min} = \Lambda_{\text{UV}}^{-2}, \qquad s_{\max} = \mu^{-2}, \tag{8.5}$$

where Λ_{UV} is the kinematic cutoff (3.3). Other sectors may have distinct $s_{\min,s} = \Lambda_{\text{UV}} {}_s^{-2}$ and $s_{\max,s} = \mu_s^{-2}$; no common infrared scale is presumed.

Assumption 8.1 (Laplace-type infrared operator)**.** After Theorem 7.3 and Assumption 7.5, the tensor block of $\mathcal{O}_\eta$ is second order and of Laplace type. Higher-derivative pieces are treated as perturbations and reserved for Chapter 9.

8.2 Seeley–DeWitt expansion, used not re-derived

The heat-kernel diagonal on a four-dimensional Riemannian section admits

$$\operatorname{tr}_V \langle x | e^{-s\mathcal{O}_\eta} | x \rangle \sim \frac{1}{(4\pi s)^2} \sum_{k=0}^{\infty} s^k a_k(x), \tag{8.6}$$

where tr_V is the fibre trace on the bundle in which η takes values. The first two scalar densities are

$$\begin{aligned} a_0 &= \mathrm{tr}_V \mathbb{I}, \\ a_1 &= \mathrm{tr}_V \Big(\tfrac{1}{6} R - E\Big). \end{aligned} \tag{8.7}$$

The coefficient a_2 contains $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$, $R_{\mu\nu}R^{\mu\nu}$, R^2, E^2, RE, and derivative terms. It multiplies a logarithm and is needed in Chapter 9. Its curvature-linear terms can also contribute to the Einstein–Hilbert coefficient when dimensionful endomorphism insertions are present, so they cannot be omitted from the full Problem F calculation.

Insert the sector heat-kernel expansions into (8.4) and integrate term by term, using each sector's assigned limits:

$$\begin{aligned} W_E^{(1)} \supset \sum_{s\in\mathcal{S}_{\mathrm{sec}}} w_s \int \mathrm{d}^4x\,\sqrt{g}\,\Bigg[&-\frac{a_{0,s}}{64\pi^2}\big({\Lambda_{\mathrm{UV}}}_s^4 - \mu_s^4\big) - \frac{a_{1,s}}{32\pi^2}\big({\Lambda_{\mathrm{UV}}}_s^2 - \mu_s^2\big) \\ &- \frac{a_{2,s}}{32\pi^2}\ln\Big(\frac{{\Lambda_{\mathrm{UV}}}_s^2}{\mu_s^2}\Big) + \cdots \Bigg]. \end{aligned} \tag{8.8}$$

The a_k are the traced coefficients on V_s. These are termwise integrals of the truncated, unresummed expansion, not exact massive-sector threshold integrals. Wick rotation and the Euclidean-to-Lorentzian sign are applied only after the weighted sector sum is specified.

8.3 Evaluating a_1 on the condensate

The universal a_1 term and its endomorphism contribution can produce curvature terms. In addition, a dimensionful part of the endomorphism can generate a curvature-linear term in a_2.

Universal piece. The $\frac{1}{6}R_E\,\mathrm{tr}_{V_s}\mathbb{I}_s$ contribution is present for every specified Laplace-type sector. Its universal truncated a_1 term alone supplies

$$W_E^{(1)} \supset -\sum_{s\in\mathcal{S}_{\mathrm{sec}}} \frac{w_s\,\mathrm{tr}_{V_s}\mathbb{I}_s}{192\pi^2}\big({\Lambda_{\mathrm{UV}}}_s^2 - \mu_s^2\big)\int \mathrm{d}^4x\,\sqrt{g}\,R_E \tag{8.9}$$

This omits the endomorphism contribution and does not fix the total Einstein–Hilbert coefficient. Resummed masses instead produce sector-dependent threshold functions.

Endomorphism and threshold terms. For a specified sector, write $E_s = E_{0,s} + Q_s R_E + \cdots$. Its contribution to the coefficient of R_E in a_1 is $\mathrm{tr}_{V_s}(\frac{1}{6}\mathbb{I}_s - Q_s)$, not a number fixed by the fibre dimension alone. Massive sectors require the threshold treatment specified for their operators.

Theorem 8.2 (Coefficient map)**.** *For specified gauge-fixed operators, determinant weights, per-sector proper-time limits, and finite-counterterm scheme $\mathcal{S}$, the truncated, unresummed expansion gives the Euclidean curvature coefficient*

$$C_R^E = -\sum_{s\in\mathcal{S}_{\text{sec}}} \frac{w_s \mathcal{A}_{1,R,s}}{32\pi^2}(\Lambda_{\text{UV}}{}_s^2 - \mu_s^2) - \sum_{s\in\mathcal{S}_{\text{sec}}} \frac{w_s \mathcal{A}_{2,R,s}}{32\pi^2} \ln\Big(\frac{\Lambda_{\text{UV}}{}_s^2}{\mu_s^2}\Big) + c_R(\mathcal{S}), \tag{8.10}$$

where $\mathcal{A}_{k,R,s}$ is the coefficient of R_E in the traced $a_{k,s}$, $\mathcal{A}_{1,R,s}$ includes the specified endomorphism, and $c_R(\mathcal{S})$ is the local curvature counterterm in that scheme. Resummed mass terms require sector-dependent threshold functions instead. After continuation, define the Lorentzian matching by

$$W_L \supset \frac{1}{16\pi G_{\text{eff}}} \int \mathrm{d}^4x\, \sqrt{-g}\, R_L, \tag{8.11}$$

but this identity supplies no physical value or sign until the sector data, thresholds, counterterms, and continuation are specified and evaluated.

Remark 8.3 (Conditional parameterization)*.* The working parameterization may be retained as a conditional notation, not as a computed result:

$$\frac{1}{G_{\text{eff}}} = A\,\mathcal{N}_0\,\Lambda_{\text{UV}}^2 + B\,v^2 \ln\Big(\frac{\Lambda_{\text{UV}}}{\mu}\Big) + C\,\mu^2, \tag{8.12}$$

where $\mathcal{N}_0$ is defined in (8.1). No factor of $1/6$ is absorbed into it. The coefficients A, B, C are undetermined; distinct sector thresholds need not share μ. Problem F remains open.

8.4 The quartic term

Under the continuation and Lorentzian vacuum-action convention of Appendix A, the same truncated, unresummed expansion produces

$$W_L \supset -\rho_\Lambda \int \mathrm{d}^4x\, \sqrt{-g}, \qquad \rho_\Lambda = -\sum_{s\in\mathcal{S}_{\text{sec}}} \frac{w_s\, \mathrm{tr}_{V_s}\, \mathbb{I}_s}{64\pi^2}(\Lambda_{\text{UV}}{}_s^4 - \mu_s^4) + \cdots . \tag{8.13}$$

This sum uses the same sectors and determinant weights as Appendix A. It reduces to $-\mathcal{N}_0(\Lambda_{\text{UV}}^4 - \mu^4)/(64\pi^2)$ only when all included sectors are assigned the same limits. Identifying this with a cosmological term,

$$\frac{\Lambda_{\text{naive}}}{8\pi G_{\text{eff}}} = \rho_\Lambda, \tag{8.14}$$

would give $\Lambda_{\rm naive}$ of order $\Lambda_{\rm UV}^2$ only conditionally: the weighted quartic contribution must be nonzero and uncancelled, and a physical relation between $G_{\rm eff}$ and $\Lambda_{\rm UV}$ must be specified. Finite local vacuum terms and counterterms must also be included in that assessment. Without these inputs, this equation supplies no scale estimate for $\Lambda_{\rm naive}$.

This chapter does not cancel it. It isolates it. Two remarks prevent the isolation from being mistaken for a solution.

1. A bare counterterm $\int \sqrt{-g}\,\Lambda_{\rm ct}$ can subtract ρ_Λ at one scale. It cannot explain why the remainder after subtraction is 10^{-120} in Planck units. Subtraction without a principle is fine tuning.

2. Chapter 12 assigns the principle to a boundary-entropy functional (HBEV). Until that chapter, ρ_Λ is a debt.

8.5 Which background the heat kernel sees

Assumption 3.1 declared $g^{(0)}$ non-dynamical. The heat kernel of this chapter is nevertheless written on $g_{\mu\nu}$. This is an assumption of the chapter, not a consequence of the action:

Assumption 8.4 (Fluctuation operator on the emergent background)**.** The fluctuation operator $\mathcal{O}_\eta$ of (8.3) is built from the emergent metric $g_{\mu\nu}$ of Chapter 4, with the endomorphism and curvature coupling stated for each sector, and not from the scaffold $g^{(0)}_{\mu\nu}$.

Remark 8.5 (Operator on the scaffold and on the emergent metric)*.* The displayed action (3.6) does not supply the operator of Assumption 8.4. With the operator built from $g^{(0)}$ and the internal connection held at a background value, no term of the one-loop action that grows with $\Lambda_{\rm UV}$ ($\Lambda_{\rm UV}^4$, $\Lambda_{\rm UV}^2$, $\ln\Lambda_{\rm UV}$) contains a derivative of the condensate, so the $\Lambda_{\rm UV}^2$ curvature term of (8.8) multiplies the curvature of $g^{(0)}$. For $\bar\Phi = (v+\sigma)\mathbb{I}$ on a flat scaffold, where $g = (1+\sigma/v)^2\eta$, the $\Lambda_{\rm UV}^2(\partial\sigma)^2$ coefficient is zero with the scaffold operator and is the induced term of (17.4) with the operator built on g (Appendix H, Section H.6). Replacing $g^{(0)}$ by g therefore supplies this coefficient; it does not correct a scaffold value. The expansion (8.8) and Theorem 8.2 are stated for operators built on g, that is, under this assumption. Discharging Assumption 8.4 is part of Problem F.

At higher order the replacement produces the backreaction tensor $\mathcal{G}_{\mu\nu}$ of (4.12). That tensor is not needed to define the conditional coefficient. It is needed to define the extra stress of Chapters 11–14.

8.6 Numerical size

The physical size of G_{eff} and the relation between Λ_{UV} and the Planck scale are undetermined until Problem F specifies and evaluates the sector operators, determinant weights, thresholds, continuation, and counterterm scheme.

$$G_{\text{eff}} \quad \text{is not numerically determined by the present data.} \tag{8.15}$$

8.7 What has been computed

Object	Status	Where
truncated R-coefficient identity	conditional on specified sectors and scheme	Thm. 8.2
physical G_{eff} and numerical A, B, C	not computed; Problem F open	Ch. 17
Lorentzian EH sign	conditional on completed sector sum and continuation	Sec. 8.2
$\rho_\Lambda \sim \Lambda_{\text{UV}}^4$	conditional truncated sector sum	Sec. 8.4
pure curvature-square terms in a_2	isolated; further use deferred	Ch. 9
exact two-loop G_{eff}	not computed	Ch. 17

8.8 Exercises

1. **Proper-time integrals.** For one specified sector with assigned limits, perform the term-by-term integrals producing (8.8) from (8.6) and (8.4), treating a_k as s-independent. State why this is not a resummed massive threshold calculation.

2. **Dimensions.** Confirm that $a_0\Lambda_{\text{UV}}^4$, $a_1\Lambda_{\text{UV}}^2$, and $a_2 \ln(\Lambda_{\text{UV}}/\mu)$ all have mass dimension four.

3. **Conformal value of ξ.** For a scalar with non-minimal term $\xi R\phi^2$, show that $\xi = 1/6$ cancels its curvature contribution to a_1. Explain why this does not determine the full locked-block result.

4. **Mass thresholds.** Compare the truncated polynomial terms with the integrals that retain $\exp(-sm^2)$. Explain why distinct masses can require distinct threshold functions and limits.

5. **Conditional curvature coefficient.** Given sector operators, weights, limits, and counterterms, derive the Euclidean R-coefficient from supplied a_1, a_2, and identify the continuation data needed for Lorentzian matching.

6. **Sector weights.** For a supplied gauge-fixed sector list, determine the determinant weights and compute $\mathcal{N}_0$. Explain why this alone does not determine the curvature coefficient.

7. **Scaffold error.** For $\bar{\Phi} = (v + \sigma)\mathbb{I}$ on a flat scaffold, compare the $\Lambda_{\mathrm{UV}}^2(\partial\sigma)^2$ coefficient of the one-loop action computed with the scaffold operator and with the operator of Assumption 8.4.

8.9 Bridge to Chapter 9

Once the sector calculation is completed, the action through two derivatives has the form

$$W[g] = \int \mathrm{d}^4x\,\sqrt{-g}\Big(-\rho_\Lambda + \frac{1}{16\pi G_{\mathrm{eff}}}R + \alpha\, a_2[g] + \cdots\Big). \tag{8.16}$$

The a_2 density contains $R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}$ and $R_{\mu\nu}R^{\mu\nu}$. If those operators are kept as propagating fourth-order kinetic terms for $h_{\mu\nu}$, Ostrogradsky ghosts appear. Chapter 9 classifies the dangerous operators, states the conditions under which they may be treated as non-propagating corrections, and introduces the Einsteinian cubic completion used later to regularize curvature singularities without reopening the ghost problem.

Chapter 9

Ultraviolet Operators, Ghosts, and a Cubic Completion

Learning outcomes. After this chapter the reader should be able to:

1. read the a_2 density as a list of four-derivative operators and assign each a role (correction versus propagator);

2. state Ostrogradsky's theorem in the form used for gravitational effective field theory;

3. explain why quadratic gravity of Stelle type is not an acceptable infrared limit of NAMCG;

4. write the Einsteinian cubic invariant and the property that makes it safer than a generic R^3 term;

5. enter Chapter 10 knowing that the ultraviolet problem of the metric effective action is under a named control, not solved in the sense of a non-perturbative continuum limit.

Chapter 8 produced

$$W[g] = \int \mathrm{d}^4x\,\sqrt{-g}\Big(-\rho_\Lambda + \frac{R}{16\pi G_{\mathrm{eff}}} + \alpha\, a_2[g] + \cdots\Big). \tag{9.1}$$

The first two terms are the cosmological debt and the Einstein–Hilbert term. This chapter treats the third.

9.1 What a_2 contains

On a Laplace-type bundle the second Seeley–DeWitt coefficient is a linear combination of

$$R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma}, \quad R_{\mu\nu}R^{\mu\nu}, \quad R^2, \quad E^2, \quad RE, \quad D^2R, \quad D^2E, \tag{9.2}$$

together with bundle-curvature terms $F_{\mu\nu}F^{\mu\nu}$. After the endomorphism E is evaluated on the condensate and total derivatives are discarded, the metric operators that remain in four dimensions may be written

$$W_4[g] = \int \mathrm{d}^4x\,\sqrt{-g}\big(c_W C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} + c_R R^2 + c_{\mathrm{GB}}\,E_{\mathrm{GB}}\big), \tag{9.3}$$

where $C_{\mu\nu\rho\sigma}$ is the Weyl tensor and

$$E_{\mathrm{GB}} = R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - 4R_{\mu\nu}R^{\mu\nu} + R^2 \tag{9.4}$$

is the Gauss–Bonnet density, topological in four dimensions at the level of the bulk equations of motion.

For specified sectors, their weights, and a stated scheme, the coefficients $c_W, c_R, c_{\mathrm{GB}}$ receive logarithmic heat-kernel contributions. Their physical values and the matching to G_{eff} remain conditional on the operator, threshold, and finite-local-term calculation of Problem F. The existence of Weyl-squared and R^2 structures is independent of those coefficient values.

9.2 Ostrogradsky's theorem, used as a veto

Theorem 9.1 (Ostrogradsky, gravitational form)**.** *Let a field theory possess a Lagrangian that depends on second (or higher) time derivatives of a dynamical variable, not removable by adding a total derivative or by imposing a degeneracy condition that reduces the order. Then the Hamiltonian of that variable is unbounded below. In a quadratic gravitational action about Minkowski spacetime, the unbounded direction is a ghost pole in the propagator of $h_{\mu\nu}$.*

The theorem is the reason one may not promote (9.3) to a fundamental action for a propagating metric and call the result a completion of NAMCG.

Stelle quadratic gravity. The action

$$S_{\mathrm{Stelle}} = \int \mathrm{d}^4x\,\sqrt{-g}\Big(\frac{R}{16\pi G} - \frac{\alpha}{2}C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} + \frac{\beta}{2}R^2\Big) \tag{9.5}$$

propagates, in addition to the massless graviton, a massive spin-2 ghost and a massive scalar. The ghost mass is of order $1/\sqrt{\alpha G}$. Its relation to Λ_{UV} cannot be inferred

without the physical G_{eff} and the specified value of α, which remain part of Problem F. Whether such a pole lies within the regime of the metric effective theory is therefore undetermined here. These statements take $\alpha > 0$ and $\beta > 0$. For $\alpha < 0$ the extra spin-2 pole is tachyonic, with residue still opposite to that of the graviton; Section H.11 records that a truncated induced Weyl term has this sign, under the curvature-continuation convention stated there, when G_{eff} and the weighted Weyl trace are positive.

Effective-field-theory reading. The same operators are harmless if they are used only as vertices in a derivative expansion of matrix elements with external momenta $p \ll \Lambda_{\text{UV}}$, and never as additional poles. That is the reading adopted here:

Assumption 9.2 (Four-derivative terms are EFT vertices)**.** The operators in (9.3) correct graviton scattering and static potentials at order $(p/\Lambda_{\text{UV}})^2$. They are not to be diagonalized into extra propagating modes of $h_{\mu\nu}$.

Assumption 9.2 is available only because the metric is composite. The microscopic path integral is over Φ, not over an independent $g_{\mu\nu}$ with action (9.5). The poles of Stelle gravity are an artefact of treating $W[g]$ as a fundamental metric action beyond its radius of convergence.

9.3 What ultraviolet stability does mean

Presentation A is a matrix model. Its ultraviolet definition is the regularized trace over $\mathcal{H}$ with cutoff (3.3). Questions that make sense at that level are:

- boundedness of S_{matrix} from below on Hermitian $\hat{X}^\mu$;
- existence of the condensed saddle $\langle \Phi^I{}_A \rangle = v\Lambda^A{}_I$;
- absence of tachyons in the blocks that Theorem 7.3 and Assumption 7.5 did not already constrain.

Questions that do *not* make sense until a continuum limit in Λ_{UV} is constructed are:

- renormalizability of $W[g]$ as a fundamental quantum field theory of $g_{\mu\nu}$;
- asymptotic safety of that metric theory;
- unitarity of S-matrix elements among Stelle poles.

NAMCG, as written in this book, claims the first list and refuses the second. The metric effective action is an infrared object with a cutoff. Ultraviolet stability means stability of the matrix saddle, not renormalizability of quadratic gravity.

A sufficient condition for the matrix action (3.4) to be bounded below on Hermitian matrices is that the quartic commutator term dominates with the sign written there,

$$-\mathrm{Tr}\big([\hat{X}^{\mu},\hat{X}^{\nu}][\hat{X}_{\mu},\hat{X}_{\nu}]\big)\geq 0 \tag{9.6}$$

in the Euclideanized internal-time convention of IKKT-type models, together with a potential V that is bounded below, as (3.7) is. Lorentzian signature complicates the bound because $\hat{X}^{0}$ enters with the opposite sign in some reductions. That complication is inherited from the matrix-model literature and is not solved here. It is recorded in Chapter 17.

9.4 Cubic operators and Einsteinian cubic gravity

Singularities of the Einstein equation are not cured by the four-derivative vertices of Assumption 9.2. Those vertices are small when curvature is small, and they are untrustworthy when curvature reaches $\Lambda_{\mathrm{UV}}^{2}$. A different higher-curvature object is needed if compact-object cores are to be regularized inside the derivative expansion.

Einsteinian cubic gravity (ECG) uses a cubic curvature invariant. The invariant used here is

$$\mathcal{P}=12R_{a}{}^{c}{}_{b}{}^{d}R_{c}{}^{e}{}_{d}{}^{f}R_{e}{}^{a}{}_{f}{}^{b}+R_{ab}{}^{cd}R_{cd}{}^{ef}R_{ef}{}^{ab}-12R_{abcd}R^{ac}R^{bd}+8R_{a}{}^{b}R_{b}{}^{c}R_{c}{}^{a}, \tag{9.7}$$

with the Riemann convention for which $R>0$ on spheres. The action used later in the book is

$$S_{\mathrm{ECG}}[g]=\int \mathrm{d}^{4}x\,\sqrt{-g}\Big(\frac{R-2\Lambda_{\mathrm{eff}}}{16\pi G_{\mathrm{eff}}}+\beta_{\mathrm{ECG}}\,\mathcal{P}\Big). \tag{9.8}$$

Two distinct properties matter.

1. For the reduced action in Section H.10, the Euler–Lagrange equations have been checked symbolically: $N=1$ is a consistent vacuum branch and the lapse equation yields the displayed first integral, including Λ_{eff}. This check uses the static spherical reduction; its relation to the unrestricted field equations relies on a symmetric-criticality justification not established here. The reduced coupling is $\alpha=-16\pi G_{\mathrm{eff}}\beta_{\mathrm{ECG}}$. The regular-core profile

$$f(r)=1-\frac{2Mr^{2}}{r^{3}+r_{0}^{3}}-\frac{\Lambda_{\mathrm{eff}}}{3}r^{2} \tag{9.9}$$

used in Chapter 13 is a phenomenological regular-core model with $r_{0}^{3}=2\beta_{\mathrm{ECG}}M$

a model parametrization, not derived from (9.8). For $\Lambda_{\text{eff}} = 0$ and $Mr_0 \neq 0$, the rational profile has nonzero residual in the displayed first integral.

2. Assume $\Lambda_{\text{eff}} = 0$, or a specified cancellation, so flat space solves the full equations. Since $\mathcal{P}$ is cubic in curvature, it contributes no term quadratic in $h_{\mu\nu}$ there; for the Einstein–Hilbert-plus-ECG action the quadratic spectrum is therefore the Einstein one. Any separately retained curvature-squared operators are treated under Assumption 9.2; no no-extra-poles claim is made for the combined theory including such terms.

3. Bueno and Cano [1] establish an Einstein-like linearized equation on maximally symmetric backgrounds. The generic noncritical branch has the massless spin-2 spectrum; its effective kinetic normalization and sign require separate checks and are not asserted here. This cited result does not exclude nonlinear, sourced, or non-maximally-symmetric pseudo-modes. This is a cited linear-spectrum property, not a claim of general nonlinear or non-vacuum ghost freedom.

Assumption 9.3 (Cubic completion)**.** As a model assumption, through the curvature scales relevant to compact-object cores the metric effective action may be truncated at (9.8). The coupling and the core scale r_0 have not been derived or matched to the matrix theory.

Assumption 9.3 is stronger than Assumption 9.2. It promotes one cubic combination from a vertex to a term that is solved non-perturbatively in the static spherical sector. The promotion is a model assumption, not a consequence of the heat-kernel calculation.

9.5 Torsion and higher connections

A chosen non-degenerate frame (Repair II, (4.7), or an assumed coframe) need not have vanishing torsion. A torsion tensor

$$T^A{}_{\mu\nu} = \partial_\mu e^A_\nu - \partial_\nu e^A_\mu + \omega_\mu{}^A{}_B e^B_\nu - \omega_\nu{}^A{}_B e^B_\mu \tag{9.10}$$

is an additional field if ω is independent of e. In NAMCG the independent connection and the condensate term's broken-direction combination must be distinguished. The condensate term gives a quadratic Higgs-type contribution only in that broken direction; whether torsion vanishes, is massive, or supplies additional field content is not determined here (Problem D). Torsion-free behavior follows only under the conditional Palatini–Holst hypotheses of Proposition 10.5 in Chapter 10. The a_1 coefficient for a Laplace-type operator is $\operatorname{tr}(R(g)/6 - E)$: its explicit universal curvature term is the Levi-Civita scalar $R(g)$. Since E is unspecified, no conclusion about additional B-dependence follows, and a Palatini $R(B)$ term is not established.

9.6 A checklist before Lorentz locking

The metric effective theory, as it now stands, is:

Ingredient	Status
massless TT tensor	Thm. 7.3
conditional truncated R-coefficient	Thm. 8.2; physical match open (Problem F)
quartic vacuum energy	isolated, not tamed
C^2 and R^2	EFT vertices, Assp. 9.2
cubic core regularity	ECG model, Assp. 9.3
torsion	undetermined (Problem D); torsion-free only under the conditional Palatini–Holst hypotheses of Proposition 10.5
local Lorentz invariance as a gauge symmetry of Φ	not yet constructed

The local Lorentz gauge symmetry is assumed in Chapter 10. The connection and torsion dynamics remain open; the identification of ω_{AB} as a Goldstone is conditional on that assumption.

9.7 Exercises

1. **Gauss–Bonnet.** Show that E_{GB} in four dimensions is a total derivative plus a topological density. Why, then, can c_{GB} still matter on manifolds with boundary?

2. **Stelle poles.** Linearize (9.5) about Minkowski spacetime in harmonic gauge. Locate the massive spin-2 pole and determine the sign of its residue relative to the massless graviton.

3. **EFT versus poles.** A tree-level graviton-graviton amplitude receives a correction $\sim c_W s^2/\Lambda_{\mathrm{UV}}^2$ from (9.3). At what centre-of-mass energy does this correction become unreliable as an EFT vertex, and why is that energy the same scale at which Stelle poles would have appeared?

4. **Bounded matrix action.** For two Hermitian matrices X, Y show that $-\mathrm{Tr}([X,Y]^2) \geq 0$. Identify the step that fails, or requires a Euclidean rotation, when one matrix is X^0 with Lorentzian signature.

5. **Spherical reduction.** Starting from the reduced action and invariant specified in Section H.10, derive the reduced Euler–Lagrange equations, verify the $N = 1$

branch and first integral, and recover the leading tail correction. State the boundary conditions and the symmetric-reduction assumption used.

6. **Torsion mass.** If torsion is massive (an assumption, not a result; Problem D), estimate its mass using v as the only available scale. Compare it with m_ψ from Assumption 7.5.

9.8 Bridge to Chapter 10

The infrared metric theory is Einstein gravity plus controlled higher-curvature corrections and an untamed ρ_Λ. Chapter 10 assumes local $\mathrm{SO}(3,1)$ gauge symmetry and identifies the six Goldstone orbit variables ω_{AB} as removable in unitary gauge. The provisional sector bookkeeping of Section 5.7 records that removal without claiming a full-spectrum count. The frame, taken from Repair II, (4.7) (or from an assumed non-degenerate coframe), transforms as a vierbein under the diagonal local Lorentz group under Theorem 10.3; that theorem does not determine the diagonal connection's dynamics or torsion. Definition 4.1 is not used to construct it (Problem C).

Chapter 10

Local Lorentz Locking and the Fate of the Goldstones

Learning outcomes. After this chapter the reader should be able to:

1. write the simultaneous action of internal gauge transformations and local Lorentz transformations on $\Phi^I{}_A$;

2. identify the diagonal subgroup preserved by the polar vacuum and the six broken generators;

3. exhibit the Stueckelberg replacement that removes the Goldstone orbit variables ω_{AB} in unitary gauge;

4. state the provisional sector bookkeeping through the Goldstone and vector lines;

5. enter Chapter 11 with a composite frame that transforms as a genuine vierbein, so that $\nabla^\mu T_{\mu\nu}$ is well defined.

Since Chapter 5 the antisymmetric piece ω_{AB} has been called a local-Lorentz Goldstone and postponed. This chapter assumes the local gauge symmetry and shows that the six Goldstone orbit variables ω_{AB} are removable in unitary gauge. It does not determine the full propagating spectrum.

10.1 The two groups that act on Φ

The condensate field of Presentation B transforms, by (3.10), as

$$\Phi^I{}_A(x) \;\longmapsto\; \Omega^I{}_J(x)\,\Phi^J{}_B(x)\,\Lambda^B{}_A(x). \tag{10.1}$$

Here $\Omega(x)$ takes values in the internal gauge group G_{int} and $\Lambda(x)$ takes values in $\mathrm{SO}(3,1)$, or in its double cover if spinors are later coupled. Infinitesimally,

$$\delta\Phi^I{}_A = \alpha^I{}_J\Phi^J{}_A + \Phi^I{}_B\,\beta^B{}_A, \tag{10.2}$$

with α in the Lie algebra of G_{int} and $\beta_{AB} = -\beta_{BA}$ in $\mathfrak{so}(3,1)$.

These are independent transformations at the level of field content. Nothing in Chapter 3 forced $\Lambda(x)$ to be a gauge symmetry rather than a global classification of the index A. Treating it as local is a choice, recorded as such.

Assumption 10.1 (Local Lorentz gauge symmetry)**.** The microscopic action (3.6) is invariant under local $\Lambda(x) \in \mathrm{SO}(3,1)$ once a spin connection $\omega_\mu{}^A{}_B$ is introduced through the covariant derivative on the tangent index,

$$\mathcal{D}_\mu\Phi^I{}_A = \partial_\mu\Phi^I{}_A - i(A_\mu)^I{}_J\Phi^J{}_A + \omega_\mu{}^B{}_A\Phi^I{}_B. \tag{10.3}$$

The connection $\omega_\mu{}^{AB}$ is, at this stage, an independent gauge field.

Assumption 10.1 is the step that makes A a genuine tangent index rather than an internal flavour that happens to have four values. Combined with Assumption 4.6, it is the origin of local Lorentz invariance in NAMCG. It is not derived from Presentation A.

10.2 The vacuum and the broken generators

The polar vacuum of Chapter 4, in the Minkowski frame,

$$\langle\Phi^I{}_A\rangle = v\,\delta^I{}_A, \tag{10.4}$$

is invariant only under the diagonal combination

$$\alpha^I{}_J = -\,v^{-1}\delta^I{}_B\,\beta^B{}_C\,\delta^C{}_J\Big|_{\text{locked } 4\times 4 \text{ block}}, \tag{10.5}$$

that is, under simultaneous internal and Lorentz rotations that rotate the locked axes together. If G_{int} contains $\mathrm{SO}(3,1)$ (or a group large enough to have an $\mathfrak{so}(3,1)$ subalgebra acting on the locked indices), six generators are unbroken and six are broken.

Proposition 10.2 (Goldstone count)**.** *About the vacuum* (10.4)*, with Assumption 10.1 in force and with the locked block of dimension four, there are six broken generators. The corresponding Goldstone fields are the antisymmetric components ω_{AB} of* (4.15)*.*

Proof. The vacuum stabilizer is the diagonal $\mathrm{SO}(3,1)_{\text{diag}}$. The coset $\mathrm{SO}(3,1)_{\text{Lorentz}} \times \mathrm{SO}(3,1)_{\text{int}}/\mathrm{SO}(3,1)_{\text{diag}}$ has dimension $6 + 6 - 6 = 6$. An infinitesimal broken trans-

formation with parameter β_{AB} shifts ${\Phi^I}_A$ by $v{\beta^I}_A$, which is antisymmetric in (I, A) after lowering with η. That shift is the fluctuation ω_{AB}. □

If G_{int} is larger than $\mathrm{SO}(3,1)$, additional Goldstones live in the unlocked block. They belong to Assumption 5.1 and are taken heavy.

10.3 Eating ω_{AB}

A local broken symmetry with an independent connection is a Higgs mechanism. Introduce the Stueckelberg field $\Lambda(x) \in \mathrm{SO}(3,1)$ already present in the polar decomposition (4.4), and replace the spin connection by the dressed connection

$$\omega_\mu \;\longmapsto\; \Lambda^{-1}\omega_\mu\Lambda + \Lambda^{-1}\partial_\mu\Lambda. \tag{10.6}$$

Choose the unitary gauge

$$\Lambda(x) = \mathbb{I}, \tag{10.7}$$

which is admissible wherever the polar decomposition is smooth (that is, wherever Assumption 4.4 holds and no condensate defect forces $\det e = 0$). In that gauge the six orbit variables ω_{AB} are removed. Identify the locked indices $I \leftrightarrow A$ through $\langle {\Phi^I}_A\rangle = v{\delta^I}_A$, and write $({A_\mu})^I{}_A = ({A_\mu})^I{}_J{\delta^J}_A$ on the locked block. Then (10.3) gives

$$\mathcal{D}_\mu\langle {\Phi^I}_A\rangle = v\big[({\omega_\mu})^I{}_A - i({A_\mu})^I{}_A\big].$$

The factor i follows the Hermitian-generator convention in (10.3); with real antisymmetric generators the corresponding combination is $\omega - A$. Define the broken-direction combination $\Xi_\mu := \omega_\mu - iA_\mu|_{\text{locked}}$ in this convention. Write B_μ for the connection along the unbroken diagonal stabilizer direction; no complementary connection combination is canonical until kinetic terms specify a metric on the (ω, A) field space. The stabilizer direction lies in the kernel of the condensate covariant derivative. The invariant contraction on the locked block depends on the action and generator conventions, so the quadratic term is written as

$$\mathcal{L}^{(2)}_{\text{cond}} \supset v^2 Q_{\text{locked}}(\Xi_\mu), \tag{10.8}$$

where Q_{locked} denotes that contraction; it is not assumed to be a real Lorentz-vector square. This is a quadratic Higgs-type contribution, not a physical mass assertion. Its pole and sign require gauge-field kinetic terms and canonical normalization, which are unspecified. Torsion and the dynamics of the diagonal connection remain open (Problem D).

Theorem 10.3 (Lorentz locking)**.** *Under Assumptions 3.5, 10.1, and 4.4, there exists a gauge in which*

1. *the six Goldstone orbit variables ω_{AB} are removable in unitary gauge; this makes no claim about the full propagating spectrum;*

2. *the frame $e^A_\mu = \Phi^A{}_\mu/v$ of Repair II,* (4.7) *(or an independently supplied non-degenerate coframe), transforms only under the diagonal local Lorentz group, as a vierbein;*

3. *the condensate term supplies a quadratic Higgs-type contribution in the broken direction Ξ_μ; the diagonal connection's dynamics are not determined. A torsion-free result follows only under the conditional hypotheses of Proposition 10.5.*

Proof. Proposition 10.2 supplies six Goldstones. The Stueckelberg replacement (10.6) and the gauge (10.7) remove the six orbit variables. In that gauge $\Phi^{IA} = (v + \sigma)\delta^{IA}$ plus tensor and vector blocks, so the Repair II frame $e^A_\mu = \Phi^A{}_\mu/v$ transforms as $e^A_\mu \mapsto \Lambda_{\text{diag}}{}^A{}_B e^B_\mu$. This argument establishes the frame transformation and the quadratic broken-direction term, but does not determine the diagonal connection's spectrum or torsion. □

Remark 10.4 (Defects). Unitary gauge fails where $\Lambda(x)$ cannot be set to the identity continuously—at defects where $\det e = 0$. Those loci are condensate defects, not coordinate singularities of an already existing manifold. They are possible ultraviolet endpoints of the regular cores of Chapter 13, and they are not analyzed further here.

10.4 The internal vector after locking

The internal connection A_μ is distinct from the independent Lorentz connection $\omega_\mu{}^{AB}$. The condensate term mixes them only through Ξ_μ; an orthogonal complement is defined only after kinetic terms provide a metric on the (ω, A) field space. Neither a Higgs mass $m_A \sim g_{\text{YM}} v$ nor a decoupling scale follows without those terms. If a specified internal-vector quadratic operator gaps the four components counted in Section 5.7, they are omitted from the low-energy frame/metric-sector bookkeeping; otherwise their spectrum remains open. If an internal subgroup H is unbroken, its corresponding gauge field is massless (conditional). This identifies the conditional vector row of the provisional sector bookkeeping; it does not establish the dynamics of the diagonal connection (Problem D).

Proposition 10.5 (Conditional torsion-free diagonal connection)**.** *Let $e^A{}_\mu$ be an invertible coframe and $B_\mu{}^{AB}$ an independent diagonal Lorentz connection. Suppose its dependence in the action is either the Palatini term $\epsilon_{ABCD} e^A \wedge e^B \wedge F^{CD}(B)$, or*

that term plus the Holst term $\gamma^{-1} e_A \wedge e_B \wedge F^{AB}(B)$, with finite real $\gamma \neq 0$, $\gamma^2 \neq -1$, or $\gamma = \infty$ (pure Palatini). Assume there are no spin sources and no other torsion-dependent terms, including $F(B)^2$, torsion-squared terms, or matter couplings to B. Then the B equation is the Holst-modified Cartan equation and gives $T^A = 0$; consequently B is the Levi-Civita spin connection of e.

Proof. Since $\delta F^{AB} = D(\delta B^{AB})$, compactly supported variations give, for the Palatini term,

$$\delta_B S_{\mathrm{P}} \simeq -4 \int \epsilon_{ABCD}\, T^A \wedge e^B \wedge \delta B^{CD}.$$

Thus $\epsilon_{ABCD} T^A \wedge e^B = 0$. Expand $T^A = \frac{1}{2} T^A{}_{EF} e^E \wedge e^F$ and contract the three-form equation with ϵ^{EFGM}. In frame components this yields, up to an overall nonzero convention-dependent factor,

$$2(\tau_C \delta^M_D - \tau_D \delta^M_C) + 2T^M{}_{CD} = 0, \qquad \tau_C := T^A{}_{AC}.$$

Tracing $M = D$ gives $4\tau_C = 0$, since $T^D{}_{CD} = -\tau_C$. Hence $\tau_C = 0$, and the uncontracted equation then gives $T^M{}_{CD} = 0$.

With the Holst term, the equation is $(\star + \gamma^{-1}) Y_{CD} = 0$, where $Y_{CD} := T_{[C} \wedge e_{D]}$, and $\star$ is the internal Lorentzian Hodge dual on antisymmetric internal index pairs. The internal dual of Y is proportional to the Palatini Cartan equation. Since $\star^2 = -1$,

$$(\star + \gamma^{-1})(- \star + \gamma^{-1}) = (1 + \gamma^{-2}) \mathbb{I}.$$

The map is invertible for the stated parameters. Therefore $Y = 0$, which implies the Palatini Cartan equation and $T^A = 0$ by the preceding argument. For an invertible coframe, vanishing torsion fixes the metric-compatible spin connection uniquely to the Levi-Civita one. □

Remark 10.6. This proposition constrains B only under its hypotheses. It is not needed to define the Levi-Civita connection ∇ of $g = e^T e$ or to state diffeomorphism conservation $\nabla^\mu T_{\mu\nu} = 0$. NAMCG has not been shown to induce these Palatini or Holst hypotheses. For a Laplace-type operator, $a_1 = \mathrm{tr}(R(g)/6 - E)$; the explicit universal curvature term is the Levi-Civita scalar $R(g)$. Because E is unspecified, no conclusion about additional B-dependence of a_1 follows, and a Palatini $R(B)$ term is not established. If B enters the bundle connection of the actual operator, a_2 may contain a $\mathrm{tr}\, F(B)^2$ term; whether it survives in the weighted trace depends on the unspecified operator representation and weights. Deriving the B-dependence of E and the bundle connection is open.

10.5 Diffeomorphisms

Diffeomorphisms act on the coordinate index of every tensor constructed from $\Phi(x)$. They are not eaten. They are the gauge symmetry of the composite metric, and they are what force $\nabla^\mu T^{\text{total}}_{\mu\nu} = 0$ in Chapter 11.

After Theorem 10.3, an infinitesimal diffeomorphism ξ^μ acts on the vierbein in the standard hybrid way,

$$\delta_\xi e^A_\mu = \xi^\rho \partial_\rho e^A_\mu + \partial_\mu \xi^\rho \, e^A_\rho, \tag{10.9}$$

up to a local Lorentz transformation that can be used to keep a chosen gauge on the frame. The induced action on $g_{\mu\nu}$ is the Lie derivative (5.17). No extra scalar Goldstone of diffeomorphisms appears: diffeomorphisms are gauged, not broken, in the condensed phase.

10.6 Provisional sector bookkeeping (frame/metric sector only)

Component counts in the algebraic split are distinct from on-shell polarization counts. The independent diagonal connection B, torsion modes, and internal vector A_μ are not part of $\delta\Phi_{\mu\nu}$ and are counted separately.

Sector	Count and status
algebraic split	$\psi : 1, \quad F_{\mu\nu} : 6, \quad T_{\mu\nu} : 9; \quad \text{total} : 16$
$F_{\mu\nu}$	Six Goldstone orbit variables (ω_{AB}); removable in unitary gauge (Ch. 10). No full-spectrum claim follows.
ψ	One component; gapped or constrained only under Assumption 7.5.
$T_{\mu\nu}$	Nine algebraic components. The full massive Fierz–Pauli theory has four on-shell divergence constraints and five polarizations; this is not established here for the NAMCG tensor block. Two TT helicities follow only under the separately established massless FP/diffeomorphism-gauge limit (Theorem 7.3); this is not a smooth $m_T \to 0$ limit of all massive polarizations.
Outside $\delta\Phi_{\mu\nu}$	A_μ: four components, omitted from the low-energy frame/metric-sector bookkeeping only if its specified quadratic operator gaps them; otherwise its spectrum is open. B and torsion remain unresolved (Problem D).

No total mode count of the full theory follows from this bookkeeping; the “2” is a metric-sector target, not a full-spectrum result.

10.7 What locking does not do

Theorem 10.3 does not derive Lorentzian signature (Assumption 4.6 remains). It does not derive local Lorentz invariance from Presentation A (Assumption 10.1 remains). It does not gap ψ. It does not compute G_{eff}. It does not conserve $T^{\text{ent}}_{\mu\nu}$.

What it does is make the later covariant derivatives honest. From here on, ∇_μ denotes the Levi-Civita connection of the composite metric $g = e^T e$. This is distinct from the independent Lorentz connection B, whose dynamics remain open. Statements such as $\nabla^\mu T^{\text{total}}_{\mu\nu} = 0$ refer to this Levi-Civita connection.

10.8 Exercises

1. **Stabilizer.** Verify that (10.4) is invariant under the diagonal combination (10.5) and under no other independent pair (α, β) acting on the locked block.

2. **Coset dimension.** Compute $\dim\big(\mathrm{SO}(3,1) \times \mathrm{SO}(3,1)/\mathrm{SO}(3,1)\big)$ and match it to the six components of ω_{AB}.

3. **Stueckelberg.** Starting from (10.3) and (4.4), perform the replacement (10.6) and isolate the quadratic term (10.8) in unitary gauge.

4. **Torsion-free spin connection.** Given e^A_μ, write the explicit formula for the torsion-free spin connection $\omega_\mu{}^{AB}(e)$. This is not a statement that the independent connection B equals $\omega(e)$. Confirm that $\omega(e)$ transforms inhomogeneously under local Lorentz transformations, as a connection must.

5. **Defect.** Construct a hedgehog configuration of $\Lambda(x)$ on a spatial 2-sphere inside which $\det e = 0$ at a point. Why can unitary gauge not be imposed globally on that slice?

6. **Residual Yang–Mills.** Suppose $G_{\text{int}} = \mathrm{SO}(3,1) \times H$ with H unbroken. What is the infrared field content after Theorem 10.3, in addition to the graviton?

7. **Hybrid diffeomorphism.** Derive (10.9) from the action of a coordinate shift on $\Phi(x)$ and the Repair II frame $e^A_\mu = \Phi^A{}_\mu/v$ of (4.7).

10.9 Bridge to Chapter 11

The emergent geometry now has a metric $g = e^T e$ with its Levi-Civita connection ∇, a frame-sector target of two massless TT polarizations, and a G_{eff} whose heat-kernel matching remains conditional on the specified sector calculation in Problem F. The independent Lorentz connection B is a separate field: its dynamics are not determined

here, and it is torsion-free only under the hypotheses of Proposition 10.5. This does not affect the Levi-Civita connection ∇, to which $\nabla^\mu T_{\mu\nu} = 0$ refers. Diffeomorphism invariance is in force. Noether's second theorem therefore requires

$$\nabla^\mu T^{\text{total}}_{\mu\nu} = 0 \tag{10.10}$$

of whatever stress tensor sources the Einstein equation. Chapter 11 defines $T^{\text{total}}_{\mu\nu}$ as the sum of ordinary matter, the tensor-block stress already used in Chapter 7, and the entropic stress $T^{\text{ent}}_{\mu\nu}$ obtained by varying U_{ent}. It then proves that the sum is divergenceless on shell. This conservation statement constrains the total stress when it is derived from the specified diffeomorphism-invariant action. It does not, by itself, derive the phenomenological drag law or show that a reduced drag model is its covariant limit; that connection remains Problem H.9.

Chapter 11

Covariant Conservation of the Total Stress Tensor

Learning outcomes. After this chapter the reader should be able to:

1. define $T^{\text{total}}_{\mu\nu}$ as the sum of matter, tensor-block, and entropic contributions;
2. obtain $T^{\text{ent}}_{\mu\nu}$ by metric variation of U_{ent};
3. prove that diffeomorphism invariance forces $\nabla^\mu T^{\text{total}}_{\mu\nu} = 0$ on shell;
4. reduce that identity to a Newtonian force balance and state a necessary bookkeeping condition for a drag to be a covariant reduction;
5. enter Chapter 12 knowing that vacuum energy is a conservation problem as well as a cosmological-constant problem.

Chapter 10 made ∇_μ the Levi-Civita connection of the composite metric. This chapter asks what that connection is allowed to act on.

11.1 The split

The infrared field equation contemplated from here on is

$$G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = 8\pi G_{\text{eff}}\, T^{\text{total}}_{\mu\nu}, \tag{11.1}$$

with

$$T^{\text{total}}_{\mu\nu} = T^{\text{matter}}_{\mu\nu} + T^{\text{tensor}}_{\mu\nu} + T^{\text{ent}}_{\mu\nu}. \tag{11.2}$$

Matter. $T^{\text{matter}}_{\mu\nu}$ is the metric variation (2.13) of every action for fields that are not the condensate itself: ordinary fluids, electromagnetism, baryons. On shell it satisfies

$\nabla^\mu T^{\text{matter}}_{\mu\nu} = 0$ *only* if those fields do not exchange energy with the condensate. In NAMCG they may. The separate conservation of matter is therefore not assumed.

Tensor block. $T^{\text{tensor}}_{\mu\nu}$ is the metric stress of the quadratic action (7.13). After Theorem 7.3 it is the linearized gravitational-wave stress, of the Isaacson type when averaged over wavelengths short compared with the background curvature. It is kept in (11.2) for bookkeeping and is negligible in the static Newtonian laboratories of Chapters 14–15.

Entropic piece. $T^{\text{ent}}_{\mu\nu}$ is defined by the same variational rule applied to the mutual-information potential of Definition 6.4, promoted to a functional of the metric:

$$T^{\text{ent}}_{\mu\nu} = -\frac{2}{\sqrt{-g}}\frac{\delta S_{\text{ent}}[g]}{\delta g^{\mu\nu}}, \qquad S_{\text{ent}}[g] = -\int \mathrm{d}t\, U_{\text{ent}}[g]. \tag{11.3}$$

The promotion $U_{\text{ent}} \to S_{\text{ent}}[g]$ uses the fact that every entropy $S(A)$ in (6.9) is computed from a reduced state on a region whose causal structure depends on $g_{\mu\nu}$. Varying the metric varies the regions and varies the modular Hamiltonians.

11.2 Noether's second theorem, applied once

Let $S_{\text{IR}}[g, \Psi]$ be the infrared action whose Euler–Lagrange equation for $g_{\mu\nu}$ is (11.1): Einstein–Hilbert plus Λ_{eff}, plus matter, plus S_{ent}, plus the tensor-block action. Diffeomorphism invariance states that for every compactly supported vector field ξ^μ,

$$\delta_\xi S_{\text{IR}} = 0 \tag{11.4}$$

identically, off shell as a statement about the integrand up to a boundary term. Noether's second theorem then supplies

$$\nabla_\mu\Big(\frac{1}{8\pi G_{\text{eff}}}(G^{\mu\nu} + \Lambda_{\text{eff}} g^{\mu\nu}) - T^{\mu\nu}_{\text{total}}\Big) = 0 \tag{11.5}$$

as an identity, once the matter and condensate fields Ψ are on shell. The contracted Bianchi identity $\nabla_\mu G^{\mu\nu} = 0$ and metric compatibility $\nabla_\mu g^{\mu\nu} = 0$ reduce (11.5) to the on-shell conservation law

$$\nabla_\mu T^{\mu\nu}_{\text{total}} = 0. \tag{11.6}$$

Theorem 11.1 (Conservation of the total stress)**.** *Assume:*

1. *the infrared action S_{IR} is a scalar under diffeomorphisms of the composite metric, including the entropic functional $S_{\text{ent}}[g]$;*

2. *the Einstein equation is* (11.1)*;*

3. matter fields and condensate fluctuations are on shell with respect to S_{IR}.

Then

$$\nabla^{\mu}(T^{\mathrm{matter}}_{\mu\nu} + T^{\mathrm{tensor}}_{\mu\nu} + T^{\mathrm{ent}}_{\mu\nu}) = 0. \tag{11.7}$$

Separate conservation of any one summand is not implied.

The Einstein-plus-extra-tensor form of the infrared equation is collected in Appendix F.

Proof. Diffeomorphism invariance of S_{IR} is Assumption 1. Noether's second theorem yields (11.5). Bianchi and (11.1) yield (11.7). The last sentence is the observation that Noether's theorem constrains only the sum that couples to $g_{\mu\nu}$. □

Remark 11.2 (The real hypothesis). Everything in the proof is standard except Assumption 1 applied to S_{ent}. Mutual information (2.31) is a scalar for each fixed partition of a spatial slice. Promoting it to a diffeomorphism scalar on a dynamical spacetime requires that the assignment $(g, A, B) \mapsto I(A : B)$ commute with Lie dragging of the regions. That is true if A and B are defined geometrically (causal domains, outermost trapped surfaces, constant-mean-curvature balls of specified volume) and fails if they are defined by coordinate inequalities such as $r \leq R$ in a preferred chart. The book uses geometric regions.

11.3 Structure of the entropic stress

A first variation of the static entanglement potential produces a conservative interaction. A time-dependent configuration or first variation may introduce additional terms, but does not by itself establish a velocity-odd or dissipative stress. The split

$$T^{\mathrm{ent}}_{\mu\nu} = T^{\mathrm{ent,rev}}_{\mu\nu} + T^{\mathrm{ent,irrev}}_{\mu\nu} \tag{11.8}$$

is phenomenological unless its second term is identified by a dynamical response calculation.

On two worldlines γ_A, γ_B in a nearly Minkowski region one has, at leading post-Newtonian order,

$$U_{\mathrm{ent}} = -\lambda I\big(\ell_A, \ell_B, d(\gamma_A, \gamma_B)\big), \tag{11.9}$$

and the four-force on body A is

$$f^{\mu}_{A} = -\frac{D}{D\tau_A}\Big(\frac{\partial L_{\mathrm{ent}}}{\partial \dot{x}_{A\mu}}\Big) + \frac{\partial L_{\mathrm{ent}}}{\partial x_{A\mu}}, \tag{11.10}$$

with $L_{\mathrm{ent}} = -U_{\mathrm{ent}}$ along the worldlines. The static dependence on d gives the conservative interaction. Terms involving $\dot{d}$ or changing ℓ_A, ℓ_B may occur in a more

general effective action, but their presence alone does not establish irreversibility. Whether they produce dissipation must be determined from the dynamics and its retarded response.

A phenomenological Newtonian drag, for a reduced mass μ and a separation d, is of the form

$$\mathbf{a}_{\rm irrev} = -\eta(d, \ell_A, \ell_B)\, \frac{\mathbf{v}}{d^n}. \tag{11.11}$$

Neither the sign of the dissipative coefficient nor the exponent n in (11.11) follows from positivity of mutual information. The source notes use $n = 4$ as a working exponent, not as a consequence of differentiating a static tail. Such differentiation does not, by itself, derive a velocity-linear dissipative force. Chapter 15 adopts

$$\mathbf{F}_{\rm diss} = -\eta\, \frac{\mu}{d^4}\, \mathbf{v} \tag{11.12}$$

as a laboratory assumption, not as a derived multipole formula; deriving it requires the dynamical response calculation specified in Problem H.9.

Proposition 11.3 (Necessary condition for a covariant drag reduction)**.** *Suppose a two-body equation*

$$\mu \ddot{\mathbf{r}} = -\nabla U_{\rm N} - \nabla U_{\rm ent}^{(\rm rev)} + \mathbf{F}_{\rm diss} \tag{11.13}$$

is claimed to follow from a covariantly conserved microscopic stress. In a non-radiative reduced subsystem, a necessary bookkeeping condition, to the post-Newtonian order retained, is that there exist a functional $E_{\rm ent}$ of the microscopic state, specified independently of $\mathbf{F}_{\rm diss}$ and of the reduced equation of motion (11.13) itself — *in particular, not any quantity derivable purely from the reduced two-body flow, such as (but not limited to) a first integral of* (11.13) *or the accumulated dissipation $-W_{\rm diss}$ along a trajectory — such that, writing*

$$E_{\rm mech} = \tfrac{1}{2}\mu v^2 + U_{\rm N} + U_{\rm ent}^{(\rm rev)},$$

$$\frac{\rm d}{{\rm d}t}(E_{\rm mech} + E_{\rm ent}) = O(c^{-2}), \qquad \frac{{\rm d}E_{\rm ent}}{{\rm d}t} = -\mathbf{F}_{\rm diss}\cdot\mathbf{v} + O(c^{-2}). \tag{11.14}$$

A microscopic candidate for $E_{\rm ent}$ must be defined from degrees of freedom of the microscopic theory (e.g. a specified functional of the reduced density matrix or of $I(A:B)$), not from $(\mathbf{r}, \mathbf{v})$ and $\mathbf{F}_{\rm diss}$ alone. We additionally require any such candidate to be bounded below, since an unbounded reservoir would let the balance accommodate arbitrary drag; this is a physical requirement imposed here, not a consequence of covariant conservation. The microscopic derivation must also account for the force on the second body and, when radiation is present, the corresponding energy flux. If the proposed $E_{\rm ent}$ is claimed to arise from mutual information, that relation must be derived from the microscopic theory.

This proposition is a necessary condition for a claimed covariant reduction, and it has content only because E_{ent} is required to come from outside the reduced two-body flow; that general exclusion is stated in the proposition itself. Two instances of a construction internal to the flow illustrate why the exclusion is needed, not an exhaustive list of what it rules out. First, $E_{\text{ent}} = -W_{\text{diss}} + E_{\text{ent}}(0)$, with $W_{\text{diss}} = \int \mathbf{F}_{\text{diss}} \cdot \mathbf{v}\, \mathrm{d}t$ a path-dependent functional of the trajectory, satisfies (11.14) identically for *any* $\mathbf{F}_{\text{diss}}$. Second, because a smooth non-vanishing vector field admits a local first integral $C(\mathbf{r}, \mathbf{v})$ with $\dot{C} = 0$ along its flow (a flow-box argument), $E_{\text{ent}} := C - E_{\text{mech}}$, and indeed any function of such first integrals, also satisfies the balance identically, for any $\mathbf{F}_{\text{diss}}$, with no reference to mutual information or any microscopic quantity. None of these constructions is a microscopic state function in the required sense, and none derives E_{ent} from $I(A : B)$ or proves covariant stress conservation. For passive drag, $\mathbf{F}_{\text{diss}} \cdot \mathbf{v} \leq 0$, so a microscopic E_{ent} satisfying the proposition, if one exists, gains the energy removed from the mechanical degree of freedom.

The laboratory (11.12) is not shown here to satisfy Proposition 11.3. Chapter 15 tracks $-W_{\text{diss}}$, which is bookkeeping for the reduced ODE and is not a microscopic E_{ent} in the sense above.

11.4 Exchange with matter

Equation (11.7) permits

$$\nabla^\mu T^{\text{matter}}_{\mu\nu} = -\nabla^\mu T^{\text{ent}}_{\mu\nu} - \nabla^\mu T^{\text{tensor}}_{\mu\nu}. \tag{11.15}$$

Matter may heat or cool if entanglement of the condensate cores is changing. In a tidal disruption event the conditional reading is: as the disrupted star is sheared, the mutual information between debris parcels and the compact core may change. If the phenomenological drag were the reduction of a covariantly conserved stress, the work done by $\mathbf{F}_{\text{diss}}$ would be accounted in a microscopic E_{ent} rather than lost; that identification has not been derived and is part of Problem H.9.

This is still not a model of microphysical dissipation in a fluid with a viscosity measured in the laboratory. Whether conservation of the total stress requires such a channel, once U_{ent} is admitted as part of the action, depends on the dynamical response calculation of Problem H.9.

11.5 Vacuum energy as a conservation constraint

A term $\rho_\Lambda g_{\mu\nu}$ with constant ρ_Λ is conserved by itself:

$$\nabla^\mu(\rho_\Lambda g_{\mu\nu}) = 0. \tag{11.16}$$

A term whose prefactor varies, $\rho_\Lambda(x)g_{\mu\nu}$, is not conserved unless

$$\partial_\mu \rho_\Lambda = 0 \tag{11.17}$$

or additional current terms are supplied. The heat-kernel density (8.13) is constant in the approximation that the included sector weights and their assigned cutoffs are constant. If Chapter 12 regulates ρ_Λ by a boundary functional that depends on the metric induced on ∂M, the regulated object $\Lambda_{\text{eff}}[g]$ must still obey (11.7). That requirement is a design constraint on HBEV, not an optional aesthetic.

11.6 What this chapter has not done

Theorem 11.1 does not compute η, does not fix $n = 4$, does not evaluate $I(A : B)$ for Kerr exteriors, and does not replace quadrupole radiation. Gravitational waves remain the dissipative channel of the tensor block. Any entropic drag would be an additional channel, admissible only if it sits inside a conserved total tensor; that the phenomenological drag does so remains Problem H.9.

The theorem also does not justify writing an arbitrary function $\mathbf{a}_{\text{irrev}}(\mathbf{r}, \mathbf{v})$ and calling it NAMCG. Phenomenological chapters that introduce a drag must exhibit the E_{ent} of Proposition 11.3 or must state that they are studying a reduced laboratory.

11.7 Exercises

1. **Bianchi reminder.** From $\nabla^\mu G_{\mu\nu} = 0$ and (11.1), obtain (11.7) in three lines.

2. **Two-body check.** For a potential $U(d)$ depending only on separation, show that the Newtonian forces are equal and opposite. Which symmetry of $I(A : B)$ is the relativistic ancestor of that statement?

3. **Energy bookkeeping.** Take (11.12) with constant η and compute $\mathbf{F}_{\text{diss}} \cdot \mathbf{v}$. Show that $E_{\text{ent}} = -W_{\text{diss}} + E_{\text{ent}}(0)$ satisfies (11.14) for the reduced ODE. Why does this construction not satisfy Proposition 11.3?

4. **Illegal drag.** Consider $\mathbf{F} = -\gamma \mathbf{v}$ with constant γ, independent of d. If E_{ent} is required to be a bounded functional of $I(A : B)$, why does this fail Proposition 11.3 for two bodies that can be separated to infinity at finite cost in $I(A : B)$?

5. **Variable vacuum energy.** Let $\rho_\Lambda = \rho_\Lambda(\phi)$ for a scalar ϕ. Construct the additional stress that restores conservation. Compare with a potential $V(\phi)$ in ordinary scalar-field cosmology.

6. **Isaacson average.** Write the Isaacson stress of a high-frequency TT wave and confirm that its divergence vanishes after a short-wavelength average. That is the tensor-block contribution to (11.7) in a radiative spacetime.

7. **Coordinate regions.** Give an explicit example of two coordinate-defined regions whose mutual information is *not* a diffeomorphism scalar. This is the counterexample that forces the remark after Theorem 11.1.

11.8 Bridge to Chapter 12

Conservation is now in force for whatever vacuum energy Chapter 8 left untamed. A constant ρ_Λ is conserved and disastrous. A regulated Λ_{eff} must remain conserved after the regulation. Chapter 12 introduces the Holographic Boundary-Entropy Vacuum: a boundary functional built from $K/8\pi G_{\text{eff}}$ and an entropy density $S_{\text{ent}}(h_{ij}, \phi)$ whose stationarity sets Λ_{eff} by an infrared entanglement datum rather than by Λ_{UV}^4. The design constraint just stated—that the resulting $\Lambda_{\text{eff}} g_{\mu\nu}$ still sit inside Theorem 11.1—is the first test that construction must pass.

Chapter 12

Holographic Boundary-Entropy Vacuum and Λ_{eff}

Learning outcomes. After this chapter the reader should be able to:

1. restate the quartic vacuum-energy debt isolated in Chapter 8 as a problem about a bulk term $\rho_\Lambda\sqrt{-g}$;
2. write the HBEV boundary action built from the Gibbons–Hawking–York term and an entropy density $S_{\text{ent}}(h_{ij},\phi)$;
3. obtain Λ_{eff} from stationarity of that boundary action rather than from Λ_{UV}^4;
4. check that the regulated cosmological term still obeys Theorem 11.1;
5. enter Chapter 13 knowing which vacuum energy is being inserted into regular-core and galactic solutions.

Chapter 8 produced

$$\rho_\Lambda = -\sum_{s\in\mathcal{S}_{\text{sec}}} \frac{w_s \operatorname{tr}_{V_s}\mathbb{I}_s}{64\pi^2}\left(\Lambda_{\text{UV}}{}_s^4 - \mu_s^4\right) + \cdots . \tag{12.1}$$

This is the truncated, unresummed sector sum of Chapter 8 under its stated continuation, with the same included sectors and determinant weights. A common-limit form using $\mathcal{N}_0$ applies only when all included sectors are assigned the same limits; otherwise sector-dependent thresholds are retained. Only if the weighted quartic contribution is nonzero and remains uncancelled, including after finite local terms and counterterms are specified, must its magnitude be reconciled with the observed cosmological constant. Without those conditions no such comparison follows from (12.1). This chapter does not cancel it by a bare counterterm tuned to 120 decimal places. It replaces the bulk

cosmological term by a boundary variational problem whose solution is an infrared scale.

12.1 What would not count as a solution

Three common manoeuvres are excluded in advance.

1. **Subtraction by hand.** Adding $-\rho_\Lambda \int \sqrt{-g}$ to the action and declaring the sum zero is a renormalization condition at one scale. It does not explain the residual scale.

2. **Setting $\Lambda_{\rm UV}$ to meV.** The cutoff constraint cannot be assessed until the physical $G_{\rm eff}$ is obtained from the specified sector calculation of Problem F.

3. **A spacetime-dependent $\rho_\Lambda(x)$ with no current.** Section 11.5 forbids this unless extra stress is supplied. Regulation that breaks Theorem 11.1 is not available.

Any construction that survives must therefore (i) replace the bulk quartic by an infrared datum, (ii) use a $G_{\rm eff}$ matched from specified operators, weights, thresholds, continuation, and scheme as required by Problem F, and (iii) produce a conserved $\Lambda_{\rm eff} g_{\mu\nu}$ or a conserved completion of it.

12.2 Boundary geometry

Let M be a region of the emergent spacetime with boundary ∂M, outward normal n^μ, induced metric

$$h_{ij} = g_{\mu\nu} e_i^\mu e_j^\nu, \tag{12.2}$$

and extrinsic curvature

$$K_{ij} = \nabla_{(\mu} n_{\nu)} e_i^\mu e_j^\nu, \qquad K = h^{ij} K_{ij}. \tag{12.3}$$

The Gibbons–Hawking–York term already introduced in (2.11) is

$$S_{\rm GHY} = \frac{1}{8\pi G_{\rm eff}} \int_{\partial M} \mathrm{d}^3x \, \sqrt{|h|}\, K. \tag{12.4}$$

On a timelike boundary K measures the expansion of the outward normal; on a spacelike boundary it measures the expansion of a unit-normal congruence. HBEV uses a timelike conformal boundary, or a large-radius timelike tube that will be sent to infinity after variation, according to the problem.

The boundary may also carry condensate data. Write ϕ for the pull-back of the polar radial mode σ and of any unlocked scalars that survive Assumption 5.1. The pair (h_{ij}, ϕ) is the boundary field content.

12.3 The HBEV action

Definition 12.1 (HBEV boundary action)**.** The holographic boundary-entropy vacuum functional is

$$S_{\mathrm{HBEV}}[h,\phi] = \int_{\partial M} \mathrm{d}^3x\, \sqrt{|h|}\Big(\frac{K}{8\pi G_{\mathrm{eff}}} - T_{\mathrm{bdy}} - S_{\mathrm{ent}}(h_{ij},\phi)\Big), \tag{12.5}$$

where T_{bdy} is a boundary tension (a counterterm density built from h_{ij} only) and $S_{\mathrm{ent}}(h_{ij},\phi)$ is an entanglement-entropy density of the condensate restricted to ∂M.

The first term is GHY. The second term is the standard counterterm that renders the on-shell action finite on AdS-like or asymptotically flat tubes; in the present setting it is fixed by demanding that S_{HBEV} vanish on the reference condensate $\bar{e}^A_\mu = \delta^A_\mu$, $\phi = 0$. The third term is the new ingredient. It is not the diamond entropy S_D of Chapter 6. It is a local density on ∂M whose integral tracks the entanglement of bulk condensate degrees of freedom that end on the boundary.

Assumption 12.2 (Existence of a local entropy density)**.** In the condensed phase there exists a diffeomorphism-scalar density $S_{\mathrm{ent}}(h_{ij},\phi)$, of mass dimension three, obtained from the replica construction of Section 6.4 applied to a thin tubular neighbourhood of ∂M and then localized. Its leading terms are

$$S_{\mathrm{ent}} = \alpha_0 \Lambda_{\mathrm{UV}}^3 + \alpha_1 \Lambda_{\mathrm{UV}} R^{(h)} + \alpha_2\, \phi^2 \Lambda_{\mathrm{UV}} + \cdots, \tag{12.6}$$

with dimensionless coefficients α_i of order the fibre trace.

The $\alpha_0 \Lambda_{\mathrm{UV}}^3$ piece is the boundary analogue of ρ_Λ. It will be cancelled against T_{bdy} and against the leading GHY contribution on the reference background. What is not cancelled is required to be an infrared quantity.

12.4 Stationarity and the induced cosmological term

Vary $S_{\mathrm{EH}} + S_{\mathrm{HBEV}}$ with respect to the boundary metric h_{ij}, holding the bulk fields on shell. The bulk Einstein tensor produces the usual Brown–York stress. The boundary variation produces

$$\pi^{ij} = \frac{1}{8\pi G_{\mathrm{eff}}}\big(K^{ij} - K h^{ij}\big) - T^{ij}_{\mathrm{bdy}} - \frac{2}{\sqrt{|h|}}\frac{\delta S_{\mathrm{ent}}}{\delta h_{ij}}. \tag{12.7}$$

HBEV imposes entanglement equilibrium at the boundary in the same logical role that Assumption 6.1 played in the bulk diamonds:

Assumption 12.3 (Boundary entanglement equilibrium)**.** The physical vacuum is a stationary point of S_{HBEV} at fixed boundary volume element, in the sense that the trace-adjusted boundary stress vanishes,

$$h_{ij}\pi^{ij} = 0. \tag{12.8}$$

On an asymptotically simple tube of radius r_B, with $K = 2/r_B + O(r_B^{-3})$ in asymptotically flat space and $K = 3/\ell_{\mathrm{AdS}} + O(r_B^{-2})$ in AdS, equation (12.8) reduces to a single condition on the integration constant in the bulk Einstein equation with a cosmological term. That constant is Λ_{eff}.

Theorem 12.4 (Induced cosmological constant)**.** *Under Assumptions 12.2 and 12.3, and with T_{bdy} chosen to cancel the reference $\alpha_0\Lambda_{\mathrm{UV}}^3$ piece, the bulk field equation compatible with boundary stationarity is*

$$G_{\mu\nu} + \Lambda_{\mathrm{eff}} g_{\mu\nu} = 8\pi G_{\mathrm{eff}}\, T_{\mu\nu}^{\mathrm{total}}, \tag{12.9}$$

where

$$\frac{\Lambda_{\mathrm{eff}}}{8\pi G_{\mathrm{eff}}} = \langle S_{\mathrm{ent}}^{(\mathrm{IR})} \rangle_{\partial M} - T_{\mathrm{bdy}}^{(\mathrm{IR})} \tag{12.10}$$

is determined by the infrared part of the boundary entropy density after the Λ_{UV}^3 subtraction. In particular Λ_{eff} does not scale as Λ_{UV}^2.

Proof sketch. On-shell variation of $S_{\mathrm{EH}} + S_{\mathrm{GHY}}$ with respect to a constant bulk cosmological term is proportional to the spacetime volume, which by Stokes is a boundary integral of the normal. That boundary integral is matched to $h_{ij}\pi^{ij}$ after the GHY contribution is included. Imposing (12.8) fixes the constant. The ultraviolet pieces of S_{ent} are cancelled by the choice of T_{bdy} on the reference background, which is a finite renormalization of a three-dimensional density, not a fine-tuning of a four-dimensional quartic against itself. The remainder is infrared by construction of the subtraction, and is constant on the vacuum because Assumption 12.3 is imposed uniformly on the asymptotic tube. □

Remark 12.5 (Dimensional status of the boundary-to-bulk relation)*.* In units with $\hbar = c = 1$, the left-hand side of (12.10) has mass dimension four: $[\Lambda_{\mathrm{eff}}] = 2$ and $[G_{\mathrm{eff}}] = -2$. The terms in the expansion (12.6) each have mass dimension three, as does T_{bdy} in (12.7). Thus the right-hand side of (12.10), as written, has mass dimension three, and the equality is dimensionally incomplete.

The missing input enters the proof sketch at the step that matches the on-shell bulk variation to a boundary integral and uses boundary stationarity to fix the bulk

cosmological constant. A boundary-to-bulk conversion of mass dimension one, with its normalization specified by the boundary geometry and variational prescription, is needed for that step. No such conversion is supplied here, and no length is inserted by assumption. Until it is supplied, the conclusion of Theorem 12.4 on the scaling of $\Lambda_{\rm eff}$, including its independence of $\Lambda_{\rm UV}^2$, is conditional on that conversion. Accordingly, (12.10) is schematic pending this input, and the boundary mechanism remains conditional; Problem G remains OPEN.

Remark 12.6 (What the theorem does not claim). Equation (12.10) does not compute the observed value 10^{-122} in Planck units. It changes the *origin* of the scale: from $\Lambda_{\rm UV}^4$ to an infrared boundary entropy. Predicting the numerical value requires an evaluation of $S_{\rm ent}^{\rm (IR)}$ on the actual cosmological boundary, which this book does not perform. Problem G also requires deriving the entropy density and boundary stationarity from the matrix theory; evaluating a scaling within an assumed boundary model alone remains programme-conditional (Section 17.6).

12.5 Conservation

A constant $\Lambda_{\rm eff}$ produced by Theorem 12.4 satisfies

$$\nabla^\mu(\Lambda_{\rm eff} g_{\mu\nu}) = 0 \tag{12.11}$$

automatically. The design constraint of Section 11.5 is therefore met in the vacuum.

If the boundary data ϕ are dynamical, $S_{\rm ent}$ can depend on time and $\Lambda_{\rm eff}$ can drift. In that case Theorem 11.1 requires a compensating current. The compensation is the boundary analogue of Exercise 5 in Chapter 11: the variation of $S_{\rm HBEV}$ with respect to ϕ supplies a boundary flux that, by the on-shell identity for the radial mode, matches $\partial_\mu \Lambda_{\rm eff}$. The book uses a constant $\Lambda_{\rm eff}$ in all later solutions and models and treats a rolling case as a cosmological refinement, not as the default.

12.6 Relation to the diamond entropy of Chapter 6

The two entanglement functionals in the book now have distinct jobs.

Functional	Support	Job
S_D	small bulk diamonds	kill m_T
$S_{\rm HBEV}$	asymptotic boundary	set $\Lambda_{\rm eff}$
$I(A:B)$	two compact cores	conservative potential

They are not interchangeable. Using S_D to subtract ρ_Λ would confuse a local first-law identity with a global vacuum-energy problem. Using $S_{\rm HBEV}$ to kill m_T

would confuse a boundary condition with a bulk Ward identity. The architecture is intentional.

12.7 Working value used later

Later chapters use the regular-core model

$$f(r) = 1 - \frac{2Mr^2}{r^3 + r_0^3} - \frac{\Lambda_{\text{eff}}}{3} r^2 \tag{12.12}$$

with Λ_{eff} a constant of either sign. The negative sign in f for $\Lambda_{\text{eff}} > 0$ is de Sitter-like and is the default cosmological reading; the opposite sign is a mathematical option for matching interiors. No Solar-System observable in Chapters 13–15 depends on the numerical size of Λ_{eff} at laboratory precision. In this model, $f(r) \to 1 - \Lambda_{\text{eff}} r^2/3$ as $r \to 0$.

12.8 Exercises

1. **GHY variation.** Vary (12.4) with respect to h_{ij} and recover the $K^{ij} - Kh^{ij}$ piece of (12.7).

2. **Reference subtraction.** On Minkowski space in a large sphere of radius r_B, compute K and $\sqrt{h}$. What T_{bdy} cancels S_{HBEV} at $\phi = 0$, $S_{\text{ent}} = \alpha_0 \Lambda_{\text{UV}}^3$?

3. **Why not Λ_{UV}^4.** Contrast a four-dimensional counterterm $\int_M \sqrt{-g}\, \Lambda_{\text{UV}}^4$ with a three-dimensional counterterm $\int_{\partial M} \sqrt{h}\, \Lambda_{\text{UV}}^3$. Which tuning is a cancellation of two bulk quartics, and which is a renormalization of a boundary density?

4. **Trace condition.** Reduce (12.8) on a spherically symmetric tube $r = r_B$ in Schwarzschild–de Sitter and show that it constrains Λ_{eff} rather than M.

5. **Rolling Λ.** Let S_{ent} depend on a homogeneous boundary scalar $\phi(t)$. Write the extra flux needed to keep Theorem 11.1. Compare with a slowly rolling quintessence potential.

6. **Two functionals.** In one paragraph, explain what goes wrong if one attempts to set Λ_{eff} from $\delta S_D = 0$ on a single small diamond.

12.9 Bridge to Chapter 13

The conditional infrared Einstein equation has the form

$$G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = 8\pi G_{\text{eff}}\, T_{\mu\nu}^{\text{total}}, \tag{12.13}$$

with G_{eff} to be matched by the specified operator, weight, threshold, continuation, and scheme calculation of Problem F, Λ_{eff} from Theorem 12.4, $m_T = 0$ from Theorem 7.3, and conservation from Theorem 11.1. What remains of the geometric programme is the high-curvature interior, where the Einstein truncation fails and Assumption 9.3 is used. Chapter 13 uses the static spherical sector of Einsteinian cubic gravity coupled to (12.13) to motivate the regular-core model (12.12), which is a model and not an established solution (Section H.10), and only then turns to galactic backreaction as a consequence of $T^{\mathrm{ent}}_{\mu\nu}$ in the weak- field limit—not as a substitute for the derivation just completed.

Chapter 13

Regular Cores and Galactic Backreaction

Learning outcomes. After this chapter the reader should be able to:

1. write the static, spherically symmetric reduction of the infrared equation $G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = 8\pi G_{\text{eff}} T^{\text{total}}_{\mu\nu}$ with the Einsteinian cubic term included;
2. state the regular-core profile $f(r) = 1 - 2Mr^2/(r^3 + r_0^3) - \Lambda_{\text{eff}} r^2/3$ as a phenomenological model, and state the scope of the no-go on its exactness (Section H.10);
3. reduce the reversible part of $T^{\text{ent}}_{\mu\nu}$ to a modification of Poisson's equation;
4. derive a schematic circular-velocity formula and list the observations that would distinguish it from a particulate halo;
5. enter Chapter 14 knowing that any time-dependent dissipative term would be a separate addition to the static extra potential used here.

The geometric programme of Parts II–IV is applied to two symmetric sectors: static spherical interiors and weak-field galactic exteriors. The regular-core profile below remains a model conditional on Problem J, not a consequence of the earlier theorems.

13.1 Static spherical ansatz

Take the emergent metric in Schwarzschild coordinates,

$$ds^2 = -f(r)\,dt^2 + \frac{dr^2}{f(r)} + r^2 d\Omega^2, \tag{13.1}$$

and restrict $T^{\text{total}}_{\mu\nu}$ to a static, spherical form. The Einstein tensor of (13.1) has the standard components

$$G^t{}_t = G^r{}_r = -\frac{1-f-rf'}{r^2}, \qquad G^\theta{}_\theta = -\frac{rf''+2f'}{2r}. \tag{13.2}$$

With only the Einstein–Hilbert term, vacuum solutions with constant Λ_{eff} are Schwarzschild–de Sitter,

$$f(r) = 1 - \frac{2M}{r} - \frac{\Lambda_{\text{eff}}}{3}r^2, \tag{13.3}$$

which is singular at $r = 0$ whenever $M \neq 0$.

13.2 Cubic correction in the spherical sector

Assumption 9.3 promotes the Einsteinian cubic density $\mathcal{P}$ to a term that is solved non-perturbatively in this sector. Conditional on the first-integral equation of Section H.10 being the static spherical vacuum equation of the action (9.8), that equation is a nonlinear second-order ODE for f, not an algebraic modification of the mass function. The mass function below parametrises f; it is not thereby a solution of that equation. Write

$$f(r) = 1 - \frac{2m(r)}{r} - \frac{\Lambda_{\text{eff}}}{3}r^2. \tag{13.4}$$

The regular-core profile used here is the model

$$m(r) = \frac{Mr^3}{r^3 + r_0^3}, \tag{13.5}$$

and therefore

$$f(r) = 1 - \frac{2Mr^2}{r^3 + r_0^3} - \frac{\Lambda_{\text{eff}}}{3}r^2. \tag{13.6}$$

The core scale is

$$r_0^3 = 2\beta_{\text{ECG}}M \tag{13.7}$$

as a phenomenological parametrization (up to the conventional normalization of $\mathcal{P}$); this relation is not derived from the action or the first-integral equation. The ECG asymptotic correction instead defines a characteristic tail length $r_* \sim |\alpha M|^{1/5}$, where $\alpha = -16\pi G_{\text{eff}}\beta_{\text{ECG}}$ in the stated action normalization. This does not determine r_0 or identify it with a core length. The book does not claim a measured value of r_0.

Limits. As $r \to \infty$, (13.6) returns to (13.3). As $r \to 0$,

$$f(r) \to 1 - \frac{\Lambda_{\text{eff}}}{3}r^2 - \frac{2M}{r_0^3}r^2 + O(r^5), \tag{13.8}$$

which is finite. The curvature scalars remain finite at the origin for $r_0 \neq 0$. Geodesic completeness through $r = 0$ is a separate question: the chart (13.1) is not itself complete, and a regular origin still requires a smooth matching onto a neighbourhood covered by local inertial frames of the Levi-Civita geometry of the composite metric. The frame retains local Lorentz freedom; the independent Lorentz connection B is not used here.

Remark 13.1 (Model, not an established solution). Equation (13.6) is the working profile of the notes. For the first-integral equation of Section H.10, it is not an exact vacuum solution when $\Lambda_{\text{eff}} = 0$ and $Mr_0 \neq 0$. The equation also excludes a nonzero integration constant for a smooth regular centre for any Λ_{eff}. These conclusions are conditional on the reduction and the first integral for the invariant $\mathcal{P}$ of (9.8). Whenever a later section uses (13.6), it is using this model, not a theorem that ECG produces $r^3/(r^3 + r_0^3)$.

13.3 Horizons and the core

Roots of $f(r) = 0$ are Killing horizons of (13.1). For $\Lambda_{\text{eff}} = 0$ and $r_0 > 0$ there is a range of M in which two positive roots exist, one, or none, according to the ratio M/r_0. For astrophysical masses and microscopic r_0 the exterior is indistinguishable from Schwarzschild at the photon sphere and beyond. The core is hidden behind a horizon if a horizon exists, and is naked if it does not.

NAMCG does not require every compact object to be horizonless. It requires that the Einstein singularity not be accepted as fundamental. Whether LIGO ringdowns or EHT images can see r_0 is a question for Chapter 16, not a claim of this chapter.

The Einstein-plus-extra-tensor form of the infrared equation is collected in Appendix F.

13.4 Weak-field exterior and the extra Poisson equation

Far from a core, $r \gg r_0$, the cubic term is negligible and the geometry is Einstein plus $T^{\text{ent}}_{\mu\nu}$. In the Newtonian gauge

$$ds^2 = -(1 + 2\Phi)\, dt^2 + (1 - 2\Phi)\delta_{ij} dx^i dx^j, \tag{13.9}$$

the reversible, static piece of the entropic stress reduces to an effective density

$$\rho_{\text{ent}} = T^{\text{ent,rev}}_{00}. \tag{13.10}$$

Poisson's equation becomes

$$\nabla^2\Phi = 4\pi G_{\text{eff}}(\rho_{\text{b}} + \rho_{\text{ent}}), \tag{13.11}$$

where ρ_{b} is baryonic density. Conservation, Theorem 11.1, requires that ρ_{ent} not be an arbitrary halo function. It is determined by the variation of U_{ent} for the condensate configuration sourced by ρ_{b}.

A useful schematic, sufficient for circular orbits and not sufficient for lensing tomography, is a source-tracking extra density

$$\rho_{\text{ent}}(r) = \frac{1}{4\pi G_{\text{eff}}}\nabla\cdot\big(\mu_{\text{ent}}(g_{\text{N}})\,\nabla\Phi_{\text{b}}\big), \tag{13.12}$$

where $g_{\text{N}} = |\nabla\Phi_{\text{b}}|$ is the Newtonian baryonic field and μ_{ent} is a dimensionless interpolating function fixed by the mutual-information functional. Equation (13.12) is written to make the structure visible: ρ_{ent} is slaved to baryons. It is not a six-parameter halo fitted independently of ρ_{b}.

Assumption 13.2 (Reversible extra density tracks baryons)**.** In static, weak-field systems the leading reversible entropic density is a local or mildly nonlocal functional of the baryonic Newtonian field, of the form (13.12) or a convolution thereof with kernel width of order the core scale of the constituent condensates.

Assumption 13.2 is the galactic input. It is narrower than "there exists dark matter," and it is stronger than anything proved from $I(A:B)$ in Chapter 6. Chapter 17 lists a derivation of μ_{ent} from $I(A:B)$ as open.

Assumption 13.3 (External field enters the interpolator)**.** For a subsystem whose internal Newtonian field $\mathbf{g}_{\text{N}}$ is embedded in a baryonic field $\mathbf{g}_{\text{ext}}$ that varies slowly across the subsystem, the reversible boost is evaluated on the total field,

$$\mathbf{g} = \nu_{\surd}(g_{\text{tot}})\,\mathbf{g}_{\text{N}} + \Big[\nu_{\surd}(g_{\text{ext}}) - \nu_{\surd}(g_{\text{tot}})\Big]\mathbf{g}_{\text{ext}}, \tag{13.13}$$

with $g_{\text{tot}} = |\mathbf{g}_{\text{N}} + \mathbf{g}_{\text{ext}}|$ and $\nu_{\surd}$ as in (13.15). In the external-field-dominated limit $g_{\text{N}} \ll g_{\text{ext}}$,

$$\gamma \equiv \frac{g_{\text{int}}}{g_{\text{N}}} \to \nu_{\surd}(g_{\text{ext}}), \tag{13.14}$$

not $\nu_{\surd}(g_{\text{N}})$.

Assumption 13.3 is a coefficient. It is what $U_{\text{ent}}[I(A:B)]$ must do if mutual information sees the Galactic baryons as well as the binary. It is not implied by Theorem 11.1, and it is not a solution of Problem H. The Galactic scale at the Sun, $g_{\text{gal}} \approx 2.2\times10^{-10}\,\text{m}\,\text{s}^{-2}$, already satisfies $g_{\text{gal}} > g_{\dagger}$, so $\nu_{\surd}(g_{\text{gal}}) \approx 1.1$. Isolated-RAR boosts of order 4 are then illegal for solar-neighborhood wide binaries. A null Gaia result constrains the isolated reading of Appendix D, not $m_T = 0$.

Working interpolator. Until Problem H of Chapter 17 produces $\mu_{\rm ent}$ from $I(A:B)$, galactic comparisons use

$$\nu_{\surd}(g_{\rm N}) = \frac{1}{1-\exp(-\sqrt{g_{\rm N}/g_s})}, \tag{13.15}$$

so that $v_c = V_{\rm b}\sqrt{\nu_{\surd}}$. The default scale is the published SPARC value $g_s = g_\dagger = 1.20\times10^{-10}\,{\rm m\,s^{-2}}$ [4]. This ν is the RAR interpolator written as a boost. It is not $1/(1-\alpha\exp(-g_{\rm N}/g_s))$ and it is not a per-galaxy radial profile. Appendix D records the SPARC ranking and retires the older exponential.

13.5 Circular velocities

For circular orbits in the midplane,

$$v_c^2(r) = r\partial_r\Phi = G_{\rm eff}\frac{M_{\rm b}(<r)+M_{\rm ent}(<r)}{r}, \tag{13.16}$$

with $M_{\rm ent}$ the integral of (13.10). Two regimes follow once $\mu_{\rm ent}$ interpolates from 0 at large $g_{\rm N}$ to a finite value at small $g_{\rm N}$.

Interior / high acceleration. $\rho_{\rm ent}$ is negligible and $v_c^2 \simeq G_{\rm eff}M_{\rm b}/r$. Solar-System and inner-disk data constrain how small $\mu_{\rm ent}$ must be at those accelerations.

Outer / low acceleration. If $M_{\rm ent}(<r)$ grows approximately linearly with r, v_c flattens. That growth is not mysterious once (13.12) is granted: it is the integral of an extra density that falls more slowly than $\rho_{\rm b}$.

On 175 SPARC rotmod files with fixed mass-to-light ratios, (13.15) at frozen $g_\dagger$ is statistically equivalent to the published RAR and about 0.2% behind simple MOND in χ^2, while baryons-only is an order of magnitude worse (Appendix D). That ranking is a coefficient check of Assumption 13.2. It is not a measurement of $G_{\rm eff}$, not a derivation from $I(A:B)$, and not evidence reserved for the cover. Whether a single $\nu_{\surd}$ survives clusters remains a Chapter 16 question.

13.6 What this is not

It is not a proof that dark matter particles are absent. Cluster lensing, the CMB, and colliding clusters probe $T^{\rm ent}_{\mu\nu}$ in regimes that (13.12) may fail. If those data require a stress that does not track baryons, NAMCG must either supply an additional condensate configuration that does, or yield the dark-matter interpretation in those regimes.

It is not MOND, though the interpolating function $\mu_{\rm ent}$ invites comparison. MOND is a modification of inertia or of the gravitational field equation at a fixed acceleration scale a_0. NAMCG obtains an extra conserved tensor from entanglement. If $\mu_{\rm ent}$ happens to mimic MOND in disks and fails in clusters, that is a phenomenological coincidence to be measured, not a founding principle.

It is not a replacement for Chapters 3–12. A successful rotation- curve fit that used an illegal drag, or a non-conserved extra density, would not support the framework.

13.7 Matching core to exterior

A compact object of mass M with core (13.6) matched to a weak-field exterior (13.11) raises no new consistency issue beyond ordinary matching of f and f' at an intermediate radius. The cubic term is already negligible there if r_0 is microscopic. The entropic density $\rho_{\rm ent}$ of a single compact object cannot be quantified from the mutual-information tail alone. Deriving that density and comparing its field with the Newtonian field requires varying $U_{\rm ent}$ for a specified source configuration. Galactic flattening, if it occurs, is a collective effect of many baryonic parcels; it is not established here from the tail of a single compact core.

13.8 Exercises

1. **Einstein limits.** Recover (13.3) from (13.6) as $r_0 \to 0$ at fixed $r > 0$, and as $r \to \infty$ at fixed r_0.

2. **Core curvature.** For $\Lambda_{\rm eff} = 0$, compute R at $r = 0$ from (13.8). Show it is finite for $r_0 \neq 0$.

3. **Horizon condition.** With $\Lambda_{\rm eff} = 0$, discuss the number of positive roots of $f(r) = 0$ as a function of M/r_0. A graphical argument is enough.

4. **Poisson split.** Starting from (13.11) and (13.12), write an effective equation for Φ in terms of $\Phi_{\rm b}$ only. What operator acts on $\Phi_{\rm b}$?

5. **Flat-curve kinematics.** Suppose $M_{\rm b} + M_{\rm ent} = \kappa r$ at large r. Show v_c is constant. What density profile $\rho_{\rm ent}$ produces that mass growth, and is it integrable at infinity?

6. **Solar-System bound.** Estimate an upper bound on $\mu_{\rm ent}$ at $g_{\rm N} \sim 1\,{\rm cm\,s^{-2}}$ from the requirement that perihelion precession not be spoiled at current precision. Order of magnitude is enough.

7. **Illegal halo.** Write a spherical $\rho_{\rm halo} \propto 1/(r^2 + r_s^2)$ independent of $\rho_{\rm b}$. Which assumption of this chapter does it violate, and which theorem of Chapter 11 does it risk violating if it cannot be obtained from $U_{\rm ent}$?

13.9 Bridge to Chapter 14

The extra stress used in this chapter was the reversible, static piece $T_{\mu\nu}^{\rm ent,rev}$. The working d^{-4} drag introduced in Chapter 14 is a phenomenological addition; it is not derived here from time-dependent mutual information. Its microscopic dynamical origin and compatibility with a covariantly conserved stress remain unresolved and are part of Problem H.9. Chapter 15 integrates a reduced two-body laboratory model. Neither that laboratory nor the assumed drag substitutes for quadrupole radiation.

Chapter 14

Irreversible Entropic Stress and Dissipative Drag

Learning outcomes. After this chapter the reader should be able to:

1. distinguish the conservative force from the static U_{ent} from a proposed, phenomenological dissipative component;
2. state the passivity condition under which a retarded low-frequency response gives nonnegative dissipation;
3. implement and analyze the working law $\mathbf{F}_{\mathrm{diss}} = -\eta\mu\mathbf{v}/d^4$ as a reduced laboratory model, without treating it as a microscopic derivation;
4. compare that assumed channel with quadrupole gravitational-wave losses under stated model conditions;
5. distinguish the ODE's energy-bookkeeping identity from a derivation of mutual-information energy or covariant stress conservation.

Chapter 13 used only the static extra density. A time-dependent configuration can change $I(A : B)$, but that change alone does not establish irreversible work or a velocity-odd stress. Neither the first law nor the static mutual-information potential determines a retarded response by itself. The drag discussed in this chapter is a phenomenological component unless it is derived from microscopic dynamics, as required by Problem H.9.

14.1 Reversible versus irreversible

On a static spatial slice the mutual-information potential (6.9) is a function of the instantaneous regions A and B. Its metric variation is conservative: the associated force derives from U_{ent} and reverses under $\mathbf{v} \to -\mathbf{v}$.

If the regions change in time—because two cores inspiral, because a star is sheared across a tidal radius, because a condensate defect is stretched—then in general

$$\frac{\mathrm{d}}{\mathrm{d}t} I(A:B) \neq 0. \tag{14.1}$$

A changing $I(A:B)$ records a change in the correlations of the state. It does not, by itself, establish irreversible work or a force odd under time reversal.

Definition 14.1 (Phenomenological entropic-stress split)**.** For bookkeeping, write

$$T^{\mathrm{ent}}_{\mu\nu} = T^{\mathrm{ent,rev}}_{\mu\nu} + T^{\mathrm{ent,irrev}}_{\mu\nu}. \tag{14.2}$$

The first term denotes the contribution assigned to the static potential U_{ent}; the second is a proposed remainder in the effective description. This split alone does not establish that the remainder is dissipative or odd under time reversal. Those properties require a dynamical response calculation and an appropriate passivity condition.

Until that derivation is supplied, the split is phenomenological. Change or production of mutual information along a trajectory is not, on its own, evidence of irreversibility.

14.2 Worldline reduction

Consider two compact condensed cores with worldlines γ_A, γ_B, rest masses m_A, m_B, reduced mass $\mu = m_A m_B/(m_A + m_B)$, and separation four-vector $r^\mu = x^\mu_A - x^\mu_B$, $d = \sqrt{r^\mu r_\mu}$ in a local inertial frame. The entropic action along the worldlines is

$$S_{\mathrm{ent}} = -\int \mathrm{d}\tau\, U_{\mathrm{ent}}(d, \ell_A, \ell_B, \dot{d}), \tag{14.3}$$

where ℓ_A, ℓ_B are proper core radii and dots are proper time of the reduced motion. Varying with respect to x^μ_A produces the four-force (11.10).

At leading post-Newtonian order, with ℓ_A, ℓ_B frozen,

$$\mathbf{F}_A = -\nabla_{\mathbf{x}_A} U_{\mathrm{ent}}(d) - \eta^\mu(d, \dot{d})\, \mathbf{v} + O(c^{-2}). \tag{14.4}$$

The first term is Chapter 13. The second term is a phenomenological drag; it is not obtained by varying the static U_{ent}, and the coefficient η^μ is not derived here. A covariant reduction of such a term would have to satisfy Proposition 11.3, that is, there would have to be a microscopic entanglement energy E_{ent} such that

$$\frac{\mathrm{d}}{\mathrm{d}t}(E_{\mathrm{mech}} + E_{\mathrm{ent}}) = O(c^{-2}), \tag{14.5}$$

and the force on B would have to be equal and opposite up to radiation reaction of the tensor block.

The dissipative response is defined by a conditional Kubo prescription. At fixed separation $\mathbf{r}$, choose a stationary reference state of the microscopic Hamiltonian $H(\mathbf{r})$ and define the connected force operator $\delta\hat{F}_i = -\partial H/\partial r_i - \langle -\partial H/\partial r_i\rangle_{\mathbf{r}}$. For a perturbation $\delta H(t) = -\delta r_j(t)\,\delta\hat{F}_j$, the retarded response kernel is

$$\chi^R_{ij}(t) = \frac{i}{\hbar}\,\theta(t)\,\big\langle[\delta\hat{F}_i(t), \delta\hat{F}_j(0)]\big\rangle_{\mathbf{r}}, \qquad \chi^R_{ij}(\omega) = \int_{-\infty}^{\infty} e^{i\omega t}\chi^R_{ij}(t)\,\mathrm{d}t. \tag{14.6}$$

If the reference state is reciprocal and passive and the limit exists, the low-frequency dissipative tensor is

$$\Gamma_{ij} = \lim_{\omega\downarrow 0} \frac{\operatorname{Im}\chi^R_{(ij)}(\omega)}{\omega}, \qquad \mathbf{F}_{\mathrm{diss},i} = -\Gamma_{ij}v_j, \tag{14.7}$$

where $\chi^R_{(ij)}$ is the symmetric part and Γ_{ij} is positive semidefinite by passivity. This is a conditional prescription: the NAMCG force correlator, its stationary reference state, and its passivity have not been calculated. The causal family $\chi^R(\omega) = \chi_0/(1 - i\omega\tau)$, $\tau > 0$, has the same static value χ_0 for every τ but low-frequency slope $\tau\chi_0$, so a static susceptibility does not determine the dissipative coefficient. Positivity of $I(A:B)$ for each state likewise establishes neither an ordering between adiabatic and non-adiabatic mutual informations nor the sign of Γ; Problem H.9 states the calculation required.

14.3 The working exponent

Section 6.7 gave the conformal large-separation tail

$$I(A:B) \;\sim\; c_{p_*}\left(\frac{\ell_A\ell_B}{d^2}\right)^{p_*}, \qquad d \gg \ell_A, \ell_B, \tag{14.8}$$

with p_* and c_{p_*} and the associated assumptions as in (6.10); for a free massless scalar in $3+1$ dimensions $p_* = 2$ and the static tail falls as d^{-4}. This static scaling concerns mutual information only and does not determine the low-frequency dissipative response. The scalar relation

$$\Gamma(d) = \eta\,\frac{\mu}{d^4} \tag{14.9}$$

is therefore a working isotropic laboratory choice, not a deduction from the static tail or a calculation of the microscopic response tensor. The coefficient η has not been derived from a microscopic force-response function. The isotropic scalar form also requires the longitudinal and transverse components of Γ_{ij} to agree, or an explicit restriction to radial motion.

Assumption 14.2 (Working dissipative law)**.** In the reduced two-body laboratory of this book,

$$\mathbf{F}_{\rm diss} = -\eta\,\frac{\mu}{d^4}\,\mathbf{v}, \tag{14.10}$$

with $\eta \geq 0$ constant on a given inspiral and with $E_{\rm ent}$ defined so that (14.5) holds identically.

Assumption 14.2 is not a multipole theorem. A different static tail of $I(A:B)$ does not, by itself, imply a different drag exponent. The response exponent remains undetermined until the dynamical calculation in Problem H.9 is supplied. Chapter 16 treats the working exponent as falsifiable; Chapter 15 integrates the specified laboratory law, not a derived microscopic response, and no other drag.

For the laboratory bookkeeping, define the accumulated work

$$W_{\rm diss}(t) = \int_0^t \mathbf{F}_{\rm diss}\cdot\mathbf{v}\,{\rm d}t'. \tag{14.11}$$

Then set

$$E_{\rm ent}(t) = -W_{\rm diss}(t) + E_{\rm ent}(0). \tag{14.12}$$

This definition makes $E_{\rm mech}+E_{\rm ent}$ constant at the retained order when $\dot{E}_{\rm mech} = \mathbf{F}_{\rm diss}\cdot\mathbf{v}$. It is bookkeeping for the reduced laboratory, not a derivation of an entanglement energy from $I(A:B)$. The integrator in Chapter 15 displays $E_{\rm mech}(t) - W_{\rm diss}(t)$ as this conserved combination.

14.4 Comparison with quadrupole losses

The Isaacson flux of the tensor block for a weak, slow binary is the Peters–Mathews power

$$P_{\rm GW} = \frac{32}{5} G_{\rm eff}\mu^2 d^4\Omega^6 \langle \text{inclination factors}\rangle \tag{14.13}$$

in the usual quadrupole units, up to the precise harmonic decomposition. The entropic power is

$$P_{\rm ent} = -\mathbf{F}_{\rm diss}\cdot\mathbf{v} = \eta\mu\frac{v^2}{d^4}. \tag{14.14}$$

For a circular Newtonian orbit, $v^2 = G_{\rm eff}M/d$ and $\Omega^2 = G_{\rm eff}M/d^3$, so

$$\frac{P_{\rm ent}}{P_{\rm GW}} \;\sim\; \frac{\eta}{G_{\rm eff}^3 M^2\mu}\left(\frac{d}{G_{\rm eff}M}\right)^k \tag{14.15}$$

with a positive power of d that makes the ratio *grow* toward wide separations if η is held fixed. That conclusion is an artefact of treating η as constant down to laboratory distances. If η is instead suppressed at wide separations, the growth need not occur;

this book does not calculate that dependence, and a sufficiently small η is required for consistency with Solar-System tests. The intended domain of (14.10) is:

- late inspirals of two compact condensed cores, where d is not astronomically large compared with $\ell_A + \ell_B$;
- tidal disruption events, where a stellar core is sheared at $d \sim r_{\rm t}$ and $\dot{d}$ is not small;
- numerical laboratories that isolate the extra channel.

In LIGO-band vacuum inspirals of two objects whose condensate cores are hidden behind horizons, quadrupole radiation remains the default dissipative channel. NAMCG does not replace the Peters formula by (14.10) in that regime unless η is independently constrained to be large there. It is not.

14.5 Tidal disruption

A star of radius R_* and mass m_* approaching a compact object of mass M is disrupted at

$$r_{\rm t} \sim R_* \Big(\frac{M}{m_*}\Big)^{1/3}. \tag{14.16}$$

Across that radius the baryonic body ceases to be a single region A and becomes a collection of debris parcels. The mutual information between the compact core and the star changes over a dynamical time

$$t_{\rm dyn} \sim \sqrt{r_{\rm t}^3 / G_{\rm eff} M}. \tag{14.17}$$

The non-adiabatic character of this process may motivate testing the phenomenological drag model near $r_{\rm t}$, but it does not establish a dissipative response or its magnitude. The existence and size of any additional drag require the microscopic dynamics, reference state, and response calculation specified in Problem H.9. Observable consequences such as changes to fallback rates or extra heating are therefore conditional phenomenological possibilities, not predictions derived in this chapter. They belong to Chapter 16 and require hydrodynamics that this text does not contain.

The book does not simulate debris streams.

14.6 Illegal and legal integrators

An integrator that evolves

$$\mu \ddot{\mathbf{r}} = -\frac{G_{\rm eff} M \mu}{d^2} \hat{\mathbf{r}} - \gamma \mathbf{v} \tag{14.18}$$

with constant γ and no $E_{\rm ent}$ is not a NAMCG calculation, even if its trajectories look like inspirals. It dissipates energy into nowhere.

An integrator that evolves

$$\mu\ddot{\mathbf{r}} = -\nabla U_{\rm N}(d) - \nabla U_{\rm ent}^{\rm rev}(d) - \eta\frac{\mu}{d^4}\mathbf{v} \tag{14.19}$$

and accumulates (14.11) is the reduced laboratory implementation of Assumption 14.2. Accumulating $W_{\rm diss}$ makes $E_{\rm mech} - W_{\rm diss}$ constant for the modeled ODE, up to numerical error. This bookkeeping identity does not establish a microscopic mutual-information energy or prove that the ODE is the reduction of a covariantly conserved NAMCG stress. Chapter 15 implements (14.19) with a symplectic Verlet step on the conservative forces and an explicit treatment of the drag. The gravitational constant used there is a toy value, not a physical $G_{\rm eff}$; that matching remains open in Problem F.

14.7 Exercises

1. **Passivity and the sign of drag.** For the reciprocal low-frequency response tensor $\Gamma_{ij}(\mathbf{r}; \ell_A, \ell_B)$ defined in Problem H.9, state the passivity condition. Show that positive semidefiniteness implies $F_i^{\rm diss} v_i \le 0$. Explain why $I(A:B) \ge 0$ alone implies neither an ordering between adiabatic and non-adiabatic states nor a positive drag coefficient. Which of the assumptions needed for this Kubo conclusion have actually been established for NAMCG?

2. **Static versus dynamical scaling.** Explain why a power law for static $I(A:B)$ does not determine the separation dependence of the low-frequency drag tensor. Identify the dynamical response data needed to calculate that dependence.

3. **Energy identity.** For (14.10) with circular motion at instantaneous radius $d(t)$, compute $P_{\rm ent}$. Compare with $\mathrm{d}U_{\rm N}/\mathrm{d}t$ along a Peters inspiral and estimate a separation where the two powers could be comparable for a chosen η.

4. **Equal and opposite.** Show that (14.10), written in the reduced-mass frame, corresponds to equal-and-opposite three-forces on A and B. What radiation-reaction term would have to be added to break that balance, and which block of $T_{\mu\nu}^{\rm total}$ supplies it?

5. **Constant-γ counterexample.** Take (14.18) and send $d \to \infty$. What happens to the dissipated energy per unit time, and why is that incompatible with a mutual-information tail that vanishes at infinity?

6. **Tidal time.** Evaluate (14.17) for a solar-type star at the tidal radius of a $10^6 M_\odot$ compact object. Why is the adiabatic approximation of Chapter 13 unreliable there?

14.8 Bridge to Chapter 15

The ordinary-differential laboratory is now specified: Newtonian gravity, optional reversible extra potential, drag (14.10), and an accumulated work (14.11). Chapter 15 integrates that system with a symplectic Verlet step, plots the trajectory and the energy balance, and states, in the first paragraph, that the figure is a reduced model. The Python code that produces the figure lives in `code/` and is not a derivation of Einstein gravity. After the figure, Chapter 16 collects every named assumption in the book and lists the observations that would kill the framework.

Chapter 15

A Reduced Inspiral Laboratory

Learning outcomes. After this chapter the reader should be able to:

1. write the ordinary-differential system of Assumption 14.2 in a form a computer can integrate;

2. implement a velocity-Verlet step on the conservative force with an explicit drag kick;

3. accumulate the dissipated work W_{diss} and verify the bookkeeping identity numerically;

4. read a trajectory plot as a laboratory output, not as a derivation of general relativity;

5. enter Chapter 16 with every phenomenological extra in the book already named.

This chapter is a reduced Newtonian laboratory. It does not derive Einstein gravity, does not compute G_{eff} from Theorem 8.2, and does not replace quadrupole radiation. It integrates Assumption 14.2 and checks that the energy bookkeeping identity of Chapter 14 can be made visible on a plot. That identity is not a check of Proposition 11.3.

15.1 The ordinary-differential system

Work in the reduced-mass frame with relative coordinate $\mathbf{r}(t)$ and $d = |\mathbf{r}|$. The laboratory equation is (14.19) without a reversible extra potential, which is optional at this order:

$$\mu\ddot{\mathbf{r}} = -\frac{G_{\mathrm{toy}} M \mu}{d^3}\mathbf{r} - \eta\frac{\mu}{d^4}\dot{\mathbf{r}}. \tag{15.1}$$

Here G_{toy} is a numerical constant chosen so that an orbit is visible in a few thousand steps. It is not G_{N} and not G_{eff}. The source-note values used below are

$$G_{\mathrm{toy}} = 36, \qquad \eta = 0.006, \qquad M = 1, \qquad \mu = 1 \tag{15.2}$$

in unspecified code units, together with a bound initial condition that would be an ellipse if $\eta = 0$.

Mechanical energy and angular momentum of the conservative problem are

$$E_{\mathrm{mech}} = \frac{1}{2}\mu\dot{\mathbf{r}}^2 - \frac{G_{\mathrm{toy}} M \mu}{d}, \qquad \mathbf{L} = \mu\, \mathbf{r} \times \dot{\mathbf{r}}. \tag{15.3}$$

The dissipated work is

$$W_{\mathrm{diss}}(t) = \int_0^t \Big(-\eta \frac{\mu}{d^4}\dot{\mathbf{r}}\Big) \cdot \dot{\mathbf{r}}\, dt' = -\int_0^t \eta\mu \frac{v^2}{d^4}\, dt'. \tag{15.4}$$

The reduced ODE implies that

$$\mathcal{E}(t) \equiv E_{\mathrm{mech}}(t) - W_{\mathrm{diss}}(t) \tag{15.5}$$

is constant up to integrator error. The combination $\mathcal{E}$ is the laboratory stand-in for $E_{\mathrm{mech}} + E_{\mathrm{ent}}$ with $E_{\mathrm{ent}} = -W_{\mathrm{diss}}$ accumulated from the initial time. This is bookkeeping: it does not construct a microscopic E_{ent}. If $\mathcal{E}$ drifts systematically, the implementation is inconsistent with the bookkeeping identity of Section 14.6.

15.2 Integrator

A velocity-dependent force is not Hamiltonian. A symplectic Verlet step is still used on the conservative piece because it conserves a nearby energy of the $\eta = 0$ problem over long times. The drag is applied as an explicit kick.

Let $\mathbf{a}_{\mathrm{N}}(\mathbf{r}) = -G_{\mathrm{toy}} M \mathbf{r}/d^3$ and $\mathbf{a}_{\mathrm{d}}(\mathbf{r}, \mathbf{v}) = -\eta \mathbf{v}/d^4$. One step of size Δt is

$$\begin{aligned} \mathbf{v}_{1/2} &= \mathbf{v}_n + \tfrac{1}{2}\Delta t\, \big(\mathbf{a}_{\mathrm{N}}(\mathbf{r}_n) + \mathbf{a}_{\mathrm{d}}(\mathbf{r}_n, \mathbf{v}_n)\big), \\ \mathbf{r}_{n+1} &= \mathbf{r}_n + \Delta t\, \mathbf{v}_{1/2}, \\ \mathbf{v}_{n+1} &= \mathbf{v}_{1/2} + \tfrac{1}{2}\Delta t\, \big(\mathbf{a}_{\mathrm{N}}(\mathbf{r}_{n+1}) + \mathbf{a}_{\mathrm{d}}(\mathbf{r}_{n+1}, \mathbf{v}_{1/2})\big). \end{aligned} \tag{15.6}$$

The last evaluation of the drag uses $\mathbf{v}_{1/2}$ rather than an implicit $\mathbf{v}_{n+1}$. The resulting error is $O(\Delta t^2)$ in the dissipative piece, which is adequate for the plots below. The work increment on the step is

$$\Delta W = \mathbf{F}_{\mathrm{diss}}(\mathbf{r}_n, \mathbf{v}_n) \cdot \mathbf{v}_n\, \Delta t, \tag{15.7}$$

or a trapezoidal average of the two endpoints. Either choice converges to (15.4).

15.3 Initial data

A bound ellipse of the conservative problem is specified by

$$\mathbf{r}(0) = (d_0, 0), \qquad \mathbf{v}(0) = (0, v_0), \tag{15.8}$$

with $v_0^2 < 2G_{\mathrm{toy}}M/d_0$. The runs shown use a nearly circular start,

$$d_0 = 1.0, \qquad v_0 = \sqrt{G_{\mathrm{toy}}M/d_0}, \tag{15.9}$$

in code units, together with $\Delta t = 10^{-3}$ and $N = 1.2 \times 10^4$ steps. Those numbers are inherited from the corrected notebook `dmcg5.py`. They are not physical units.

15.4 What the figures are allowed to show

Two panels are produced.

Trajectory. The curve $\mathbf{r}(t)$ in the plane. For $\eta = 0$ it is a closed ellipse (a circle, for (15.9)). For $\eta > 0$ the radius shrinks. That shrinkage is the laboratory image of P_{ent} in (14.14). It is not a plot of a relativistic inspiral and must not be overplotted on LIGO waveforms.

Energy. Three curves: $E_{\mathrm{mech}}(t)$, $W_{\mathrm{diss}}(t)$, and $\mathcal{E}(t)$. The first falls, the second is negative and falls, and the third is flat within integrator noise if the implementation is correct.

If a previously generated PNG from the notes is used instead, it belongs at this caption and nowhere else in the book.

15.5 Code

The integrator lives in `code/namcg_inspiral.py`. A minimal version sufficient to produce Figure 15.1 is the following.

Listing 15.1: Python implementation of the reduced inspiral laboratory velocity-Verlet integrator.

```
import numpy as np
import matplotlib.pyplot as plt

def run_namcg_laboratory(
```




Figure 15.1: Laboratory trajectory and energy balance for (15.1) with the toy parameters (15.2). The figure is a reduced model.

```
    G_toy=36.0,
    eta=0.006,
    M=1.0,
    mu=1.0,
    d0=1.0,
    dt=1.0e-3,
    num_steps=12000,
):
    r = np.array([d0, 0.0])
    v = np.array([0.0, np.sqrt(G_toy * M / d0)])
    traj = np.zeros((num_steps + 1, 2))
    E_mech = np.zeros(num_steps + 1)
    W_diss = np.zeros(num_steps + 1)
    traj[0] = r

    def a_N(r):
        d = np.linalg.norm(r)
        return -G_toy * M * r / d**3

    def a_d(r, v):
        d = np.linalg.norm(r)
        return -eta * v / d**4

    def mechanical_energy(r, v):
        d = np.linalg.norm(r)
        return 0.5 * mu * np.dot(v, v) - G_toy * M * mu / d

    E_mech[0] = mechanical_energy(r, v)
    for n in range(num_steps):
```

```
        F_d = mu * a_d(r, v)
        W_diss[n + 1] = W_diss[n] + np.dot(F_d, v) * dt
        v_half = v + 0.5 * dt * (a_N(r) + a_d(r, v))
        r = r + dt * v_half
        v = v_half + 0.5 * dt * (a_N(r) + a_d(r, v_half))
        traj[n + 1] = r
        E_mech[n + 1] = mechanical_energy(r, v)
    return traj, E_mech, W_diss

def plot_laboratory(traj, E_mech, W_diss, filename):
    E_bal = E_mech - W_diss
    fig, axes = plt.subplots(1, 2, figsize=(8.4, 3.6))
    axes[0].plot(traj[:, 0], traj[:, 1], lw=0.8)
    axes[0].set_aspect("equal")
    axes[0].set_xlabel("x")
    axes[0].set_ylabel("y")
    axes[0].set_title("Laboratory trajectory")
    t = np.arange(len(E_mech))
    axes[1].plot(t, E_mech, label=r"$E_{\mathrm{mech}}$")
    axes[1].plot(t, W_diss, label=r"$W_{\mathrm{diss}}$")
    axes[1].plot(t, E_bal, label=r"$\mathcal{E}$")
    axes[1].set_xlabel("step")
    axes[1].set_title("Energy balance")
    axes[1].legend(fontsize=8)
    fig.tight_layout()
    fig.savefig(filename)
    plt.close(fig)

if __name__ == "__main__":
    traj, E_mech, W_diss = run_namcg_laboratory()
    plot_laboratory(
        traj, E_mech, W_diss,
        "figs/namcg_gr_simulation_corrected.pdf",
    )
```

Run the file from the project root after creating `figs/`. The script writes a vector PDF suitable for the print interior. A PNG of the same figure, if already produced by an earlier notebook, may be kept for draft viewing but should not be the print original.

15.6 How to misread the figure, and how not to

The shrinking orbit is not a measurement of η in nature. The value 0.006 was chosen so that decay is visible in 1.2×10^4 steps. This book does not calculate η, so no

realistic value at $d \gg \ell_A + \ell_B$ is established here.

The flatness of $\mathcal{E}(t)$ is not a proof of Theorem 11.1. It is a check of the reduced ODE's bookkeeping identity, not of a covariant reduction. If $\mathcal{E}$ is flat and the force law is not (14.10), the check has been passed for the wrong theory.

Comparing the laboratory decay time with a LIGO inspiral time is not a test of NAMCG. The laboratory has no gravitational waves, no horizons, and no relativity. Section 14.4 already stated the regime in which quadrupole losses dominate.

15.7 Exercises

1. **Zero drag.** Set $\eta = 0$ and confirm that E_{mech} and $|\mathbf{L}|$ are constant at the level of Verlet error. What is the size of that error as a function of Δt?

2. **Balance.** With $\eta = 0.006$, plot $\mathcal{E}(t)$ for $\Delta t \in \{10^{-3}, 5 \times 10^{-4}, 2 \times 10^{-3}\}$. Does the drift shrink with Δt?

3. **Illegal drag.** Replace d^{-4} by a constant γ and rerun. Track W_{diss} as d decreases. Why is the late behaviour a counterexample to Exercise 5 of Chapter 14?

4. **Angular momentum.** The drag (14.10) is antiparallel to $\mathbf{v}$. Show that it exerts a torque and compute $\mathrm{d}\mathbf{L}/\mathrm{d}t$. Confirm the identity on the numerical trajectory.

5. **Units.** Restore SI symbols in (15.1) and assign dimensions to η. What combination of ℓ_A, ℓ_B, c and a response time could supply those dimensions?

15.8 Bridge to Chapter 16

Every extra structure used in applications now has a name: the regular core (13.6), the baryon-tracking extra density of Assumption 13.2, the working drag of Assumption 14.2, and the laboratory of this chapter. Chapter 16 collects the full list of assumptions from Chapters 3–15, restates the derivation chain of Chapter 1 as a single page, and writes the observations that would kill the framework rather than merely constrain η.

Chapter 16

Assumptions, Predictions, and Falsifiers

Learning outcomes. After this chapter the reader should be able to:

1. restate the derivation chain of Chapter 1 with the theorem numbers now attached;
2. separate assumptions that kill NAMCG if they fail from coefficients that can be adjusted;
3. name observations that would refute the framework rather than merely bound η or r_0;
4. refuse to treat a successful laboratory plot as evidence for Theorems 7.3–12.4;
5. enter Chapter 17 knowing which open calculations are optional refinements and which are critical-path derivations.

A framework that cannot say what would kill it is not a framework. This chapter is the scorecard promised in Section 1.4.

16.1 The chain, with theorem numbers

The argument of the book is the following sequence, and no shorter sequence.

1. A matrix order parameter $\Phi^I{}_A$ with potential (3.7) admits a frame-locking vacuum (Assumptions 3.5, 3.6).
2. A composite vierbein and metric exist in the condensed phase (Repair II, (4.7); Definition 4.3, Assumption 4.4).
3. Fluctuations split into ψ, A_μ, ω_{AB}, and $T_{\mu\nu}$ (Chapter 5).

4. Entanglement equilibrium on emergent diamonds is a property of the condensate vacuum (Assumption 6.1).

5. A Fierz–Pauli tensor block is incompatible with that equilibrium unless $m_T = 0$ (Theorem 7.3).

6. The Euclidean curvature coefficient obeys the conditional, truncated heat-kernel identity of Theorem 8.2 for operators built on the emergent metric (Assumption 8.4); physical matching requires specified operators, weights, thresholds, continuation, and scheme (Problem F).

7. Four-derivative metric operators are treated as EFT vertices, not extra poles, under Assumption 9.2; this is not a claim about a non-perturbative completion.

8. Under the local Lorentz gauge assumption, the six Goldstone orbit variables ω_{AB} are removable in unitary gauge (Theorem 10.3); the diagonal connection's dynamics remain open (Problem D).

9. The conformal mode is heavy (Assumption 7.5).

10. The total stress, including $T^{\text{ent}}_{\mu\nu}$, is covariantly conserved (Theorem 11.1).

11. Λ_{eff} is set by boundary entanglement after an ultraviolet subtraction (Theorem 12.4).

Items 5, 6, 8, 10, and 11 are theorems *inside* the list of assumptions that precede them. If an earlier assumption fails, the later theorem is idle. Phenomenology in Chapters 13–15 sits after the whole list. It cannot substitute for any line.

16.2 Inventory of named assumptions

The named hypotheses of Parts II–V are collected here. "Load- bearing" means that failure returns the infrared theory to analog gravity, scalar-tensor gravity, or an inconsistent extra force. "Coefficient" means that failure changes a number or a regime of validity without destroying the graviton.

Tag	Kind	Content
3.1	setup	$g^{(0)}$ is a scaffold
3.5	load-bearing	vacuum $\langle \Phi^I{}_A \rangle = v\delta^I{}_A$ exists
3.6	load-bearing	commutator / covariant-derivative dictionary
4.4	load-bearing	condensed frame is non-degenerate
4.6	load-bearing	signature is Lorentzian
5.1	coefficient	extra internal scalars are heavy
6.1	load-bearing	microscopic entanglement equilibrium
7.5	load-bearing	conformal mode is gapped
8.1	setup	infrared operator is Laplace type
8.4	load-bearing	fluctuations propagate on the emergent metric
9.2	load-bearing	C^2, R^2 are vertices, not poles
9.3	model	cubic truncation in the static spherical sector
10.1	load-bearing	local $\mathrm{SO}(3,1)$ is gauged
12.2	load-bearing	local boundary entropy density exists
12.3	load-bearing	boundary stationarity sets Λ_{eff}
13.2	coefficient	static extra density tracks baryons; disks use $\nu_{\surd}$ of Appendix D
13.3	coefficient	external field saturates ν at $\nu(g_{\mathrm{ext}})$, not $\nu(g_{\mathrm{N}})$
14.2	coefficient	working laboratory law $\mathbf{F}_{\mathrm{diss}} \propto -\mathbf{v}/d^4$

Two comments on the table.

First, "coefficient" is not a synonym for "unimportant." Assumption 13.2 is how galaxies enter the book. If it fails, galactic claims fail. The graviton claims need not.

Second, Assumption 3.6 is the most important unproved microscopic bridge. If the infrared dictionary from Presentation A to Presentation B cannot be constructed, the heat kernel of Chapter 8 is a calculation in a different theory than the matrix action of Chapter 3.

Here "load-bearing" classifies an assumption as a premise in the derivation chain; it does not classify the priority of the open problem associated with that premise. Chapter 17 lists the critical-path problems for a completed matrix-derived derivation. Assumptions 3.6, 4.6 and 9.2 are the targets of critical-path Problems A, E and N, and frame existence is part of Problem D1. Assumption 8.4 is a premise of Theorem 8.2 that Problem F must discharge. An assumption may remain in a scope-limited model, but that does not derive or close the corresponding problem.

16.3 What would kill the framework

The following outcomes are incompatible with NAMCG as written, not merely with a choice of η.

A light extra tensor or a ghost below the cutoff. If precision gravity, pulsar timing, or gravitational-wave propagation requires a second massless or light spin-2 field, or a ghost pole below Λ_{UV}, Theorem 7.3 plus Assumption 9.2 has failed.

A light conformal partner. If Solar-System or binary-pulsar tests require a Brans–Dicke parameter that cannot be produced by a heavy ψ, Assumption 7.5 has failed and the infrared theory is scalar-tensor gravity. That is the ceiling of Chapter 1, reopened.

Non-conservation of the extra stress. If an extra acceleration is demanded by data and cannot be written as $\nabla^{\mu}T^{\mathrm{ent}}_{\mu\nu}$ with Theorem 11.1 intact—a drag into nowhere, a preferred-frame force that cannot be improved to a tensor—then the geometric reading has failed. Analog dissipation remains possible. Geometrogenesis of the extra channel has not occurred.

Failure of Lorentz locking. If the unitary-gauge removal of the six Goldstone orbit variables ω_{AB} fails under the theorem's assumptions, Theorem 10.3 has failed. No later chapter can hide that failure.

A cutoff that cannot serve Newton and Lorentz together. If the only way to obtain $G_{\mathrm{eff}} = G_{\mathrm{N}}$ forces Λ_{UV} into a regime where Lorentz-violating $\Theta^{\mu\nu}$ corrections are already excluded by current bounds, the coefficient map and the kinematic algebra cannot coexist.

An induced Einstein–Hilbert term of the wrong sign. If a complete heat-kernel evaluation on the locked bundle produces a repulsive Einstein term that cannot be flipped by fermions or by the Lorentzian continuation, the infrared force is not gravity.

None of these killers is a rotation curve. None of them is a pretty laboratory inspiral.

16.4 What would only constrain a coefficient

The following outcomes bound parameters or restrict regimes. They do not, by themselves, refute Theorems 7.3–12.4.

- SPARC ranking of a global $\nu(g_{\rm N})$ against simple MOND and the published RAR (Appendix D). Frozen $\nu_{\vee}$ at $g_\dagger$ is equivalent to the RAR and about 0.2% behind simple MOND in χ^2; baryons-only is not competitive. That is a passed coefficient check on disks, not a theorem. A later sample that rejects every high-g-Newtonian, low-g-square-root boost would restrict Assumption 13.2 to a smaller class of systems, or kill the galactic claim only.

- Wide binaries in the solar neighbourhood. Isolated $\nu_{\vee}(g_{\rm N})$ predicts $\gamma \approx 4$ at $g_{\rm N} = 10^{-11}\,{\rm m\,s^{-2}}$. Assumption 13.3 predicts $\gamma \to \nu_{\vee}(g_{\rm gal}) \approx 1.1$. A quality-controlled Gaia sample consistent with Newton constrains the isolated reading and is compatible with Assumption 13.3. A surviving $\gamma \approx 1.4$ is not explained by that saturation. Neither outcome touches Theorems 7.3–12.4.

- Cluster lensing or the bullet-cluster mass offset. Same status: the extra tensor may fail as a complete replacement for particulate dark matter while the graviton derivation stands.

- Absence of an extra drag in LIGO-band vacuum inspirals. Expected if η is negligible at those separations; Section 14.4 already said so.

- Absence of a visible core scale r_0 in EHT images or ringdowns. Expected if r_0 is microscopic; bounds $\beta_{\rm ECG}$.

- A measured $\Lambda_{\rm eff}$ that does not match a particular guess for $S_{\rm ent}^{\rm (IR)}$. Theorem 12.4 changes the origin of the scale. It does not compute the number.

- A different static mutual-information tail does not, by itself, determine the drag exponent. The response exponent must be calculated from the dynamical force response. The d^{-4} law remains a falsifiable laboratory assumption (Assumption 14.2).

16.5 Observational checklist

A short list of current and near-term measurements, with the question each actually asks of the book.

Solar System and pulsar timing. Do they tolerate a massive ψ, and, if torsion is massive at some as-yet-undetermined scale (Problem D), that torsion mode, along with a $\mu_{\rm ent}$ negligible at high acceleration? Whether torsion is massive is a condition to test, not an established result. If the scalar or tracking assumption fails, the corresponding Assumption 7.5 or 13.2 fails. If they survive, nothing about diamonds has been tested.

Gravitational waves. Is the propagating mode two TT polarizations at the speed of light, with no extra light tensor? That is a test of Theorem 7.3 plus Assumption 9.2, and it is already passed at present precision. A measured extra polarization would be a killer. A measured inspiral-time deviation at LIGO frequencies is only a bound on η.

Black-hole images and ringdowns. Is there a visible deviation from Kerr at the photon ring or in quasi-normal modes? That bounds r_0. A perfectly Kerr object is compatible with microscopic r_0.

Galaxies, wide binaries, and clusters. Do disks accept a single $\nu_{\surd}(g_{\mathrm{N}})$ with $\nu \to 1$ at Solar-System acceleration? Appendix D says yes, at RAR accuracy. Do solar-neighbourhood wide binaries accept the isolated evaluation of that law? No: Assumption 13.3 saturates them at $\nu_{\surd}(g_{\mathrm{gal}})$. Do merging clusters still require a stress that does not track baryons? That remains an open constraint on Assumption 13.2, not on Theorem 7.3, and the external field is not a substitute for the lensing offset.

Cosmology. Is Λ_{eff} constant enough for a Friedmann equation, and is the ultraviolet quartic absent from the expansion history? This tests Theorem 12.4 as a mechanism, not as a number. A detected bulk Λ_{UV}^4 vacuum energy would kill the HBEV subtraction.

Tidal disruptions. Is there extra heating or altered fallback that hydrodynamics plus GR tides cannot account for? A possible window on Assumption 14.2. A null result bounds η at r_{t} and nothing else.

16.6 Claims the book has not made

The following sentences do not appear as theorems in this text, and must not appear on a cover, in an abstract, or in a chapter heading as if they did.

- Dark matter has been explained.
- The cosmological-constant problem has been solved numerically.
- Singularities have been proved absent in dynamical collapse.
- LIGO inspirals are driven by a $1/d^4$ drag.
- Lorentzian signature has been derived from a Euclidean matrix model.
- Presentation A has been reduced to Presentation B without Assumption 3.6.
- The rationals A, B, C in the coefficient map are unique.

- Isolated $\nu_{\surd}$ has been derived, and wide binaries have been predicted without an external field.

A sale-ready monograph that prints any of those sentences as results has left this book and entered advertising.

16.7 How to use Chapter 15 in public

Figure 15.1 may be shown as a numerical bookkeeping check of an assumed drag in a Newtonian laboratory. It may not be shown as "NAMCG predicting an inspiral." The caption must retain the words "reduced model." The toy value $G_{\mathrm{toy}} = 36$ must not be silently replaced by G_{N} in a talk slide.

16.8 Exercises

1. **Sort the list.** Take any popular article on emergent gravity and assign each of its claims to "killer," "coefficient," or "not a claim of this book."

2. **One-page chain.** Write the eleven-line chain of Section 16.1 from memory, without looking. Where you stall is the section to reread.

3. **Design a killer.** Propose one measurement, not already listed, that would refute Theorem 7.3 without passing through η. State the null and the fatal outcome.

4. **Design a non-killer.** Propose one measurement that looks dramatic and would only bound r_0.

5. **Abstract hygiene.** Rewrite a five-sentence abstract of this book that contains no sentence from Section 16.6.

16.9 Bridge to Chapter 17

The killers and the coefficients are now distinguished. What remains is a research agenda: the calculations that would turn Assumption 3.6 into a lemma, recompute A, B, C, evaluate $S_{\mathrm{ent}}^{(\mathrm{IR})}$, derive μ_{ent} from $I(A:B)$, including the external-field argument of Assumption 13.3, and decide whether Lorentzian signature can be more than Assumption 4.6. Chapter 17 writes that agenda at the thickness of a reader who has read the book and wants to work, not at the thickness of a grant advertisement.

Chapter 17

Open Calculations and a Research Agenda

Learning outcomes. After this chapter the reader should be able to:

1. distinguish a load-bearing assumption from a critical-path problem and an optional refinement;
2. pick a first calculation that actually moves the framework, rather than a calculation that only decorates Chapter 15;
3. state what would have to be true before NAMCG may be described as finished rather than as a programme;
4. use the appendices as tools, not as a substitute for the open problems listed here.

Chapter 16 said what would kill the theory. This chapter says what would complete it. The two lists are not the same.

17.1 What is closed conditionally inside the book

The following are closed *as arguments from named hypotheses to named conclusions.* They are not closed as derivations from empty premises.

- Composite metric from a non-degenerate condensate frame (Repair II; Definition 4.3, Assumption 4.4). The Repair I formula of Definition 4.1 is not a constructed frame (Problem C).
- Algebraic split of $\delta\Phi_{\mu\nu}$ (Chapter 5).
- $m_T = 0$ from entanglement equilibrium plus Fierz–Pauli (Theorem 7.3).

- The conditional, truncated heat-kernel identity for specified sectors and scheme (Theorem 8.2); physical G_{eff} remains open in Problem F.
- Lorentz locking in unitary gauge (Theorem 10.3).
- Conservation of the total stress (Theorem 11.1).
- Replacement of a bulk quartic by an infrared boundary datum (Theorem 12.4).
- A legal Newtonian laboratory for Assumption 14.2 (Chapter 15).

A reader who has worked every exercise through Chapter 16 can reproduce that list. Reproducing it is not the same as removing the hypotheses.

17.2 Status vocabulary and completion labels

Status words are *OPEN* and *CLOSED CONDITIONALLY*. Closed conditionally means proved from named hypotheses to named conclusions; it does not remove those hypotheses. Completion words are *DERIVED* and *PROGRAMME-CONDITIONAL*. Derived means that the problem's target has no remaining assumed premise in the claimed chain. Programme-conditional means an assumption, input, reduced scope, or unresolved branch remains.

> Keeping an assumption, treating a quantity as an input, narrowing the scope, or choosing a scope that leaves a branch unresolved documents a limitation and keeps the programme conditional. It does not close the problem or satisfy the finished-derivation criterion of Section 17.7. Closure of a critical-path problem means DERIVED. A reduced-scope result is stated as such and the problem stays OPEN. A critical-path problem is met for the matrix-derived label only by a derivation from the matrix theory; an external justification, model input, imposed premise, or reduced scope supports at most a scope-limited Presentation-B model.

OPTIONAL means that a problem may remain open within a stated scope provided the claim given up is stated. Optional does not mean solved. Priority words are CRITICAL PATH and OPTIONAL. In Chapter 16, a *load-bearing assumption* is a premise used by the chain. The critical-path problems are those listed in Section 17.7. For the matrix-derived label, each load-bearing assumption must be discharged by one of them; a scope-limited model may retain it.

Label	Meaning
Completed infrared derivation (matrix-derived)	The only label for meeting the criterion in Section 17.7; every critical-path problem listed there is DERIVED for NAMCG from the matrix theory, so no load-bearing assumption of Chapter 16 remains assumed in their chains.
Scope-limited Presentation-B infrared model	A separately stated model with its imposed premises named, including any retained load-bearing assumption such as Assumption 3.6, 4.6 or 9.2. It is not matrix-derived, does not close critical-path problems, and does not satisfy the criterion of Section 17.7.
Programme	Any other status; one or more critical-path premises or targets remain unresolved.

17.3 Critical-path problems I: B, C, and D

The critical-path problems are those listed in Section 17.7. Problems B, C and D are treated here. The remaining critical-path problems, together with the optional problems H, I and J, are treated in Section 17.4. The order of work in Section 17.6 follows dependencies, not project size.

Problem B: entanglement equilibrium from the path integral. *CRITICAL PATH; OPEN.* The fixed-algebra first-law identity is established, but not the required stationarity. Assumption 6.1 is the physical engine of Theorem 7.3. Derive $\delta S_D = 0$ at fixed volume as a Ward identity of the microscopic action (3.6), or of (3.4) after Problem A, without importing Jacobson's hypothesis from a theory that already has a metric. A replica computation on small diamonds, with the polar vacuum as background, is the obvious route. A result that holds only for conformal fluctuations must be flagged as such, because the tensor block is not conformal. Closure requires fixed-volume variations of the tensor (non-conformal) block from the microscopic action, a specified map $g^{(0)} \mapsto g$, and no use of an already-existing metric. A conformal-only derivation is a sub-case and leaves B OPEN.

Problem C: the conformal mode. *CRITICAL PATH; OPEN.* The radial potential curvature is known, but the physical scalar operator and coupling are not. Assumption 7.5 is the remaining scalar-tensor ceiling. Compute the full quadratic scalar operator on the locked vacuum in a constructed frame: the canonically normalised kinetic coefficient of the pure-trace scalar (including the induced Einstein–Hilbert

contribution), its mixing with $T_{\mu\nu}$, its non-minimal curvature coupling, and its matter coupling. The potential (3.7) alone fixes only the radial curvature λv^2. From the operator, produce the pole mass, or a ratio m_ψ/v, together with the coupling that can be compared with Cassini bounds. If m_ψ is not parametrically of order $\sqrt{\lambda}\,v$, the assumption is false and the infrared theory must be rewritten as scalar-tensor NAMCG. Closure requires the operator, scalar coupling, and phenomenological comparison that define and test either the safe-mass or scalar-tensor outcome. Declaring a scalar-tensor scope without those data leaves Problem C open and the result programme-conditional.

Lemma (tree-level radial mass). On the locked vacuum $\langle \Phi^I{}_A \rangle = v\delta^I{}_A$, write the polar field $\Phi = (v+\sigma)\Lambda$ with $\Lambda^\dagger\Lambda = \mathbb{I}$ and $\langle\sigma\rangle = 0$. Then $\Phi^\dagger\Phi = (v+\sigma)^2\mathbb{I}$, and (3.7) collapses to a singlet potential. With $N = \mathrm{Tr}(\mathbb{I})$ locked directions,

$$V = \frac{\lambda}{4} N\big[(v+\sigma)^2 - v^2\big]^2 = N\lambda v^2\,\sigma^2 + O(\sigma^3). \tag{17.1}$$

The kinetic term in (3.6), in the signature $(-,+,+,+)$, contributes $-N\,(\partial\sigma)^2$. Canonical normalisation $\sigma_c = \sigma\sqrt{2N}$ puts the quadratic action in the form $-\frac{1}{2}(\partial\sigma_c)^2 - \frac{1}{2}m_\sigma^2\sigma_c^2$, with

$$m_\sigma^2 = \lambda v^2, \qquad m_\sigma = v\sqrt{\lambda}. \tag{17.2}$$

The would-be Goldstones in Λ remain massless at this order. They are the modes Chapter 10 is required to eat. They are not ψ.

Lemma (radial potential curvature in the pure-trace sector). Work in the pure-trace sector: $T_{\mu\nu} = 0$, $F_{\mu\nu} = 0$, and $\omega_{\mu\nu} = 0$. Two repairs of the frame must be kept distinct.

Repair II, unitary gauge. Chapter 10 permits $\Lambda = \mathbb{I}$ wherever Assumption 4.4 holds. The vacuum is then the frame, $e^A{}_\mu = \Phi^A{}_\mu/v$, so

$$g_{\mu\nu} = (1+\sigma/v)^2\eta_{\mu\nu}, \qquad \psi \equiv \eta^{\mu\nu}h_{\mu\nu} = 8\sigma/v + O(\sigma^2). \tag{17.3}$$

The conformal mode and the radial singlet are the same field in this gauge. The potential (3.7) gives the radial singlet the curvature $V'' \propto \lambda v^2$, which with the microscopic kinetic normalisation of Lemma 17.3 is $m_\sigma^2 = \lambda v^2$. This is a statement about the potential only. It is not yet the mass of a canonically normalised pole of ψ (see the identification paragraph below).

Repair I. Definition 4.1 does not define a vierbein (Appendix H, Section H.3), so ψ has no definition in terms of Λ and no Hessian on (σ,ψ) is computed for Repair I. What survives is that (3.7) depends only on $\Phi^\dagger\Phi = (v+\sigma)^2\mathbb{I}$, so it gives curvature only to σ and none to the $\mathfrak{so}(3,1)$-valued directions of Λ.

Identification $g^{(0)} = g[e]$. For the metric (17.3), $g = \Omega^2\eta$ with $\Omega = 1+\sigma/v$, the

induced Einstein–Hilbert density is $\sqrt{-g}\,R = -6\,\Omega\,\Box\Omega$, which equals $6(\partial\Omega)^2$ up to a total derivative. With the curvature convention of Chapter 2, Section 2.1 (identified in Chapter 9 as the one for which $R > 0$ on spheres), this adds to the Lagrangian $+\,3(\partial\sigma)^2/(8\pi G_{\rm eff} v^2)$ if the Einstein–Hilbert term has the form of (8.11) with $G_{\rm eff} > 0$; that sign of $G_{\rm eff}$ is a declared condition (Chapter 8, Problem F), not a derived result. In the signature $(-,+,+,+)$ a healthy scalar enters as $-\frac{1}{2}Z(\partial\sigma)^2$ with $Z > 0$, so this term has the conformal-factor (ghost) sign relative to the microscopic term $-N(\partial\sigma)^2$ of Lemma 17.3. Conditional on the pure-trace truncation, on adding the induced term to the microscopic one, and on $G_{\rm eff} > 0$ in the Lorentzian continuation, the quadratic action $-\frac{1}{2}Z(\partial\sigma)^2 - \frac{1}{2}V''\sigma^2$ has

$$Z = 2N - \frac{3}{4\pi G_{\rm eff} v^2} = 2N\Big(1 - \frac{3M_{\rm P}^2}{Nv^2}\Big), \qquad V'' = 2N\lambda v^2, \qquad M_{\rm P}^2 = \frac{1}{8\pi G_{\rm eff}}. \tag{17.4}$$

If $Nv^2 > 3M_{\rm P}^2$ then $Z > 0$ and a pole of squared mass $m^2 = V''/Z = \lambda v^2(1 - 3M_{\rm P}^2/(Nv^2))^{-1}$ exists in this truncation; it tends to (17.2) only when $3M_{\rm P}^2/(Nv^2) \to 0$, i.e. $G_{\rm eff}Nv^2 \to \infty$. For $Z < 0$ (i.e. $Nv^2 < 3M_{\rm P}^2$) the truncated mode has a wrong-sign kinetic term and, with $\lambda > 0$, also $m^2 = V''/Z < 0$: in this truncation it is both ghost-like and tachyonic, so the kinetic sign is not the only instability. The pure-trace truncation therefore cannot support Assumption 7.5 there; this signals an instability of the chosen background and truncation and is not a statement about the full scalar spectrum. At $Nv^2 = 3M_{\rm P}^2$ one has $Z = 0$: the quadratic kinetic term of this truncation degenerates, and the formula establishes no canonically normalized finite-mass pole there. Since $v/M_{\rm P}$ requires $G_{\rm eff}$, this ties Problem C to Problem F. The mixing with $T_{\mu\nu}$, the non-minimal curvature coupling, and the matter coupling are not derived here, and no Cassini conclusion follows.

Scalar exchange in the pure-trace truncation. Assume in addition that matter couples to $g = \Omega^2\eta$ of (17.3) only through the metric, with no further direct σ–matter coupling; this is an assumption, not a result of Chapters 11–12. Then to first order $\delta g_{\mu\nu} = 2(\sigma/v)\eta_{\mu\nu}$ and the matter Lagrangian contains $(\sigma/v)\,T^{\mu}{}_{\mu}$. For $Z > 0$ of Equation (17.4), nonrelativistic sources of positive mass exchange the canonical scalar through the attractive Yukawa potential

$$V_\sigma(r) = -\frac{m_1 m_2}{4\pi Z v^2}\frac{e^{-mr}}{r}.$$

Relative to the nominal Einstein-sector Newtonian potential

$$V_N(r) = -\frac{G_{\rm eff} m_1 m_2}{r},$$

its strength is βe^{-mr}, where

$$\beta = \frac{2M_{\rm P}^2}{Zv^2} = \frac{M_{\rm P}^2}{Nv^2 - 3M_{\rm P}^2}, \qquad M_{\rm P}^2 = \frac{1}{8\pi G_{\rm eff}}.$$

The physical tensor normalization is not established, so this Einstein-sector comparison is not identified with a measured Newton constant. As $Z \to 0^+$, β diverges but $m^2 = V''/Z$ also diverges, so at fixed separation $\beta e^{-mr} \to 0$; for $G_{\rm eff} N v^2 \gg 1$, $\beta \sim M_{\rm P}^2/(Nv^2)$. This concerns the unmixed truncation only. Mixing with $T_{\mu\nu}$, the non-minimal coupling, $G_{\rm eff} > 0$, and the metric-only coupling assumption remain unestablished, so no Cassini bound follows.

Cassini, revised. The potential (3.7) gives the radial singlet the curvature λv^2 in the Repair II pure-trace sector and produces no massless scalar by itself. The bound $\gamma - 1 = (2.1 \pm 2.3) \times 10^{-5}$ [5] constrains a massless or long-range scalar partner. Whether the physical scalar of this sector is short-range on an astronomical unit depends on the total kinetic coefficient, the non-minimal coupling and the matter coupling, none of which is computed. This is not a Cassini bound on NAMCG.

Status of Assumption 7.5. Not promoted. What is established is the potential curvature $m_\sigma^2 = \lambda v^2$ (Lemma 17.3) and its identification with the conformal mode in the Repair II pure-trace sector, under Assumptions 3.5 and 4.4. The physical scalar spectrum, the mixing with $T_{\mu\nu}$ and the matter coupling remain open (Appendix H, Section H.3). Loop corrections and the wrong-sign conformal kinetic term of the heat kernel remain Chapter 8 bookkeeping.

Problem D: local Lorentz invariance from Presentation A. *CRITICAL PATH; OPEN.* Generator counting is closed conditionally, while local Lorentz symmetry is imposed. Assumption 10.1 puts $\mathrm{SO}(3,1)$ into Presentation B by hand. Either derive a local Lorentz gauge symmetry from the matrix algebra and condensate (D1), or state that NAMCG is a theory of a condensate field on an auxiliary spacetime and stop calling Presentation A ontologically prior (D2). D1 closure requires the ω, A_μ, B, and ghost operators and the resulting field content to be specified. D1 closure also requires that the locked frame exist and be non-degenerate on the claimed domain, derived from the same microscopic data rather than taken from Assumptions 3.5 and 4.4. Under D2, even with these operators specified, the result is only a scope-limited Presentation-B model: D remains OPEN and CRITICAL PATH, and a full matrix-derived spectrum is not established. The Chapter 10 generator count identifies six Goldstone orbit variables removable in unitary gauge; it does not determine the dynamics of the independent diagonal connection B or whether torsion vanishes, propagates, or is massive. Torsion-free B follows only under the conditional Palatini–Holst hypotheses

of Proposition 10.5; NAMCG has not been shown to induce them. For a Laplace-type operator, $a_1 = \mathrm{tr}(R(g)/6 - E)$; its explicit universal curvature term is the Levi-Civita scalar $R(g)$, while unspecified E leaves additional B-dependence open and does not establish a Palatini $R(B)$ term. If B enters the actual bundle connection, a_2 may contain $\mathrm{tr}\, F(B)^2$; its survival in the weighted trace depends on unspecified operators and weights. Deriving the connection and endomorphism dependence or counting B and torsion as additional field content remains open; Problem D is unresolved. Within D2, reduced claims may be limited to Levi-Civita metric geometry of $g = e^T e$ and conservation of the total stress, without using B; unitary-gauge Lorentz locking removes the six Goldstone orbit variables, while frame/metric bookkeeping remains provisional. D2 does not establish a full-spectrum account, the dynamics or masslessness of B, torsion-free B, or the quadratic operator of A_μ. Problem F then applies only to an explicitly stated frame/metric truncation omitting B and A_μ, conditional on that truncation and with the corresponding ghost sector specified.

17.4 Open problems by scope and priority

17.4.1 Critical-path problems A and E

Problem A: the infrared dictionary. *OPEN; CRITICAL PATH for the matrix-derived label.* A scope-limited Presentation-B model may retain Assumption 3.6; it then claims no derivation from the matrix model, and Chapter 8 remains a Presentation-B calculation. Assumption 3.6 is the map

$$[\hat{X}_\mu, \Phi] \longleftrightarrow i\Theta_{\mu\nu} D^\nu \Phi \tag{17.5}$$

plus higher-derivative corrections. A derivation must specify the Hilbert space, Weyl map, continuum limit, and error at each order in Θ.

Problem E: Lorentzian signature. *OPEN; CRITICAL PATH for the matrix-derived label.* A scope that retains Assumption 4.6 gives up the claim that Lorentzian signature is emergent or derived. Euclidean positivity and a formal Wick rotation do not select it; rotating because the infrared world is Lorentzian is not a derivation.

17.4.2 Critical-path problems F, G and N

Problem F: the rationals A, B, C. *CRITICAL PATH; OPEN.* The conditional heat-kernel identity does not compute the physical coupling. The target is to evaluate the weighted a_1 and the curvature-linear part of a_2 on the locked block, including diffeomorphism and local-Lorentz ghosts and the radial-mode threshold, with a scheme stated. The finite-dimensional fibre count alone does not determine these coefficients:

the quadratic operator, its connection and endomorphism, the determinant weights, and finite counterterms enter.

Verified heat-kernel bookkeeping (not closure). For the proper-time convention of Appendix A,

$$W_E = -\frac{1}{2}\int_{\Lambda_{\rm UV}^{-2}}^{\mu^{-2}} \frac{\mathrm{d}s}{s}\,\mathrm{Tr}\,e^{-s\mathcal{O}},$$

termwise integration of the truncated, unresummed small-s expansion gives the a_0, a_1, a_2 factors $-1/(64\pi^2), -1/(32\pi^2), -1/(32\pi^2)$, respectively, with powers $\Lambda_{\rm UV}^4 - \mu^4$, $\Lambda_{\rm UV}^2 - \mu^2$, and $\ln(\Lambda_{\rm UV}^2/\mu^2)$. These are not exact finite-cutoff integrations of massive determinants; resumming mass exponentials produces sector-dependent threshold functions. The standard curvature-square part of a_2 for $\mathcal{O} = -D^2 + E$ is

$$\frac{1}{180}\left(R_{\mu\nu\rho\sigma}R^{\mu\nu\rho\sigma} - R_{\mu\nu}R^{\mu\nu} + \frac{5}{2}R^2\right),$$

not the previously displayed $7R^2/2$.

For one minimally coupled real bosonic Laplace-type field with $E_R = 0$, the universal a_1 contribution to the Euclidean coefficient of $\int \sqrt{g}\, R_E$ is

$$C_{R,E}^{(a_1)} = -\frac{N_0}{192\pi^2}(\Lambda_{\rm UV}^2 - \mu^2),$$

where N_0 is the fibre trace of this single sector and the displayed limits are assigned to it. For a specified collection, the corresponding Euclidean contribution is the sum over the explicitly weighted sectors and their assigned limits. This is not the full locked-block result and does not establish a ghost-subtracted trace. It makes no claim about the Lorentzian Einstein–Hilbert sign or a physical $G_{\rm eff}$; those require the specified sector sum and an explicit action/curvature continuation convention.

If $E = E_0 + QR + \cdots$, the a_1 curvature coefficient is $\mathrm{tr}(\frac{1}{6}\mathbb{I} - Q)$. For a scalar with $E = m^2 + \xi R$, the mass-dependent curvature-linear term in a_2 is $(\xi - \frac{1}{6})m^2 R$, vanishing at $\xi = 1/6$. Thus the conformal radial scalar alone does not establish a nonzero logarithmic B; the additional frame operator must be specified and evaluated.

Missing step. Specify the gauge-fixed quadratic Hessian for every locked-block field, including bundle connections, endomorphisms, diffeomorphism and local-Lorentz ghost operators and weights, the radial threshold prescription, finite counterterms, and continuation convention. Then compute the weighted a_1 and relevant a_2 terms in that scheme and match them to A, B, C. The present manuscript does not specify those data. The asserted net $+2$ from the on-shell TT count is not an off-shell heat-kernel trace, and no replacement ghost count can be inferred without the actual gauge fixing and ghost operators. The values $A = 1/\pi$, $B = 8\pi/3$, and $C = 0$ are not established.

The quadratic Φ–Φ Hessian of the displayed action (3.6) at fixed $A_\mu = 0$ (Appendix H, Problem F) is an input to, not a specification of, these operators. Problem F remains open.

Problem G: infrared boundary entropy. *CRITICAL PATH; mechanism CLOSED CONDITIONALLY, derivation OPEN.* Evaluate $S_{\rm ent}^{(\rm IR)}$ on an explicit cosmological or large-tube boundary and produce a candidate $\Lambda_{\rm eff}$ from (12.10). A symbolic formula in terms of a late-time entanglement scale is more useful than a claim to have obtained 10^{-122}.

G completion levels. For G-model, a scaling or value computed within a stated assumed boundary model is PROGRAMME-CONDITIONAL and does not satisfy the matrix-derived completion criterion. For G-external, independently justified boundary entropy and stationarity may support only a scope-limited Presentation-B model; G remains open for the matrix-derived label. G is DERIVED only when the entropy density and stationarity relation are derived from the matrix theory and $\Lambda_{\rm eff}$ follows from them. A scaling selected to reproduce a target remains an input. The required entropy density is the one in Assumption 12.2, derived by the replica construction from the matrix theory, and boundary stationarity Assumption 12.3 must likewise be derived rather than retained.

Problem N: control of the effective expansion. *CRITICAL PATH for the matrix-derived label; OPEN.* Assumption 9.2 states that the four-derivative terms are vertices of an expansion in $p/\Lambda_{\rm UV}$ and are not diagonalized into extra propagating modes of $h_{\mu\nu}$. Chapter 9 justifies it only by observing that the metric is composite. The target is to derive from the matrix path integral, and not from Stelle's action (9.5), that the graviton two-point function has no propagating Ostrogradsky ghost for $p < \Lambda_{\rm UV}$, for the values of α and $G_{\rm eff}$ computed in Problem F (here α is the Weyl-squared coupling of (9.5), whose $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$ coefficient is $-\alpha/2$; with the continuation convention of Section H.11, $c_W = -\alpha/2$ in (9.3); it is not any other coupling denoted α in this book), with the region of those values in which it holds stated. The target is no stronger than the requirement of Chapter 1, item 6, that no propagating Ostrogradsky ghosts remain in the infrared; it claims no nonperturbative ultraviolet completion. A statement valid only for a truncated or imposed expansion leaves Assumption 9.2 in place and yields at most a scope-limited model.

17.4.3 Optional problems continued

Problem H: interpolating function. *OPEN; OPTIONAL research within the theory.* The claim given up is a derivation of $\nu_{\surd}$ from modular data; current empirical

comparisons remain coefficient checks. Derive $\nu_{\surd}$ in (13.15), or a function that beats it on a SPARC hold-out, from the modular data of Chapters 6 and 12. The target is

$$\nu_{\surd}(g_{\rm N}) = \frac{1}{1-\exp(-\sqrt{g_{\rm N}/g_s})}, \tag{17.6}$$

which is identically $1 + n_{\rm BE}(\sqrt{g_{\rm N}/g_s})$. Appendix D already shows that this law, frozen at the published $g_\dagger$, is equivalent to the RAR and about 0.2% behind simple MOND, while baryons-only fails. That ranking is a coefficient check. It does not solve Problem H.

The first finite computation is the replica free energy of a free field on a weak-field diamond whose infrared boundary is specified as part of the calculation. Chapter 12 supplies $\Lambda_{\rm eff}$ but does not define a stationary boundary geometry or its surface gravity $\kappa_{\rm IR}$; a placeholder $\kappa_{\rm IR}$ with $\Lambda = \Lambda_{\rm eff}$ must be constructed and normalized there. Read the source-dependent modular excitation gap $\Delta k(g_{\rm N}, \kappa_{\rm IR})$ when a Newtonian source of acceleration $g_{\rm N}$ is introduced; the absolute lowest eigenvalue is not used, because it depends on an additive normalization (Section H.8). Problem H is solved if the response to the extra stress, computed from the weighted multimode spectral sum of Section H.8 (or from a justified single-ladder response), reproduces (17.6) with no SPARC input. In the single-ladder case this requires

$$\Delta k = \sqrt{\frac{g_{\rm N}}{g_s}}, \qquad g_s = c_\star\, \kappa_{\rm IR}, \tag{17.7}$$

with the order-one constant $c_\star$ obtained from the calculation and not from galaxy data. The values $c_\star = 1$ and $c_\star = 1/(2\pi)$ both appear in this manuscript; neither is adopted, and a scaling $\Delta k \propto \sqrt{g_{\rm N}/\kappa_{\rm IR}}$ alone does not fix $c_\star$. A computed spectrum that does not reproduce (17.6) does not solve Problem H. A linear gap $\Delta k \propto g_{\rm N}/\kappa_{\rm IR}$ reproduces the retired $\nu_{\rm exp}$ and kills Assumption E.1.

Alternatively, prove that no local $\mu_{\rm ent}(g_{\rm N})$ exists and write the nonlocal kernel instead. Refitting α, r_s, or g_s to rotmod files is not a solution. Identifying g_s with $g_\dagger$ inside the replica calculation is not allowed; compare $\kappa_{\rm IR}$ to $g_\dagger$ only after the source-dependent gap and response are calculated.

Problem I: dynamical origin of the drag. *OPEN; OPTIONAL falsifiable phenomenological extension.* It becomes critical only if the drag is claimed to be derived or predicted. Otherwise the microscopic exponent and rate remain open, and Assumption 14.2 is a working law. Replace Assumption 14.2 by a low-frequency response calculation of $\Gamma_{ij}(\mathbf{r}; \ell_A, \ell_B)$ from specified microscopic dynamics and a specified stationary reference state, including the finite-size factors ℓ_A, ℓ_B and the longitudinal and transverse components. Report the exponent and the coefficient. Then, and only

then, estimate η at tidal radii and at LIGO-band separations. The static tail and its assumptions are recorded in Problem H.9; no static-tail power determines the retarded response.

Problem J: spherical ECG solutions. *OPEN; OPTIONAL research within the stated model scope.* The claim given up is singularity resolution derived from the matrix theory; reduced results remain conditional on symmetric reduction and the ECG model assumption. For the explicit invariant $\mathcal{P}$ in (9.7), used in the action (9.8), Section H.10 records checked reduced Euler–Lagrange identities, the resulting first integral including Λ_{eff}, its conditional regular-centre constraint for every Λ_{eff}, the maximally symmetric curvature relation, and the large-radius tail. These results are conditional on the static spherical reduction capturing the full field equations; that symmetric-criticality step is not established here. The Einstein-like linearized spectrum on maximally symmetric backgrounds is cited separately and does not establish nonlinear or sourced stability. Problem J remains open: justify the reduction against the full field equations, establish the cubic truncation and its microscopic matching, and construct a regular massive core from specified sourced or modified equations, including its entanglement stress and residual relative to the phenomenological profile (13.6).

17.5 Problems that look like work and are not

The following activities are useful as exercises and do not move the framework.

- Producing further Newtonian integrators with new colour maps of Figure 15.1.
- Fitting SPARC with an unconstrained μ_{ent}, a per-galaxy (α, r_s), or a new ν that is not (17.6) and is not derived from $I(A:B)$. Appendix D already did the legitimate coefficient ranking.
- Writing $\mathbf{a}_{\text{irrev}}$ with additional free exponents and calling each a prediction.
- Replacing G_{toy} by G_{N} in a slide.
- Adding adjectives ("holographic," "non-Abelian," "entropic") to an abstract without adding a lemma.

Chapter 15 exists so that a legal laboratory is on the record. It does not exist so that the laboratory can become the theory.

17.6 A recommended order of work

The critical-path dependency order is:

1. Problem D1: derive local $SO(3,1)$ from the matrix algebra and condensate, together with the existence and non-degeneracy of the locked frame; specify the ω, A_μ, B and ghost operators needed by Problem F.

2. Problem C: compute the tree-level scalar operator, including kinetic coefficient, mixing, mass and testable matter coupling.

3. Problem F: compute A, B, C with the specified operators, ghosts, thresholds and scheme, and derive G_{eff}.

4. Problem N: using the α and G_{eff} of Problem F, control the effective expansion and exclude propagating Ostrogradsky ghosts for $p < \Lambda_{\text{UV}}$.

5. Problem B: derive the fixed-volume tensor-block stationarity from the microscopic action, not only for conformal variations.

6. Problem G: derive $S_{\text{ent}}^{(\text{IR})}$ and boundary stationarity from the matrix theory, then derive the Λ_{eff} scaling or value.

Problems A and E are not placed in this sequence. Each removes a premise (Assumption 3.6 and Assumption 4.6) that several items above use, so either may be attacked at any point. A derivation of B or F that still invokes Assumption 3.6 is DERIVED only after Problem A is.

Alternate scope: model work outside the critical-path sequence. Taking Presentation B with imposed local Lorentz symmetry (D2), an externally justified boundary result (G-external), or G_{eff} as an input may make a scope-limited Presentation-B model calculable and testable, for example by specifying full quadratic and ghost operators and computing Problem F for a stated truncation. Such work leaves Problems D, F, G and the matrix-derived completion criterion open, must name its imposed premises, and is not a step in the sequence above.

Problems H, I, and J are optional within their stated scopes. Appendix D records the coefficient-level check for Problem H; further galactic work should not be described as a derivation from $I(A:B)$ without the modular calculation.

17.7 What "finished" would mean

The label *Completed infrared derivation (matrix-derived)* applies only if all the critical-path problems below are DERIVED. DERIVED means no remaining assumed premise in the claimed chain, so a chain that still uses Assumption 3.6, 4.6, 3.5, 4.4, 7.5 or 9.2 does not qualify.

1. Problem A: Assumption 3.6 is derived, or the chains of the other items are re-derived without invoking it. Renaming the same map does not meet this item.

2. Problem B is a lemma: Assumption 6.1 is no longer an assumption, fixed-volume variations include the non-conformal tensor block, and the derivation does not invoke Assumption 3.6.

3. Problem C has a computed scalar operator, coupling, and phenomenological comparison defining and testing either a sufficiently heavy m_ψ or the scalar-tensor outcome; Assumption 7.5 is discharged.

4. Problem D1: an $SO(3,1)$ gauge symmetry is derived from the matrix algebra and condensate, with the ω, A_μ, B and ghost operators specified and the field content stated. The locked frame and its non-degeneracy on the claimed domain are derived, not assumed (Assumptions 3.5 and 4.4).

5. Problem E: Lorentzian signature is derived rather than retained (Assumption 4.6).

6. Problem F has A, B, C and the sign/scheme dependence of G_{eff} computed from the specified operators; G_{eff} is not an input.

7. Problem G-derived: the infrared boundary entropy density and stationarity relation are derived from the matrix theory, and the Λ_{eff} scaling or value follows from them. G-model and G-external do not meet this item.

8. Problem N: Assumption 9.2 is discharged; there is no propagating Ostrogradsky ghost for $p < \Lambda_{\text{UV}}$ in the derived graviton two-point function, for the values of α and G_{eff} computed in Problem F, with the region of those values in which it holds stated.

If local Lorentz symmetry is instead imposed (D2), even with the full quadratic and ghost content specified, the result is a scope-limited Presentation-B infrared model. It may be studied and tested as such, but Problem D stays OPEN and CRITICAL PATH and the criterion of this section is not met. No label of "completed infrared derivation" is allowed for that scope. Any critical-path item met only by an imposed premise, input, or reduced scope likewise yields at most a scope-limited model or programme, never the matrix-derived label.

Problems H, I, and J may remain open in a completed derivation within the stated scopes; the claims given up are listed above. Problem I is a prerequisite only if drag is claimed to be derived or predicted.

Until all the critical-path problems are derived, the honest public description is the subtitle of the book, read as a programme: *an attempt* to derive general relativity and

massless spin-2 dynamics from a matrix condensate held at entanglement equilibrium. The attempt is written so that it can fail in public. That is the point of Chapter 16.

17.8 Exercises

1. **Status report.** Copy the eleven-line chain of Section 16.1 and mark each line "lemma," "theorem from assumptions," or "assumption." Do not use a mark you cannot defend from the text.

2. **Proposal.** Write a two-page project proposal for Problem C or Problem F only. Include the first calculation you would perform and the result that would make you stop.

3. **Not a proposal.** Take any project that Section 17.5 would reject and explain, in six sentences, why it does not change the status of Theorem 7.3.

4. **Public sentence.** Write one sentence suitable for an Amazon description that survives Section 16.6 and the last paragraph of Section 17.7.

17.9 Bridge to the appendices

The main text is complete. Appendix A records the Seeley–DeWitt algorithm at the working thickness used in Chapter 8. Appendix B records the replica formula and the modular Hamiltonian at the thickness used in Chapter 6. Appendix C records the Verlet laboratory and the energy identity used in Chapter 15. Appendix D records the SPARC ranking of $\nu_{\surd}$ against the RAR, simple MOND, and baryons-only. None of the appendices introduces a new physical hypothesis. If an appendix appears to do so, the hypothesis belongs in the main text or it does not belong in the book.

Appendix A

Seeley–DeWitt Coefficients at Working Thickness

This appendix records the heat-kernel formulae used in Chapter 8. It introduces no new physical hypothesis. Conventions follow Section 2.6: Euclidean section first, Lorentzian continuation afterwards, signature of the main text restored only when an Einstein–Hilbert term is isolated.

A.1 Laplace-type operators

Let $V \to M$ be a vector bundle over a smooth four-manifold with Riemannian metric $g_{\mu\nu}$. A second-order differential operator on sections of V is of Laplace type if it can be written

$$\mathcal{O} = -g^{\mu\nu} D_\mu D_\nu + E, \tag{A.1}$$

where $D_\mu = \nabla_\mu + \mathcal{A}_\mu$ is a connection on V (including the Levi-Civita connection on tensor factors) and E is a bundle endomorphism. The curvature of D is

$$\mathcal{F}_{\mu\nu} = [D_\mu, D_\nu]. \tag{A.2}$$

On the locked condensate block of Chapter 5, V is the bundle of tensors $\delta\Phi_{\mu\nu}$ together with whatever ghosts are included in a given computation.

The heat operator $e^{-s\mathcal{O}}$ is defined for $s > 0$ by the usual functional calculus after $\mathcal{O}$ has been made positive by a Wick rotation. Its integral kernel $K(s; x, y)$ satisfies

$$\partial_s K + \mathcal{O}_x K = 0, \qquad K(0; x, y) = \delta(x, y). \tag{A.3}$$

A.2 Small-time expansion

On the diagonal in four dimensions,

$$\operatorname{tr}_V K(s;x,x) \sim \frac{1}{(4\pi s)^2} \sum_{k=0}^{\infty} s^k a_k(x), \qquad s \to 0^+. \tag{A.4}$$

Each a_k is a local scalar density of mass dimension $2k$, built from $R_{\mu\nu\rho\sigma}$, $\mathcal{F}_{\mu\nu}$, E, and their covariant derivatives. The first three coefficients, up to total derivatives that do not enter the bulk effective action, are

$$a_0 = \operatorname{tr}_V \mathbb{I}, \tag{A.5}$$

$$a_1 = \operatorname{tr}_V \Big(\tfrac{1}{6} R - E \Big), \tag{A.6}$$

$$\begin{aligned} a_2 = \operatorname{tr}_V \Bigg(& \frac{1}{180} (R_{\mu\nu\rho\sigma} R^{\mu\nu\rho\sigma} - R_{\mu\nu} R^{\mu\nu} + \tfrac{5}{2} R^2) + \frac{1}{2} E^2 - \frac{1}{6} R E \\ & + \frac{1}{12} \mathcal{F}_{\mu\nu} \mathcal{F}^{\mu\nu} + \frac{1}{30} D^2 R - \frac{1}{6} D^2 E \Bigg). \end{aligned} \tag{A.7}$$

Different references redistribute the $D^2 R$ and $D^2 E$ terms by integration by parts. For the bulk action of Chapter 8 those terms are discarded.

The numerical coefficients in (A.7) are the standard DeWitt values for a Laplace-type operator. They do not determine the coefficients A, B, C in the conditional parameterization (8.12). Their evaluation requires the specified operators, fibre traces, determinant weights, and thresholds; this remains part of Problem F in Chapter 17.

A.3 Proper-time integrals

For the sectors actually included in a gauge-fixed calculation, let $\mathcal{S}_{\mathrm{sec}}$ be their index set, w_s their determinant weights relative to a real bosonic determinant, and $\mathcal{O}_s$ their operators. Define

$$W_E^{(1)} = \frac{1}{2} \sum_{s \in \mathcal{S}_{\mathrm{sec}}} w_s \operatorname{Tr}_{V_s} \ln \mathcal{O}_s, \qquad \mathcal{N}_0 := \sum_{s \in \mathcal{S}_{\mathrm{sec}}} w_s \operatorname{tr}_{V_s} \mathbb{I}_s. \tag{A.8}$$

The corresponding proper-time representation is

$$W_E^{(1)} = -\frac{1}{2} \sum_{s \in \mathcal{S}_{\mathrm{sec}}} w_s \int_{s_{\min,s}}^{s_{\max,s}} \frac{\mathrm{d}s}{s} \operatorname{Tr}_{V_s} e^{-s\mathcal{O}_s}. \tag{A.9}$$

The expressions below are term-by-term integrals of the truncated, unresummed small-s heat-kernel expansion through a_2, not exact finite-cutoff integrations of massive determinants. If E_s contains m_s^2, resumming $\exp(-sm_s^2)$ replaces polynomial cutoff terms by sector-dependent threshold functions. A common pair of limits applies only to sectors explicitly assigned those limits; otherwise the sector-dependent limits are retained. Inserting the heat-kernel expansion and integrating term by term gives

$$W_E^{(1)} \supset \sum_{s\in\mathcal{S}_{\rm sec}} w_s \int \mathrm{d}^4x\,\sqrt{g}\,\Bigg[-\frac{a_{0,s}}{64\pi^2}(\Lambda_{\mathrm{UV}_s}^4-\mu_s^4)-\frac{a_{1,s}}{32\pi^2}(\Lambda_{\mathrm{UV}_s}^2-\mu_s^2) - \frac{a_{2,s}}{32\pi^2}\ln\Big(\frac{\Lambda_{\mathrm{UV}_s}^2}{\mu_s^2}\Big)\Bigg], \tag{A.10}$$

where $a_{k,s}$ are the traced coefficients on V_s. With common limits explicitly assigned to every included sector, the a_0 term is $-\mathcal{N}_0(\Lambda_{\rm UV}^4-\mu^4)/(64\pi^2)$.

For the Lorentzian vacuum-sign statement adopt $S_E=-iS_L$ under $t=-i\tau$ and define the Lorentzian vacuum term by $S_L\supset-\rho_\Lambda\int\mathrm{d}^4x\,\sqrt{-g}$. For one real scalar, $a_0=1$, and only under these conventions, $W_E/\mathrm{Vol}\supset-(\Lambda_{\rm UV}^4-\mu^4)/(64\pi^2)$ corresponds to $\rho_\Lambda=-(\Lambda_{\rm UV}^4-\mu^4)/(64\pi^2)$. The common-limit sector sum gives $\rho_\Lambda=-\mathcal{N}_0(\Lambda_{\rm UV}^4-\mu^4)/(64\pi^2)$ only when all displayed sectors share the assigned limits, within the same truncated, unresummed expansion. The Lorentzian sign is conditional on this continuation/action convention.

A.4 Worked example: a non-minimal scalar

For a single real scalar with

$$\mathcal{O}=-\nabla^2+\xi R+m^2, \tag{A.11}$$

one has $E=\xi R+m^2$, $\mathcal{F}_{\mu\nu}=0$, and $\mathrm{tr}_V\mathbb{I}=1$. Then

$$a_1=(\tfrac{1}{6}-\xi)R-m^2. \tag{A.12}$$

The curvature piece of a_1 cancels at $\xi=1/6$. That cancellation removes this scalar's contribution to the curvature coefficient through a_1. The NAMCG radial mode is not assumed to sit at $\xi=1/6$; Exercise 3 of Chapter 8 is the corresponding warning.

For one real scalar the Euclidean a_0 contribution is $-(\Lambda_{\rm UV}^4-\mu^4)/(64\pi^2)$. Under the continuation and Lorentzian vacuum-action convention stated above, this corresponds to $\rho_\Lambda=-(\Lambda_{\rm UV}^4-\mu^4)/(64\pi^2)$. For a collection of sectors the total density includes their determinant weights and sector-dependent thresholds.

A.5 Fibre traces on the locked block

If V is the space of symmetric traceless tensors on a four-manifold, $\mathrm{tr}_V \mathbb{I} = 9$. If the algebraic split of Chapter 5 is retained before gauge fixing, the sixteen components of $\delta\Phi_{\mu\nu}$ give $\mathrm{tr}_V \mathbb{I} = 16$. Ghosts for diffeomorphisms contribute with the opposite sign and with their own Laplace-type operators. Any local-Lorentz ghost sector must be specified together with the gauge fixing of the independent Lorentz connection; Theorem 10.3 does not supply that operator.

$\mathcal{N}_0$ is the weighted fibre trace over the sectors actually included, not automatically the locked-block dimension or a physical-mode count. A total locked-block result must specify the gauge-fixed fluctuation, Goldstone, diffeomorphism-ghost and local-Lorentz-ghost operators, determinant weights, and assigned thresholds. No factor of $1/6$ is absorbed into $\mathcal{N}_0$; the complete calculation is Problem F.

A.6 What this appendix will not do

It will not list a_3 and a_4. Those coefficients generate six- and eight-derivative operators that the book treats as suppressed by further powers of $\Lambda_{\mathrm{UV}}^{-1}$.

It will not prove the coefficients (A.5)–(A.7). Standard derivations (coincidence limits of the Synge world function, or the recursive DeWitt transport equations) are in the references collected in `bib/namcg.bib`. The book uses the coefficients; it does not redo the recursion.

It will not evaluate a_2 on the Einsteinian cubic background of Chapter 13. That evaluation is part of Problem J if it is needed, not part of the working thickness of Chapter 8.

A.7 Exercises

1. **Dimensions.** Assign mass dimensions to s, a_0, a_1, a_2, and $\mathrm{tr}_V K(s; x, x)$. Confirm that each term in (A.10) has dimension four.

2. **Conformal scalar.** Set $\xi = 1/6$ and $m = 0$ in (A.12). What remains of a_1? What remains of the $\Lambda_{\mathrm{UV}}^2 R$ term in W_E?

3. **Gauss–Bonnet combination.** Rewrite the curvature part of (A.7) in the basis $\{C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}, R^2, E_{\mathrm{GB}}\}$. Which combination multiplies a topological density in four dimensions?

4. **Ghost sign.** A diffeomorphism ghost in the background-field gauge is a complex vector with a Laplace-type operator. What factor multiplies its contribution to a_0 relative to a real scalar, and why is the sign opposite to a physical field?

5. **Conditional curvature coefficient.** Given specified sector operators, determinant weights, proper-time limits, and a finite-counterterm scheme, derive the truncated Euclidean curvature coefficient from a_1 and a_2. Identify the additional data needed for its Lorentzian continuation and comparison with G_{eff}.

Appendix B

Modular Hamiltonians and the Replica Formula

This appendix records the entanglement formulae used in Chapters 6, 11, and 12. It introduces no new physical hypothesis. The working identity of the main text remains Proposition 6.5. What follows is the language in which that identity was written.

B.1 States, reductions, and von Neumann entropy

Let $\mathcal{H}$ be the Hilbert space of the microscopic theory and ρ a density operator on $\mathcal{H}$. For a von Neumann algebra $\mathcal{A}_D$ of observables associated with a region D, the reduced state ρ_D is the restriction of ρ to $\mathcal{A}_D$. In a tensor-product approximation $\mathcal{H} \simeq \mathcal{H}_D \otimes \mathcal{H}_{D^c}$,

$$\rho_D = \mathrm{Tr}_{\mathcal{H}_{D^c}} \rho. \tag{B.1}$$

Continuum quantum field theory does not have that tensor-product factorization. The algebraic definition (restriction to $\mathcal{A}_D$) is the honest one; equation (B.1) is the formula used in every explicit replica computation in the book.

The von Neumann entropy and modular Hamiltonian of ρ_D are

$$S_D = -\mathrm{Tr}(\rho_D \ln \rho_D), \qquad H_D = -\ln \rho_D, \tag{B.2}$$

so that

$$S_D = \langle H_D \rangle_{\rho_D}. \tag{B.3}$$

If ρ_D has kernel, $\ln \rho_D$ is defined on the support of ρ_D and S_D may be infinite. In quantum field theory S_D is ultraviolet divergent. Differences δS_D at fixed cutoff, and mutual informations, are the finite objects used in the main text.

B.2 Relative entropy and mutual information

For two states ρ and σ on the same algebra,

$$S(\rho\|\sigma) = \mathrm{Tr}(\rho \ln \rho) - \mathrm{Tr}(\rho \ln \sigma) \tag{B.4}$$

is non-negative and vanishes if and only if $\rho = \sigma$ (when the expression is finite). Mutual information of two regions A and B is the relative entropy of ρ_{AB} with respect to $\rho_A \otimes \rho_B$,

$$I(A:B) = S(\rho_{AB}\|\rho_A \otimes \rho_B) = S(A) + S(B) - S(A \cup B), \tag{B.5}$$

the second equality holding when the three von Neumann entropies are regulated in the same scheme. Positivity of $I(A:B)$ is therefore positivity of relative entropy. Under Definition 6.4, for $\lambda \geq 0$, this implies $U_{\mathrm{ent}} \leq 0$. That concerns the chosen potential convention; it does not establish an ordering between adiabatic and non-adiabatic states or the sign of a dynamical drag coefficient. A nonnegative low-frequency drag requires a separate passivity condition on the retarded response, as discussed in Chapter 14.

Strong subadditivity,

$$S(A \cup B) + S(B \cup C) \geq S(B) + S(A \cup B \cup C), \tag{B.6}$$

is recorded for completeness. The main text does not use it beyond the positivity of $I(A:B)$.

B.3 The replica formula

For integer $n \geq 1$,

$$\mathrm{Tr}\,(\rho_D)^n = Z_n / Z_1^n, \tag{B.7}$$

where Z_n is the partition function of the microscopic theory on an n-sheeted cover branched over the edge $\mathcal{B} = \partial D \cap \Sigma$. Analytic continuation in n and the identity

$$S_D = \lim_{n \to 1} (1 - n\partial_n) \ln \mathrm{Tr}\,(\rho_D)^n \tag{B.8}$$

produce the von Neumann entropy. Differentiating at $n = 1$ inserts a single modular Hamiltonian: the first variation $\delta S_D = \delta \langle H_D \rangle$ used in Chapter 6 is the replica formula at first order, not a second definition.

In Presentation B the cover is a branched cover of the auxiliary manifold. Identifying it with the emergent spacetime requires a specified map from the auxiliary metric to the emergent metric, which the displayed action does not supply (see Assumption 8.4).

In Presentation A the same construction requires Problem A of Chapter 17.

B.4 Bisognano–Wichmann and small diamonds

On Minkowski space, let W be a Rindler wedge. The Bisognano–Wichmann theorem identifies the modular Hamiltonian of the vacuum restricted to W with the boost generator that fixes the edge,

$$H_W = \frac{2\pi}{\kappa} \int_{\Sigma \cap W} \mathrm{d}\Sigma^\mu \, \xi^\nu T_{\mu\nu}, \tag{B.9}$$

where ξ^μ is the boost Killing vector and κ its surface gravity (equal to 1 in standard Rindler coordinates, restored here for matching (6.5)).

A small causal diamond in a slowly curved emergent geometry is, to leading order in Riemann normal coordinates (2.27), a compact analogue of two opposing Rindler wedges glued at a 2-sphere. The working formula (6.5) is that analogue: the same boost-energy integral plus an area term H_{area} whose first variation is $\delta A/(4G_{\mathrm{eff}})$ after Chapter 8 has named G_{eff}. The error is $O(\ell^3)$ in the diamond radius, which is the order dropped in Proposition 6.5.

The identification is exact for free fields in Rindler space. It is an operator-product approximation for interacting condensate fluctuations. The book does not claim a Bisognano–Wichmann theorem for the full matrix potential (3.7). For free scalar fields and coherent states, the relative entropy between the vacuum and the excited state on a bifurcate Killing horizon is identified with the horizon energy flux in [2]. That result is for a prescribed metric and says nothing about the interacting matrix condensate.

B.5 Fixed volume

Jacobson's entanglement-equilibrium condition is imposed at fixed spatial volume of the maximal slice. In Riemann normal coordinates the volume of a small ball of radius ℓ is (6.2). A pure conformal variation $\delta g_{\mu\nu} = \chi \eta_{\mu\nu}$ changes that volume at order ℓ^3. Holding V fixed removes that breathing from the comparison and is why the conformal block ψ and the tensor block $T_{\mu\nu}$ do not enter the first law on the same footing.

Fixed volume is a constraint on the variations admitted in Assumption 6.1. It is not a statement that ψ is massive. That statement is Assumption 7.5.

B.6 Boundary entropy versus diamond entropy

Chapter 12 uses a different functional: a local density $S_{\mathrm{ent}}(h_{ij}, \phi)$ on ∂M. The replica construction of that density is the same formula, (B.8), applied to a thin tubular

neighbourhood of the boundary and then localized as in Assumption 12.2. The two functionals must not be substituted for each other. Section 12.6 already assigned them different jobs.

B.7 What this appendix will not do

It will not develop Tomita–Takesaki theory beyond the definition $H_D = -\ln \rho_D$. It will not compute S_D for the matrix potential (3.7). It will not prove Assumption 6.1. Those are Problem B of Chapter 17 and the physical input of Chapter 6, respectively.

B.8 Exercises

1. **Replica derivative.** From (B.8) and $\operatorname{Tr} \rho^n = \exp(\ln \operatorname{Tr} \rho^n)$, recover $S = \langle H \rangle$ at $n = 1$.

2. **Relative entropy and Klein's inequality.** For finite-dimensional density matrices, prove $S(\rho \| \sigma) \geq 0$. Deduce $I(A : B) \geq 0$.

3. **Rindler temperature.** In standard Rindler coordinates, show that the Unruh temperature seen by an accelerated observer is $\kappa / 2\pi$. Match the factor in (B.9).

4. **Area divergence.** For a free scalar in Minkowski space, the unregulated S_D diverges as $\operatorname{Area}(\mathcal{B}) / \epsilon^2$. Why do mutual information and first variations δS_D at fixed cutoff remain useful?

5. **Two functionals.** Write four sentences explaining why a small-diamond S_D cannot be used as the boundary density of Definition 12.1.

Appendix C

Verlet Laboratory and Energy Identity

This appendix records the integrator and the energy identity used in Chapter 15. It introduces no new physical hypothesis. The force law is Assumption 14.2. The bookkeeping reference is Proposition 11.3, which this laboratory does not satisfy in the microscopic sense; the integrator tracks only W_{diss}. The source file is `code/namcg_inspiral.py`.

C.1 Reduced equation

In the reduced-mass frame the laboratory system is

$$\ddot{\mathbf{r}} = \mathbf{a}_{\mathrm{N}}(\mathbf{r}) + \mathbf{a}_{\mathrm{d}}(\mathbf{r}, \mathbf{v}), \tag{C.1}$$

with

$$\mathbf{a}_{\mathrm{N}} = -\frac{G_{\mathrm{toy}} M}{d^3}\mathbf{r}, \qquad \mathbf{a}_{\mathrm{d}} = -\frac{\eta}{d^4}\mathbf{v}, \qquad d = |\mathbf{r}|. \tag{C.2}$$

The conservative force is Newtonian. The drag is (14.10) with μ cancelled. Toy parameters are (15.2).

C.2 One step

Let $\Delta t > 0$. Given $(\mathbf{r}_n, \mathbf{v}_n)$,

$$\begin{aligned} \mathbf{v}_{1/2} &= \mathbf{v}_n + \tfrac{1}{2}\Delta t\big(\mathbf{a}_{\mathrm{N}}(\mathbf{r}_n) + \mathbf{a}_{\mathrm{d}}(\mathbf{r}_n, \mathbf{v}_n)\big), \\ \mathbf{r}_{n+1} &= \mathbf{r}_n + \Delta t\, \mathbf{v}_{1/2}, \\ \mathbf{v}_{n+1} &= \mathbf{v}_{1/2} + \tfrac{1}{2}\Delta t\big(\mathbf{a}_{\mathrm{N}}(\mathbf{r}_{n+1}) + \mathbf{a}_{\mathrm{d}}(\mathbf{r}_{n+1}, \mathbf{v}_{1/2})\big). \end{aligned} \tag{C.3}$$

When $\eta = 0$ this is velocity Verlet for a position-dependent force and is second-order accurate and symplectic on the conservative phase space. When $\eta \neq 0$ the last drag evaluation uses $\mathbf{v}_{1/2}$ rather than $\mathbf{v}_{n+1}$, so the dissipative piece is explicit and the scheme is no longer symplectic. The local error in the drag kick remains $O(\Delta t^2)$.

An implicit alternative replaces $\mathbf{a}_{\mathrm{d}}(\mathbf{r}_{n+1}, \mathbf{v}_{1/2})$ by $\mathbf{a}_{\mathrm{d}}(\mathbf{r}_{n+1}, \mathbf{v}_{n+1})$ and solves a linear equation for $\mathbf{v}_{n+1}$. The published script does not do that. The difference is visible only if one drives Δt up to the orbital period.

C.3 Mechanical energy, work, and balance

Define

$$E_{\mathrm{mech}} = \frac{1}{2}\mu v^2 - \frac{G_{\mathrm{toy}} M \mu}{d} \tag{C.4}$$

and the accumulated work

$$W_{\mathrm{diss}}(t_{n+1}) = W_{\mathrm{diss}}(t_n) + \mathbf{F}_{\mathrm{diss}}(\mathbf{r}_n, \mathbf{v}_n) \cdot \mathbf{v}_n \, \Delta t, \tag{C.5}$$

with $\mathbf{F}_{\mathrm{diss}} = \mu \mathbf{a}_{\mathrm{d}}$ and $W_{\mathrm{diss}}(0) = 0$. Along exact solutions of (C.1),

$$\frac{\mathrm{d}E_{\mathrm{mech}}}{\mathrm{d}t} = \mathbf{F}_{\mathrm{diss}} \cdot \mathbf{v}, \tag{C.6}$$

because the Newtonian force is minus the gradient of $U_{\mathrm{N}} = -G_{\mathrm{eff}}$-less toy potential $-G_{\mathrm{toy}} M \mu / d$. Therefore

$$\mathcal{E}(t) \equiv E_{\mathrm{mech}}(t) - W_{\mathrm{diss}}(t) \tag{C.7}$$

is constant on exact solutions. On discrete solutions it is constant up to the quadrature error in (C.5) and the Verlet error in (C.3). A systematic drift that does not shrink with Δt means the implementation does not obey (C.6) and is illegal in the sense of Section 14.6.

A trapezoidal quadrature,

$$\Delta W = \frac{1}{2}\Big(\mathbf{F}_{\mathrm{diss},n} \cdot \mathbf{v}_n + \mathbf{F}_{\mathrm{diss},n+1} \cdot \mathbf{v}_{n+1}\Big)\Delta t, \tag{C.8}$$

reduces that drift by one order in Δt. The published script uses the left-endpoint rule (C.5) because it is enough to make $\mathcal{E}$ look flat on the scale of Figure 15.1.

C.4 Angular momentum

The Newtonian force is central and exerts no torque. The drag is antiparallel to $\mathbf{v}$, so

$$\frac{\mathrm{d}\mathbf{L}}{\mathrm{d}t} = \mathbf{r} \times \mathbf{F}_{\mathrm{diss}} = -\eta\mu \frac{\mathbf{r} \times \mathbf{v}}{d^4}. \tag{C.9}$$

Angular momentum is therefore not conserved when $\eta \neq 0$. That is required: an inspiral that conserved $|\mathbf{L}|$ at fixed mass would be incompatible with a shrinking circular orbit. The identity (C.9) is a useful code check (Exercise 4 of Chapter 15).

C.5 Initial data and units

The published run uses

$$\mathbf{r}(0) = (1, 0), \qquad \mathbf{v}(0) = (0, \sqrt{G_{\text{toy}} M}), \tag{C.10}$$

with $G_{\text{toy}} = 36$, $\eta = 0.006$, $\Delta t = 10^{-3}$, and 1.2×10^4 steps. These are code units. They are chosen so that decay is visible, not so that η matches a tidal-disruption estimate from Problem I of Chapter 17.

Restoring dimensions, η in (C.2) has dimension length^4/time, since $\mathbf{a} = -\eta \mathbf{v}/d^4$. The static tail (14.8) does not supply this coefficient; deriving it requires the response calculation of Problem H.9, which this appendix does not attempt.

C.6 How to run the script

From the project root, after `figs/` exists:

```
python code/namcg_inspiral.py
```

The script writes `figs/namcg_gr_simulation_corrected.pdf`. Point Figure 15.1 at that PDF for the print interior. A legacy PNG from an earlier notebook may be kept for draft viewing.

If `matplotlib` is unavailable, the integrator still runs and can write `traj`, `E_mech`, and `W_diss` to text; the figure is then optional.

C.7 What this appendix will not do

It will not introduce gravitational waves, horizons, or a relativistic two-body Hamiltonian. It will not treat G_{toy} as G_{N}. It will not fit η to data. Those refusals are the content of Sections 15.6 and 16.7.

C.8 Exercises

1. **Conservative order.** With $\eta = 0$, measure the secular drift of E_{mech} over 10^4 steps at three values of Δt. Confirm second-order scaling.

2. **Quadrature.** Replace (C.5) by (C.8) and compare the drift of $\mathcal{E}$.

3. **Implicit drag.** Solve the linear equation for $\mathbf{v}_{n+1}$ when the last drag evaluation uses $\mathbf{v}_{n+1}$. Does $\mathcal{E}$ improve enough to justify the extra algebra?

4. **Torque check.** Integrate (C.9) in lockstep with the trajectory and compare with $\mathbf{L}_{n+1} - \mathbf{L}_n$.

5. **Illegal control.** Set $\mathbf{a}_{\mathrm{d}} = -\gamma \mathbf{v}$ with constant γ and watch W_{diss} as d decreases. This is Exercise 3 of Chapter 15, recorded here so the appendix can be used alone as a lab sheet.

Appendix D

SPARC Rotation Curves: A Coefficient-Level Consistency Check

D.1 What is being tested

The function

$$\nu_{\surd}(g_{\rm N}, g_s) = \frac{1}{1 - \exp(-\sqrt{g_{\rm N}/g_s})} \tag{D.1}$$

is the published Radial Acceleration Relation interpolator of McGaugh, Lelli & Schombert (2016). Appendix H records that a derivation of this (or any other) interpolator from the mutual-information functional remains open (Problem H). The present appendix asks only whether the same function, with the single scale fixed at the published value $g_s = g_\dagger = 1.20 \times 10^{-10}\,\mathrm{m\,s^{-2}}$, is compatible with the SPARC rotation-curve sample.

The predicted circular velocity is

$$V = V_{\rm b}\sqrt{\nu_{\surd}(g_{\rm N}, g_s)}, \tag{D.2}$$

where $V_{\rm b}$ is the Newtonian baryonic contribution computed from the SPARC mass models. No additional galactic parameters are introduced.

D.2 SPARC comparison

The comparison uses the public SPARC rotmod files (175 galaxies, 3391 radial points). Three models are scored by χ^2 and AIC:

- baryons only ($\nu = 1$);

- simple MOND with one free acceleration scale a_0;
- $\nu_{\surd}$ with g_s fixed at $g_\dagger$ (zero free parameters) and, for reference, with g_s free.

Table D.1: Global χ^2 and AIC on 175 SPARC galaxies (3391 points). Lower AIC is better.

Model	Free params	χ^2	AIC	Scale ($\mathrm{m\,s^{-2}}$)
$\nu_{\surd}$ (fixed $g_s = g_\dagger$)	0	1.695×10^5	1.695×10^5	1.20×10^{-10}
$\nu_{\surd}$ (free g_s)	1	1.695×10^5	1.695×10^5	1.20×10^{-10}
Simple MOND (free a_0)	1	1.691×10^5	1.691×10^5	1.16×10^{-10}
Baryons only	0	1.960×10^6	1.960×10^6	—

The fixed-g_s and free-g_s versions of $\nu_{\surd}$ are statistically indistinguishable; the free fit returns $g_s = 1.20 \times 10^{-10}\,\mathrm{m\,s^{-2}}$. Relative to the best one-parameter MOND fit the χ^2 penalty is approximately 0.2%. Both modified-gravity models improve on baryons-only by more than an order of magnitude in χ^2.

D.3 Numerical implementation

Listing D.1: Core evaluation functions.

```
import numpy as np

KMS2_PER_KPC_TO_SI = 1.0e6 / 3.085677581491367e19
G_DAGGER = 1.20e-10

def gn_si(vb, r):
    r = np.maximum(np.asarray(r, float), 1e-6)
    return (np.asarray(vb, float)**2 / r) * KMS2_PER_KPC_TO_SI

def nu_sqrt(gn, gs):
    gs = max(float(gs), 1e-18)
    x = np.clip(np.asarray(gn, float)/gs, 0.0, 1e12)
    s = np.sqrt(x)
    denom = 1.0 - np.exp(-s)
    nu = np.empty_like(denom)
    tiny = denom < 1e-12
    nu[tiny] = 1.0 / np.maximum(s[tiny], 1e-30)
    nu[~tiny] = 1.0 / denom[~tiny]
    return nu

def v_namcg(r, vb, gs=G_DAGGER):
```

```
    return vb * np.sqrt(np.clip(nu_sqrt(gn_si(vb, r), gs), 1.0,
1e6))
```

D.4 What the comparison is and is not

At low Newtonian acceleration, $\nu_{\surd}$ approaches the square-root map $g_{\rm obs} \approx \sqrt{g_{\rm N} g_\dagger}$. On the SPARC disks this keeps V from falling as fast as $V_{\rm b}$. The numerical result shows that this particular function, frozen at the published $g_\dagger$, is compatible with the sample at the same level as a one-parameter MOND fit.

The comparison does not:

- derive $\nu_{\surd}$ from $I(A : B)$ or from the modular spectrum (Problem H remains open);
- fix $g_\dagger$ from the condensate;
- constrain $m_T = 0$, the heat-kernel coefficients, or the conservation identity;
- rule out particle dark-matter halos on other scales.

It is a coefficient-level consistency check: the infrared equation of the monograph can wear the published RAR interpolator without contradiction on this data set. That is the strongest statement the SPARC sample supports.

Appendix E

A Provisional Bridge from Mutual Information to the RAR

This appendix states the strongest logical contact between NAMCG and the SPARC radial acceleration relation that the book can currently defend. It does not solve Problem H. It does not derive $g_\dagger$ from $\Phi^I{}_A$. It replaces an unspecified $\mu_{\rm ent}$ with a single modular conjecture whose failure would leave Appendix D as a coefficient ranking and nothing more.

E.1 What the two sides already share

NAMCG, after Theorems 7.3–11.1, is Einstein gravity plus a conserved extra tensor obtained by varying an entanglement functional of baryonic regions. In a static disk that statement has three immediate consequences, none of which uses SPARC.

1. The extra acceleration is slaved to the baryonic Newtonian field. It cannot be an independent halo function of r (Assumption 13.2).

2. At Solar-System and inner-disk accelerations the extra piece must vanish, or Cassini bounds fail: $\nu \to 1$ as $g_{\rm N} \to \infty$.

3. The only scalar available to mark the transition is an acceleration built from the same modular data that defined the diamonds of Chapter 6 and the infrared boundary of Chapter 12.

A fourth consequence is needed once the subsystem is not isolated. The entanglement functional is a functional of the baryonic configuration, not of an excised disk. A slowly varying external Newtonian field $\mathbf{g}_{\rm ext}$ therefore enters the argument of the

boost (Assumption 13.3). In the external-field-dominated limit $g_{\rm N} \ll g_{\rm ext}$,

$$\gamma \equiv \frac{g_{\rm int}}{g_{\rm N}} \to \nu_{\surd}(g_{\rm ext}), \tag{E.1}$$

not $\nu_{\surd}(g_{\rm N})$. At the Sun, $v_\odot^2/R_0 \approx 2.2 \times 10^{-10}\,{\rm m\,s^{-2}} \approx 1.8\,g_\dagger$, so $\nu_{\surd}(g_{\rm gal}) \approx 1.1$. An isolated evaluation of (E.2) at $g_{\rm N} = 10^{-11}\,{\rm m\,s^{-2}}$ wants $\gamma \approx 4$ and is illegal for a solar-neighborhood binary.

The RAR is an empirical map of exactly that type [4, 6]:

$$g_{\rm obs} = \nu_{\rm RAR}(x)\, g_{\rm N}, \qquad x = \frac{g_{\rm N}}{g_\dagger}, \qquad \nu_{\rm RAR}(x) = \frac{1}{1 - e^{-\sqrt{x}}}, \tag{E.2}$$

with $g_\dagger = (1.20 \pm 0.02) \times 10^{-10}\,{\rm m\,s^{-2}}$ and residuals that do not track radius or Hubble type. Appendix D showed that (E.2), frozen at $g_\dagger$, is the disk law NAMCG is presently allowed to wear.

The shared content is therefore the *logical type* of the law, not the algebra of the interpolator and not the numerical value of $g_\dagger$.

E.2 Why the first NAMCG-shaped interpolator failed

Chapter 6 records the large-separation scaling

$$I(A:B) \;\sim\; c_{p_*} \left(\frac{\ell_A \ell_B}{d^2} \right)^{p_*}, \qquad d \gg \ell_A, \ell_B, \tag{E.3}$$

with p_* and c_{p_*} defined as in Eq. (6.10): p_* is the smallest sum of scaling dimensions of operators whose product has vacuum quantum numbers. Cardy [3] obtains this form for the Rényi mutual information of compact regions in a conformal vacuum, with the von Neumann limit shown explicitly for free-scalar spheres; its use for NAMCG cores is an assumption. The first NAMCG-shaped interpolator nevertheless converted a gradient of this static tail into a local boost, producing the working family

$$\nu_{\rm exp} = \frac{1}{1 - \alpha \exp(-g_{\rm N}/g_s)}. \tag{E.4}$$

That conversion is not derived: a static separation gradient is not a retarded response or a local acceleration interpolator. On the same 175 SPARC files that family lost to simple MOND by a wide AIC gap. The transition was too sharp in $g_{\rm N}$ and needed a second amplitude α that disks do not want.

The failure is useful. It shows that "mutual information implies a boost" is not enough. The *argument of the exponential* matters. SPARC selects $\sqrt{g_{\rm N}}$ in that argument, not $g_{\rm N}$.

E.3 An algebraic identity, not a derivation

Write $y = \sqrt{x} = \sqrt{g_{\rm N}/g_\dagger}$. Then

$$\nu_{\rm RAR}(x) = 1 + \frac{1}{e^y - 1} = 1 + n_{\rm BE}(y), \tag{E.5}$$

where $n_{\rm BE}$ is the Bose–Einstein occupation of a single mode with dimensionless energy y. This is an identity. It does not, by itself, mean that galactic baryons are a Bose gas. It does mean that any microscopic argument that produces a thermal factor $e^{-\sqrt{g_{\rm N}/g_s}}$ in a screen occupation, rather than $e^{-g_{\rm N}/g_s}$, lands on the published interpolator automatically.

The linear-exponential (E.4) is the Boltzmann factor of a scale set by $g_{\rm N}$ itself. The RAR is the Bose factor of a scale set by $\sqrt{g_{\rm N}}$. That is the whole algebraic distinction.

E.4 A modular conjecture

Chapter 6 already assigns a diamond an Unruh temperature $T_{\rm Unruh} = \kappa/2\pi$ and a modular Hamiltonian H_D. Chapter 12 already assigns the cosmology an infrared boundary scale through $\Lambda_{\rm eff}$. Milgrom's coincidence [7],

$$2\pi g_\dagger \;\sim\; cH_0 \;\sim\; c^2\sqrt{\Lambda/3}, \tag{E.6}$$

says that the SPARC scale sits, to a factor of order 2π, on the same infrared accelerations NAMCG already possesses as diamond and boundary data.

Assumption E.1 (Square-root modular response)**.** For a slowly varying disk of condensed parcels, the leading reversible variation of $U_{\rm ent}$ with respect to a local Newtonian acceleration is thermal in the variable

$$y = \sqrt{\frac{g_{\rm N}}{g_s}}, \qquad g_s \sim \frac{\kappa_{\rm IR}}{2\pi}, \tag{E.7}$$

where $\kappa_{\rm IR}$ is a putative surface gravity of an infrared boundary, not a new galactic parameter. The manuscript does not supply a specific boundary geometry from which it is defined, and the order-one factor in $g_s \sim \kappa_{\rm IR}/(2\pi)$ is a working convention of this assumption, not fixed here; both are part of Problem H (Section H.8). The occupation that weights the extra stress is $n_{\rm BE}(y)$. Then $\nu = 1 + n_{\rm BE}(y)$ is (E.2). For the empirical comparison, setting $g_s = g_\dagger$ reproduces (E.2); this identification is phenomenological and does not determine $g_s/\kappa_{\rm IR}$.

Assumption E.1 is the missing sentence of Problem H. It is not implied by Theorem 7.3. It is not implied by the conformal tail (E.3). It is the simplest contact that turns the book's modular objects into the interpolator SPARC already selected.

Why a square root rather than a linear Unruh factor $T \propto g_{\rm N}$ is not derived here. A linear factor reproduces (E.4) and has already failed. A justification would have to come from the spectrum of H_D on a diamond whose size is set by the local baryonic curvature radius, or from a two-horizon sharing between the local diamond and the HBEV boundary. That calculation is the content of Problem H, restated.

E.5 What this bridge excludes

Verlinde's 2016 emergent-gravity formula for finite disks depends explicitly on radius as well as on $g_{\rm N}$. Lelli, McGaugh and Schombert found that those residuals correlate with radius and that acceptable fits require stellar mass-to-light ratios in tension with population synthesis [8]. NAMCG cannot take that route. Assumption 13.2 plus the RAR residuals already require a map of $g_{\rm N}$ only.

A cosmological running $g_s(z) \propto H(z)$ is likewise not licensed by this appendix. Equation (E.6) is a present-day coincidence used to name g_s. Promoting it to $g_s(z)$ is extra physics and would have to survive high-redshift rotation curves before it is written as a NAMCG result.

Isolated application of (E.2) to wide binaries is likewise not licensed. Assumption 13.3 saturates the boost at $\nu_{\surd}(g_{\rm ext})$. A Gaia sample consistent with Newton constrains the isolated reading, not Theorem 7.3. A reported $\gamma \approx 1.4$ that survives quality cuts is not explained by $\nu_{\surd}(g_{\rm gal}) \approx 1.1$; that residual, if real, is still open.

E.6 How the bridge would be confirmed or killed

Confirmation. The published diamond first law recovers Einstein gravity at first order in the metric, hence $\nu \to 1$. An interpolator requires the modular spectrum of K_D for a weak-field diamond together with a putative infrared boundary datum $\kappa_{\rm IR}$. Neither the manuscript's Chapters 6 and 12 nor this appendix specifies a stationary boundary geometry from which $\kappa_{\rm IR}$ is defined or normalized; constructing it is part of Problem H. Problem H is solved if the weighted multimode response of the extra stress (Section H.8), or a justified single-ladder response, reproduces the target interpolator of Chapter 17 with no SPARC input. In the single-ladder case this requires $\Delta k = \sqrt{g_{\rm N}/g_s}$ with $g_s = c_\star \kappa_{\rm IR}$ and $c_\star$ fixed by the calculation. Then g_s is not a SPARC fit, and Appendix D becomes a consistency check. A computed spectrum that fails to reproduce the target does not solve Problem H. A linear gap $\Delta k \propto g_{\rm N}/\kappa_{\rm IR}$ would reproduce the retired $\nu_{\rm exp}$ and kill Assumption E.1.

The same computation must grow an external-field argument before wide binaries are a prediction. Until it does, Assumption 13.3 is a coefficient in the same sense as Assumption 13.2: it is the QUMOND external-field limit written with $\nu_{\surd}$, not a lemma from $I(A:B)$.

The explicit modular Hamiltonian of de Sitter diamonds [9] is the operator to use for the gap test (17.7). The Euclidean Schwarzschild–de Sitter action is minus the sum of the two horizon entropies, not a geometric mean of surface gravities; that instanton does not by itself produce $\sqrt{g_{\mathrm{N}}/\kappa_{\mathrm{IR}}}$. The numerical coincidence $cH_0/2\pi \sim g_\dagger$ may be recorded after Δk is computed, not used to fix g_s.

Failure. The same computation produces (E.4), a radius-dependent kernel, or a light ψ. Then disks may still follow the RAR, but not as a NAMCG prediction. Assumption 13.2 remains a coefficient; Problem H remains open; Appendix D is unchanged.

E.7 One paragraph for Chapter 13

The working disk law is (E.2). The reason NAMCG is allowed to use it on an isolated disk is Assumption 13.2. The reason a subsystem embedded in the Galaxy must not use the isolated evaluation is Assumption 13.3. The reason the book might eventually own either sentence is Assumption E.1. Until the last is a lemma, the first two are borrowed, and the galactic sector is the RAR with the external-field effect.

E.8 Exercises

1. Prove (E.5) in three lines.

2. Expand ν_{RAR} and ν_{exp} for $x \ll 1$ and $x \gg 1$. Where do they first differ by order one?

3. Using $H_0 = 70\,\mathrm{km\,s^{-1}\,Mpc^{-1}}$, compute cH_0 and $cH_0/2\pi$ in $\mathrm{m\,s^{-2}}$. Compare with $g_\dagger$. This is a coincidence check, not a derivation.

4. Why does an explicit r in ν, as in Verlinde's finite-disk formula, threaten the residual structure reported by [6]?

5. **External field.** Take $g_{\mathrm{ext}} = 2.2 \times 10^{-10}\,\mathrm{m\,s^{-2}}$ and $g_{\mathrm{N}} = 10^{-11}\,\mathrm{m\,s^{-2}}$. Compute $\nu_{\surd}(g_{\mathrm{N}})$ and $\nu_{\surd}(g_{\mathrm{ext}})$. Which one is the legal boost under Assumption 13.3, and which published wide-binary claim does each contact?

Appendix F

Einstein Gravity Plus a Conserved Extra Tensor

This appendix writes the infrared field equation of Theorems 7.3–11.1 next to Einstein's equation. It introduces no new hypothesis.

F.1 Einstein

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = 8\pi G\, T^{\text{matter}}_{\mu\nu}. \tag{F.1}$$

Bianchi identities force $\nabla^{\mu} T^{\text{matter}}_{\mu\nu} = 0$ on shell.

F.2 NAMCG infrared equation

After the composite metric, $m_T = 0$, the heat-kernel map, Lorentz locking, and conservation are granted,

$$G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = 8\pi G_{\text{eff}} \big(T^{\text{matter}}_{\mu\nu} + T^{\text{tensor}}_{\mu\nu} + T^{\text{ent}}_{\mu\nu}\big). \tag{F.2}$$

$G_{\mu\nu}$ is the Einstein tensor of the composite metric $g_{\mu\nu} = \eta_{AB} e^{A}{}_{\mu} e^{B}{}_{\nu}$. Λ_{eff} is the coefficient of Theorem 12.4; G_{eff} is to be matched by the conditional sector calculation of Problem F. The extra source is

$$T^{\text{ent}}_{\mu\nu} = -\frac{2}{\sqrt{-g}} \frac{\delta U_{\text{ent}}[I(A:B)]}{\delta g^{\mu\nu}} = T^{\text{ent,rev}}_{\mu\nu} + T^{\text{ent,irrev}}_{\mu\nu}. \tag{F.3}$$

Chapter 11 requires

$$\nabla^{\mu} \big(T^{\text{matter}}_{\mu\nu} + T^{\text{tensor}}_{\mu\nu} + T^{\text{ent}}_{\mu\nu}\big) = 0. \tag{F.4}$$

In the infrared, $T^{\text{tensor}}_{\mu\nu}$ is absorbed into the Einstein tensor of $g_{\mu\nu}$. What can appear as extra gravity in static disks is $T^{\text{ent,rev}}_{\mu\nu}$.

F.3 Weak field

Newtonian gauge, static source:

$$\nabla^2\Phi = 4\pi G_{\rm eff}(\rho_{\rm b} + \rho_{\rm ent}), \qquad \rho_{\rm ent} = T_{00}^{\rm ent,rev}. \tag{F.5}$$

Assumption 13.2 slaved $\rho_{\rm ent}$ to $\rho_{\rm b}$. The interpolator $\nu_{\surd}$ of Appendix D is a coefficient choice for that assumption, not a consequence of (F.2).

F.4 What this does and does not change

High acceleration: $T_{\mu\nu}^{\rm ent,rev} \to 0$, so (F.2) returns to (F.1) with $(G_{\rm eff}, \Lambda_{\rm eff})$ in place of (G, Λ). Low acceleration: $\rho_{\rm ent}$ may dominate $\rho_{\rm b}$ without a particle halo, if and only if Assumption 13.2 holds. Clusters, the CMB, and colliding systems test that assumption, not $G_{\mu\nu}$. Gravitational waves test $m_T = 0$, not (F.3). The irreversible drag of Chapter 14 is a phenomenological addition, not a static extra potential and not derived from (F.3).

F.5 One-line summary

Einstein equates curvature to ordinary stress. NAMCG equates the same curvature to ordinary stress plus a divergenceless entanglement stress of baryonic regions.

Appendix G

Heat-Kernel Area Term on a Causal Diamond

This appendix records the standard heat-kernel contribution to the entanglement entropy of a causal diamond. It does not derive the equilibrium condition $\delta S_D = \delta\langle H_D\rangle$ and does not recover the Einstein equation from the matrix sector alone.

G.1 Replica formula and cone expansion

The von Neumann entropy of the reduced state on a diamond D is obtained from the replica trick,

$$S_D = (1 - n\partial_n) \ln Z(n)\Big|_{n=1}, \tag{G.1}$$

where $Z(n)$ is the partition function on the n-sheeted cover branched along the entangling surface Σ. For a Laplace-type fluctuation operator on the emergent metric (Assumption 8.4) the heat-kernel expansion on the cone produces the area term

$$S_D = \frac{\mathrm{Area}(\Sigma)}{4G_{\mathrm{eff}}} + S_{\mathrm{pot}}, \tag{G.2}$$

with the prefactor given by the Seeley–DeWitt coefficient that multiplies the deficit. This is the standard cone computation for operators whose endomorphism contains no explicit curvature coupling, on a replica geometry on which the connection and the endomorphism stay bounded. For $E = E_0 + QR$ with the conical delta of R not continued into E, the replica entropy differs from $\mathrm{Area}(\Sigma)/(4G_{\mathrm{eff}})$, with G_{eff} the curvature coefficient of Theorem 8.2, by a contact term proportional to $\mathrm{tr}\, Q$; if the delta is continued into E, the difference vanishes. For operators built from a composite metric the curvature coefficient depends on the independent variables chosen (Section H.2). Chapter 6 and Appendix B use $1/(4G_{\mathrm{eff}})$ as the name of the

coefficient of δA. The conditional heat-kernel identity of Theorem 8.2 does not fix the numerical value of G_{eff}; the coefficients A, B, C and physical matching remain open in Problem F.

G.2 What is not derived

- The equilibrium condition $\delta S_D = \delta\langle H_D\rangle$ under fixed volume. It is the hypothesis of Chapters 6–7 that forces $m_T = 0$. It is not obtained as a path-integral Ward identity from the replica variation alone.
- An evaluation of the fluctuation operator of the locked NAMCG condensate (including diffeomorphism and local-Lorentz ghosts) that would determine the precise endomorphism contribution to the area coefficient.
- Recovery of the infrared equation. Once the area law and the equilibrium condition are both assumed, the Einstein equation with a matrix source follows by the usual Jacobson argument; that assumption is not removed.

G.3 Status

The area term in S_D is the standard heat-kernel term for the operators of Section G.1. The appendix adds no new derivation of the Ward identity, no scheme-fixed coefficient, and no alteration to the open-problem ledger of Appendix H.

Appendix H

Status of the Open Problems

This appendix records the precise status of Problems A–J and N as of the present text. It collects standard supporting calculations that are already rigorous, states the exact missing step for each problem, and does not convert any open problem into a lemma. The research agenda of Chapter 17 remains the authoritative list; this appendix is a working ledger. A status of OPEN is never paired with an unqualified completion claim; a reduced-scope result is labelled as such.

H.1 Problem A: matrix-to-modular map

Ledger label. Status: OPEN. Priority: CRITICAL PATH for the matrix-derived label. Completion: PROGRAMME-CONDITIONAL. The matrix-to-modular map remains unproved. A scope-limited Presentation-B model may retain Assumption 3.6 and claims no derivation from the matrix model.

For a constant-Θ Moyal realization, the commutator of a coordinate symbol x_μ with a field is exact, with no higher-derivative correction; see Remark 3.7. That result is conditional on this realization and is not a general expansion for fluctuating $\hat{X}_\mu$. The remark also records a metric-matching obstruction for a real constant Θ. Neither result supplies the map from $(\hat{X}, \Phi)$ to the causal-diamond modular Hamiltonian H_D or the mutual-information functional $U_{\text{ent}}[I(A:B)]$ before the composite metric is defined.

Missing step. An error-controlled map from the matrix order parameter (or from $[\hat{X}, \Phi]$) into the modular Hamiltonian H_D of a causal diamond, or into the mutual-information functional $U_{\text{ent}}[I(A:B)]$, that holds before the composite metric is defined. Until that map is a lemma, Chapter 8 computes a heat kernel in a theory whose relation to Chapter 3 is still Assumption 3.6.

H.2 Problem B: replica Ward identity

Ledger label. Status: OPEN. Priority: CRITICAL PATH. Completion: PROGRAMME-CONDITIONAL. The first-law identity is established, but the stationarity condition for the tensor block is not derived.

For a differentiable family of normalized reduced states on a fixed diamond algebra, the first variation of the entropy obeys the standard first-law identity

$$\delta S_D = \mathrm{Tr}(\delta\rho_D H_D^{(0)}) = \delta\langle H_D^{(0)}\rangle, \qquad H_D^{(0)} = -\ln \rho_D^{(0)},$$

where the modular Hamiltonian is held fixed at the reference state. Under the usual path-integral assumptions, the replica construction expresses this variation as a response of the microscopic action insertion on the branched cover. These identities do not imply $\delta S_D = 0$.

Missing step. Derive from the microscopic dynamics the entanglement-stationarity condition $\delta S_D = 0$ for the allowed fixed-volume variations, and establish the area contribution and its relation to the *emergent* metric without presupposing that metric or the fluctuation split of Chapter 5. For variations of the auxiliary scaffold, the replica insertion computes a response to $g_{\mu\nu}^{(0)}$; relating it to the emergent metric still requires a specified map between them. Problem B therefore remains open, although the fixed-algebra first-law identity is established.

Which first law is established (conditional; not closure). The identity above is a statement on a fixed diamond algebra: only the state varies, and $H_D^{(0)}$ is held fixed. As stated in the ledger, it does not imply $\delta S_D = 0$. The first law used in Chapter 6, Equation (6.7), also contains the area term $\delta A/(4G_{\mathrm{eff}})$, which arises when the diamond itself varies with the emergent metric. That is a variation of the algebra, not of the state on a fixed algebra, so the fixed-algebra identity does not establish it. Obtaining it requires the map between variations of the scaffold diamond and of the emergent diamond noted above. In Chapter 6 the coefficient of δA is only named $1/(4G_{\mathrm{eff}})$; its identification with the heat-kernel G_{eff} is deferred to Chapter 8 and depends on the open Problem F. The form of Assumption 6.1 stated in Chapter 6, Equation (6.8), is an assumption that goes beyond the fixed-algebra identity. This paragraph records that distinction; it does not derive the area term or the stationarity condition, and Problem B remains open.

Replica insertion identity (conditional; not closure). Let $Z_n(\varepsilon)$ be the Euclidean partition function on a regulated n-sheeted cover. Assume the usual path-integral assumptions and analytic continuation in n from integer replica number to a

neighborhood of $n = 1$. As ε varies, hold fixed the regulated geometry, branch locus, conical-locus data, and integration measure. Include the conical-locus contributions in $S_{E,n}(\varepsilon)$, and define

$$W_n(\varepsilon) = \ln Z_n(\varepsilon) - n \ln Z_1(\varepsilon), \qquad S_D(\varepsilon) = (1 - n\partial_n) W_n(\varepsilon)\big|_{n=1}, \qquad \mathcal{I}_n := \partial_\varepsilon S_{E,n}.$$

Here $\mathcal{I}_n$ is the total action insertion over the whole n-sheeted cover, not an insertion per sheet. Differentiating the normalized replica formula and using $\partial_\varepsilon \ln Z_n = -\langle \mathcal{I}_n \rangle_n$ gives

$$\partial_\varepsilon S_D = \partial_n \langle \mathcal{I}_n \rangle_n|_{n=1} - \langle \mathcal{I}_1 \rangle_1.$$

Thus $\partial_\varepsilon S_D = 0$ is equivalent, under these assumptions, to

$$\partial_n \langle \mathcal{I}_n \rangle_n|_{n=1} = \langle \mathcal{I}_1 \rangle_1.$$

The replica formula alone does not establish this equality. If a deformation changes the geometry, branch locus, conical-locus data, or integration measure, the corresponding variations must also be included. Applying the identity to non-conformal tensor-block variations still requires a specified source or background deformation representing that block in the replica action, together with the map from the auxiliary scaffold $g^{(0)}$ to the emergent metric and the corresponding area variation. Those inputs are not supplied by the displayed microscopic action. Problems B, C, D, F and G remain open.

Replica cone coefficient of the Poisson-limit operators and the area term of the first law (conditional; not closure). Keep the setting of the Presentation A Poisson-limit paragraphs of Section H.6: four matrix coordinates, the Poisson limit, the Euclidean section $\eta = \delta$, the scalar operator $\mathcal{O}_s$, the connection operator $\mathcal{O}_v$ acting on the internal index of a^μ in background gauge, the ghost operator of weight -2, $G = \rho\, g_\rho$ with $\sqrt{|G|} = \rho^2$, Assumption 8.4, and one proper-time limit $s_{\min,s} = \Lambda_{\mathrm{UV},s}^{-2}$ per sector, shared by the effective action and the replica entropy. Only the $\Lambda_{\mathrm{UV},s}^2$ term is compared, for constant $\Lambda_{\mathrm{UV},s}$; neither a value of G_{eff} nor the relation $S = A/4G$ is assumed.

Replica on the matrix chart. Let Z_n be the partition function on the n-sheeted cover of the chart x, branched over a two-surface Σ, with every background field (Θ, hence G, g_ρ, ρ, the connections and the endomorphisms) pulled back. Then G and g_ρ are conical with the same angle $2\pi n$ and ρ is a regular density of the chart: the cone is on G and g_ρ together. For a Laplace-type sector whose connection and endomorphism stay bounded on the cover, the leading surface term of the heat trace is the flat-cone

term [10],

$$\operatorname{Tr} e^{-s\mathcal{O}}\big|_n - n \operatorname{Tr} e^{-s\mathcal{O}}\big|_1 = \operatorname{tr}\mathbb{I}\, \frac{1}{12}\Big(\frac{1}{n} - n\Big)\frac{A_G(\Sigma)}{4\pi s} + O(s^0),$$

independent of the connection and of the endomorphism. The fibre index of $\mathcal{O}_v$ is internal, so the fibre is not rotated around Σ by a Levi-Civita holonomy. With $S = (1 - n\partial_n)(\ln Z_n - n \ln Z_1)|_{n=1}$ and $\ln Z_n = \frac{1}{2}\sum_s w_s \int \mathrm{d}s\, s^{-1} \operatorname{Tr} e^{-s\mathcal{O}_s}$,

$$S = \frac{A_G(\Sigma)}{48\pi} \sum_s w_s \operatorname{tr}\mathbb{I}_s\, \Lambda^2_{\mathrm{UV},s}$$

(the μ_s^2 terms of Theorem 8.2 subtracted). The same limits in Equation (8.9) give, for the coefficient of $\int \sqrt{G}\, R_G$ when the endomorphisms carry no explicit curvature, $1/(16\pi G) = \sum_s w_s \operatorname{tr}\mathbb{I}_s\, \Lambda^2_{\mathrm{UV},s}/(192\pi^2)$. Hence $S = A_G(\Sigma)/(4G)$ with that G, in the variables $(G, \rho, \partial\Theta)$ of the contact-term paragraph below: for these operators the coefficient of δA in Equation (6.7) is the Λ^2_{UV} coefficient of Theorem 8.2, namely the coefficient $A = 1/(12\pi)$ of Equation (8.12) per unit weighted fibre trace, as in the sector-sum paragraph of Section H.6, and it is determined by the operators and the cutoff. This holds for the replica and variables just stated; the scope of the whole block is given at the end of the second paragraph below.

Contact term. If an endomorphism is $E_s = E_{0,s} + Q_s R_G$ and R_G is continued onto the cone with its conical delta, the heat trace gains $-(1-n)\operatorname{tr} Q_s\, A_G/(4\pi s)$ and the entropy changes by $-\frac{A_G}{8\pi}\sum_s w_s \operatorname{tr} Q_s \Lambda^2_{\mathrm{UV},s} = -2\pi A_G \sum_s w_s \operatorname{tr} Q_s \langle\phi^2\rangle_s$, $\langle\phi^2\rangle_s = \Lambda^2_{\mathrm{UV},s}/(16\pi^2)$. This is the contact term of Susskind and Uglum [11], $S_{\mathrm{BH}} = S_{\mathrm{ent}} - Q$ in Equations (322)–(323) of [10], with S_{BH} built from the coefficient $\operatorname{tr}(\frac{1}{6} - Q_s)$ of Theorem 8.2. With $(G, \rho, \partial\Theta)$ as independent variables, the scalar endomorphism $E_s = -\frac{1}{2}\Box_G \ln\rho + \frac{1}{4}|\partial \ln\rho|^2_G$, the connection endomorphism with $\operatorname{tr} E_v = 4E_s + \operatorname{tr}(B_\sigma B^\sigma)$, $\operatorname{tr} B_\sigma = 0$, $\operatorname{tr}(B_\sigma B^\sigma) = -X$, and the ghost operator $-\mathcal{O}_s$ all have $Q = 0$ and are bounded on the pulled-back cover, so the contact term vanishes sector by sector.

Dependence on the independent variables. The Weitzenböck identity quoted in the paragraph on the torsion invariant gives $E_s = \frac{1}{4}R_G + \frac{1}{16}T_{\lambda\mu\nu}T^{\lambda\mu\nu} + \frac{1}{8}T_{\lambda\mu\nu}T^{\mu\lambda\nu}$ and $\operatorname{tr}(B_\sigma B^\sigma) = -T_{\lambda\mu\nu}T^{\lambda\mu\nu}$ (the first checked at random two-jets of a closed ω, for both signatures). With the coframe torsion in place of $\partial\Theta$ as independent variable, the scalar operator has $Q_s = \frac{1}{4}$, and its coefficient $\mathcal{A}_{1,R,s}$ is $\frac{1}{6} - \frac{1}{4} = -\frac{1}{12}$ per component instead of $\frac{1}{6}$, of opposite sign. The coefficient $\mathcal{A}_{1,R,s}$ of Theorem 8.2 is therefore not a function of these operators alone. With $N_{\mathrm{tot}} = N_s + 2$ as in the Poisson-limit paragraph, the weighted coefficient c of $\sqrt{G}\, R_G$ in $\operatorname{tr}(R_G/6 - E)$ is:

sector (weight)	c in $(G, \rho, \partial\Theta)$	c in $(G, \text{torsion})$
N_s scalar components	$N_s/6$	$-N_s/12$
connection, 4 components	$2/3$	$-1/3$
ghost, weight -2	$-1/3$	$1/6$
total	$N_{\text{tot}}/6$	$-N_{\text{tot}}/12$

Which geometry carries the cone. The first column is the replica entropy above, $S = (\Lambda_{\text{UV}}^2 A_G/8\pi)\, c$, with ρ regular. The cone on G can instead be carried by the Poisson structure: for $\Theta^{12} = 1/f(x^1, x^2)$, $\Theta^{34} = 1$, one has $G_\perp = f^2\delta$ and $\rho = f$, and $f = |z|^{n-1}$ makes G conical of angle $2\pi n$ with $\rho \propto r^{n-1}$ singular, g_ρ singular, and torsion unbounded, $|T|^2 \propto (n-1)^2/u^2$ at proper distance u from Σ. Then E_s contains $-\frac{1}{2}\Box_G \ln\rho = \pi(1-n)\delta_\Sigma$, a contact term with $Q = \frac{1}{4}$, and $\mathcal{O}_s$ reduces to a Bessel problem of index $\sqrt{m^2 + ((n-1)/2)^2}/n$. With the branch regular at $m = 0$, the first-order coefficient is that of the second column, for the scalar and ghost and for the components of a^μ in this block class (a statement conditional on that branch and not used below; the factor $-\frac{1}{2}$ is also what the local a_1 coefficient with the delta function, $Q = \frac{1}{4}$, gives). The two choices differ by the factor $-\frac{1}{2}$ in every sector (for the branch stated). The first is the replica of the same state, since $\operatorname{Tr}\rho_D^n$ is built on the cover of the same background; the second changes the background with n and corresponds to a smooth-coframe reading of the induced action. Only the first is used in the identification above.

Normalisation of the tensor kinetic term against the area coefficient (conditional; not closure). Assume Equation (6.8) for every small diamond, every unit timelike direction of ξ and the fixed-volume variations of the tensor block, with the geometry of Equation (7.9) in the form of [12], in which $\delta A|_V$ is proportional to $G_{00}^{(1)}[h]$, and write $1/(4G_A)$ for the coefficient of δA. A symmetric tensor whose contraction with every timelike unit vector vanishes is zero, so the condition is the tensor equation $G_{\mu\nu}^{(1)}[h] = 8\pi G_A\, \delta\langle T_{\mu\nu}^{\text{op}}\rangle[h]$ up to a term proportional to $\eta_{\mu\nu}$. For a tensor block with the Fierz–Pauli structure of Equation (7.7) this requires both $m_T = 0$ and $G_T = G_A$, where $1/(16\pi G_T)$ is the coefficient with which the quadratic action whose metric response defines T^{op} carries the linearised Einstein form. The second condition is not implied by the first, by the identification of the area prefactor with the coefficient of Theorem 8.2, or by Proposition 6.5: it equates the coefficient of the quadratic term of the block with the coefficient of the area term.

In the class of the Poisson-limit paragraphs, about a constant $\bar{\Theta}$, the induced Λ_{UV}^2 quadratic form on the two modes with $\delta\rho = 0$ is $-\frac{\Lambda_{\text{UV}}^2}{32\pi^2}\bar{\rho}^2 p_\eta^2\,(1 - N_{\text{tot}}/24)\,\mathcal{M}$, with $\mathcal{M}$ the tree-level kernel of Remark 3.8. A term $c\,\rho R[g_\rho]$ has eigenvalue $-c/4$ on these modes, so the form is that of an Einstein term with $c_T = N_{\text{tot}}/6 - 4$, whereas the area coefficient above, for the replica of the first paragraph, is $c_A = N_{\text{tot}}/6$. With a

common cutoff $1/G_T = (N_{\rm tot} - 24)\Lambda_{\rm UV}^2/(12\pi)$ and $1/G_A = N_{\rm tot}\Lambda_{\rm UV}^2/(12\pi)$; with the connection sector at its own limit $\Lambda_{{\rm UV},v}$, $1/G_A - 1/G_T = 2\Lambda_{{\rm UV},v}^2/\pi$, independent of N_s. The difference is the term $\rho^2 X$ of the connection sector, which is bounded on the replica cover and contributes no entropy but contributes to the quadratic form; for scalar content alone $G_T = G_A$. In this class (Poisson limit, Euclidean section, background Feynman gauge, one loop, $\Lambda_{\rm UV}^2$ term, the two modes with $\delta\rho = 0$) and for the replica of the first paragraph, the induced term therefore has the Einstein structure but not the coefficient that $G_T = G_A$ requires, for every N_s and every $\Lambda_{{\rm UV},v} > 0$.

These statements are for the Poisson-limit operators of the preceding paragraphs at one loop, for the $\Lambda_{\rm UV}^2$ term, the replica above and the background gauge stated. They do not derive Assumption 6.1, fixed-volume stationarity of the tensor block, or the identification of G with the emergent metric (Assumption 8.4); they do not treat the tree-level term (of lower order in momentum and not of Einstein form), fermions or additional scalars, Moyal corrections at the cutoff, the mixing of Φ with a^μ, edge modes or the interpretation of the ghost-weighted sum as an entropy, or the logarithmic terms; the off-shell quadratic form is gauge dependent; and they give no value or sign of $G_{\rm eff}$. Problems B, C, D, F and G remain open.

Gauge parameter and gauge function in the comparison of G_T with G_A (conditional; not closure). Keep the setting, the class and the normalisation of the two preceding paragraphs. Write e_i for the contribution of a sector of weight w_i to the bracket $1 - N_{\rm tot}/24$ of the preceding paragraph, so that $\sum_i w_i e_i$ is the net eigenvalue on the two modes with $\delta\rho = 0$ and $c_T = -4\sum_i w_i e_i$; a scalar component has $e = -\frac{1}{24}$ there and $-\frac{1}{8}$ on the mode with $\delta\rho \neq 0$. Let $D_\mu = \{x_\mu, \cdot\}$, anti-self-adjoint on $L^2(\rho\,{\rm d}^4x)$, $({\rm Gr}\varphi)_\nu = D_\nu\varphi$, and $\mathfrak{J}_\nu = D^\mu\Theta_{\mu\nu}$, the Poisson limit of the tree-level equation of motion; for the perturbation of the quadratic-form paragraph $\mathfrak{J}_\nu = -\epsilon\cos(p\cdot x)\,(Mu)_\nu$, with $M = q^2\eta - qq$ the kernel of Remark 3.8. The background gauge $\frac{1}{2\alpha}\rho\,(D_\mu a^\mu)^2$, $\alpha > 0$, $\kappa = 1 - 1/\alpha$, leaves the ghost operator unchanged and replaces $\mathcal{O}_v$ by the non-minimal operator $\mathcal{O}_\kappa = \mathcal{O}_v - \kappa\,{\rm Gr}{\rm Gr}^\dagger$, with principal symbol $q^2\mathbb{I} - \kappa\,qq^T$; one proper-time cut-off on $\mathcal{O}_\kappa$ serves the effective action and the replica, and the normalisation of the gauge-fixing weight is local and carries no two-derivative term.

For every Poisson bivector ${\rm Gr}^\dagger{\rm Gr} = \mathcal{O}_s$ and $\mathcal{O}_v{\rm Gr} = {\rm Gr}\mathcal{O}_s - \{\mathfrak{J}_\nu, \cdot\}$: the commutator terms are fixed by $[D_\mu, D_\nu] = \{\Theta_{\mu\nu}, \cdot\}$ and the Leibniz property of D_μ on the bracket, and the $\{\Theta_{\mu\nu}, \cdot\}$ coupling of $\mathcal{O}_v$ cancels them up to $\mathfrak{J}$ (checked in exact arithmetic for a nonlinear bivector). Hence for $\mathfrak{J} = 0$, $\mathcal{O}_\kappa{\rm Gr} = (1-\kappa){\rm Gr}\mathcal{O}_s$, the longitudinal sector is the scalar operator with proper time rescaled by $1 - \kappa$, and up to zero modes ${\rm Tr}\,e^{-s\mathcal{O}_\kappa} = {\rm Tr}\,e^{-s\mathcal{O}_v} + {\rm Tr}\,e^{-s(1-\kappa)\mathcal{O}_s} - {\rm Tr}\,e^{-s\mathcal{O}_s}$. The integrated $\Lambda_{\rm UV}^2$ density of the connection sector is then $\frac{3+\alpha}{6}\,\rho R[g_\rho] + \rho^2 X$ on such backgrounds, with the coefficient 1 of $\rho^2 X$ independent of α. On the replica cover with constant $\bar\Theta$ the same intertwining, with the domain at Σ for which the longitudinal sector is the image

of the regular scalar domain, gives the area coefficient $\frac{3+\alpha}{6}$ for the connection sector in the units c of the table above, equal to the coefficient of $\rho R[g_\rho]$. This depends on that choice of domain; other choices are not examined.

About constant $\bar\Theta$, with $\mathfrak{J} \neq 0$, a numerical second-order evaluation of the heat trace (momentum-space Duhamel expansion, Gauss–Hermite quadrature, contour extraction of the p^4 coefficient; it reproduces the scalar value above, the $\alpha = 1$ values $\frac{5}{6}$ and $\frac{1}{2}$ of the connection sector, and the weight $3 + \alpha^2$ of the Λ^4_{UV} term) gives for the connection sector the values, matched by the rational forms, $\frac{5}{6} - \frac{\alpha-1}{24}$ on the two modes with $\delta\rho = 0$ and $\frac{1}{2} - \frac{\alpha-1}{8} + \frac{\alpha^2-1}{4}$ on the mode with $\delta\rho \neq 0$ ($\alpha = \frac{1}{3}, \frac{1}{2}, \frac{2}{3}, 1, 2, 3$ in one background, $\frac{1}{2}, 1, 2$ in another; rational fits to 2×10^{-3}, not derived; $\alpha = \frac{1}{3}, \dots, 3$ only). The two modes with $\delta\rho = 0$ thus show the α-dependence of the $\mathfrak{J} = 0$ result, and the remainder, $\frac{\alpha^2-1}{2}\rho|\nabla \ln \rho|^2_{g_\rho}$, acts on the mode with $\delta\rho \neq 0$ only. With scalar and ghost sectors independent of α, $c_T(\alpha) = \frac{N_s+\alpha-1}{6} - \frac{11}{3}$ and the replica coefficient is $c_A(\alpha) = \frac{N_s+\alpha+1}{6}$, so $1/G_A - 1/G_T = 2\Lambda^2_{\mathrm{UV}}/\pi$ for the tested range $\frac{1}{3} \leq \alpha \leq 3$; on backgrounds with $\mathfrak{J} = 0$ and under the conditions above the α-dependence cancels in the difference for every $\alpha > 0$. The TT value agrees with the exact lemma on $\mathfrak{J} = 0$ backgrounds. For $\alpha \neq 1$ the form is not spanned by $\rho R[g_\rho]$ and $\rho^2 X$; the third-mode to $\delta\rho = 0$ ratio of a term $c\,\rho R[g_\rho]$ is 3, and the net form attains this ratio at $\alpha = 3$, where the gap is unchanged, so that ratio does not diagnose the coefficient of $\rho^2 X$ outside the Feynman gauge. A gauge function $D_\mu a^\mu + c\,\mathfrak{J}_\mu a^\mu$ ($\alpha = 1$, ghost operator $-D_\mu D_\mu - c\mathfrak{J}_\mu D_\mu$) adds $c^2\rho^2\mathfrak{J}_\nu\mathfrak{J}_\nu$ to the density of the connection and ghost sectors together at quadratic order about $\bar\Theta$, that is $c^2q^2/(2p^2_\eta)$ to the eigenvalue on all three modes (two backgrounds, $c = \frac{1}{2}, 1, 2$); this lowers c_T at fixed c_A and depends on the direction of p unless $\bar\Theta$ is isotropic. Gauge functions with other first-derivative vectors are not examined.

Fermionic content and the connection-sector cone in the comparison of G_T with G_A (conditional; not closure). The displayed action is bosonic. Add n_D Dirac fields with action $\int \rho\,\bar\Psi\gamma^\mu D_\mu\Psi$, the Poisson limit of $\mathrm{Tr}\,\bar\Psi\Gamma^\mu[\hat X_\mu, \Psi]$, and the connection gauge $\alpha = 1$; the Yukawa couplings $\Gamma^i[\hat X_i, \Psi]$ to the N_s transverse matrices vanish on the background and are dropped. The operator $\mathcal{O}_f = -(\gamma^\mu D_\mu)^2 = \mathcal{O}_s - \frac{1}{2}\gamma^{\mu\nu}\{\Theta_{\mu\nu}, \cdot\}$ has the form (8.3) on G with $\omega_\sigma = -\frac{1}{2}\partial_\sigma \ln\rho\,\mathbb{I} + \beta_\sigma$, $\beta^\sigma = \frac{1}{4}\gamma^{\mu\nu}\Theta^{\rho\sigma}\partial_\rho\Theta_{\mu\nu}$, the spinor lift of $\mathcal{B}^\sigma$, and $\mathrm{tr}\,\beta_\sigma\beta^\sigma = -\frac{1}{2}X$ over four components (the vector value is $-X$); it contains no explicit curvature, so $\sqrt{|G|}\,\mathrm{tr}(\frac{1}{6}R_G - E_f) = \frac{4}{6}\rho R[g_\rho] + \frac{1}{2}\rho^2 X$, with eigenvalues $\frac{1}{3}, \frac{1}{3}, 0$ on the three modes (two backgrounds). The statistical weight is -1 per Dirac field ($\Gamma = -\frac{1}{2}\ln\det\mathcal{O}_f$ over four complex components). The coefficient of $\rho^2 X$ of the whole content is therefore $1 - n_D/2$: the weight of a field is its index under the rotation generated by $\mathcal{B}_\sigma$, and it vanishes for $n_D = 2$, the sixteen real components of the ten-dimensional maximally supersymmetric matrix model [13] with the Poisson background along four coordinates and $N_s = 6$. No symmetry of the displayed action

forces this. For n_D fields $c_T = \frac{N_s+\alpha-1}{6} + \frac{4n_D}{3} - \frac{11}{3}$. With the antiperiodic polar-frame condition, which gives the Cartesian-frame components the phase $(-1)^{n+1}$, the area coefficient of a Dirac field is that of two scalar components, $+\frac{1}{3}$ (Equations (126)–(127) of [10]; the zeta-function evaluation of the twisted cone gives $-\frac{1}{2}$ times the scalar value per component on the branch $n < 1$). Then $c_A = \frac{N_s+\alpha+1}{6} + \frac{n_D}{3}$ and $c_A - c_T = 4 - n_D$. This gluing transports the spinor index with the replica rotation; it is not the unrotated pull-back gluing used above for a^μ. With unrotated periodic gluing of the Cartesian components the area coefficient of a Dirac field is $-\frac{2}{3}$, equal to its coefficient of $\rho R[g_\rho]$, and $c_A - c_T = 4 - 2n_D$, which vanishes at $n_D = 2$ with $c_T = c_A = 0$: an equality of vanishing Λ^2_{UV} coefficients, not a derivation. The $\rho^2 X$ coefficient vanishes at $n_D = 2$; for antiperiodic gluing $c_A - c_T = 2$ there and equality needs $n_D = 4$, where $\rho^2 X$ has coefficient -1. The gluing selected by a quasi-free state with rotational modular flow is stated below. For $n_D = 2$, $N_s = 6$ one has $N_s + 2 - 4n_D = 0$, $c_T = 0$ and $c_A = 2$: the Λ^4_{UV} and Λ^2_{UV} tensor coefficients vanish, and the area coefficient is 2 for antiperiodic and 0 for periodic gluing.

If the internal index of a^μ is rotated with the replica angle, as for a vacuum invariant under the combined rotation, the area coefficient of the connection sector at $\alpha = 1$ is $-\frac{1}{3}$ on the branch $n < 1$ and $\frac{5}{3}$ on $n > 1$ instead of $\frac{2}{3}$; with the ghosts these are $-\frac{2}{3}$, which is $(D_1(4) - 6)/6$ of Equation (147) of [10], and $\frac{4}{3}$, instead of $\frac{1}{3}$; this function of n is not analytic at $n = 1$, and the gap is $3 - n_D$ or $5 - n_D$. These statements are for the Poisson-limit operators, the Λ^2_{UV} term, one loop, the quadratic form about constant $\bar{\Theta}$ on the two modes with $\delta\rho = 0$, the gauge families and the replica prescriptions stated. They do not derive the content, the fermion coupling, the gluing or the rotation of the internal index, which belong to Problem A, Assumption 6.1 or the modular flow; they do not treat Moyal corrections, gauge functions beyond those stated, Φ–a mixing, or the Kaluza–Klein sector. Problems B, C, D, F and G remain open.

Fermion replica gluing for $\mathrm{Tr}\,\rho^n$ and the supertrace character (conditional; not closure). Let the state of the Poisson-limit fermions on the wedge be quasi-free, and let its modular flow be the rotation by $2\pi t$ of the transverse plane of Σ, acting on every field through its representation of the rotation group (for integer spin, the identity at integer n). The ungraded trace $\mathrm{Tr}\,\rho^n$ then glues a Grassmann-odd field with the rotation by $2\pi n$ and one sign from the cyclicity of the trace: in the polar frame the spinor is antiperiodic under the angle $2\pi n$, and the Cartesian components of transverse spin $\sigma = \pm\frac{1}{2}$ acquire the monodromy $-e^{2\pi i n\sigma}$, which is $(-1)^{n+1}$ for integer n [14, 15]. Diagonalising the replica matrix gives the twist phases $e^{2\pi i k/n}$, $k = -(n-1)/2, \ldots, (n-1)/2$ [14]. For a block of two components with twist a, $|a| < \frac{1}{2}$, and the less singular self-adjoint extension of the first-order operator $\gamma^\mu D_\mu$, whose square is $\mathcal{O}_f$, the conical heat-trace constant is $a^2 + \frac{1}{6}$: the componentwise-regular value $B_2(|a|)$, $B_2(x) = x^2 - x + \frac{1}{6}$, plus $|a|$ from the reversed sign of the one Bessel order that

the first-order operator fixes. This extension is the one giving an analytic continuation: the componentwise-regular domain gives $\sum_k |a_k|$ equal to $n/4$ or $(n^2-1)/(4n)$ for even or odd n, a parity-dependent and non-analytic $\operatorname{Tr}\rho^n$. With the less singular extension, $\sum_k a_k^2 = (n^2-1)/(12n)$, analytic in n, equal to the zeta-function value of the polar-frame cone on the branch $n<1$, with $(1-n\partial_n)$ equal to $-\frac{1}{6}$ per block at $n=1$. The area coefficient of a Dirac field (two blocks, weight -1) is therefore $+\frac{1}{3}$, as used above for antiperiodic gluing.

The unrotated periodic gluing above has the phase $+1$ for all n: its twists are those of $\operatorname{Tr}\rho^n$ for odd n and those of the graded trace $\operatorname{Tr}[(-1)^F\rho^n]$ for even n, and with the componentwise-regular domain it gives -2 times the Dirac-domain value at odd n. It is not $\operatorname{Tr}\rho^n$, and for the stated extension its values $-\frac{2}{3}$ and $4-2n_D$ do not apply to $\operatorname{Tr}\rho^n$. The graded trace has the twist $a=\frac{1}{2}$ at $n=1$ and is not normalised there. The rotation by $2\pi n$ is the identity on the vector index for integer n, so the integer-n traces of the connection sector are those of the unrotated gluing; if the continuation from integer n is unique, the rotated branches $3-n_D$ and $5-n_D$ above do not reproduce them and are not continuations of $\operatorname{Tr}\rho^n$. With $c_A = \frac{N_s+\alpha+1}{6}+\frac{n_D}{3}$ and c_T as above, $c_A - c_T = 4(1-\frac{n_D}{2})+n_D$, the sum of the weight of $\rho^2 X$ and the excess $+1$ per Dirac field of the cone value $+\frac{1}{3}$ over its coefficient $-\frac{2}{3}$ of $\rho R[g_\rho]$. For $n_D=2$, $N_s=6$, $\alpha=1$: $c_T=0$, $c_A=2$.

In the character Q_{10} of [16], Eq. (3.7), with the flux along the four Poisson directions, the vector, six inert scalars, the ghosts and n_D Dirac fields in place of the sixteen-component spinor, the coefficient of order α^0 is $N_{\mathrm{tot}} = 8-4n_D$ and that of order α^2 is $-b\,(x^2+y^2)$ with $b = 1-n_D/2$, where $\pm\mathrm{i}x, \pm\mathrm{i}y$ are the eigenvalues of the flux. These are the weighted component count and the weighted index of the preceding paragraphs, so that $c_T = N_{\mathrm{tot}}/6 - 4b$ vanishes at $n_D=2$, $N_s=6$ because the orders α^0 and α^2 of the character vanish. At $n_D=2$ the order α^4 is $(x^2-y^2)^2/16$, which vanishes for self-dual or anti-self-dual flux. This concerns the character only, not the a_2 coefficient of the operators and not the area side. These statements are conditional on the quasi-free state, the rotational modular flow, the choice of self-adjoint extension, the uniqueness of the continuation from integer n, the domain of the first-order operator, the Λ_{UV}^2 order and the class of the preceding paragraphs; they do not derive the state, the flow, the content, or a value or sign of G_{eff}, and they do not compute the next order of either side. Problems B, C, D, F and G remain open.

Spin term, Lichnerowicz contact and the smooth first-order cone in the comparison of G_T with G_A (conditional; not closure). Keep the setting of the paragraphs on fermionic content and on fermion replica gluing: Poisson limit, Euclidean section, $\alpha=1$, the Λ_{UV}^2 term, N_s scalars, the connection with its ghosts and n_D Dirac fields, $N_{\mathrm{tot}} = N_s+2-4n_D$, and the coframe $e^A{}_\mu = \eta^{AB}\omega_{\mu B}$ with torsion T

of the paragraph on the torsion invariant. The *fixed-state replica* is the pulled-back replica of the paragraph on the replica cone coefficient (ρ regular, cone on G) with the antiperiodic Dirac gluing of the paragraph on fermion replica gluing. At random two-jets of a closed ω, the operator $\mathcal{O}_f$ of the fermionic-content paragraph satisfies $\frac{1}{4}\operatorname{tr} E_f = \frac{1}{4}R_G - \frac{1}{16}T_{\lambda\mu\nu}T^{\lambda\mu\nu} + \frac{1}{8}T_{\lambda\mu\nu}T^{\mu\lambda\nu}$, with no further term in the representation by R_G, T^2 and TT'; the scalar identity differs only in the sign of the first torsion term, so that $Q_f = Q_s = \frac{1}{4}$ in the torsion reading, whereas in the variables $(G, \rho, \partial\Theta)$ one has $Q_f = 0$. The coefficient of R_G in $\operatorname{tr}(\frac{1}{6}R_G - E_f)$ is then $-\frac{1}{3}$ over four components, and with weight -1 it gives $+\frac{1}{3}$ per Dirac field, the antiperiodic cone value, instead of $-\frac{2}{3}$. In the normalisation of the paragraph on fermion replica gluing, the n-derivative at $n = 1$ of the heat-trace constant of a two-component block is $\frac{1}{6}$ for the antiperiodic gluing with the less singular extension and $-\frac{1}{3}$ for the unrotated gluing with the componentwise-regular domain; the difference $\frac{1}{2} = 2Q_f$ equals the contact term of $Q_f R_G$ continued with its conical delta. This is an equality of n-derivatives at $n = 1$, not an identity in n. The excess $+1$ per Dirac field of the antiperiodic over the unrotated area coefficient is therefore the Lichnerowicz contact. It is already contained in c_A, it enters the area side only and raises c_A, and adding to it the contact of $Q_f R_G$, which at the weight -1 shifts c_A by $+1$ per Dirac field, would count it twice. For a Dirac field with the Levi-Civita spin connection the black-hole and entanglement entropies coincide [17]; for the torsion reading the corresponding equality (both $+\frac{1}{3}$ per Dirac field) is the computation above.

Smooth first-order cone. For $\omega = \bar{\omega} + \epsilon\,\mathrm{d}A$, with any constant antisymmetric invertible $\bar{\omega}$ and any A, one has to first order in ϵ $R_G^{(1)} = -2\Box_{\bar{G}}\,\delta\ln\rho$ and $\bar{\rho}\,R[g_\rho]^{(1)} = -\frac{1}{2}\bar{\rho}^2 R_G^{(1)}$. These were checked symbolically for a generic $\bar{\omega}$ and follow from the Weitzenböck identities of the paragraph on the torsion invariant. A conical deficit $2\pi(1-n)$ at first order in $n-1$ of a smooth Poisson structure about constant $\bar{\Theta}$ therefore requires $\ln\rho$ to carry the flux $\oint \nabla\ln\rho$ through Σ, that is a singular ρ; the fixed-state replica is not a limit of smooth Poisson structures at this order. This does not by itself exclude the fixed-state replica as the definition of $\operatorname{Tr}\rho^n$. For a regularised smooth family the first-order local density of every sector has $Q = \frac{1}{4}$, the factor $-\frac{1}{2}$ of the paragraph on the replica cone coefficient then carries no branch dependence, and the area coefficient is $c_A^{\mathrm{sm}} = -N_{\mathrm{tot}}/12$, the factor $-\frac{1}{2}$ times the coefficient of $\rho R[g_\rho]$ in every sector including the Dirac fields, for which it coincides with the antiperiodic value $+\frac{1}{3}$ without any gluing prescription. The branch issue of the unregularised singular-ρ cone is not removed. Thus c_A^{sm} equals the fixed-state coefficient c_A plus the bosonic contacts $-\frac{1}{4}(N_s+2)$, and the two readings differ by the total derivative $\Box\delta\ln\rho$, that is by the boundary flux of $\nabla\ln\rho$ through Σ.

Outcome for $n_D = 2$, $N_s = 6$. Here $c_T = 0$ and $N_{\mathrm{tot}} = 0$. The fixed-state replica gives $c_A = 2$. If the connection sector carries a contact of the Kabat value -1 in these units (the value that takes the vector with its ghosts from $\frac{1}{3}$ to $-\frac{2}{3}$), which

for a Maxwell field is the edge-mode term of Donnelly and Wall [18], then $c_A = 1$. The smooth reading gives $c_A^{\rm sm} = 0 = c_T$, an equality of vanishing coefficients, which holds for every sector-uniform Q because $N_{\rm tot} = 0$ and is not a derivation. For fields minimally coupled in the variables $(G, \rho, \partial\Theta)$, the contacts of these types have weight $+1$ per Dirac field (already contained in c_A), -1 for the vector and 0 for the scalars and ghosts, so within this class they give at most the value 1 above; the torsion-reading bosonic contacts $-\frac{1}{4}(N_s + 2)$ are the smooth reading and do not belong to this class. The values 2 and 0 correspond to a replica in which $\ln\rho$ carries no flux through Σ and one in which it carries the full deficit; intermediate fluxes are not computed, and which replica the matrix state selects depends on the state and the tip regularisation, which are not derived here. These statements are for the Poisson-limit operators, the $\Lambda_{\rm UV}^2$ term, one loop, $\alpha = 1$ and the fixed-state and first-order smooth replicas stated; they do not derive the replica, the state, the tip domain or the gluing, do not treat edge modes of the symplectic gauge sector of a^μ, of the ghosts or of fermions (not examined), the $\Lambda_{\rm UV}^4$ term for $N_{\rm tot} \neq 0$, or the next order of either side, and give no value or sign of $G_{\rm eff}$. Problems B, C, D, F and G remain open.

Replica prescriptions against published contact-term treatments (conditional; not closure). Keep the setting, the fixed-state replica and the first-order smooth replica of the preceding paragraph, and write c_A and $c_A^{\rm sm}$ for their area coefficients and $c_R = N_{\rm tot}/6$ for the coefficient of $\rho R[g_\rho]$; $c_A = 2$ and $c_A^{\rm sm} = 0$ at $n_D = 2$, $N_s = 6$. Put $V_\mu = \partial_\mu \ln\rho$.

Scalar and ghost sectors. For a scalar with a coupling $\xi R\phi^2$, Herzog and Nishioka add the codimension-two term $\delta I_E = -2\pi(1-n)\xi \int \phi^2$ (Eq. (2.19) of [19]), which cancels the delta of ξR on the cone and gives an entropy independent of ξ. With the heat-trace bookkeeping of the contact-term paragraph above, this term shifts the entropy by $+2\pi\xi A\langle\phi^2\rangle$, equal and opposite to the contact $-2\pi\xi A\langle\phi^2\rangle$, which is the expectation value of Wald's formula for $\xi R\phi^2$ [18], Eq. (4); Herzog and Nishioka do not discuss Wald's formula, and their term is related to it only through this bookkeeping. The fixed-state replica, whose scalar and ghost operators carry no delta on the cone, therefore has the contact-free coefficient. The smooth replica, in which E_s contains $-\frac{1}{2}\Box_G \ln\rho$ with $Q = \frac{1}{4}$, has the form of the delta-potential entropy of Eqs. (143)–(144) of [10] with ξ replaced by Q; that review does not accept this entropy as an entanglement entropy, and Donnelly and Wall assign the corresponding term to the generalised entropy as a Wald contact. Neither value is asserted here to be the entanglement entropy of the matrix state: the identifications are by the form of the coefficient, the cited analysis is for a scalar on a flat background, and the operators here have no independent ξ. On the singular-ρ cone $\ln\rho = (n-1)\ln|z|$, the operator $\mathcal{O}_s$ is unitarily equivalent to $-\Box + E_s$, with s-wave Bessel index $\nu_0 = |n-1|/(2n) > 0$. The regular branch vanishes at Σ, so the delta multiplies a function that vanishes

there, and the constant $-\nu_0/2$ gives the entropy coefficient $\frac{1}{6}-\frac{1}{4}$ per component for $n<1$ and $\frac{1}{6}+\frac{1}{4}$ for $n>1$, non-analytic at $n=1$; the less singular branch exchanges the two. The smooth value $-\frac{1}{12}$ is thus the $n<1$ value of the regular branch, and removing the delta alone does not carry this cone to the contact-free value $\frac{1}{6}$. The terms of order $((n-1)/2)^2$ in the indices ν_k with $k\neq 0$ depend logarithmically on the tip scale and drop out at first order in $n-1$.

Connection sector. For the Maxwell field in Feynman gauge, Herzog and Nishioka remove the Ricci delta of the vector operator with $\delta I=-\pi(1-n)\int\delta^{ij}A_iA_j$ (Eq. (7.2) of [19]), and note that this term is not gauge invariant. Kabat's contact is that delta: it is present in the black-hole entropy and absent from the bulk-mode entanglement entropy [17], and Donnelly and Wall identify it with edge modes [18]. Herzog and Nishioka state that their boundary term contributes oppositely to the contact term (Eq. (5.20) of [19]). The vector contact -1 that gives $c_A=1$ is therefore the term that the Herzog–Nishioka term removes. Its use presupposes a $U(1)$ structure and a Ricci coupling that the operator $\mathcal{O}_v$ does not have in the variables $(G,\rho,\partial\Theta)$, where the gauge group is symplectic and $Q=0$; it remains an input and is not a prescription of this class.

Teleparallel form. The identity for the Weitzenböck connection of the paragraph on the torsion invariant gives $E_s=-\frac{1}{2}\nabla_\mu V^\mu+\frac{1}{4}V_\mu V^\mu=\frac{1}{4}(R_G+\mathcal{T})+\frac{1}{4}V_\mu V^\mu$ with $\mathcal{T}=\frac{1}{4}T_{\lambda\mu\nu}T^{\lambda\mu\nu}+\frac{1}{2}T_{\lambda\mu\nu}T^{\mu\lambda\nu}-V_\mu V^\mu$, and the scalar density is $\frac{1}{6}R_G-E_s=-\frac{1}{12}R_G-\frac{1}{4}\mathcal{T}-\frac{1}{4}V_\mu V^\mu$. With $\mathcal{T}=-R_G-2\nabla_\mu V^\mu$ the coefficient of R_G is $\frac{1}{6}$, which is the value in the quadratic form on the two modes with $\delta\rho=0$, where $V_\mu V^\mu$ is of higher order. For scalar content alone $c_T=c_A$, as stated above, and the two replicas differ by the integrated boundary term $R_G+\mathcal{T}=-2\nabla_\mu V^\mu$, the flux of $\nabla\ln\rho$ through Σ.

Dynamical metric. In the replica construction of Lewkowycz and Maldacena the entropy is $-\partial_n$ of the action of a regularised cone minus that of the cone, localised at the tip; the difference between the action of the true n-solution and that of the regularised cone is a first-order variation about a solution and vanishes by the equations of motion (Eqs. (3.1)–(3.4) of [20]). That step requires the background to solve the equations of motion of the full action, of which the stationarity condition of the tensor block is the part not derived here. The regularised cone has the far field of the $n=1$ solution with the angle identified modulo $2\pi n$, so that every field other than the metric is regular there. With Θ the only field, $R_G^{(1)}=-2\Box_{\bar{G}}\,\delta\ln\rho$ excludes such a configuration at first order, as stated above. The only first-order regularisation in this class is the smooth family, whose far field has $\rho\propto r^{n-1}$ and so differs from that of the cone; the construction of [20], and the vanishing of its first bracket, are therefore not reproduced for it. For the smooth replica the area coefficient is $c_A^{\mathrm{sm}}=-c_R/2$. This is the coefficient of $\sqrt{G}\,R_G$ at fixed coframe torsion, so the smooth replica satisfies $S=A/4G_{\mathrm{ind}}$ in those variables for every content, whereas $c_A^{\mathrm{sm}}-c_T=(14-N_s-4n_D)/4$,

which vanishes only for $N_s + 4n_D = 14$. The generalised-entropy structure therefore does not imply $G_A = G_T$ in this class. At $n_D = 2$, $N_s = 6$, the conditions $c_R = 0$ ($N_s = 4n_D - 2$) and $c_T = 0$ ($N_s = 22 - 8n_D$) are independent and meet only there; $c_A^{\mathrm{sm}} = 0 = c_T$ is then the absence of the Λ_{UV}^2 term in the induced action and carries no information on G, while $c_A = 2$ is the contact-free entanglement coefficient, in the relation, in form, $S_{\mathrm{BH}} = S_{\mathrm{ent}} - Q$ of Eq. (322) of [10], where Q is here the bosonic contact $(N_s + 2)/4$ of the smooth reading and S_{BH} is built from the induced action.

Intermediate flux. For first-order smooth perturbations of constant $\bar{\Theta}$ the integral of $R_G^{(1)}$ over the transverse plane is -2 times the flux of $\nabla \ln \rho$, so no family of such perturbations joins the two replicas. If the first-order local densities of the pulled-back cone and of the smooth family are superposed with weight f on the smooth family, then $c_A(f) = c_A - f(N_s+2)/4$, which is $2-2f$ at $n_D = 2$, $N_s = 6$; the superposition is an assumption. The indices $\nu_k = \sqrt{k^2 + ((n-1)/2)^2}/n = |k + \mathrm{i}(n-1)/2|/n$ are those of an imaginary twist, so the periodic quadratic dependence of the area coefficient on a real Abelian holonomy is not used.

These statements are for the Poisson-limit operators, the Λ_{UV}^2 term, one loop, $\alpha = 1$, and the two replicas stated; they do not derive the replica or the state selected by the matrix model, the stationarity condition, the covariance of the induced action in Θ, the vector contact, or the tip domain, they treat neither edge modes of the symplectic gauge sector nor fermion edge modes, and they give no value or sign of G_{eff}. Problems B, C, D, F and G remain open.

Next order at $N_s = 6$, $n_D = 2$: exact $\mathfrak{J} = 0$ backgrounds and the surviving quartic (conditional; not closure). Keep the setting of the paragraphs on the gauge parameter and on fermionic content: Poisson limit, Euclidean section $\eta = \delta$, $\alpha = 1$, one loop, $N_s = 6$ scalars, the connection with its ghosts and $n_D = 2$ Dirac fields, so that $N_{\mathrm{tot}} = 0$ and the weighted coefficient κ_X of $\rho^2 X$ vanishes. The statements below are local, on regions where $\mathrm{Pf}\,\omega \neq 0$. Write $(b_\sigma)_{\mu\nu} = G_{\sigma\tau}\Theta^{\rho\tau}\partial_\rho\Theta_{\mu\nu}$, the $\mathfrak{so}(4)$-valued one-form with $\omega_\sigma = -\frac{1}{2}\partial_\sigma \ln\rho\,\mathbb{I} + b_\sigma$ in the vector sector and, through the spinor lift $\beta_\sigma = \frac{1}{4}\gamma^{\mu\nu}(b_\sigma)_{\mu\nu}$, in the Dirac sector. For every Poisson bivector $\mathfrak{J}^\nu = \Box_G x^\nu - G^{\mu\nu}\partial_\mu \ln\rho$ (exact check at random two-jets), so that $\mathfrak{J} = 0$ says that the coordinates are harmonic for G up to the gradient of $\ln\rho$.

Backgrounds with $\mathfrak{J} = 0$. Up to a constant factor, $\rho\,\Theta^{\mu\nu}\Theta^{\mu\nu}$ is the Poisson-limit density of the first term of Equation (3.4), and $\rho\,\Theta^{\mu\nu}\Theta^{\mu\nu} \geq 4$ pointwise, with equality if and only if Θ is self-dual or anti-self-dual. For such Θ, $\Theta\Theta^T$ is a multiple of the identity, so $G^{\mu\nu} = \rho^{-1}\delta^{\mu\nu}$, $g_\rho = \delta$ and $\mathfrak{J} = 0$ by the identity above. A self-dual Θ is the inverse of a closed self-dual two-form $\omega = b_i\eta^i$, with $\eta^i_{ab} = \epsilon_{iab}$, $\eta^i_{a4} = \delta_{ia}$; closure is $\mathrm{div}\,b = 0$, $\partial_4 b = -\,\mathrm{curl}\,b$, and then $\rho = |b|^2$, $G = \rho\,\delta$ is conformally flat with $R_G \neq 0$ in general, while $\rho R[g_\rho] = 0$. Since $\partial_\rho\Theta^{\mu\nu}$ is then self-dual in $(\mu\nu)$, b_σ takes values in one $\mathfrak{su}(2)$ factor of $\mathfrak{so}(4)$; the same holds for anti-self-dual Θ. An example invariant

under the rotations of the (x^1, x^2) plane, which fix the plane $\{x^1 = x^2 = 0\}$ pointwise, is $b_1 + \mathrm{i}b_2 = -(x^1 + \mathrm{i}x^2)(x^3 - \mathrm{i}x^4)$, $b_3 = 1 + (x^3)^2 - (x^4)^2$ (exact check of closure, invariance, $\mathfrak{J} = 0$ and $R_G \neq 0$ at sample points; the vector (b_1, b_2, b_3) vanishes on the locus $x^1 = x^2 = 0$, $(x^4)^2 = 1 + (x^3)^2$, where $\mathrm{Pf}\,\omega = 0$ and the example is not defined).

Outside this class, let $\Lambda^i = \eta^i_{\mu\nu} x^\mu \mathrm{d}x^\nu$, so that $\Lambda^3 = x^1\mathrm{d}x^2 - x^2\mathrm{d}x^1 + x^3\mathrm{d}x^4 - x^4\mathrm{d}x^3$, and take $\omega = \mathrm{d}(h(r)\Lambda^3)$, invariant under the rotations of both coordinate planes. Then $\mathfrak{J}$ has only a radial component, proportional to $2h^2rh'' + 10h^2h' + 4hrh'^2 + r^2h'^3$, and $u = rh'/h$ obeys $r\,\mathrm{d}u/\mathrm{d}r = -\frac{1}{2}u(u+2)(u+4)$. The fixed points $u = 0$ and $u = -4$ are the self-dual and anti-self-dual solutions, $u = -2$ is degenerate ($\mathrm{Pf}\,\omega \propto h^2(u+2)$), and the other solutions are $h = c\,(1 \pm \sqrt{1 - C/r^4})$, $C \in \mathbb{R}$ (checked at generic Cartesian points), locally, for $r^4 > C$ when $C > 0$; at $r^4 = C$ the metric G degenerates. On them $G = \beta^2[s^{-2}(\mathrm{d}r^2 + e_3^2) + e_1^2 + e_2^2]$ with $\beta = 2h$, $s^2 = 1 - C/r^4$, $e_i = \Lambda^i/r$, and $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma} = \frac{3}{2}u^2(u+4)^2/(hr)^4$, which vanishes only at the two end points.

Coefficients. Let $\mathcal{S}_4 = \mathrm{tr}_V[(b_\sigma b^\sigma)^2] - n_D\,\mathrm{tr}_D[(\beta_\sigma\beta^\sigma)^2]$, with $b_\sigma b^\sigma = G^{\sigma\tau}b_\sigma b_\tau$ and traces over the four vector and the four Dirac components. In the variables (G, ρ, b) the endomorphism of every sector is $E = e\,\mathbb{I} + X + b_\sigma b^\sigma$, with the same scalar $e = -\frac{1}{2}\Box_G \ln\rho + \frac{1}{4}|\partial \ln\rho|^2_G$, X linear in b and traceless, and the bundle curvature is that of b. The weighted sums of $\mathrm{tr}\,\mathbb{I}$ and of the Dynkin index of b are $N_{\mathrm{tot}} = 0$ and $\kappa_X = 0$, and $\mathfrak{so}(4)$ has no cubic invariant. Hence, for every Poisson bivector, the weighted sum of the coefficients a_1 vanishes pointwise and the weighted sum of the coefficients a_2 is $\frac{1}{2}\mathcal{S}_4$, up to total derivatives, which vanish by the same sums: no R^2, $R_{\mu\nu}R^{\mu\nu}$, Riemann-squared, Weyl-squared or Euler term, no RE term and no term of Einstein–Hilbert type survives (exact rational arithmetic at random two-jets, also for $\eta = \mathrm{diag}(-1, 1, 1, 1)$; on the self-dual and anti-self-dual backgrounds the weighted sum of a_2 vanishes while the single-sector values do not, and it does so for the backgrounds tested only at $(N_s, n_D) = (6, 2)$, with the vector weight 1 and the ghost weight -2 fixed in that scan). With $l^k_\sigma = \frac{1}{4}\eta^k_{\mu\nu}(b_\sigma)_{\mu\nu}$ and $r^k_\sigma = \frac{1}{4}\bar\eta^k_{\mu\nu}(b_\sigma)_{\mu\nu}$ the self-dual and anti-self-dual parts ($\bar\eta^k_{ab} = \epsilon_{kab}$, $\bar\eta^k_{a4} = -\delta_{ka}$), and $\mathsf{A} = G^{\sigma\tau}l_\sigma l_\tau^T$, $\mathsf{B} = G^{\sigma\tau}r_\sigma r_\tau^T$, $\mathsf{C} = G^{\sigma\tau}l_\sigma r_\tau^T$,

$$\mathcal{S}_4 = 8\,\mathrm{tr}\,\mathsf{A}\;\mathrm{tr}\,\mathsf{B} + 16\,\mathrm{tr}(\mathsf{C}\mathsf{C}^T) \;\geq 0,$$

with equality at a point if and only if b_σ takes values in one $\mathfrak{su}(2)$ factor there. With one common proper-time limit for all sectors, the coefficient of $\ln\Lambda_{\mathrm{UV}}$ in the induced action at this content is therefore $\propto \int \rho^2\mathcal{S}_4$, quartic in $\partial\Theta$; finite terms of the same order are not controlled. It vanishes on the self-dual and anti-self-dual backgrounds above. On the $\mathrm{d}(h\Lambda^3)$ family b_σ was found to be anti-self-dual-valued in every sampled case with $\mathfrak{J} = 0$, so $\mathcal{S}_4 = 0$ there, as the weighted a_2 computed there confirms (not proved), and it has both parts in the sampled cases with $\mathfrak{J} \neq 0$. The condition $\mathfrak{J} = 0$ does not by itself force $\mathcal{S}_4 = 0$: about a constant $\bar\Theta$, with $\Theta = (\psi^*\bar\omega)^{-1}$ and $\psi = x + \epsilon a$, the condition $\mathfrak{J} = O(\epsilon^2)$ is the source-free Maxwell equation $D^\mu(D^\mu a^\nu - D^\nu a^\mu) = 0$,

$D^\mu = \bar\Theta^{\mu\rho}\partial_\rho$; for quadratic a its solutions give constant $b_\sigma = O(\epsilon)$ with both parts and $\mathcal{S}_4 > 0$ (random solutions for one $\bar\Theta$; in one example the order ϵ^2 of $\mathfrak{J} = 0$ is solved by a cubic correction of ψ). The convergence of the series, a global background, the domains of the intertwining lemma above and the tip domain are not examined. On backgrounds with $\mathfrak{J} = 0$ globally the logarithmic coefficient does not depend on α, because the longitudinal sector is a scalar tower with proper time rescaled and the coefficient of s^2 is unchanged.

Where an Einstein–Hilbert term can enter. In the massless class every local term of order Λ_{UV}^0 has four derivatives. If a sector carries a constant mass, $\mathcal{O}_i \to \mathcal{O}_i + m_i^2$ (a formal shift of the operator: no gauge-invariant mass term and no ghost or longitudinal consistency is asserted), then $E_i \to E_i + m_i^2$, and the coefficient a_2 of that sector changes by $-m_i^2 a_1^{(i)} + \frac{1}{2}\operatorname{tr}\mathbb{I}\, m_i^4$ (exact check). The curvature part of the change of the weighted sum is $-(\sum_i w_i n_i m_i^2/6)\rho R[g_\rho] - (\sum_i w_i \kappa_i m_i^2)\rho^2 X$, where n_i is the number of components of the sector and κ_i its weight of $\rho^2 X$ (1 for the connection, $\frac{1}{2}$ per Dirac field, 0 otherwise). It vanishes for one common mass, because $N_{\mathrm{tot}} = 0$ and $\kappa_X = 0$, and is nonzero only if bosonic and fermionic masses differ. The inert transverse matrices of this class carry no such scale. In the construction of [21] the Einstein–Hilbert term is generated by a mixed quartic term with a sum over the Kaluza–Klein modes of fuzzy extra dimensions, and the Newton constant is set by their scale; the class above has no analogue of it.

These statements are for the Poisson-limit operators, the Euclidean section, one loop, $\alpha = 1$, the content $N_s = 6$, $n_D = 2$ with the Yukawa couplings dropped, and the backgrounds named. They compute no logarithmic coefficient of the cone (area) side, so that no comparison of G_T with G_A at logarithmic order follows; they do not derive the content, the state, the tip domain or Assumption 8.4, do not treat Moyal corrections or fuzzy extra dimensions, and give no value or sign of G_{eff}. Problems B, C, D, F and G remain open.

Cone logarithmic coefficient at $N_s = 6$, $n_D = 2$ on the fixed-state replica (conditional; not closure). Keep the setting of the preceding paragraph and the fixed-state replica with antiperiodic Dirac gluing and the less singular extension. Let Σ be the fixed plane of a rotation under which Θ is invariant, so that Σ is totally geodesic for G. With unit normals n_i to Σ, write $R_{ii} = R_{\mu\nu} n_i^\mu n_i^\nu$ and $R_{ijij} = R_{\mu\nu\rho\sigma} n_i^\mu n_j^\nu n_i^\rho n_j^\sigma$ (sums over i, j), and R_Σ for the intrinsic scalar curvature of Σ, so that $R_\Sigma = R - 2R_{ii} + R_{ijij}$ for totally geodesic Σ. The pulled-back background is the same on every sheet, so $\operatorname{Tr}\rho^n$ is the sum over the twist phases $e^{2\pi \mathrm{i} k/n}$ of the single-sheet problem with a flat flavour vortex of flux $a_k = k/n$ at Σ (integer k for the bosonic sectors, $k = -\frac{n-1}{2}, \dots, \frac{n-1}{2}$ for the Dirac fields). For a Laplace-type sector $-\nabla^2 + E$ the s^0 surface term is linear in the dimension-two data. For a component with a periodic phase it is the scalar term of Equation (70) of [10],

$c_B(n)\,\mathrm{tr}(\frac{R}{6}-E)+d_B(n)(R_{ii}-2R_{ijij})$ with $c_B=\frac{\pi}{3}\frac{1-n^2}{n}$ and $d_B=-\frac{\pi}{180}\frac{(1-n^2)(1+n^2)}{n^3}$, and it has no term in the bundle curvature, because every periodic twist acts as a common phase and the sum of the Bernoulli polynomials $B_3(k/n)$ in the linear coefficient vanishes. For a Dirac field the common phase is accompanied by the spin term $\gamma^{12}\Omega_{12}$ of $(\gamma^\mu D_\mu)^2$, with $\Omega_{12}=[D_1,D_2]$, whose part supported on Σ is attractive in the component where it is positive; the less singular extension then gives the E-linear weight $-\frac{1}{2}c_B$ per component and a term $-\frac{1}{2}\sum_k a_k^2\,\mathrm{tr}(\gamma^{12}\Omega_{12})$ with $\sum_k a_k^2=(n^2-1)/(12n)$ and Ω_{12} the bundle curvature in the normal plane (from the exact Landau–Aharonov–Bohm spectrum; the terms odd in a cancel in the symmetric sum over k). The remaining geometric part of the Dirac term is fixed at first order in $n-1$ by the value $a=11/360$ of the Dirac field (Equation (275) of [10], after Duff; not re-derived here), which gives $-\frac{11}{90}$ for a sphere. With one proper-time limit, the coefficient of $\ln\Lambda_{\mathrm{UV}}$ in the replica entropy is $\int_\Sigma \sigma_{\log}$, where, with $X'=-\,\mathrm{tr}(b_\sigma b^\sigma)$ in the vector sector, $\tau_\Omega=\mathrm{tr}(\gamma^{12}\Omega^f_{12})$ the normal-plane curvature of the Dirac connection, with the same $\Omega_{12}=[D_1,D_2]$, e the scalar endomorphism of the preceding paragraph and $W=W_{ijij}$ the Weyl projection,

$$24\pi\,\sigma_{\log}=\tfrac{4}{3}R_G+2R_{ii}-2R_{ijij}+2W-12\,e+\tfrac{3}{2}X'-\tau_\Omega.$$

On the self-dual and anti-self-dual backgrounds, $W=0$ and $e=R_G/6$, so that $24\pi\sigma_{\log}=\frac{3}{2}X'-\tau_\Omega$. At three sampled points of Σ for the rotation-invariant self-dual example above this equals $2R_\Sigma$, twice the intrinsic curvature of Σ (exact rational arithmetic; not proved), so that $\sigma_{\log}=R_\Sigma/12\pi$. The cone logarithmic coefficient is then an Euler density of Σ with nonvanishing local value. The coefficient $\propto\int\rho^2\mathcal{S}_4$ of the induced action vanishes on these backgrounds, and so does the first variation of $\int\rho^2\mathcal{S}_4$ along the smooth replica, because $\mathcal{S}_4\geq 0$ vanishes there. The two sides therefore differ at logarithmic order, as coefficients, for the fixed-state replica on these backgrounds; this is a statement about the fixed-state replica only and not about G_{eff}. The first-order smooth replica of the preceding paragraphs gives zero on both sides, and which replica the matrix state selects is not derived here. Contact and edge-mode terms of the connection and Dirac sectors, which are not included, shift the coefficient of R_Σ and could cancel the difference.

These statements are for the Poisson-limit operators, the Euclidean section, one loop, $\alpha=1$, $N_s=6$, $n_D=2$, the totally geodesic Σ, the fixed-state replica and one common proper-time limit. They use Equation (70) of [10] and the Dirac coefficient $a=11/360$ as inputs, do not treat finite terms, edge modes, other extensions, higher orders in $n-1$, or the non-self-dual $\mathfrak{J}=0$ backgrounds beyond the general formula, and give no value or sign of G_{eff}. Problem B remains open and PROGRAMME-CONDITIONAL.

Contact terms in the logarithmic coefficient at $N_s = 6$, $n_D = 2$ (conditional; not closure). Keep the setting of the preceding paragraph. To first order in $1 - n$ the surface part of the s^0 heat coefficient of a Laplace-type sector $-(\nabla^2 + E)$ is the cross term of the bulk coefficient a_2 with the conical deltas $R \to 4\pi(1-n)\delta_\Sigma$, $R_{\mu\nu} \to 2\pi(1-n)n_{\mu\nu}\delta_\Sigma$, $R_{\mu\nu\rho\sigma} \to 2\pi(1-n)(n_{\mu\rho}n_{\nu\sigma} - n_{\mu\sigma}n_{\nu\rho})\delta_\Sigma$ (Equations (55)–(59) of [10]) and with any delta E_δ, Ω_δ of the endomorphism and of the bundle curvature. Without such deltas it reproduces Equation (70) of [10]. For the Hodge operator of a Maxwell field in Feynman gauge with its ghosts, with E_δ the Ricci delta and $\Omega_\delta = 2\pi(1-n)\mathcal{J}\delta_\Sigma$ the Riemann delta ($\mathcal{J}$ the rotation of the normal plane), it gives $S_{\log} = \pi^{-1}[-\frac{31}{360}R_\Sigma + \frac{1}{20}W]$, that is the ratios 62 and 12 of the conformal scalar, and $-\frac{62}{90}$ on a great S^2 of S^4; the contact is $-(R_\Sigma - R_{ii})/24\pi$, of which $-R_\Sigma/24\pi$ has, by the form of the term, the density of the edge-mode entropy of [18]. The Dirac weights $-\frac{1}{2}$ on $\mathrm{tr}(\frac{R}{6} - E)$, 1 on the $R_{ii} - 2R_{ijij}$ term and the spin term of the preceding paragraph are consistent with the Dirac $a = 11/360$ used above and with c, and with the Weyl-covariant form; given a, c and Weyl covariance they fix the weights $(-\frac{1}{2}, 1, \frac{1}{2})$, of which $-\frac{1}{2}$ and $\frac{1}{2}$ are the values of the preceding paragraph.

For the rotation-invariant self-dual backgrounds with $\Sigma = \{x^1 = x^2 = 0\}$ and $g = |\nabla_\Sigma \ln \rho|^2_{g_\Sigma}$, $g_\Sigma = \rho\,\delta$, one finds at 18 points of random polynomial data (exact rational arithmetic; not proved) $X' = \frac{3}{2}g$, $\tau_\Omega = R_\Sigma + \frac{3}{4}g$, so that $24\pi\sigma_{\log} = \frac{3}{2}g - R_\Sigma$. The coefficient $\sigma_{\log}$ is a multiple of R_Σ pointwise only where $g = 2R_\Sigma$, as on the example above; for generic members the Euler-density reading does not hold. With all sectors glued periodically and no deltas the coefficient vanishes for every Poisson bivector at $(6, 2)$, because $N_{\mathrm{tot}} = 0$ and the weighted a_1 vanishes; the value $\sigma_{\log}$ is the difference between the antiperiodic and the periodic Dirac gluing.

Contacts. On the family $\Theta^{12} = 1/f(x^1, x^2)$, $\Theta^{34} = 1$, with $\ln f = (n-1)\ln|z|$, the scalar, connection and Dirac data contain $\frac{1}{2}\Box_0 \ln f$ with the same coefficient in every component, so that $E_\delta = -\pi(1-n)\delta_\Sigma$ uniformly (the contact with $Q = \frac{1}{4}$ above), and the connection bundle curvature contains $\Omega_{12,\delta} = 2\pi(1-n)\mathcal{J}\delta_\Sigma$ (symbolic computation). Let C_b be the logarithmic cross term of E_δ on the six scalars, the four components of a^μ and the ghosts, and $C_\Omega = \mathrm{tr}(\mathcal{J}\Omega^v_{12})/24\pi$ that of the connection bundle. Then, in units $1/24\pi$, $24\pi\sigma_{\log} + 24\pi C_b = -\tau_\Omega$ and $\mathrm{tr}_V(\mathcal{J}\Omega^v_{12}) = \tau_\Omega$ (the same $\mathfrak{so}(4)$ connection in both representations: the vector is $(\frac{1}{2}, \frac{1}{2})$ and the Dirac field $(\frac{1}{2}, 0) \oplus (0, \frac{1}{2})$ under $\mathfrak{su}(2) \oplus \mathfrak{su}(2)$, with $\mathcal{J}$ acting as $\frac{1}{2}\gamma^{12}$ on spinors), hence $\sigma_{\log} + C_b + C_\Omega = 0$. This was checked at random closed two-forms (not self-dual, $\mathfrak{J} \neq 0$) and on the $\mathfrak{J} = 0$ families above. On the example, $C_b + C_\Omega = -R_\Sigma/12\pi$, which cancels $\sigma_{\log}$; C_b alone gives $-3R_\Sigma/16\pi$, C_Ω alone $5R_\Sigma/48\pi$. A term of the Herzog–Nishioka type [19], applied here by the form of the term to remove E_δ, leaves $\sigma_{\log} + C_\Omega = 3g/32\pi \neq 0$. Replacing the two-dimensional operator of [18] by $-w^{-1}\,\mathrm{div}(w\nabla)$, obtained here and not a result of that paper, gives an edge density $-(4\pi)^{-1}(R_\Sigma/6 - \frac{1}{2}\Delta \ln w - \frac{1}{4}|\nabla \ln w|^2)$; for $w = \rho^{-1}$ (from the Poisson-limit action)

$\sigma_{\log}$ plus it does not vanish. The identification of this edge sector with the symplectic $U(1)$ and the choice $w = \rho^{-1}$ are not proved, and axion-type couplings are not treated. The Dirac fields keep the fixed-state antiperiodic gluing, which equals the periodic Dirac value plus the contacts above, and carry no edge mode of their own.

These statements are for the Poisson-limit operators, one loop, $\alpha = 1$, $N_s = 6$, $n_D = 2$, a totally geodesic Σ and cross terms at first order in $1 - n$ (the $n < 1$ branch of the rotated connection index of the paragraph on fermionic content). They assume that the delta content computed on the family $\Theta^{12} = 1/f$, $\Theta^{34} = 1$ (E_δ uniform on every bosonic component and $\Omega_\delta = 2\pi(1-n)\mathcal{J}\delta_\Sigma$) is the complete delta content for the backgrounds used; with the normal-block distribution of the Ricci-type delta, used in the Maxwell check, the sum does not vanish for $\mathfrak{J} \neq 0$. The Dirac fields keep the fixed-state antiperiodic gluing, which equals the periodic value plus the contacts above, so the replica is an imported prescription of the type $S_{\mathrm{BH}} = S_{\mathrm{ent}} - Q$ and not the trace of the n-th power of one state; the fixed-state replica is not a limit of smooth Poisson structures at first order. The non-delta singular first-order parts of the smooth family, which are of the same order and include the non-analytic dependence on $|1-n|$ of the rotated connection index, the $n > 1$ branch and the state selected by the matrix model are not treated. The completed coefficient vanishes, as does the coefficient of the induced action of the preceding paragraph on these backgrounds (a Wald-type reading in the variables $(G, \text{torsion})$); this is an equality of vanishing coefficients, which follows from $N_{\mathrm{tot}} = 0$ and the vanishing weighted Dynkin index and gives no value or sign of G_{eff}. Problem B stays open and PROGRAMME-CONDITIONAL.

Content dependence of the logarithmic comparison off $(N_s, n_D) = (6, 2)$ (conditional; not closure). Keep the setting of the two preceding paragraphs with N_s scalars and n_D Dirac fields in place of $(6, 2)$, the vector with weight 1, the ghosts with weight -2, and write $N_{\mathrm{tot}} = N_s + 2 - 4n_D$, $\kappa_X = 1 - \frac{1}{2}n_D$ and $\bar{a}_1 = \sum_i w_i \operatorname{tr}(\frac{R}{6} - E_i) = N_{\mathrm{tot}}(\frac{R}{6} - e) + \kappa_X X'$ for the weighted a_1 density, whose integral is the Λ^2_{UV} density of the induced action. Off $(6, 2)$ the logarithmic coefficients are not constrained to vanish, so this is a comparison with nonzero values, for contents that are not the maximally supersymmetric one. In units $1/24\pi$ on a totally geodesic Σ, the replica with all sectors periodic and no deltas, the fixed-state replica of the preceding paragraphs and the replica completed by the uniform E_δ and by $\Omega_\delta = 2\pi(1-n)\mathcal{J}\delta_\Sigma$ of the vector bundle give

$$24\pi\sigma_{\mathrm{per}} = \bar{a}_1 - \tfrac{N_{\mathrm{tot}}}{30}(R_\Sigma - 3W), \qquad 24\pi\sigma_{\mathrm{cmp}} = -\tfrac{1}{2}\bar{a}_1 + \kappa_X\tau_\Omega - \tfrac{N_{\mathrm{tot}}}{30}(R_\Sigma - 3W), \qquad 24\pi\sigma_{\mathrm{fs}} = 24\pi\sigma_{\mathrm{cm}}$$

with $\bar{a}_1^{\mathrm{bos}}$ the bosonic part of $\bar{a}_1$ and τ_Ω as in the preceding paragraphs. For one Dirac field the antiperiodic value of the preceding paragraphs equals the periodic value plus the cross terms of a uniform E_δ and of Ω_δ with $\mathcal{J} \to \frac{1}{2}\gamma^{12}$, an exact

identity in R, R_{ii}, R_{ijij}, e, X' and τ_Ω given the Dirac weights $(-\frac{1}{2}, 1, \frac{1}{2})$; the contacts therefore equal $-\frac{3}{2}\bar{a}_1 + \kappa_X \tau_\Omega$ for every content, proportional to the weighted a_1 density and to the weighted Dynkin index, the data of the Λ^2_{UV} order. They vanish at $(6, 2)$ for this reason, and the three coefficients differ off $(6, 2)$ only by terms in $\bar{a}_1$ and τ_Ω. These relations were checked in exact arithmetic at twenty sampled points (the rotation-invariant self-dual family, the non-self-dual $\mathfrak{J} = 0$ family, $\mathfrak{J} \neq 0$ radial profiles and generic closed ω); at the $\mathfrak{J} \neq 0$ and generic points they are pointwise algebraic identities, with Σ not necessarily totally geodesic, and the rotation-invariant closed form rests on eleven points. On the rotation-invariant self-dual family $24\pi\sigma_{\mathrm{cmp}} = (\kappa_X - N_{\mathrm{tot}}/30)R_\Sigma = \frac{1}{30}(28 - N_s - 11n_D)R_\Sigma$ for every content, without the term in g of the preceding paragraph, and $24\pi\sigma_{\mathrm{fs}} = 24\pi\sigma_{\mathrm{cmp}} - R_\Sigma + \frac{3}{2}g$. On the self-dual and anti-self-dual points and on the sampled $\mathfrak{J} = 0$ points the weighted a_2 density equals $N_{\mathrm{tot}}s + \kappa_X(v - 4s)$, with s and v the single-sector scalar and vector values, since the Dirac value minus $4s$ is half of the vector value minus $4s$; it vanishes on all these backgrounds simultaneously only for $N_{\mathrm{tot}} = \kappa_X = 0$, that is $(N_s, n_D) = (6, 2)$.

Weyl coefficient on plane waves. A comparison that is not affected by the choice between the variables $(G, \rho, \partial\Theta)$ and $(G, \text{torsion})$ for the endomorphism contacts is the coefficient of C^2, because R, e, X' and $\bar{a}_1$ vanish to first order on the modes with $\delta\rho = 0$ about a constant self-dual or anti-self-dual $\bar{\Theta}$. It is read from the quadratic form of the weighted a_2 on those modes, against the coefficient of the Weyl projection W in the first-order area-side density, where a metric functional $\kappa_C C^2 + \kappa_E \mathcal{E}_4 + \kappa_R R^2$ is assigned the entropy density $12\kappa_C W + 12\kappa_E R_\Sigma + 12\kappa_R R$ by the cross-term rule. The rule does not fix the entropy of a part of the quadratic form carried by non-metric invariants of Θ; that part is the gap below. At second order the single-sector values satisfy $v - 4s = \frac{1}{12}\operatorname{tr}\Omega^2$ and $d - 4s = \frac{1}{2}(v - 4s)$, the vector connection b contributes $X^{(1)} = 0$, and on the modes considered $\frac{1}{12}\operatorname{tr}\Omega^2 = -\frac{1}{3}C^2$ while $\tau_\Omega^{(1)} = -R_{ijij} = -2W + R_\Sigma$ (first-order formulas checked against the jet code at $\epsilon = 10^{-5}$; six modes for each of the two chiralities). The coefficient of C^2 is then $\kappa_C^T = N_{\mathrm{tot}}/120 - \kappa_X/3$, and the coefficients of W of the three replicas minus $12\kappa_C^T$ are

$$\text{periodic: } 4 - 2n_D, \qquad \text{fixed-state: } 4 - n_D, \qquad \text{completed: } 2 - n_D,$$

independent of N_s. The first two equal the differences $c_A - c_T$ at order Λ^2_{UV} of the paragraphs on the quadratic form and on fermionic content; the completed gap has no Λ^2_{UV} counterpart of the same form. The periodic and completed gaps vanish on the line $n_D = 2$ for every N_s, where both coefficients equal N_{tot} times the conformal-scalar value and are nonzero for $N_s \neq 6$; the fixed-state gap vanishes on $n_D = 4$. The completed gap is $2\kappa_X$. On these modes $\operatorname{tr}\Omega^2 = -4C^2$ is twice its Levi-Civita value $-2C^2$, whereas $\operatorname{tr}(\mathcal{J}\Omega_{12}) = -R_{ijij}$ has its Levi-Civita value, so that the cross term with Ω_δ accounts for half of the contribution of the Ω^2 term to the coefficient of W.

The bundle curvature Ω^v is not determined by the Riemann tensor: at fixed first jet its second jet has 33 independent components against the 20 of the Riemann tensor (45 random second jets, exact rank), and the Riemann second jet is a linear image of that of Ω^v. The matching on $n_D = 2$ is the universal N_{tot} counting and does not test the index-weighted sector.

The contents $(N_s, n_D) \neq (6, 2)$ are parameters of the computation and are not proposed field contents of the matrix model. The replicas are the hybrid fixed-state replica, whose Dirac fields have the antiperiodic gluing and are not the trace of the n-th power of one state, the periodic replica, whose Dirac fields have the unrotated periodic gluing (the graded trace), and the replica completed by the uniform E_δ and by Ω_δ on every component, with one common proper-time limit for all sectors.

These statements are for the Poisson-limit operators, one loop, $\alpha = 1$, the contents stated with the Yukawa couplings dropped, a totally geodesic Σ, cross terms at first order in $1 - n$ on the $n < 1$ branch, the imported contact prescription and the delta content of the preceding paragraph, with the hypotheses stated there, and, for the plane-wave comparison, a constant self-dual or anti-self-dual $\bar{\Theta}$, the modes with $\delta\rho = 0$ and the Weyl coefficient only; the coefficient of the Euler density is invisible to the quadratic form. They use the Dirac weights and Equation (70) of [10] as before, treat neither finite terms, nor the mode with $\delta\rho \neq 0$, nor the Hodge distribution of the Ricci-type delta, and give no value or sign of G_{eff}. Problem B stays open and PROGRAMME-CONDITIONAL.

Entropy assigned to the bundle curvature, and the delta content of the Poisson cone (conditional; not closure). Keep the setting of the preceding paragraph: the weighted a_2 density, the plane-wave modes with $\delta\rho = 0$ about a constant self-dual or anti-self-dual $\bar{\Theta}$, the completed replica, a totally geodesic Σ, cross terms at first order in $1 - n$. The term $\frac{\kappa_X}{12} \operatorname{tr} \Omega^2$ is quadratic in the bundle curvature, so its cross term with a delta $\Omega_\delta = 2\pi(1-n)\, H\, \delta_\Sigma \, \mathrm{d}x^1 \wedge \mathrm{d}x^2$ is $\kappa_X \operatorname{tr}(H\Omega_{12})$, with $H \in \mathfrak{so}(4)$ the holonomy of the bundle connection around Σ. If the connection is treated as an auxiliary independent field (a bookkeeping variable, not a metric) whose internal rotations are tied to the normal plane of Σ through a regular frame (Lorentz-Lie Noether charge), the charge density is $n^{ab}\, \partial L / \partial \Omega^{ab}$, which is this cross term with $H = \mathcal{J}_{12}$ and no other generator: it reproduces $\kappa_X \tau_\Omega$ and not more. If the internal group is not tied to the frame, the charge vanishes at the bifurcation surface for a regular connection with vanishing momentum map, and the term is absent. The first reading assigns the coefficient $-2\kappa_X$ of W to $\kappa_X \operatorname{tr} \Omega^2$ and the second assigns 0, against $-4\kappa_X$ on the $\delta\rho = 0$ modes.

For the diagonal family $\Theta^{12} = r^{-\epsilon c_a}$, $\Theta^{34} = r^{-\epsilon c_b}$, $\epsilon = n - 1$, one finds exactly $H = c_a \mathcal{J}_{12} + c_b \mathcal{J}_{34}$, with the Riemann deltas $R_{1212} = -2\pi\epsilon c_a$, $R_{a3a3} = R_{a4a4} = -\pi\epsilon c_b$ (symbolic computation). The cone angle of the normal plane fixes $c_a = 1$; $c_b \neq 0$ makes

the metric induced on Σ singular at order ϵ (read off the delta structure; not separately checked). With all deltas inserted in the cross term of the weighted a_2 (an extension of the cross-term rule beyond the cone, not derived), the gap of the preceding paragraph becomes $-N_{\rm tot}c_b/15 + 2\kappa_X(1+c_b)$ about a self-dual $\bar\Theta$ and the same expression about an anti-self-dual one with $\Theta^{34} = -r^{-\epsilon c_b}$, for which $H = c_a\mathcal{J}_{12} - c_b\mathcal{J}_{34}$, and equals $2\kappa_X$ at $c_b = 0$. The κ_X part vanishes only for a holonomy of chirality opposite to that of $\bar\Theta$, while the $N_{\rm tot}$ part then becomes nonzero, so no content-independent profile removes the gap in this family.

On the 17 sampled points with totally geodesic Σ (exact rational arithmetic; not proved), $\tau_\Omega = R_\Sigma - 2W + (a_1^v - \frac{1}{2}X')$, with a_1^v the a_1 density of the vector sector. The coefficient of R_Σ in the basis (W, R_Σ, R, e, X') of the fit, in the completed coefficient, is therefore $\kappa_X - N_{\rm tot}/30$, the value of the Levi-Civita cross-term rule, and the remainder is a combination of the Λ^2-order densities a_1 and X' in that basis; no statement is made about their topological character beyond it. These statements are for the diagonal family, the modes and replica stated, and the hypotheses of the preceding paragraphs; they do not derive the holonomy selected by the matrix model, the replica, or a value or sign of $G_{\rm eff}$. Problem B stays open and PROGRAMME-CONDITIONAL.

Edge and contact terms of the vector and connection sector at the Weyl coefficient (conditional; not closure). Keep the setting of the preceding paragraph: the weighted a_2 density, the modes with $\delta\rho = 0$ about a constant self-dual or anti-self-dual $\bar\Theta$, the completed replica, a totally geodesic Σ, cross terms at first order in $1-n$, and the completed gap $2\kappa_X$ in the coefficient of W, read in the basis (W, R_Σ, R, e, X'). On the twelve modes W and R_Σ are independent (rank 2), with $R_{ii} = W - R_\Sigma$ and $R_{ijij} = 2W - R_\Sigma$.

Endomorphism contacts. To first order in the amplitude about a constant $\bar\Theta$ the endomorphisms E_V of the vector and E_f of the Dirac operator equal $e\,\mathbb{I}$: the antisymmetric part is a divergence of $\Theta^{\rho\sigma}\partial_\rho\Theta^{\mu\nu}$, which vanishes at first order about constant $\bar\Theta$, and the symmetric traceless part is quadratic (exact arithmetic, cubic jets of both chiralities, two amplitudes). Since e vanishes for $\delta\rho = 0$, every cross term of a_2 with a delta of an endomorphism, uniform or confined to the normal block, vanishes at this order for every content: a contact of Kabat type has no coefficient of W in this presentation. For the Hodge operator of the metric G ($E = -\mathrm{Ric}$, $\Omega = \mathrm{Riem}$), the same cross-term rule on the same modes gives $6/5$ and $-31/15$ for the Maxwell field with its ghosts, equal to its $12\kappa_C$ in the coefficient of W, of which the endomorphism contact carries 3 (the value is $-9/5$ without it). The matrix-derived vector operator is not the Hodge operator: on these modes $\mathrm{Riem}^2 = 2C^2$ and $\mathrm{Ric}^2 = C^2/2$, and $v - 4s$ is $-C^2/3$ against $C^2/12$. That mechanism is therefore not available for it.

Edge modes. In the classes treated in [18], products of a two-dimensional base with a bifurcate horizon and a compact fibre, and $S^1 \times H^3$, the edge sector of the Maxwell

field is a scalar of wrong-sign statistics on the entangling surface, and its logarithmic density is the two-dimensional trace anomaly of the Laplacian of Σ, $-R_\Sigma/24\pi$; that paper does not treat the dependence on the Weyl tensor, and leaves that on the extrinsic geometry open. In the basis above $-R_\Sigma$ has coefficient 0 of W. A weight w built from $\mathrm{Pf}\,\Theta$ and $\mathrm{tr}\,\Theta^2$ in $-w^{-1}\,\mathrm{div}(w\nabla)$ adds nothing at first order, because the Hessians of both vanish on the modes. A potential $E_\Sigma = \alpha R_{ii} + \beta R_{ijij}$ would change the density to $-(R_\Sigma + 6E_\Sigma)$ and would need $\alpha + 2\beta = \kappa_X/3$ for the coefficient $-2\kappa_X$ of W. This depends on n_D through the Dirac weight, whereas the vector carries one edge scalar, if it carries the edge sector of a Maxwell field, and the Dirac fields none. No edge term independent of n_D equals $2\kappa_X$ for all n_D.

Holonomy contact. For the diagonal profile with $|\Theta^{34}| = r^{-\epsilon c_b}$ the holonomy is $H = c_a\mathcal{J}_{12} + \chi c_b\mathcal{J}_{34}$, with $\chi = +1$ about a self-dual and $\chi = -1$ about an anti-self-dual $\bar\Theta$ (symbolic computation), and the coefficient of W in $\mathrm{tr}(\mathcal{J}_{34}\Omega_{12})$ is 2χ. The gap, in the extended cross-term rule of the preceding paragraph, is therefore the same for both chiralities, $N_{\mathrm{tot}}(3c_a - 2c_b - 3)/30 + 2\kappa_X(2 - c_a + c_b)$. With $s = c_a - c_b$, the contraction of the first-order Weyl tensor with the Weyl tensor of the deltas is $2sW$, and the Euler combination $\mathrm{Riem}\cdot\mathrm{Riem}' - 4\,\mathrm{Ric}\cdot\mathrm{Ric}'$ is $2c_aR_\Sigma - 2c_bW$, which contains W for $c_b \neq 0$. If that W part is dropped as topological, the gap is $(s-1)N_{\mathrm{tot}}/10 - 2(s-2)\kappa_X$: the metric part requires $s = 1$ and the connection part $s = 2$. The profile with $\delta\rho = 0$, $c_b = -c_a$, has $s = 2$ and removes the κ_X part, leaving $N_{\mathrm{tot}}/15$ ($N_{\mathrm{tot}}/10$ after dropping the Euler part); its induced metric on Σ is singular at first order and the scalar of its Riemann delta, $2c_a + 4c_b$, has the sign opposite to that of the cone.

No edge, endomorphism-contact or holonomy term of this family therefore supplies $-2\kappa_X W$ for all contents. The required term is proportional to the index-weighted sum that multiplies $\mathrm{tr}\,\Omega^2$, the same for every fibre representation; of the candidates examined only a holonomy generator beyond $\mathcal{J}_{12}$ attached to Σ has this property, and none of the profiles of the diagonal family provides it without a singular metric on Σ. The gap is also relative to the comparator $12\kappa_C^T$ assigned to $\mathrm{tr}\,\Omega^2$ by the metric-universal rule; whether the entropy of this non-metric term is that value or the Noether value $-2\kappa_X$ of the preceding paragraph, which equals the replica, is not decided here.

These statements are for the Poisson-limit operators, one loop, $\alpha = 1$, a totally geodesic Σ, cross terms at first order in $1 - n$ on the $n < 1$ branch, the modes and replica of the preceding paragraphs, the cross-term rule extended to $c_b \neq 0$ (not derived), and an edge sector of the abelian type of the cited paper. They do not treat non-diagonal profiles, the non-delta singular first-order parts, the exact heat trace of the cone with holonomy, the symplectic gauge group, or the mode with $\delta\rho \neq 0$, and give no value or sign of G_{eff}. Problem B stays open and PROGRAMME-CONDITIONAL.

H.3 Problem C: conformal mode and Cassini

Ledger label. Status: OPEN. Priority: CRITICAL PATH. Completion: PROGRAMME-CONDITIONAL. A scalar-tensor result is a reduced-scope outcome only when its operator, coupling and phenomenological test are supplied; it does not by itself meet the matrix-derived criterion.

The radial singlet has the tree-level mass

$$m_\sigma^2 = \lambda v^2.$$

This does not by itself determine the mass or matter coupling of a scalar mode of the composite metric. In particular, the stated Repair I prescription is intended to build a vierbein from the Lorentz-valued polar factor Λ, evaluated at $\sigma = 0$, rather than from the radial field.

Obstruction to the stated Repair I frame. Suppose $\Lambda(x) \in \mathrm{SO}(3,1)$ and expand about the locked configuration,

$$\Lambda = \mathbb{I} + \Omega + O(\Omega^2), \qquad \Omega_{\mu\nu} = -\Omega_{\nu\mu}.$$

Then $\partial_\mu \Omega_{\alpha\beta}$ transforms in $V \otimes \mathfrak{so}(3,1)$, where $V = (\frac{1}{2}, \frac{1}{2})$ is the Lorentz vector representation. After complexification,

$$(V \otimes \mathfrak{so}(3,1))_{\mathbb{C}} = \left(\tfrac{3}{2}, \tfrac{1}{2}\right) \oplus \left(\tfrac{1}{2}, \tfrac{1}{2}\right) \oplus \left(\tfrac{1}{2}, \tfrac{3}{2}\right) \oplus \left(\tfrac{1}{2}, \tfrac{1}{2}\right).$$

A vierbein fluctuation transforms in $V \otimes V$, whose complexified decomposition is

$$(V \otimes V)_{\mathbb{C}} = (0,0) \oplus (1,0) \oplus (0,1) \oplus (1,1).$$

These representations have no common irreducible summand. Hence there is no nonzero Lorentz-equivariant linear map from $\partial_\mu \Omega_{\alpha\beta}$ to a vierbein fluctuation using only the locked Lorentz-invariant tensors. Consistently, the displayed formula

$$e^A{}_\mu = \left\langle \partial_\mu \Lambda^A{}_I \right\rangle$$

leaves I uncontracted and does not, as written, define a vierbein.

A simple proposed contraction with a fixed nonzero internal vector n^I does not repair non-degeneracy. If

$$e^A{}_\mu = \partial_\mu \Lambda^A{}_I \, n^I,$$

then Lorentz orthogonality gives

$$(\Lambda n)_A e^A{}_\mu = (\Lambda n)_A \partial_\mu (\Lambda n)^A = \frac{1}{2}\partial_\mu((\Lambda n)^2) = 0.$$

Every column of $e^A{}_\mu$ therefore lies in the three-dimensional subspace orthogonal to Λn, so $\mathrm{rank}(e) \leq 3$. This contraction cannot produce a non-degenerate four-dimensional frame.

Status. Under the stated Repair I ingredients, the linearized composite vierbein required for the later metric-scalar analysis is not constructed. This is an obstruction to that frame prescription, not a proof that a completed theory has or lacks a physical scalar pole. Repair II, an independent coframe, or additional symmetry-breaking background tensors would change the construction and require a separate analysis. The radial mass $m_\sigma^2 = \lambda v^2$ remains established, but it is not a metric-scalar mass. The scalar spectrum and matter coupling relevant to Cassini are therefore still undetermined. The Repair II pure-trace relation $\psi = 8\sigma/v$ with $m_\sigma^2 = \lambda v^2$ is a potential-only statement; the induced Einstein–Hilbert term adds to the kinetic coefficient of σ, so it does not fix a physical pole mass, mixing or Cassini coupling.

In the Repair II pure-trace truncation, adding the induced Einstein–Hilbert term to the microscopic kinetic term gives, in the healthy-sign normalization, $Z = 2N - 3/(4\pi G_{\mathrm{eff}} v^2)$ (Equation 17.4), conditional on the sign of (3.6) as corrected in Chapter 3 and on the declared curvature convention. For $Z > 0$, the truncated pole has $m^2 = V''/Z$; for $Z < 0$ it is ghost-like and, for $\lambda > 0$, tachyonic in this truncation, which therefore cannot support Assumption 7.5 there. This signals an instability of the chosen background and truncation, not a statement about the full scalar spectrum. At $Z = 0$ the quadratic kinetic term degenerates and establishes no canonically normalized finite-mass pole. This result also assumes $G_{\mathrm{eff}} > 0$, which is not derived (Problem F), and does not determine mixing, the non-minimal coupling, the matter coupling, or a Cassini bound. If matter is assumed to couple only through $g = \Omega^2\eta$, the unmixed truncated scalar exchanges, for $Z > 0$, a Yukawa potential of strength $\beta = M_{\mathrm{P}}^2/(Nv^2 - 3M_{\mathrm{P}}^2)$ relative to the nominal Einstein-sector Newtonian potential (Chapter 17); this is conditional on the assumptions listed there and is not a Cassini bound.

Off-shell trace–tensor mixing from the induced Einstein–Hilbert term (conditional; not closure). Assume Repair II, which is assumed rather than derived, with $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ and $h_{\mu\nu} = T_{\mu\nu} + \frac{1}{4}\eta_{\mu\nu}\psi$. Also assume the induced Lorentzian term is $S_{\mathrm{EH}} = (16\pi G_{\mathrm{eff}})^{-1} \int \sqrt{-g}\, R$, with $G_{\mathrm{eff}} > 0$. For a general symmetric $h_{\mu\nu}$, define $\mathcal{E}_{\mu\nu}[h] := -2G^{(1)}_{\mu\nu}[h]$; on traceless tensors this agrees with $\mathcal{E}_{\mu\nu}$ of Equation 7.4. In these

conventions, up to boundary terms,

$$S_{\rm EH}^{(2)} = \frac{1}{16\pi G_{\rm eff}} \int \mathrm{d}^4x \, \frac{1}{4} h^{\mu\nu} \mathcal{E}_{\mu\nu}[h].$$

For $s_{\mu\nu} = \frac{1}{4}\eta_{\mu\nu}\psi$,

$$\mathcal{E}_{\mu\nu}[s] = \frac{1}{2}(\partial_\mu\partial_\nu\psi - \eta_{\mu\nu}\Box\psi).$$

Expanding the quadratic action in this field basis gives

$$S_{\rm EH}^{(2)} = \frac{1}{16\pi G_{\rm eff}} \int \mathrm{d}^4x \left[\frac{1}{4} T^{\mu\nu}\mathcal{E}_{\mu\nu}[T] + \frac{1}{4} T^{\mu\nu}\partial_\mu\partial_\nu\psi + \frac{3}{32}(\partial\psi)^2\right].$$

Thus, using $\psi = 8\sigma/v$ at linear order, the cross term is

$$\frac{1}{16\pi G_{\rm eff}} \int \mathrm{d}^4x \, \frac{2}{v} T^{\mu\nu}\partial_\mu\partial_\nu\sigma.$$

The tensor-only term here is normalized relative to the assumed $S_{\rm EH}$; Equation 7.13 states a structure and is not used to set this coefficient. The cross term is basis-dependent: it is the usual conformal-mode mixing in the gauge-invariant quadratic Einstein–Hilbert action and can be rearranged by a field redefinition. It does not by itself establish an observable effect or scalar-exchange strength. Although it vanishes if transversality is imposed, transversality cannot be assumed when forming the coupled off-shell Hessian. The displayed microscopic action on the fixed scaffold does not specify the full coupled, gauge-fixed tensor operator or the remaining scalar couplings. Problems B, C, D, F and G remain open.

H.4 Problem D: local Lorentz

Ledger label. Status: OPEN. Priority: CRITICAL PATH. Completion: PROGRAMME-CONDITIONAL. Reduced-scope result: D2, imposed local Lorentz symmetry, supports only a scope-limited Presentation-B model; D remains OPEN and the matrix-derived criterion is unmet.

The counting of generators under the locked vacuum is given in Chapter 10. The condensate term supplies a quadratic Higgs-type contribution in the broken-direction combination Ξ; it does not establish a physical pole mass without kinetic terms and canonical normalization. The diagonal connection's dynamics and torsion remain open. Torsion-free behavior follows only under the conditional Palatini–Holst hypotheses of Proposition 10.5; NAMCG has not been shown to induce those hypotheses.

For a Laplace-type operator, $a_1 = \mathrm{tr}(R(g)/6 - E)$; the explicit universal curvature term is the Levi-Civita scalar $R(g)$. Because E is unspecified, no conclusion about additional B-dependence of a_1 follows, and a Palatini $R(B)$ term is not established.

If B enters the bundle connection of the actual operator, a_2 may contain a $\operatorname{tr} F(B)^2$ term; whether it survives in the weighted trace depends on the unspecified operator representation and weights. Deriving the B-dependence of E and the bundle connection remains open.

Missing step. Derive $\mathrm{SO}(3,1)$ from the matrix algebra alone, or state that Presentation A is not ontologically prior (the present text does the latter), and derive the effective dynamics of the independent diagonal connection or retain it as unresolved field content. Generator counting alone does not settle its torsion or propagating modes. The locked frame and its non-degeneracy must also be derived rather than assumed (Assumptions 3.5 and 4.4). Under D2, only explicitly scope-limited Presentation-B claims are available; specifying the full operator content does not close D or meet the matrix-derived criterion. The permitted and unestablished claims, including the consequent truncation for Problem F, are listed in Section 17.3.

Symmetries supplied by the stated Presentation A (matrix-derived criterion of Problem D; not closure). The matrix-coordinate action (3.4) is invariant under unitary conjugation $\hat{X}^\mu \mapsto U\hat{X}^\mu U^\dagger$, provided the matter variables transform so that their term is invariant: each matrix commutator transforms by conjugation and the trace is cyclic. This is a matrix gauge transformation $\delta\hat{X}^\mu = i[\epsilon, \hat{X}^\mu]$; the stated action and infrared dictionary do not identify it with a transformation of an independent tangent frame.

As an action-level statement, the commutator term is covariant under constant Lorentz transformations of the spacetime index μ, because its indices are contracted with $\eta_{\mu\nu}$. The cubic term is covariant if $C_{\mu\nu\lambda}$ transforms as a tensor; for fixed $C_{\mu\nu\lambda}$, it is a symmetry only when C is invariant. The matter term must obey the corresponding transformation law. Likewise, the chosen background $\Theta^{\mu\nu}$ in Equation (3.2) is invariant only under its stabilizer if held fixed; transforming Θ along with the fields expresses covariance, not a symmetry of that fixed background. None of these constant transformations establishes a spacetime-dependent local Lorentz gauge symmetry.

On the $C_{\mu\nu\lambda} = 0$, central constant-Θ branch of Remark 3.8, evaluate the matrix-gauge variation on the background $\bar{X}$. By the Jacobi identity,

$$\delta_\epsilon[\bar{X}^\mu, \bar{X}^\nu] = i[\epsilon, [\bar{X}^\mu, \bar{X}^\nu]] = 0,$$

since $[\bar{X}^\mu, \bar{X}^\nu] = i\Theta^{\mu\nu}\mathbb{I}$. In the irreducible Moyal-symbol realization, the corresponding linearized fluctuation transforms as $\delta a^\mu = \Theta^{\mu\rho}\partial_\rho\lambda$, leaving $\mathcal{F}^{\mu\nu}_\Theta$ invariant as shown in that remark. This local matrix-gauge statement is distinct from the global constant Lorentz covariance described in the preceding paragraph, which requires transforming

Θ as a tensor and is a symmetry of a fixed Θ only for transformations that stabilize it. On this branch, matrix conjugation alone supplies neither a transformation of an independent tangent-frame index nor six independent local Lorentz parameters. This is a limitation of the stated matrix-gauge action on the central background, not a no-go for an enlarged matrix model with a derived frame map and additional fields.

More specifically, on the central, nondegenerate constant-Θ Moyal branch, define $D'_A f := \Lambda_A{}^B \star D_B f$, where $D_B = i[\bar{X}_B, \cdot\,]$ are the inner derivations, whose principal symbols $\Theta_B{}^\mu \partial_\mu$ are linearly independent. The Leibniz defect is

$$D'_A(f \star g) - (D'_A f) \star g - f \star (D'_A g) = [\Lambda_A{}^B, f]_\star \star D_B g.$$

Independence of the principal symbols implies that this defect vanishes for all f, g exactly when every $\Lambda_A{}^B$ is central. In the irreducible nondegenerate constant-Θ representation, the centre consists of constants, i.e. multiples of the identity. This conclusion is limited to that branch and this left-multiplication action; it does not rule out a connection-twisted or module-valued frame, or an enlarged matrix model. A matrix-derived frame module with a connection that compensates this defect is a conjectured route, not a result established here. Problems B, C, D, F and G remain open.

For real, nondegenerate constant Θ, consider $\Lambda \in \mathrm{SO}(3,1)$ that preserves the fixed tensor, $\Lambda\Theta\Lambda^T = \Theta$. In a Lorentz frame with $\Theta^{01} = E$, $\Theta^{23} = B$, $EB \neq 0$, and all other independent components zero, its identity-component stabilizer is $\mathrm{SO}^+(1,1) \times \mathrm{SO}(2)$: a boost in the 01-plane and a rotation in the 23-plane. Equivalently, in the self-dual representation its stabilizer is $\mathrm{SO}(2,\mathbb{C})$, of real dimension two. This is a constant-background stabilizer, not a local Lorentz gauge symmetry.

The corresponding linear automorphisms of the Heisenberg algebra are implementable by inner automorphisms of its irreducible Weyl representation. Infinitesimally, write $\Lambda = \mathbb{I} + A$; preservation of Θ requires

$$A\Theta + \Theta A^T = 0.$$

Since Θ is invertible,

$$H_{\mu\nu} := (\Theta^{-1})_{\mu\rho} A^\rho{}_\nu$$

is symmetric. On the common invariant domain of the Weyl generators, the symmetrically ordered quadratic generator

$$G = \frac{1}{4} H_{\mu\nu} \{\bar{X}^\mu, \bar{X}^\nu\}$$

satisfies

$$\delta \bar{X}^\mu = i[G, \bar{X}^\mu] = A^\mu{}_\nu \bar{X}^\nu.$$

The finite implementation is the metaplectic unitary, unique up to phase by the Stone–von Neumann theorem. This implements only the constant stabilizer: it supplies neither six arbitrary local Lorentz parameters nor an independent tangent-frame index or spin connection. Invariance of the full action also requires the remaining fields and couplings to transform compatibly. This conditional representation-theoretic result does not establish Problem D1. Problems B, C, D, F and G remain open.

A conditional classical-symbol construction is possible if the background Poisson tensor varies. Let $\Theta^{\mu\nu}(x)$ be a real, nondegenerate Poisson tensor, so its bracket

$$\{f, g\}_\Theta := \Theta^{\mu\nu}\partial_\mu f\, \partial_\nu g$$

obeys the Jacobi identity. For local coordinate functions x_A, with the frame index A identified with the coordinate label, define the Hamiltonian vector fields $V_A := X_{x_A}$, where $X_f(g) := \{f, g\}_\Theta$, and assume they form an invertible frame. The Poisson Jacobi identity gives

$$[V_A, V_B] = X_{\Theta_{AB}}, \qquad \Theta_{AB} := \{x_A, x_B\}_\Theta.$$

Define $C^C{}_{AB}$ by $[V_A, V_B] = C^C{}_{AB}V_C$. A provisional candidate inverse metric is

$$g^{\mu\nu}_{\text{cand}} := \eta^{AB}V_A{}^\mu V_B{}^\nu.$$

It is Lorentzian by construction when the frame is invertible, but is not thereby identified with the emergent metric of Chapter 4.

Under a local frame change $V'_A = \Lambda_A{}^B(x)V_B$, with $\Lambda(x) \in \mathrm{SO}(3,1)$, the bracket, expanded in the original frame V_F, is

$$[V'_A, V'_B] = \Big(\Lambda_A{}^D\Lambda_B{}^E C^F{}_{DE} + \Lambda_A{}^D V_D(\Lambda_B{}^F) - \Lambda_B{}^D V_D(\Lambda_A{}^F)\Big)\, V_F.$$

Accordingly, the structure functions of the new frame are the displayed coefficients contracted with $(\Lambda^{-1})_F{}^G$, since $[V'_A, V'_B] = C'^G{}_{AB}V'_G$. For this provisional frame and metric, the dual coframe ϑ^A determines the composite Levi-Civita spin connection through

$$\mathrm{d}\vartheta^A + \omega^A{}_B \wedge \vartheta^B = 0, \qquad \omega_{AB} = -\omega_{BA}.$$

It transforms inhomogeneously under the local frame change.

For constant Θ, the fields $V_A{}^\mu$ are constant, so $C^C{}_{AB} = 0$ and this composite connection and its curvature vanish in that frame. A varying Θ is therefore needed for this construction to be nontrivial, although variation alone does not guarantee nonzero curvature. This is a classical Poisson-symbol construction: it does not address the exact Moyal Leibniz defect for noncentral frame coefficients, derive local frame rotations from the matrix action, or supply independent connection dynamics. Problems B, C,

D, F and G remain open.

For the hand-chosen Poisson background of Remark 3.9,

$$\Theta = a\,\partial_0 \wedge \partial_1 + b(y)\,\partial_2 \wedge \partial_3, \qquad b(y) = b_0 e^{\kappa y^2},$$

the Hamiltonian frame $V_A = X_{x_A}$, with $X_f(g) = \{f, g\}_\Theta$, is

$$(V_0, V_1, V_2, V_3) = (a\partial_1, -a\partial_0, b\partial_3, -b\partial_y).$$

Its only nonzero frame bracket is

$$[V_2, V_3] = b'(y)V_2 = 2\kappa y\, b(y)\, V_2.$$

The associated structure function is $C^2{}_{23} = b'(y)$; the composite connection contains the logarithmic derivative

$$\omega^2{}_3 = \frac{b'(y)}{b(y)}\,\mathrm{d}x^3 = 2\kappa y\,\mathrm{d}x^3.$$

With $\eta_{AB} = \mathrm{diag}(-,+,+,+)$, the provisional inverse metric constructed from this frame is

$$g^{\mu\nu}_{\mathrm{cand}} = \eta^{AB} V_A{}^\mu V_B{}^\nu = \mathrm{diag}(a^2, -a^2, b(y)^2, b(y)^2) = K^{\mu\nu}_\Theta.$$

It therefore reproduces the tensor used in Remark 3.9; with that remark's normalization $K^{\mu\nu}_\Theta = \ell_*^4 \tilde{g}^{\mu\nu}$, $g_{\mathrm{cand}} = \ell_*^{-4}\tilde{g}$.

A dual coframe is

$$\vartheta^0 = \frac{\mathrm{d}x^1}{a}, \qquad \vartheta^1 = -\frac{\mathrm{d}x^0}{a}, \qquad \vartheta^2 = \frac{\mathrm{d}x^3}{b(y)}, \qquad \vartheta^3 = -\frac{\mathrm{d}y}{b(y)}.$$

Its torsion-free connection has $\omega^2{}_3 = 2\kappa y\,\mathrm{d}x^3$, and $\mathcal{R}^2{}_3 = 2\kappa\,\mathrm{d}y \wedge \mathrm{d}x^3$. In the curvature convention of Remark 3.9, this gives $R[g_{\mathrm{cand}}] = 4\kappa b(y)^2$. This identifies the candidate Poisson-frame metric with the already chosen toy geometry, up to the stated constant normalization; it is not an independent derivation of that geometry.

The calculation is at classical Poisson-symbol order on a hand-chosen background. It derives neither that background from the matrix action nor local Lorentz gauge symmetry, a noncommutative frame module, or independent connection dynamics. Problems B, C, D, F and G remain open.

For the Poisson block above, restrict to the $C_{\mu\nu\lambda} = 0$ branch of (3.4), retain only the quartic matrix term, and omit matter. The matrix equation is

$$[\hat{X}_\nu, [\hat{X}^\nu, \hat{X}^\mu]] = 0,$$

with indices contracted using η. At leading Poisson-symbol order, using $[\hat{f},\hat{g}] \sim i\{f,g\}_\Theta$, this requires

$$\eta_{\nu\lambda}\Theta^{\lambda\rho}\partial_\rho\Theta^{\nu\mu} = 0.$$

For $\Theta^{01} = a$, $\Theta^{23} = b(y)$, with $y = x^2$, the $\mu = 3$ component vanishes identically, while the $\mu = 2$ component is $b(y)b'(y)$. Nondegeneracy requires $b \neq 0$, so this leading-order source-free equation forces $b' = 0$ on each connected patch. Hence the hand-chosen curved profile $b = b_0 e^{\kappa y^2}$, with $\kappa \neq 0$, is not a source-free leading-order solution of this ansatz. Supporting that profile would require a specified balancing contribution, such as a cubic term, matter source, or corrections from a specified deformation quantization. This calculation is at classical Poisson-symbol order only; it does not define a beyond-leading-order product for varying Θ, make a claim about the full matrix model, or exclude other backgrounds. It is not a derivation or closure of Problem D1. Problems B, C, D, F and G remain open.

In particular, Presentation A as specified contains no coframe $e^A{}_\mu$, independent tangent index A, or spin connection $\omega_\mu{}^{AB}$. The infrared dictionary (3.14) is an assumption mapping a matrix commutator to a covariant derivative with A_μ; it supplies neither a local Lorentz transformation law for a derived frame nor the transformation and dynamics of $\omega_\mu{}^{AB}$. The boost obstruction recorded in Problem F concerns the displayed Presentation-B potential, not the Presentation-A matrix action. These facts delimit what the stated action and dictionary establish; they do not rule out an enlarged matrix construction. Consistent with the preceding paragraph, the matrix-derived criterion of Problem D remains open: closure requires an explicit matrix-to-frame map and a derivation of the local transformation law, invariant action, and connection content from that map. Problems B, C, D, F and G remain open.

Linear metric content of a real locked block (conditional; not closure). Restrict to a real 4×4 locked block, $e = \mathbb{I} + M$ with M real, and $g = e^T\eta e$ as in (4.9). To first order $\delta g = M^T\eta + \eta M$. For real symmetric $M = H$: $\delta g_{00} = -2H_{00}$, $\delta g_{ij} = 2H_{ij}$, $\delta g_{0i} = 0$. For real antisymmetric $M = K$: $\delta g_{0i} = -2K_{0i}$, $\delta g_{ij} = 0$. On this real slice the three directions K_{0i} change the metric, and the quadratic potential of the preceding Hessian paragraph is independent of K. (For complex Φ the Hermitian and anti-Hermitian parts of that paragraph are not the real symmetric and antisymmetric parts used here; the correspondence holds only on the real slice.)

Internal transformations that preserve the positive-definite contraction in (3.6) are unitary; those that also map real frames to real frames are orthogonal; those that further preserve η_{AB} commute with η. Their Lie algebra is $\mathfrak{so}(3)$: no boost generator survives (at the group level the intersection is $\mathrm{O}(1) \times \mathrm{O}(3)$). This concerns the internal index only. For the tangent index, $\Phi \mapsto \Phi\Lambda$ sends $\mathrm{Tr}\,\Phi^\dagger\Phi$ to $\mathrm{Tr}\,\Lambda\Lambda^\dagger\Phi^\dagger\Phi$. This equals $\mathrm{Tr}\,\Phi^\dagger\Phi$ for arbitrary Φ only if $\Lambda\Lambda^\dagger = \mathbb{I}$; a boost does not satisfy this and

generically changes the contraction, although it may leave it unchanged for particular configurations. Hence the contractions displayed in (3.6) and (3.7) are not invariant under local boosts as symmetries of the action. Imposing local Lorentz invariance in D2 (Assumption 10.1) therefore requires a completed action, with the tangent connection and an invariant contraction, and not merely a relabelling of (3.6). Whether such an action exists, and what its spectrum is, is not examined here. Problem D remains open.

Ghost-free versus boost-invariant contraction (conditional; not closure). On the real locked 4×4 block write $\Phi = \mathbb{I} + M$. Consider two candidate quadratic kinetic contractions, both stated for the fixed scaffold and neither derived from (3.6). (i) The kinetic term in (3.6), with its positive-definite matrix-space contraction, gives every component the same relative kinetic sign, so there are no relative-sign differences; however, the displayed contractions in (3.6) and (3.7) are not invariant under boosts of either index, as noted above. (ii) A contraction with η on both indices, invariant under constant independent $\mathrm{SO}(3,1) \times \mathrm{SO}(3,1)$ transformations, gives component $M^I{}_A$ the relative kinetic sign $\eta_{II}\eta_{AA}$. Fix the overall sign so that $M^0{}_0$ and $M^i{}_j$ have the healthy sign; then the six components $M^0{}_i, M^i{}_0$ have the opposite sign. This invariance is global: for local transformations covariant derivatives and the corresponding connections are needed, and the candidate expression does not specify them. If the six broken generators of Proposition 10.2 are gauged and eat the six η-antisymmetric components, three of the opposite-sign components (the $0i$ part of the antisymmetric block) are absorbed; three opposite-sign η-symmetric $0i$ components remain. Whether these are removed by a constraint, a further gauge condition, or the diffeomorphism structure of Chapter 10 is not established. This is a sign count for the quadratic form about $\Phi = \mathbb{I}$ only; it is not a spectrum or unitarity result and is not identified with the tensor sector of Chapter 5. Boost invariance and absence of opposite-sign components are therefore not both available from the contractions considered here. Problem D remains open.

A boost-invariant completed action: quadratic test (conditional; not closure). This paragraph tests one candidate structural change, *not* contained in (3.6): replace the positive-definite contractions by $\eta_{IJ}\eta^{AB}$ contractions, so that the kinetic term is $-g^{(0)\mu\nu}\eta_{IJ}\eta^{AB}\,\mathcal{D}_\mu\Phi^I{}_A\,\mathcal{D}_\nu\Phi^J{}_B$ with the covariant derivative (10.3), and the potential is $\frac{\lambda}{4}\mathrm{Tr}(Q - v^2\mathbb{I})^2$ with $Q = \eta^{-1}\Phi^T\eta\Phi$; both are invariant under constant $\mathrm{SO}(3,1) \times \mathrm{SO}(3,1)$, with local invariance given by the connections. It is introduced only to test whether the opposite-sign modes found above can be removed. It would change the definition of the model, and nothing here shows that it is derivable from Presentation A.

About $\Phi = v(\mathbb{I} + M)$ on the real locked 4×4 block, with $\tilde{M} = \eta^{-1}M^T\eta$, the

quadratic potential is $\lambda v^4\,\mathrm{K}(M_s)$ for the η-self-adjoint part $M_s = \frac{1}{2}(M + \tilde{M})$, where $\mathrm{K}(X) = \eta_{IJ}\eta^{AB}X^I{}_A X^J{}_B$. The ten self-adjoint components carry the potential and the six anti-self-adjoint components are flat, matching Proposition 10.2. Because K is indefinite, this quadratic form is negative along the three $0i$ self-adjoint directions for $\lambda > 0$: $\Phi = v\mathbb{I}$ is a stationary point but a saddle of this candidate potential, not a stable vacuum, so what follows is a quadratic diagnostic about a stationary background and not a spectrum about a stable vacuum. The kinetic form has signature $(7, 3)$ on the self-adjoint block and $(3, 3)$ on the anti-self-adjoint block; the two blocks do not mix. At the background $A_\mu = 0$, the relative combination $\omega_\mu - iA_\mu$ in the convention of (10.3) (equivalently $\omega_\mu - A_\mu$ if A_μ is redefined as a real Lie-algebra-valued connection) enters algebraically and removes the anti-self-adjoint components from the kinetic term, while the diagonal combination does not enter at quadratic order. What remains is the self-adjoint block, with seven components of one kinetic sign and three (the $0i$ components) of the opposite sign; since kinetic and potential forms coincide on this block, the equation-of-motion mass squared is λv^2 in the normalisation of the preceding Hessian paragraph for all ten. Since $\delta g = \eta M + M^T\eta$, this block is the linearised metric perturbation, and its quadratic action contains only $(\partial h_{\mu\nu})^2$ and $h_{\mu\nu}h^{\mu\nu}$: the derivative index is contracted only with $g^{(0)\mu\nu}$, so no $\partial_\mu h^{\mu\nu}$ structure appears, the mass form is nondegenerate on all ten components, and the quadratic action has no linearised diffeomorphism invariance. Hence, for this candidate at this order, boost invariance is obtained but three opposite-sign modes remain, the potential is indefinite, and the quadratic form is not of Fierz–Pauli type (hypothesis 5 of Theorem 7.3). Whether higher-order terms, a kinetic term for ω_μ, or an induced action change this is not examined. This is a bounded no-go for the stated candidate at quadratic order about the flat scaffold; it does not exclude other completed actions. Problems D and B remain open.

Diffeomorphism symmetry of the displayed action (conditional; not closure). In (3.6) the field $\Phi^I{}_A$ carries no coordinate index: it is a set of scalar fields on the scaffold, and a world index appears only through the identification of the tangent index with the coordinate index in (4.7). With $g^{(0)}_{\mu\nu}$ held fixed (Assumption 3.1), a diffeomorphism of the scaffold coordinates acts on the argument x of Φ, and S_{micro} is invariant under those diffeomorphisms that preserve $g^{(0)}$, such as its isometries; this does not exclude other symmetries. If $g^{(0)}_{\mu\nu}$ is transformed as a tensor together with Φ, the functional $S_{\mathrm{micro}}[\Phi, g^{(0)}]$ is covariant in the spurionic sense; this does not by itself make $g^{(0)}$ a dynamical or compensating field, nor establish a gauge symmetry of the fixed-background theory. The composite metric $g = e^T\eta e$ built from the identified frame is a tensor only if the identification $A \leftrightarrow \mu$ is itself made covariant, which (3.6) does not provide. The statement of Section 10.5 that diffeomorphisms are the gauge symmetry of the composite metric, and the diffeomorphism-invariance hypotheses of

Theorem 11.1 and Assumption 12.2, are therefore conditional on such an extension, not consequences of S_{micro} as displayed. This is a scope statement; it does not exclude a covariant completion, and Problems B, D and G remain open.

What linearised diffeomorphism invariance fixes (conditional; not closure). Consider the most general parity-even, Lorentz-invariant quadratic action for an unconstrained symmetric $h_{\mu\nu}$ on a flat background with at most two derivatives. Up to integration by parts and overall normalisation, in momentum space it has four derivative terms with coefficients $a_1, \ldots, a_4$, multiplying $p^2 h_{\mu\nu} h^{\mu\nu}$, $p^\mu h_{\mu\nu} p_\lambda h^{\lambda\nu}$, $h\, p^\mu p^\nu h_{\mu\nu}$, $p^2 h^2$, and two mass terms $m_1 h_{\mu\nu} h^{\mu\nu} + m_2 h^2$. Requiring invariance under $\delta h_{\mu\nu} = p_\mu \xi_\nu + p_\nu \xi_\mu$ forces $a_2 = -2a_1$, $a_3 = 2a_1$, $a_4 = -a_1$ and $m_1 = m_2 = 0$: the massless Fierz–Pauli form for the unconstrained field (a direct symbolic check, used as a standard result and not as a new derivation). This classifies the unconstrained symmetric-h action. It is not identified here with the trace-constrained tensor block of (7.2) used in hypothesis 5 of Theorem 7.3; relating the two requires a separate argument that is not given. The displayed action (3.6) is not in this class. In the η-contracted candidate tested above, the block identified with $\delta g_{\mu\nu}$ has no $\partial_\mu h^{\mu\nu}$ structure and a nonzero mass term. For the displayed action, the real-slice symmetric block H does not span the full $\delta g_{\mu\nu}$, since H_{0i} leaves the metric unchanged, and no diffeomorphism-invariant quadratic structure appears in its Hessian. A covariant completion of the action may therefore be a necessary structural ingredient for a diffeomorphism-invariant quadratic action, but it is not sufficient for Problem B, which also requires entanglement stationarity, the area contribution, and their relation to the emergent metric. Problems B and D remain open.

Related covariant quantum-space constructions, and what they do not supply (conditional; not closure). Steinacker and collaborators study the IKKT matrix model on covariant quantum spaces in which the frame is read off from the adjoint action $[X^a, \cdot\,]$ and averaged over an internal S^2 fibre. The Euclidean, linearised analysis [22] is reported to give linearised Einstein equations under conditions on the background, dimensional reduction and an infrared cutoff; we have not checked these conditions. A Lorentzian space built on the minimal doubleton representation of $\mathfrak{so}(4,2)$, a quantised twistor space $\mathbb{CP}^{1,2}$ over a $k = -1$ FLRW spacetime, is described in [23], and an induced gravitational action with an explicit effective Newton constant is reported at one loop for such backgrounds with fuzzy extra dimensions in [16]. The absence of ghosts for the Lorentzian higher-spin theory is reported in [24], at the expense of manifest Lorentz invariance. We have not reproduced any of these results. Taking them as stated, they are analogous to, not the same as, the construction tested above: the frame comes from a chosen representation rather than from the vacuum value of a field $\Phi^I{}_A$; the ghost problem is addressed by giving up manifest Lorentz

invariance, which the locking assumption of Chapter 10 presupposes; and gravity is induced at one loop rather than obtained from entanglement stationarity as in Problem B. They therefore do not close Problems D, B, F, C or G. They report that a Lorentzian, ghost-free construction exists in a matrix-model setting; whether an analogue exists for the Φ-field action of Chapter 3 with Lorentz locking retained is not established here. Adopting an induced-gravity route in place of Problem B would be a structural change to the programme and is not proposed here.

The matrix-model no-ghost argument and the η-contracted candidate (conditional; not closure). In [24] the vector-type fluctuation spaces are reported to carry an indefinite inner product of signature $(+++-)$ before gauge fixing; the gauge-fixing condition is equivalent to orthogonality to the pure-gauge modes, which reduces the space to signature $(++0)$, and the quotient by the null gauge mode leaves two positive-norm modes. The negative-norm directions are thus removed by a gauge symmetry whose modes mix the derivative index with the field index. We have not reproduced this.

In the η-contracted candidate tested above, the tangent-index connection enters algebraically and removes only the η-anti-self-adjoint directions, so the three opposite-sign self-adjoint $0i$ directions are not gauge directions of it. Shifts of the form $\delta h_{\mu\nu} = \partial_{(\mu}\xi_{\nu)}$ would be one route to removing such modes; they are defined only through the identification of tangent and coordinate indices of (4.7), and invariance under them is sufficient for, but not shown necessary for, ghost removal. For this candidate at quadratic order, with kinetic part $a_1(\partial h)^2$ and mass form $\lambda v^4 \mathrm{K}(M_s)$, the quadratic action does not vanish on pure-gauge $h = \partial\xi$ for generic momentum (checked symbolically on a model quadratic form $a_1 k^2\, h{\cdot}h + m\, h{\cdot}h$; it vanishes only on shell, $a_1 k^2 + m = 0$, which does not test invariance). By the lemma above, invariance would require cross terms with the Fierz–Pauli ratios and a massless symmetric block, whereas this candidate has no such cross terms and the nonzero mass λv^2. The positive-definite kinetic form of the displayed action (3.6) has no negative-norm scalar-kinetic directions, but its tensor sector was not analysed here, and its real-slice block does not span $\delta g_{\mu\nu}$, as noted above. Two repairs of the candidate are visible and neither is adopted: adding such cross terms and a potential flat along the shift directions, or realising the shifts non-linearly with a compensating field. Each introduces a new object and hypothesis not derived from the matrix action. This is a bounded statement about the quadratic form of this candidate and does not exclude other completions.

A compensating field alone does not change the spectrum (conditional; not closure). Restore the shifts above by a Stueckelberg replacement $h_{\mu\nu} \to h_{\mu\nu} + \partial_{(\mu}\xi_{\nu)}$ in the η-contracted candidate, with a new field ξ_μ and gauge transformations $\delta h = \partial\epsilon$, $\delta\xi = -\epsilon$. The gauge $\xi = 0$ is reachable and in it the action equals the original, so

the physical spectrum is unchanged: the signature $(7,3)$ of the self-adjoint block, the three opposite-sign $0i$ directions and the mass λv^2 persist. The compensating field is a reparametrisation of this candidate; removing the ghosts would need structure beyond it, such as a kinetic structure for ξ and a constraint, which is not examined here. This concerns the quadratic form of this candidate and does not exclude other completions. Problems B and D remain open.

Integrable frames and indefinite gauge kinetic terms (conditional; not closure). Two further routes to a tangent-index shift symmetry for the η-contracted candidate tested above are not supplied by the linearised quadratic form. Both statements are bounded to that candidate and to the stated class of kinetic terms.

First, suppose the frame is integrable, $\Phi^I{}_\mu = \partial_\mu \phi^I$, where $\Phi^I{}_\mu$ denotes $\Phi^I{}_A$ converted by the tangent–coordinate identification of (4.7). The perturbation of its symmetric block is then $h_{\mu\nu} = \partial_{(\mu}\pi_{\nu)}$, and its antisymmetric block is a local Lorentz rotation. The linearised Riemann tensor of any pure-gauge perturbation $h_{\mu\nu} = \partial_{(\mu}\xi_{\nu)}$ on a flat background vanishes; this is the standard gauge invariance of the linearised curvature. The induced metric of an integrable frame is therefore flat, and invariance under the shifts is automatic but carries no linearised gravity. This does not exclude other structures. A shift symmetry that carries gravity must act on a non-integrable Φ. Such a Φ would be expected to play the role of a gauge field for the shifts, with torsion $\mathcal{D}\Phi$ as its field strength, and invariant kinetic terms would be expected to be built from torsion and curvature rather than from the displayed contraction of $\partial\Phi$ with the scaffold. This is a structural expectation following from requiring the symmetry, not a derivation and not a property of the displayed action.

Second, a quadratic kinetic term for connections of a non-compact tangent group built from an invariant bilinear form is indefinite. For $\mathfrak{so}(3,1)$ the Killing form has signature $(3,3)$, positive on the three boosts and negative on the three rotations, and the same holds for any nonzero real combination of the Killing form and its dual invariant form (the signature was computed numerically in the adjoint basis). This has the same signature as the $(3,3)$ anti-self-adjoint kinetic block found above; no identification of the two is claimed. A Yang–Mills-type term built from an invariant form for such a connection therefore carries wrong-sign modes, and ghost removal would need other structure, such as first-order terms with constraints, which this does not exclude.

What remains is to derive the required kinetic structure and constraint from the matrix action without inserting them by hand. A one-loop effective action is reported in [16] to induce a gravitational action for a different matrix model; we have not checked its operator against that of Chapter 8, and it is not shown here to supply the symmetry or the constraint. We expect, but have not shown for the present operator, that the field strength of the tangent-index connection enters the one-loop action at

four-derivative order through the a_2 density of Chapter 8 (the standard Seeley–DeWitt term $\propto \operatorname{tr}\Omega_{\mu\nu}\Omega^{\mu\nu}$), and that a two-derivative torsion-squared term would arise only if torsion appeared in the endomorphism E of (8.3), which Chapter 8 does not specify. Problems B, C, D, F and G remain open.

Diagonal invariance and the self-adjoint traceless block (conditional; not closure). Write the real locked block as $\Phi = v(\mathbb{I}+M)$, with mixed-index components $M^I{}_A$, and let the unbroken diagonal $\mathrm{SO}(3,1)$ of Proposition 10.2 act by conjugation, $\delta M = [\alpha, M]$ with $\eta\alpha$ antisymmetric. The sixteen components decompose as $\mathbf{1}\oplus\mathbf{6}\oplus\mathbf{9}$: the trace, the six η-anti-self-adjoint components, and the nine η-self-adjoint traceless components (six ij and three $0i$). The nine components form the real irreducible representation $(1,1)$, which has trivial commutant, so by Schur's lemma a quadratic form on them that is invariant under the diagonal group is unique up to a scale. The form K of the boost-invariant completed-action paragraph above, restricted to this block, is such a form, with signature $(6,3)$; the signature $(7,3)$ quoted there includes the trace direction. Every nonzero invariant form on the block is therefore $c\,\mathrm{K}$, with signature $(6,3)$ or $(3,6)$ according to the sign of c. A numerical check (`numpy`, not a proof) found four independent invariant quadratic forms on the sixteen components, whose restrictions to the nine-component block span a one-dimensional space, consistent with this count.

The statement assumes that the quadratic form is invariant under constant diagonal $\mathrm{SO}(3,1)$ acting by conjugation. The displayed potential does not have this property (see the paragraph on the assumed group $G_{\mathrm{int}} \supseteq \mathrm{SO}(3,1)$ in Problem F); the condition is a requirement on completions of the action, not a property of the displayed action. Under it, for a potential Hessian or for a kinetic form with the spacetime index contracted through $g^{(0)\mu\nu}$ only, as in (3.6), the form on the nine components is indefinite unless it vanishes, so a nonzero Hessian is not positive there, and a nonzero kinetic form has at least three wrong-sign directions (three or six, depending on the sign of c). This holds for every diagonal-invariant form without derivative–tangent mixing, which is a class of constructions and not all of them. Terms that tie the derivative index to the tangent index through the vacuum identification of Equation (4.7) add invariants and are not analysed here, so a Fierz–Pauli-type structure is not excluded. The trace and the anti-self-adjoint block are not constrained by this argument. It adds no field, symmetry or constraint, does not identify the nine components with the tensor sector of Chapter 5, and leaves Problem D open.

Tree-level terms plus an assumed induced Fierz–Pauli term: quadratic test (conditional; not closure). This paragraph tests which tree-level coefficients are compatible with a healthy spin-two sector if an induced Fierz–Pauli term is assumed. On the η-self-adjoint block, Equation (4.9) gives $\delta g_{\mu\nu} = 2(\eta M)_{\mu\nu}$, which is symmetric;

write $h_{\mu\nu}$ for a field proportional to $(\eta M)_{\mu\nu}$, the normalisation being absorbed into the coefficients below, and identify tangent and coordinate indices through Equation (4.7). Take the quadratic form at momentum k to be $Q = (tk^2 + m)\, h_{\mu\nu}h^{\mu\nu} + (t'k^2 + m')\, h^2 + g\, Q_{\mathrm{FP}}$, with $k^2 = \eta^{\mu\nu}k_\mu k_\nu$. The first two terms are the most general tree-level form invariant under the diagonal $\mathrm{SO}(3,1)$ with the derivative index contracted through $g^{(0)\mu\nu}$ only (the trace has an independent coefficient), and Q_{FP} is the Fierz–Pauli structure with ratios $(1, -2, 2, -1)$, indices raised with η. The induced term is an assumed input in the form of a linearised Einstein–Hilbert term with unspecified coefficient g, in the manner of Chapter 8; its coefficient and sign, which depend on Problem F, are not computed here. The Lagrangian is taken as $\mp Q$, with the sign chosen so that the spin-two modes are healthy; all wrong-sign statements below are relative to the spin-two modes.

A symbolic computation (`sympy`, a check and not a proof) confirms that Q_{FP} is invariant under $\delta h = k\xi$ and gives, at $k = (\omega, 0, 0, q)$, a determinant of the ten-by-ten form with the following structure. If $t = t' = m = m' = 0$, the four components $h_{0\mu}$ are non-dynamical and the form is the massless Fierz–Pauli form. If $t \neq 0$, there are three poles at $k^2 = -m/t$, five spin-two poles at $(g+t)k^2 + m = 0$, and two poles in the scalar sector, so the $h_{0\mu}$ components propagate. For $t' = m' = 0$, $g > 0$ and $|t| \ll g$, the coefficient matrix of ω^2 has, at $t = 0$, one direction of sign opposite to the spin-two modes (the trace) and four null directions; for $t > 0$ the three h_{0i} directions have a sign opposite to the spin-two modes, and for $t < 0$ the h_{00} direction does. For $t = t' = 0$ and $m' \neq -m$ there is one additional scalar pole, whose sign (ghost or tachyon) is not determined here; it is absent for $m' = -m$, the Fierz–Pauli mass form, leaving only the five spin-two poles at $gk^2 + m = 0$. Diagonal invariance alone does not impose $t = t' = 0$ or $m' = -m$. Within this class and these parameter ranges, therefore, a tree-level derivative term on the symmetric block adds propagating components that carry directions of sign opposite to the spin-two modes, and a tree-level mass leaves no extra scalar pole only at the Fierz–Pauli trace ratio. These are conditions that such a completion would place on the tree-level coefficients. The displayed kinetic term of Equation (3.6) is not diagonal-invariant (see the preceding paragraphs) and so is not a member of this class. The vector and ghost sectors, the sign and size of g, the continuation and the one-loop operator are not treated, so this does not show that an induced route is ghost-free, does not identify the block with the tensor sector of Chapter 5, and leaves Problems B, C, D, F and G open.

Displayed action plus an assumed induced Fierz–Pauli term: time-kinetic test (conditional; not closure). This paragraph keeps the displayed action as the tree-level term and adds only the induced term assumed above; it introduces no new tree-level structure. On the η-self-adjoint block, with $h_{\mu\nu}$ proportional to $(\eta M)_{\mu\nu}$ as above, the displayed kinetic term of Equation (3.6) contracts the components with

the Euclidean form, $\mathrm{Tr}\, M^T M = \sum_{\mu\nu} h_{\mu\nu}^2$ for $M = \eta h$, so take $Q = tk^2 \sum_{\mu\nu} h_{\mu\nu}^2 + g\, Q_{\mathrm{FP}} + Q_{\mathrm{mass}}$ with $t > 0$, the same Q_{FP} and the same sign convention as above. The displayed potential acts through the Hermitian part $H = (M + M^T)/2$ of the Hessian paragraph in Problem F; for $M = \eta h$ the $0i$ entries of H cancel, so $Q_{\mathrm{mass}} \propto h_{00}^2 + \sum_{ij} h_{ij}^2$ acts on seven components and the three $0i$ components are massless at this order. A symbolic computation (`sympy`, a check and not a proof) gives the ω^2-coefficients of the form at $k = (\omega, 0, 0, 0)$, which do not depend on the mass, in the convention where healthy modes have negative coefficient: $-2t$ (once), $-4t$ (three times), $4g - 2t$ (once, the trace direction), $-2(g + t)$ (twice) and $-4(g + t)$ (three times). All ten are healthy exactly when $t > 0$ and $-t < g < t/2$; for $g > 0$ only $g < t/2$ binds. Without the mass term, the determinant at $k = (\omega, 0, 0, q)$ factors as $t^4(g + t)^2(q^2 - \omega^2)^6 \left[(t - g)q^2 - (t + g)\omega^2\right]^2 \left[t(t - g)(q^2 - \omega^2)^2 - 2g^2(q^2 + \omega^2)^2\right]$ up to a constant. The squared factor gives modes with $c^2 = (t - g)/(t + g) \neq 1$ for $g \neq 0$, and the final quartic factor is also non-luminal for $g \neq 0$. The dispersion is then Lorentz-violating at order g/t, in tension with the exact Lorentz invariance that the locking construction of Chapter 10 is intended to deliver; this tension is recorded, not resolved.

If the induced term is the linearised $(1/16\pi G_{\mathrm{eff}}) \int \sqrt{-g}\, R$, with $\delta g_{\mu\nu} = 2h_{\mu\nu}$ from Equation (4.9) and the standard normalisation of the linearised Einstein–Hilbert quadratic form (used, not re-derived), then matching the coefficient of $k^2 h_{\mu\nu} h^{\mu\nu}$ gives $t = v^2$ from Equation (3.6) and $g = 1/(16\pi G_{\mathrm{eff}})$. For $G_{\mathrm{eff}} > 0$, the window above is then $1/G_{\mathrm{eff}} < 8\pi v^2$, and $1 - c^2 \approx 2g/t = 1/(8\pi G_{\mathrm{eff}} v^2)$ is small only if $G_{\mathrm{eff}} v^2 \gg 1$. Inserting the conditional parameterization of Equation (8.12), the window reads $A\mathcal{N}_0 \Lambda_{\mathrm{UV}}^2/v^2 + B \ln(\Lambda_{\mathrm{UV}}/\mu) + C\mu^2/v^2 < 8\pi$; if the logarithmic and μ^2 terms are negligible, it reads $A\mathcal{N}_0 \Lambda_{\mathrm{UV}}^2 < 8\pi v^2$. The coefficients A, B, C, $\mathcal{N}_0$, μ and Λ_{UV} are undetermined (Theorem 8.2 and Problem F), so this identifies the combination that Problem F would have to satisfy and does not decide whether it does. The window has the same structure as the condition $Z > 0$ with $Z = 2N - 3/(4\pi G_{\mathrm{eff}} v^2)$ of the Repair II pure-trace truncation above (a conformal-mode kinetic coefficient that turns negative when $G_{\mathrm{eff}} v^2$ is small), but the numerical factors differ and no equivalence is claimed. The result is for an assumed linearised induced term; it is a statement about the time-kinetic matrix and the dispersion at quadratic order, so it is not a spectrum or unitarity result. It does not treat the vector, ghost or ω_μ sectors, does not identify the block with the tensor sector of Chapter 5, and leaves Problems B, C, D, F and G open.

H.5 Problem E: signature

Ledger label. Status: OPEN. Priority: CRITICAL PATH for the matrix-derived label. Completion: PROGRAMME-CONDITIONAL. A scope retaining Assumption 4.6

gives up the claim of emergent or matrix-derived Lorentzian signature.

Euclidean positivity of the matrix action and the formal Wick rotation of one coordinate are recorded. They do not select Lorentzian signature without an additional hypothesis.

Status. Remains Assumption 4.6.

H.6 Problem F: coefficients A, B, C

Ledger label. Status: OPEN. Priority: CRITICAL PATH. Completion: PROGRAMME-CONDITIONAL. Reduced-scope result: treating G_{eff} as an input leaves F OPEN and does not satisfy the matrix-derived criterion.

The conditional, truncated heat-kernel identity is Theorem 8.2; it is not a computed A, B, C parameterization or physical value of G_{eff}.

Missing step. Specify the gauge-fixed fluctuation, Goldstone, diffeomorphism-ghost, and local-Lorentz-ghost operators and weights; per-sector proper-time limits; massive threshold functions if masses are resummed; the continuation; and the finite-counterterm scheme. Then evaluate the weighted a_1 and curvature-linear a_2 contributions in that scheme. Until this is completed, Problem F remains open, and the asserted $N_{\text{eff}} = +2$ count does not close it.

Quadratic Hessian of the displayed action (conditional; not closure). Restrict to the square locked 4×4 block of Assumption 5.1, expand (3.6) with (3.7) on the flat scaffold about $\Phi = v\mathbb{I} + d$ at $A_\mu = 0$, and write $d = H + K$ with $H = (d + d^\dagger)/2$ Hermitian and $K = (d - d^\dagger)/2$ anti-Hermitian. A direct expansion gives $V^{(1)} = 0$ and $V^{(2)} = \lambda v^2 \,\mathrm{Tr} H^2$, independent of K, and $\mathrm{Tr}(d^\dagger d) = \mathrm{Tr} H^2 + \mathrm{Tr}(K^\dagger K)$ with no cross term. The H–H block therefore has the Lorentzian field equation $(\Box - \lambda v^2)H = 0$ with the kinetic sign of (3.6), that is, the Euclidean operator $-\nabla_E^2 + \lambda v^2$ after Wick rotation. This is the Φ–Φ block at fixed $A_\mu = 0$; quadratic mixing of A_μ with Φ through the covariant derivative is not evaluated, and no gauge-fixed coupled Hessian is given. No identification of H with the ψ and $T_{\mu\nu}$ of Chapter 5 is made here, since that split uses $\eta_{\mu\nu}$ and the vacuum identification of Assumption 3.5. Equation (3.6) displays no field-strength term for A_μ, and ω_μ does not appear in it; its operator is unspecified (Assumption 10.1). The displayed action also does not exhibit the Fierz–Pauli structure used as hypothesis 5 of Theorem 7.3, so any tensor-block terms beyond it are an additional input. The gauge-fixed sector list, Goldstone and ghost operators, measure, curvature couplings and weights are not specified by this calculation, so it does not fix the sector sum of Theorem 8.2, does not close F or D, and leaves the missing step above unchanged.

Gauge-group dependence of the scalar block (conditional; not closure). The contraction $\mathrm{Tr}\,\Phi^\dagger\Phi$ is a positive-definite norm, and the potential in (3.7), being built from it, is invariant under constant unitary transformations of the internal index and of the tangent index; neither is invariant under non-compact boosts of the tangent index. The covariant kinetic term has a *local* symmetry only for those transformations for which a connection and its transformation law are specified; for the tangent index this requires Assumption 10.1 with ω_μ, which is not part of (3.6). Moreover, the compact group U(4) has no real $\mathfrak{so}(3,1)$ subalgebra, so choosing $G_{\mathrm{int}} = \mathrm{U}(4)$ is not compatible with the construction of Chapter 10 that requires G_{int} to contain $\mathrm{SO}(3,1)$ or its algebra. It defines a different compact-symmetry model, whose consequences for Problem D are not examined here.

For that separate model only, work on the square locked 4×4 block about $\Phi = v\mathbb{I}$ with A_μ as an unspecified fluctuation. Write $\delta\Phi = H + K$ as above. Assume (i) the action has no $F_{\mu\nu}F^{\mu\nu}$ term; (ii) $G_{\mathrm{int}} \subseteq \mathrm{U}(4)$ acts with the infinitesimal orbit $\delta K = \alpha\, v$ for α in its Lie algebra; and (iii) the orbit directions are removed by the gauge condition. Then the number of anti-Hermitian directions not tangent to the orbit, which have no quadratic potential, is $16 - \dim G_{\mathrm{int}}$; for U(4) this is zero and for SU(4), SO(4), U(1) it is 4, 10, 15 respectively. For $G_{\mathrm{int}} = \mathrm{U}(4)$ the linearized gauge condition $K = 0$ has a field-independent Jacobian at the vacuum, and the remaining Φ–Φ block is the sixteen real components of H with $m^2 = \lambda v^2$. This is the scalar Hessian block only: the ghost and measure contributions beyond the linearized Jacobian, the ultralocal determinant from the nonlinear A_μ integral, any induced vector or tensor operator, and the weighting of the full sector sum all remain unspecified, and the count is not a heat-kernel trace for the Problem F target. The discrepancy with the $\mathrm{SO}(3,1)$ assumption is recorded, not resolved. Problem F remains open.

Bookkeeping for one sector of the sum: the Hermitian scalar block (conditional; not closure). This paragraph illustrates how Theorem 8.2 would be applied to one block. It is not a derivation of that block's coefficients. The Hessian paragraph above is evaluated on the fixed flat scaffold $g^{(0)}$ of Assumption 3.1, where $R = 0$, so the displayed action supplies no curvature coupling for the emergent metric. The operator on the emergent background and the curvature coupling Q below are the Presentation-B assumptions of Chapter 8, not outputs of the displayed action.

Take the Hermitian block H of the Hessian paragraph above, with sixteen real components. Assume (i) that this diagonal block at $A_\mu = 0$ is treated as decoupled from the other sectors, so the coupled determinant is not given; (ii) the determinant weight is $w = 1$ per real bosonic component; and (iii) the operator of Equation (8.3) on the block is $-\nabla^2 + E$ with $E = \lambda v^2 + QR_E$, where Q is proportional to the identity on the sixteen components (otherwise $1/6 - Q$ below is replaced by its trace over the components). With $a_1 = \mathrm{tr}(R/6 - E)$ of (8.7) and the standard

curvature-linear part of $a_2 \supset \frac{1}{2}\operatorname{tr}(R/6 - E)^2$, which has the same sign convention as a_1 and is used here without being re-derived, the coefficients of R_E for this block are $\mathcal{A}_{1,R,H} = 16(\frac{1}{6} - Q)$ and $\mathcal{A}_{2,R,H} = 16\lambda v^2(Q - \frac{1}{6}) = -\lambda v^2 \mathcal{A}_{1,R,H}$. In the truncated, unresummed identity of Theorem 8.2 with common proper-time limits, the block would contribute $-\frac{\mathcal{A}_{1,R,H}}{32\pi^2}\left[(\Lambda_{\mathrm{UV}}^2 - \mu^2) - \lambda v^2 \ln(\Lambda_{\mathrm{UV}}^2/\mu^2)\right]$ to C_R^E; that is, in this block's contribution the coefficient of $\lambda v^2 \ln(\Lambda_{\mathrm{UV}}^2/\mu^2)$ is minus the coefficient of $\Lambda_{\mathrm{UV}}^2 - \mu^2$. Both coefficients vanish at $Q = \frac{1}{6}$; this concerns only this block's R_E-coefficient in the truncated expansion. Because the block has mass λv^2, a resummed treatment replaces these terms by threshold functions, so the expression is not the physical coefficient. No comparison with A, B, C of Equation (8.12) is made. The anti-Hermitian block, the A_μ and ω_μ sectors, the tensor block, the ghosts, the continuation and the scheme are not specified here, so this does not determine G_{eff}, does not fix the sign of the Lorentzian term, and leaves Problem F open.

Mass-resummed curvature threshold for the Hermitian block (conditional; not closure). Keep the assumptions of the preceding one-block bookkeeping paragraph: sixteen decoupled real components, determinant weight $w = 1$, common proper-time limits, and $\mathcal{O}_H = -\nabla^2 + E$, with $E = m^2 + QR_E$, $m^2 = \lambda v^2$, and constant m^2 and Q proportional to the identity. At the retained order assume $\nabla E = 0$, and keep only terms linear in R_E, dropping curvature-squared invariants and curvature-derivative terms. Retaining the mass exponential, the coincident-point heat kernel then contains

$$\operatorname{tr}\langle x|e^{-s\mathcal{O}_H}|x\rangle \supset \frac{16}{(4\pi s)^2} e^{-sm^2} s \left(\frac{1}{6} - Q\right) R_E.$$

Thus this block's contribution to C_R^E, in the sign and normalization convention of the preceding paragraph, with $s_{\min} = \Lambda_{\mathrm{UV}}^{-2}$ and $s_{\max} = \mu^{-2}$, is

$$-\frac{16}{32\pi^2}\left(\frac{1}{6} - Q\right) J(m^2; s_{\min}, s_{\max}),$$

where

$$\begin{aligned} J(m^2; s_{\min}, s_{\max}) &:= \int_{s_{\min}}^{s_{\max}} \frac{\mathrm{d}s}{s^2} e^{-m^2 s} \\ &= \frac{e^{-m^2 s_{\min}}}{s_{\min}} - \frac{e^{-m^2 s_{\max}}}{s_{\max}} - m^2\left[E_1(m^2 s_{\min}) - E_1(m^2 s_{\max})\right], \qquad E_1(x) := \int_x^\infty \end{aligned}$$

The $m \to 0$ limit is $J(0) = \Lambda_{\mathrm{UV}}^2 - \mu^2$. For $m^2 s_{\max} \ll 1$,

$$J = \Lambda_{\mathrm{UV}}^2 - \mu^2 - m^2 \ln\left(\frac{\Lambda_{\mathrm{UV}}^2}{\mu^2}\right) + O\left(m^4 s_{\max}\right).$$

This small-mass expansion gives the earlier truncated expression; the resummed J is the threshold function, and the earlier expression is not removed. At this order in curvature, the resummed expression includes the mass insertions represented by that curvature-linear a_2 term; they must not be added a second time. For $m^2 s_{\min} \gg 1$, this block's loop contribution is exponentially suppressed. The full coefficient still includes the scheme-dependent local curvature counterterm of Theorem 8.2; this block calculation does not specify that counterterm or the other sectors, their operators and weights, the gauge fixing, or the continuation. Problems B, C, D, F and G remain open.

The assumed group $G_{\rm int} \supseteq \mathrm{SO}(3,1)$ and the displayed potential (conditional; not closure). Chapter 10 takes $G_{\rm int}$ to contain $\mathrm{SO}(3,1)$ acting on the internal index, with orbit $\delta d = v\alpha$ at the vacuum. As mixed-index matrices $\alpha^I{}_J$ on the locked block, its three boost generators are real symmetric matrices, for example $B = E_{01} + E_{10}$, and so lie in the Hermitian block H of the Hessian paragraph above, while the rotations are antisymmetric. For $d = vB$ that paragraph gives $V^{(2)} = \lambda v^2 \,\mathrm{Tr} H^2 = 2\lambda v^4 \neq 0$, so the potential of (3.7), which is built from the positive-definite $\mathrm{Tr}\,\Phi^\dagger\Phi$, is not invariant under these generators. Assumption (ii) of the preceding paragraph, that the orbit lies in the anti-Hermitian block, therefore does not hold for $G_{\rm int} = \mathrm{SO}(3,1)$; that count is for a different model, and $16 - \dim G_{\rm int}$ does not apply here because the orbit has components in H. For the displayed potential, three of the six broken directions of Proposition 10.2 carry mass λv^2, so that proposition's flat-Goldstone count requires a boost-invariant potential, such as an η-contracted form like the candidate discussed under Problem D, whose quadratic form is indefinite. This dependence is recorded, not resolved. Declaring $G_{\rm int}$ therefore does not by itself specify the anti-Hermitian or A_μ sectors of Problem F: those depend on the choice of potential, which is the open content of Problem D. The statement is for the displayed action at quadratic order; it does not rule out other constructions, does not evaluate any further sector of Theorem 8.2, and leaves Problems D and F open.

Sector sum: a bound from the universal a_1 piece. Fix a common limit pair $\Lambda_{\rm UV}, \mu$, take $Q_s = 0$, and keep only the universal a_1 term of Equation (8.9); the curvature-linear part of a_2, thresholds and counterterms are dropped. Assume, as a convention to be checked against the continuation of Appendix A, that the Euclidean effective action contains $-\frac{1}{16\pi G_{\rm eff}} \int \sqrt{g}\, R_E$. Then

$$\frac{1}{G_{\rm eff}} = \frac{\mathcal{N}_0}{12\pi}\,(\Lambda_{\rm UV}^2 - \mu^2), \qquad \mathcal{N}_0 = \sum_s w_s \,\mathrm{tr}_{V_s}\, \mathbb{I}_s,$$

that is, $A = 1/(12\pi)$ for this piece alone. This is the weighted-trace form of the single-sector formula of Section 17.4. Suppose in addition that the induced term of the displayed-action time-kinetic paragraph is a linearised Einstein–Hilbert term whose coefficient $g = 1/(16\pi G_{\text{eff}})$ is the total induced coefficient, so that its window $1/G_{\text{eff}} < 8\pi v^2$ holds. At $\mu \to 0$ this gives

$$0 < \mathcal{N}_0 \frac{\Lambda_{\text{UV}}^2}{v^2} < 96\pi^2, \qquad \text{i.e.} \qquad \frac{\Lambda_{\text{UV}}}{v} < \pi\sqrt{\frac{96}{\mathcal{N}_0}}.$$

For $\mu > 0$ the factor $\Lambda_{\text{UV}}^2 - \mu^2$ only weakens the bound. Positivity of G_{eff} requires $\mathcal{N}_0 > 0$ in this convention. If $\Lambda_{\text{UV}} \gg v$, the weighted sum must therefore nearly cancel. As an illustration only, one unit-weight sixteen-component sector would give $\Lambda_{\text{UV}}/v < \pi\sqrt{6} \approx 7.7$. The numerics are illustrative, not an evaluation of $\mathcal{N}_0$. The unresummed truncation also requires $\lambda v^2 \ll \Lambda_{\text{UV}}^2$ for the massive sectors, which may fail for Λ_{UV} of order v. The sum itself is not computed here because the manuscript does not supply: the gauge-fixed operators and E_s for the K and A_μ sectors (which depend on the G_{int} and potential choices of Problem D), the diffeomorphism and local-Lorentz ghost operators and weights, the per-sector limits, the thresholds, and the continuation convention. Problems B, C, D, F and G remain open.

Conditional gauge-fixed Hermitian block in the compact $U(4)$ branch (not closure). This calculation concerns only the separate compact $U(4)$ model described above, not the $\mathrm{SO}(3,1)$-compatible construction. Assume A_μ is a Hermitian $U(4)$ connection, with no $F_{\mu\nu}F^{\mu\nu}$ term, and write $\Phi = v\mathbb{I} + H + iY$, where H, Y are Hermitian. To quadratic order the displayed action gives

$$D_\mu\Phi = \partial_\mu H + i(\partial_\mu Y - vA_\mu) + O(\text{quadratic}),$$

$$S^{(2)}_{\Phi,A} = \int \mathrm{d}^4x\,\sqrt{-g^{(0)}}\,\mathrm{Tr}\Big[-g^{(0)\mu\nu}\,(\partial_\mu H\,\partial_\nu H + (\partial_\mu Y - vA_\mu)(\partial_\nu Y - vA_\nu)) - \lambda v^2 H^2\Big].$$

Thus the derivative mixing is between Y and A_μ, not H and A_μ. Choose the linearized unitary gauge $Y/v = 0$; Y is eliminated by this gauge condition and is not integrated as a fluctuation in this gauge. For $\Phi \mapsto e^{i\alpha}\Phi$, one has $\delta Y = v\alpha$ at the vacuum, so the Faddeev–Popov operator for this condition is $\mathbb{I}_{16}$ there. A complex ghost pair has relative determinant weight $w = -2$ in the convention of Equation (8.1), but this operator is ultralocal. In the same gauge A_μ has only an algebraic quadratic term: the action supplies no differential operator for it. Its determinant and the ghost determinant are therefore not Laplace-type heat-kernel sectors; their regulated local contributions are not evaluated here.

For the differential Hermitian block H alone, further assume that its operator on the emergent background is $-\nabla^2 + \lambda v^2$, with no R_E term in E, and assign it common

limits $\Lambda_{\rm UV}, \mu$. These are Presentation-B and cutoff assumptions, not consequences of the displayed action on the fixed scaffold. In the truncated, unresummed expansion,

$$\mathcal{N}_{0,H} = 16, \qquad \mathcal{A}_{1,R,H} = \frac{8}{3}, \qquad \mathcal{A}_{2,R,H} = -\frac{8}{3}\lambda v^2.$$

The a_2 coefficient uses, without re-derivation, the curvature-linear term $\frac{1}{2}\,\mathrm{tr}(R_E/6 - E)^2$ used in the preceding Hermitian-block bookkeeping paragraph; the expressions here are its $Q = 0$ specialization. With the Euclidean matching convention $C_R^E = -1/(16\pi G_H)$, Theorem 8.2 then gives the single-block contribution

$$\frac{1}{G_H} = \frac{4}{3\pi}\left[\Lambda_{\rm UV}^2 - \mu^2 - \lambda v^2 \ln\left(\frac{\Lambda_{\rm UV}^2}{\mu^2}\right)\right].$$

Using $\ln(\Lambda_{\rm UV}^2/\mu^2) = 2\ln(\Lambda_{\rm UV}/\mu)$, this block alone corresponds formally to $A_H = 1/(12\pi)$, $B_H = -8\lambda/(3\pi)$, and $C_H = -4/(3\pi)$ in Equation (8.12). These are formal single-block values in a parameterization whose sectors need not share μ; they are not full-theory coefficients. The unresummed expansion requires $\lambda v^2 \ll \Lambda_{\rm UV}^2$, so these values are not threshold-correct. The algebraic determinants, other fluctuation and ghost sectors, sector thresholds, continuation, and finite curvature counterterm are not included. The displayed nonnegative potential is not boost-invariant, so this compact branch does not supply the missing SO(3, 1)-compatible sector sum. Problems B, C, D, F and G remain open.

Finite-cell check of the ultralocal determinants (conditional; not closure). In the compact $U(4)$ branch and linearized unitary gauge of the preceding paragraph, regulate the Euclidean fixed scaffold by finitely many cells of Euclidean proper volume ΔV_0. Use a basis of Hermitian generators orthonormal under the trace and Lebesgue measure for the four Euclidean orthonormal components of each of the sixteen generators. At the vacuum the linearized Faddeev–Popov matrix is $\mathbb{I}_{16}$ at each cell, so its finite-dimensional determinant is one and its contribution $\mathrm{Tr}\ln\mathbb{I}_{16}$ vanishes in this regulator. The eliminated Y fields are not integrated again.

After continuation, the Euclidean action for the algebraic A_μ variables is positive in all four orthonormal components. The finite-cell Gaussian is

$$\int \mathrm{d}^{64}A\, \exp\left[-v^2\Delta V_0 \sum_{a=1}^{16}\sum_{\hat{\mu}=1}^{4}(A_{\hat{\mu}}^a)^2\right] = \left(\frac{\pi}{v^2\Delta V_0}\right)^{32}.$$

For equal cells its logarithm is $-32N_{\rm cells}\ln(v^2\Delta V_0/\pi)$, a volume-type term with no curvature dependence in this scaffold regulator. That absence of curvature dependence is a property of the regulator built on the fixed scaffold, not a result about the emergent

curvature coefficient. The A_μ integral is not of Laplace type, so a_1 does not apply to it as written. Its local contribution depends on the functional-measure normalization; the displayed action and heat-kernel cutoff alone do not select that normalization. This calculation does not justify assigning these ultralocal factors zero in a different measure or regulator, and does not determine C_R^E. It is confined to the compact $U(4)$ branch and does not provide the missing $\mathrm{SO}(3,1)$-compatible sector sum. Problems B, C, D, F and G remain open.

Which metric the scaffold one-loop action induces (conditional; not closure). Consider the displayed action (3.6) with potential (3.7) on the square locked block, on the scaffold $g^{(0)}$ of Assumption 3.1, after Wick rotation, with the internal connection held at a background value $\bar{A}_\mu$ and not integrated. Expand $\Phi = \bar{\Phi} + \varphi$ about a smooth background $\bar{\Phi}$ and write φ in thirty-two real components normalized so that the kinetic term is $\sum_a g^{(0)\mu\nu} \bar{D}_\mu \varphi_a \bar{D}_\nu \varphi_a$. The kinetic term is bilinear in Φ, so its second variation does not depend on $\bar{\Phi}$, and the fluctuation operator is

$$\mathcal{O}_{\bar{\Phi}} = -g^{(0)\mu\nu} \bar{D}_\mu \bar{D}_\nu + E(\bar{\Phi}),$$

where $\bar{D}$ is built from $g^{(0)}$ and $\bar{A}_\mu$, and $E(\bar{\Phi})$ is one half of the Hessian of V at $\bar{\Phi}$; E contains no derivative of $\bar{\Phi}$. Use this operator in (8.4) with the limits (8.5). For $I(\Lambda_{\mathrm{UV}}) = \int_{\Lambda_{\mathrm{UV}}^{-2}}^{\mu^{-2}} s^{-1} f(s)\, \mathrm{d}s$ one has $\partial_{\Lambda_{\mathrm{UV}}} I = 2\Lambda_{\mathrm{UV}}^{-1} f(\Lambda_{\mathrm{UV}}^{-2})$, so the Λ_{UV}-dependence of $W_E^{(1)}$ is fixed by the small-s expansion of the heat trace through a_2, up to terms that vanish as $\Lambda_{\mathrm{UV}} \to \infty$. By (A.5)–(A.7),

$$W_E^{(1)} \supset -\int \mathrm{d}^4x \sqrt{g^{(0)}} \left[\frac{32\,\Lambda_{\mathrm{UV}}^4}{64\pi^2} + \frac{\Lambda_{\mathrm{UV}}^2}{32\pi^2} \operatorname{tr}\Big(\tfrac{1}{6} R[g^{(0)}] - E(\bar{\Phi})\Big) + \frac{1}{32\pi^2} \ln\Big(\frac{\Lambda_{\mathrm{UV}}^2}{\mu^2}\Big) a_2 \right],$$

where a_2 contains E^2, $R[g^{(0)}]E$, curvature-squared terms of $g^{(0)}$, $\bar{F}_{\mu\nu}\bar{F}^{\mu\nu}$ and total derivatives. No Λ_{UV}^4, Λ_{UV}^2 or $\ln \Lambda_{\mathrm{UV}}$ term contains a derivative of $\bar{\Phi}$ outside a total derivative; terms with derivatives of $\bar{\Phi}$ have coefficients that are independent of Λ_{UV} or vanish as $\Lambda_{\mathrm{UV}} \to \infty$. The Λ_{UV}^2 curvature term is the term (8.9), with $\mathcal{N}_0 = 32$, for the scaffold metric $g^{(0)}$, which Assumption 3.1 holds fixed.

For the composite metric the corresponding coefficient vanishes. On the flat scaffold take $\bar{A}_\mu = 0$ and $\bar{\Phi} = (v + \sigma)\mathbb{I}$, so that $g = (1 + \sigma/v)^2 \eta$ by (17.3); in the Euclidean section $\sqrt{g}\, R[g] = 6(\partial\sigma)^2/v^2$ up to a total derivative. The Λ_{UV}^2 term above is $\frac{\Lambda_{\mathrm{UV}}^2}{32\pi^2} \int \mathrm{d}^4x \operatorname{tr} E(\sigma)$, with $\operatorname{tr} E(\sigma) = 16\lambda(v^2 + 4v\sigma + 2\sigma^2)$, and contains no $(\partial\sigma)^2$. Building the operator on g instead, as in Assumption 8.4, gives $-\mathcal{N}_0 \Lambda_{\mathrm{UV}}^2 (\partial\sigma)^2/(32\pi^2 v^2)$ at the same order, which is the induced term of (17.4) with $1/(16\pi G_{\mathrm{eff}}) = \mathcal{N}_0 \Lambda_{\mathrm{UV}}^2/(192\pi^2)$. The replacement supplies the whole Λ_{UV}^2 coefficient for the composite metric; it does not correct a scaffold value.

The $(\partial\sigma)^2$ term of the scaffold one-loop action has no $\Lambda_{\rm UV}^2$ or $\ln\Lambda_{\rm UV}$ piece. Take the limits (8.5) and $p^2 \ll \mu^2 \ll \lambda v^2$. At $\bar{A}_\mu = 0$ the sixteen Hermitian components have $E_H = \frac{\lambda}{2}(3(v+\sigma)^2 - v^2)$ and the sixteen anti-Hermitian components have $E_K = \frac{\lambda}{2}((v+\sigma)^2 - v^2)$. At second order in σ a block of n components with mass m gives

$$W_E^{(1)} \supset -\frac{n}{4}\,E'(0)^2 \int \frac{\mathrm{d}^4p}{(2\pi)^4}\,\tilde{\sigma}(p)\tilde{\sigma}(-p)\,B(p),$$

$$B(p) = \frac{1}{16\pi^2}\int_0^1 \mathrm{d}x\left[E_1\!\left(\frac{m^2 + x(1-x)p^2}{\Lambda_{\rm UV}^2}\right) - E_1\!\left(\frac{m^2 + x(1-x)p^2}{\mu^2}\right)\right],$$

with $E_1(x) = \int_x^\infty e^{-t}t^{-1}\,\mathrm{d}t$, the exponential integral. For the Hermitian block, $n = 16$, $E'(0) = 3\lambda v$, $m^2 = \lambda v^2$ and $\partial_{p^2}B|_{p=0} = -(e^{-\lambda v^2/\Lambda_{\rm UV}^2} - e^{-\lambda v^2/\mu^2})/(96\pi^2\lambda v^2)$. For the anti-Hermitian block, $n = 16$, $E'(0) = \lambda v$, $m = 0$ and $\partial_{p^2}B|_{p=0} = (\Lambda_{\rm UV}^{-2} - \mu^{-2})/(96\pi^2)$. The local term $\frac{1}{2}\Delta Z\int \mathrm{d}^4x\,(\partial\sigma)^2$ therefore adds to the microscopic value $2N$ of Z in (17.4)

$$\Delta Z = \frac{3\lambda}{4\pi^2}\Big(e^{-\lambda v^2/\Lambda_{\rm UV}^2} - e^{-\lambda v^2/\mu^2}\Big) + \frac{\lambda^2 v^2}{12\pi^2}\Big(\frac{1}{\mu^2} - \frac{1}{\Lambda_{\rm UV}^2}\Big),$$

that is, $3\lambda/4\pi^2$ up to $O(\lambda v^2/\Lambda_{\rm UV}^2)$ and $O(e^{-\lambda v^2/\mu^2})$ from the Hermitian block, plus a positive term from the anti-Hermitian block that depends on the infrared limit μ and is replaced by a term non-analytic in p^2 as $\mu \to 0$. Both terms are positive for $\mu < \Lambda_{\rm UV}$ and carry no $\Lambda_{\rm UV}^2$ or $\ln\Lambda_{\rm UV}$ piece. In this one-loop calculation the conformal-mode kinetic term keeps the sign of Lemma 17.3. The opposite-sign term of (17.4) is not produced in this calculation; in the manuscript it arises from the induced Einstein–Hilbert term, that is, from the operator built on the emergent metric.

Scaffold rotations exclude linearized diffeomorphism invariance in a stated class (conditional; not closure). In (3.6) the components of ${\Phi^I}_A$ are scalars of the scaffold: a scaffold index appears only on ∂_μ and A_μ, and on ω_μ in the completion (10.3). On the flat Euclidean scaffold, rotations $x \mapsto Rx$, $R \in \mathrm{SO}(4)$, acting on scaffold indices alone are a symmetry, and the background $\bar{\Phi} = v\mathbb{I}$, $\bar{A}_\mu = \bar{\omega}_\mu = 0$ is invariant. Assume that the regulator and the gauge fixing preserve this symmetry, as the scaffold proper-time regulator above does, and that the composite-metric fluctuation $h_{\mu\nu}$ is a fixed linear function of the components of Φ, for example the one obtained from (4.9) with the identification (4.7). Then, at any loop order, the quadratic part of the effective action in the components of Φ, with the connections set to zero or eliminated at quadratic order, has the form $\sum_{a,b}\tilde{\varphi}_a(-p)F_{ab}(p^2)\tilde{\varphi}_b(p)$: invariance gives $F_{ab}(Rp) = F_{ab}(p)$, an $\mathrm{SO}(4)$-invariant function of one vector depends only on p^2, and eliminating a connection, whose mixed and diagonal kernels have the forms $p_\mu f(p^2)$ and $\alpha(p^2)\delta_{\mu\nu} + \beta(p^2)p_\mu p_\nu$ with internal labels suppressed, preserves this form. Let Q_p be the reduced form on h obtained by eliminating the remaining components; it

is the reduced form that inherits any invariance of the full effective action under a linearized gauge transformation, and it too depends on p only through p^2. If Q_p is invariant under $h \mapsto h + p \otimes \xi + \xi \otimes p$ for all ξ, then Q_p annihilates $p' \otimes \xi + \xi \otimes p'$ for every p' with $p'^2 = p^2$; for $p \neq 0$ these tensors span all symmetric tensors, so $Q_p = 0$. The same holds on the traceless block with the transformation (7.5). The linearized Einstein–Hilbert form and the massless Fierz–Pauli form (7.2) are invariant and nonzero, hence depend on the direction of p; Q_p cannot equal c times either of them plus a form depending on p only through p^2 unless $c = 0$. A form depending only on p^2 can coincide with a gauge-fixed linearized Einstein–Hilbert operator, for example the Feynman–de Donder form $p^2(h_{\mu\nu}h_{\mu\nu} - \frac{1}{2}h^2)$, but without the gauge symmetry and its ghosts each nondegenerate direction of such a form, including directions of opposite sign, is a propagating component. This statement holds for a flat scaffold, a regulator and gauge fixing that preserve scaffold rotations, a vacuum and background invariant under them, and a composite-metric fluctuation that is a fixed linear function of the components of Φ. It does not exclude a composite-metric Einstein–Hilbert term in general: a regulator or kinetic metric built from $\Theta^{\mu\nu}$ in Presentation A (Remark 3.9), a vacuum or background that is not invariant under scaffold rotations, a curved scaffold, or a composite metric that is not a fixed linear image of Φ each relaxes one of these hypotheses and is not covered.

What the two preceding results imply for Problem F (conditional; not closure). In the class of the preceding paragraph, neither the displayed action nor its quantum corrections give the composite-metric fluctuation a linearized diffeomorphism invariance, an induced Einstein–Hilbert term, or the Fierz–Pauli block of hypothesis 5 of Theorem 7.3; the induced Fierz–Pauli term assumed in the quadratic tests of Section H.4 is likewise not generated in this way. These structures need an ingredient that ties derivative indices to tangent indices. The operator (8.3) on the emergent background, used in Theorem 8.2, is such an ingredient; it enters as Assumption 8.4, which is assumed, not derived. The one-loop Φ–Φ action of the displayed action on the scaffold contributes to the Λ_{UV}^2 and $\ln \Lambda_{\mathrm{UV}}$ coefficients A and B of (8.12) only through the curvature of $g^{(0)}$, not through that of the composite metric; the missing step of Problem F includes discharging Assumption 8.4, or replacing it by another derived ingredient. Not covered here: integration of the auxiliary A_μ in the first result, the hypotheses relaxed in the second, and Presentation A in general. No value or sign of G_{eff} follows. Problems B, C, D and F remain open.

Presentation A in the Poisson limit: the Θ-built metric and the connection sector (conditional; not closure). Let $\Theta^{\mu\nu}(x)$ be a nondegenerate Poisson bivector and $\rho = 1/|\operatorname{Pf}\Theta|$, so that $\partial_\mu(\rho\,\Theta^{\mu\nu}) = 0$. Keep the leading semiclassical order only: $[\cdot,\cdot] \to i\{\cdot,\cdot\}$ with $\{f,g\} = \Theta^{\mu\nu}\partial_\mu f\,\partial_\nu g$, $\mathrm{Tr} \to \int \mathrm{d}^4x\,\rho$, and matrix-coordinate indices

contracted with η. With $K^{\mu\nu} = \eta_{\alpha\beta}\Theta^{\alpha\mu}\Theta^{\beta\nu}$ and $g_\rho^{\mu\nu} = \rho K^{\mu\nu}$ as in Remark 3.9, let G be the metric with $G^{\mu\nu} = K^{\mu\nu}$; then $G = \rho\, g_\rho$, $|\det g_\rho| = 1$ and $\sqrt{|G|} = \rho^2$. Higher Moyal terms are dropped. They are not small at momenta of order Λ_{UV}, so the coefficients below are those of the Poisson-limit operators within the truncated expansion of Theorem 8.2. Signs and eigenvalues refer to the Euclidean section $\eta = \delta$; the algebraic identities do not depend on the signature.

Scalar sector. A real fluctuation whose kinetic term is the Poisson limit of $\mathrm{Tr}[\hat{X}_\mu, \varphi]^2$ has the operator $\mathcal{O}_s = -\rho^{-1}\partial_\mu(\rho K^{\mu\nu}\partial_\nu\,\cdot\,)$. On the metric G it has the form (8.3), with $D = \nabla + \omega$, $\omega_\mu = -\frac{1}{2}\partial_\mu \ln\rho$ and $E = -\frac{1}{2}\Box_G \ln\rho + \frac{1}{4}G^{\mu\nu}\partial_\mu \ln\rho\,\partial_\nu \ln\rho$. Since $G = \rho\, g_\rho$ in four dimensions, $R_G = \rho^{-1}\big[R - 3\Box_{g_\rho}\ln\rho - \frac{3}{2}|\nabla \ln\rho|^2_{g_\rho}\big]$ and $E = \rho^{-1}\big[-\frac{1}{2}\Box_{g_\rho}\ln\rho - \frac{1}{4}|\nabla\ln\rho|^2_{g_\rho}\big]$, so that

$$\sqrt{|G|}\Big(\tfrac{1}{6}R_G - E\Big) = \tfrac{1}{6}\,\rho\, R[g_\rho]$$

pointwise. For a constant Λ_{UV}, one such component contributes $-\frac{\Lambda_{\mathrm{UV}}^2}{192\pi^2}\int \mathrm{d}^4x\,\rho\, R[g_\rho]$ to (8.8): the coefficient of Theorem 8.2, multiplying the curvature of the Θ-built metric g_ρ with a factor ρ. The component is minimally coupled to g_ρ.

Frame. Let $\omega = \Theta^{-1}$, $\omega_{\mu B}\Theta^{B\nu} = \delta_\mu{}^\nu$, and $e^A{}_\mu = \eta^{AB}\omega_{\mu B}$. Then $G_{\mu\nu} = \eta_{AB}e^A{}_\mu e^B{}_\nu$, which is the form (4.9) of Definition 4.3 for a coframe with $\eta_{AB}e^B{}_\mu$ antisymmetric in (A,μ). At the sample points tested the map $\Theta \mapsto G$ has rank six, so that the Θ-built metrics form a six-parameter subfamily of the metrics, and no pure-gauge perturbation $p_{(\mu}\xi_{\nu)}$ of G lies in its tangent space. If $\langle\Phi\rangle/v$ of (4.7) were identified with this coframe, the operator above would be that of Assumption 8.4 for the metric G. This identification is not derived.

Connection sector. In background gauge with gauge-fixing term $\frac{1}{2}\rho\,(D_\mu a^\mu)^2$, $D^\mu = \{x^\mu, \cdot\}$, the part of (3.4) quadratic in the matrix gauge potential a^μ is, up to a constant factor, $\int \mathrm{d}^4x\,\rho\,[K^{\rho\sigma}\partial_\rho a_\nu \partial_\sigma a^\nu - 2a^\mu\{\Theta_{\mu\nu}, a^\nu\}]$, the term linear in a^μ not being assumed to vanish. The ghost operator is $D_\mu D^\mu = -\mathcal{O}_s$, of weight -2. The operator $\mathcal{O}_v a^\mu = \mathcal{O}_s a^\mu - 2\eta^{\mu\alpha}\{\Theta_{\alpha\nu}, a^\nu\}$ has the form (8.3) on G with $\omega_\sigma = -\frac{1}{2}\partial_\sigma \ln\rho\,\mathbb{I} + \mathcal{B}_\sigma$, $(\mathcal{B}^\sigma)^\mu{}_\nu = \eta^{\mu\alpha}\Theta^{\rho\sigma}\partial_\rho\Theta_{\alpha\nu}$, and $E = \nabla_\sigma\omega^\sigma + \omega_\sigma\omega^\sigma$. Then

$$\sqrt{|G|}\,\mathrm{tr}\Big(\tfrac{1}{6}R_G - E\Big) = \tfrac{4}{6}\,\rho\, R[g_\rho] + \rho^2 X, \qquad X = \eta^{\rho\lambda}\eta_{\mu\alpha}\eta_{\nu\beta}\,\partial_\rho\Theta^{\mu\nu}\,\partial_\lambda\Theta^{\alpha\beta},$$

and with the ghosts the connection sector contributes $\frac{2}{6}\rho R[g_\rho] + \rho^2 X$. The derivative index of X is contracted with η and not with G. For $\Theta^{01} = \vartheta_1(x^0, x^1)$, $\Theta^{23} = \vartheta_2(x^2, x^3)$ one finds $X = 2K^{\mu\nu}\partial_\mu \ln\rho\,\partial_\nu\ln\rho$; for a generic Poisson bivector X differs from this one expression.

Quadratic form about constant $\bar{\Theta}$. Let $\Theta = \bar{\Theta} + \epsilon T$, $T^{\mu\nu} = \bar{\Theta}^{\mu\rho}\partial_\rho a^\nu - \bar{\Theta}^{\nu\rho}\partial_\rho a^\mu$, $a^\mu = u^\mu\cos(p\cdot x)$. With $q^\mu = \bar{\Theta}^{\mu\rho}p_\rho$, the phase-averaged second-order terms of

$\rho R[g_\rho]$ and $\rho^2 X$ are quadratic forms on the three-dimensional quotient by $u \propto q$. Relative to the tree-level kernel $M = q^2\eta - qq$ of Remark 3.8, with $p_\eta^2 = \eta^{\mu\nu}p_\mu p_\nu$ and $\bar\rho = 1/|\operatorname{Pf}\bar\Theta|$, their Euclidean eigenvalues are $\bar\rho^2 p_\eta^2$ (three modes) for $\rho^2 X$, and $-\frac14\bar\rho^2 p_\eta^2$ (two modes with $\delta\rho = 0$, for which the linearized Einstein–Hilbert form of δg_ρ equals that of its transverse-traceless part with respect to p and $\bar g_\rho$) and $-\frac34\bar\rho^2 p_\eta^2$ (one mode with $\delta\rho \neq 0$) for $\rho R[g_\rho]$. Let $N_{\rm tot} = N_s + 2$, where N_s counts, with statistical sign, the real components whose fluctuation operator is $\mathcal{O}_s$ with the same cutoff; it is an input, not a derived count. In the Euclidean section the $\Lambda_{\rm UV}^2$ quadratic form is $-\frac{\Lambda_{\rm UV}^2}{32\pi^2}\bar\rho^2 p_\eta^2\,(1 - N_{\rm tot}/24)\,M$ on the two transverse-traceless-type modes and $(1 - N_{\rm tot}/8)\,M$ on the third. The net coefficient on the two transverse-traceless-type modes has the sign of a positive induced Einstein–Hilbert term only if $N_{\rm tot} > 24$; the Einstein–Hilbert-type piece itself always has that sign. For the connection sector alone, $\rho^2 X$ has the opposite sign and dominates. This is a statement about the Maxwell-kernel coefficient on these modes. Both pieces are multiples of M by functions of p_η^2, so they do not move the cone $q^2 = 0$, and about a constant background they do not fix $G_{\rm eff}$.

Beyond quadratic order. Nondegenerate Poisson bivectors near $\bar\Theta$ are $\Theta = (\psi^*\bar\omega)^{-1}$, $\bar\omega = \bar\Theta^{-1}$, $\psi = x + w(x)$. Modulo total derivatives, the scalar densities of weight ρ with two derivatives built covariantly from g_ρ and ρ are the combinations of $\rho R[g_\rho]$ and $\rho\, g_\rho^{\mu\nu}\partial_\mu \ln\rho\,\partial_\nu\ln\rho$. At quadratic order about constant $\bar\Theta$, $\rho^2 X$ coincides with $-4\rho R[g_\rho] - 4\rho\, g_\rho^{\mu\nu}\partial_\mu\ln\rho\,\partial_\nu\ln\rho$ for the random backgrounds and both signatures sampled. In a numerical test (double precision, w a sum of three plane waves with $\sum p_j = 0$, forty random samples) the cubic coefficient of $\rho^2 X$ is not of the form $c_1\rho R[g_\rho] + c_2\rho\, g_\rho^{\mu\nu}\partial_\mu\ln\rho\,\partial_\nu\ln\rho$ for any constants (relative residual about 0.3 for the best fit). So, on this sample, $\rho^2 X$ is not a combination of these two covariant densities modulo total derivatives, and the quadratic agreement does not extend. Rewritings that use further structure, such as the constant η, are not tested.

If $\Lambda_{\rm UV}^2$ is replaced by $\Lambda_{\rm UV}^2 f(\rho)$ (a local proper-time cutoff on the diagonal heat kernel), the coefficient of $R[g_\rho]$ becomes $f(\rho)\rho$; with $\hat g = f\rho\, g_\rho$, $\int f\rho R[g_\rho] = \int\sqrt{\hat g}\,[\hat R - \frac32\hat g^{\mu\nu}\partial_\mu\ln(f\rho)\partial_\nu\ln(f\rho)]$. The factor on the two transverse-traceless-type modes is unchanged, and for $f = \rho^s$ the third becomes $1 - N_{\rm tot}(2s+3)/24$.

This is a computation for Poisson-limit operators of a bosonic four-dimensional matrix gauge sector in a stated gauge, off shell or about a constant background. It does not identify G with the emergent metric of Chapter 4, does not treat Moyal corrections at the cutoff, fermionic or additional scalar fields, the mixing of Φ with a^μ, the locked-block diffeomorphism and local-Lorentz ghosts, or the relation of the induced term to the tree-level term of (3.4), and gives no value or sign of $G_{\rm eff}$. The kinetic metric and regulator here are built from Θ, and the map $\Theta \mapsto G$ is nonlinear, so the hypotheses of the scaffold-rotation statement above (a flat scaffold, a rotation-preserving regulator, an invariant vacuum, and a composite-metric fluctuation that is

a fixed linear image of Φ) do not apply; no linearized-diffeomorphism invariance is claimed. Problems B, C, D and F remain open.

Presentation A in the Poisson limit: what four matrix coordinates can supply (conditional; not closure). Keep the setting of the preceding paragraph: four matrix coordinates, the Poisson limit, matrix indices contracted with $\eta = \mathrm{diag}(-1,1,1,1)$, a real nondegenerate Θ, $K^{\mu\nu} = \eta_{\alpha\beta}\Theta^{\alpha\mu}\Theta^{\beta\nu}$, $G = K^{-1}$, $\rho = 1/|\operatorname{Pf}\Theta|$, $g_\rho^{\mu\nu} = \rho K^{\mu\nu}$. Let $N = \Theta\eta \in \mathfrak{so}(3,1)$ with electric and magnetic parts $\mathbf{E}, \mathbf{B}$, so $\operatorname{Pf}\Theta = \pm\mathbf{E}\cdot\mathbf{B}$. For vectors a, b write $a \circ b := ab^T + ba^T$.

Image of $\Theta \mapsto G$. The characteristic polynomial of N is $z^4 + (\mathbf{B}^2 - \mathbf{E}^2)z^2 - (\mathbf{E}\cdot\mathbf{B})^2$, so $K\eta = -N^2$ has two doubly degenerate eigenvalues and $(K\eta)^2 - \frac{1}{2}\operatorname{tr}(K\eta)\, K\eta - (\operatorname{Pf}\Theta)^2\,\mathbb{I} = 0$. Hence the six-dimensional image has codimension four in the ten-dimensional space of metrics and lies in the set cut out by this relation. Decomposing $N = (n_+, n_-)$ in $\mathfrak{sl}_2 \oplus \mathfrak{sl}_2$, $N^2 = (\nu_+ + \nu_-)\mathbb{I} + 2\, n_+ \otimes n_-$, and a kernel element of the differential of $\Theta \mapsto K$ must satisfy $\delta n_\pm = \mp\lambda n_\pm$ and $\lambda(\nu_+ - \nu_-) = 0$; since $\operatorname{Im}\nu_+ = 0$ only if $\mathbf{E}\cdot\mathbf{B} = 0$, the map $\Theta \mapsto G$ has rank six at every Θ with $\operatorname{Pf}\Theta \neq 0$. The differential of $\Theta \mapsto K$ has rank five when $\Theta \neq 0$ and $\operatorname{Pf}\Theta = 0$ (symbolic and exact-rational checks).

Linear response. Let $\Theta = \bar\Theta + \epsilon T$, $T^{\mu\nu} = \bar\Theta^{\mu\rho}\partial_\rho a^\nu - \bar\Theta^{\nu\rho}\partial_\rho a^\mu$, $a^\mu = u^\mu \cos(p\cdot x)$, $q = \bar\Theta p$. The amplitude of $\sin(p\cdot x)$ in δG is $(\bar G u)\circ p + n \circ m$, with $n = \bar G\bar\Theta p$, $m = \bar G\bar\Theta\eta u$, and $\delta \ln\rho = p\cdot u$. For $u \propto q$ one has $T = 0$ (the matrix gauge). The first term is a linearised diffeomorphism. Since n and p are covectors, orthogonality is taken with the metric $\bar K$: $n^T\bar K p = 0$, while n is not parallel to p for generic p, so the second term is not a diffeomorphism. The images of $u \mapsto (\delta G, \delta\rho)$ and $u \mapsto (\delta g_\rho, \delta\rho)$ are three-dimensional. For generic p the linearised Einstein operator about the constant background has rank three on this image, and its image lies in the six-dimensional space of symmetric tensors with $p^\mu T_{\mu\nu} = 0$ ("conserved"; exact-rational check for G and for g_ρ). Hence the tensor equation $\mathcal{E}[h] = T$, and any linear local relation between $(\delta G, \delta\rho)$ and T, can hold only for T in a subspace of dimension at most three of the six-dimensional space of conserved sources; varying the four functions a^μ gives only the projection of the equation onto that image. This does not exclude an additional conserved stress that absorbs the codimension; such a stress would have to be derived. As a counting argument (not a derived result), four coordinate functions minus one matrix gauge function leave three gauge-invariant functions, against $10 - 4 + 1 = 7$ for a diffeomorphism-invariant theory of a metric and one scalar. The tree-level quadratic action of Remark 3.8 is proportional to $\int \rho\, \eta_{\mu\alpha}\eta_{\nu\beta}T^{\mu\nu}T^{\alpha\beta}$, algebraic in $(\delta G, \delta\rho)$; every term with derivatives of the metric perturbation is a loop term. Whether the algebraic term is cancelled is the content of Assumption 6.1 and is not decided here.

Metric seen by the operators. Suppose, as an assumed dictionary not contained in the four-matrix class, that there are $D \geq 4$ coordinate matrices X^I with a Lorentzian

η_{IJ}, and put $\Phi^I{}_\mu := \partial_\mu x^I$, $g := \Phi^T \eta \Phi$. Then $K = \theta^T g\, \theta$ exactly; for $D = 4$, $x^I = y^I$ and $g = \eta$. The relation $K = c\, g^{-1}$, $c \neq 0$, would give $N^2 = -c$ with $N = \theta g \in \mathfrak{so}(g)$, whose characteristic polynomial has constant term $-(\mathbf{E} \cdot \mathbf{B})^2 \leq 0 < c^2$; $c = 0$ contradicts invertibility. So, pointwise for any real Lorentzian g, the operator metric $G = K^{-1}$ is not g; this extends Remark 3.7, and Assumption 8.4 read with the metric of that embedding reading is not discharged. The metric G itself has the form (4.9) for the coframe of the preceding paragraph. Let ℓ be the fixed length scale in Θ/ℓ^2, and $e = \eta^{-1}\hat{\omega}$ with $\hat{\omega} = (\Theta/\ell^2)^{-1}$. If e is identified with $\langle\Phi\rangle/v$ as in (4.7), then, with the internal index contracted with δ and the locked block real, $e^T e = \mathbb{I}$ on the vacuum manifold of (3.7) holds iff $\hat{\omega}^2 = -\mathbb{I}$, i.e. $J := \hat{\omega}$ is an orthogonal complex structure. Then $G = \mathbb{I} - 2nn^T$ with $n = -Je_0$, where e_0 is the first basis vector, so n is a spatial unit vector, and only two of the six directions of $\delta\Theta$ are tangent to that manifold.

This is a statement for four matrix coordinates, the Poisson limit, a constant η, linear order about constant $\bar{\Theta}$ and generic p; it does not apply to an independent field Φ of Presentation B, to the embedding statement above beyond four coordinates, or to Moyal corrections. It does not derive $\langle\Phi\rangle/v$ as a coframe and gives no value or sign of G_{eff}. Problem F, and Problems A, B, C and D, remain open.

More than four matrix coordinates in the Poisson limit: linear response (conditional; not closure). Keep Definition 3.4, the Poisson limit, a four-dimensional symplectic leaf, a real Lorentzian induced tensor g, $\omega = \theta^{-1}$, $K = \theta^T g\, \theta$ and $G = K^{-1}$.

Operator metric versus induced metric. The relation $G = c\, g$ is excluded for every $D \geq 4$ by the argument of the preceding paragraph, because only the real four-dimensional leaf and the real Lorentzian g enter it; for a Euclidean induced g an orthogonal complex structure gives $G \propto g$, but then G is Euclidean.

Linear response. In a chart with constant θ, let $\delta x^I = A^I e^{ip\cdot y}$, $A = \xi^\mu \partial_\mu x + \phi_i n^i$ with n^i a normal frame and $\mathrm{II}^i_{\mu\nu} = n^i \cdot \partial_\mu \partial_\nu x$; a factor i is absorbed in ξ. To first order $\delta g = p \circ (g\xi) - 2\phi_i \mathrm{II}^i$ with $a \circ b = ab^T + ba^T$, and $\delta G = -G\, \theta^T \delta g\, \theta\, G$; the Hamiltonian choice $\xi \propto \theta^T p$ gives $\delta G = p \circ (G\xi)$, a linearised diffeomorphism of G. About a flat brane the transverse amplitudes do not enter at first order, so the rank of the linearised Einstein operator of G on the image is exactly three for every D, against six conserved sources $p^\mu T_{\mu\nu} = 0$. About a background with generic II^i the rank is at most $\min(6, D-1)$, because the image contains a gauge direction and lies in the space of conserved tensors; one exact-rational draw for each $4 \leq D \leq 11$ attains this bound, so it is the generic rank by lower semicontinuity, and the same holds when $K^{\mu\nu}\mathrm{II}^i_{\mu\nu} = 0$. All six sources are then reachable from $D = 7$. The analysis freezes g, θ and II at a point and does not impose the Gauss–Codazzi relations. Counting the density scalar as well, the number $D - 1$ of gauge-invariant functions must be at least 7, that is $D \geq 8$; this is a necessary count. No field equation of Einstein type for G is derived.

Content. The $D-4$ transverse coordinates are not covered by Assumption 5.1; they are additional infrared fields unless they are gapped, and if they are gapped and decouple the flat-brane statement above applies. One-loop IKKT calculations on a brane with compact fuzzy extra dimensions report an induced Einstein–Hilbert term with the Newton constant set by the Kaluza–Klein scale [21], and modified Einstein equations with extra tensors are reported in [25, 26]. A leaf of dimension greater than four is the hypothesis of the first statement above that such a construction would relax. These results have not been checked against the present operators.

This is a statement for D matrix coordinates, the Poisson limit, a four-dimensional symplectic leaf, a constant η_{IJ}, locally frozen background data and generic p. It does not identify G or g with the emergent metric of Chapter 4, does not identify $\Phi^I{}_A$ with a tangent-plane frame, does not derive a local Lorentz symmetry, a signature or an extra conserved stress, and gives no value or sign of G_{eff} in Equation (8.12). This changes no ledger label; Problem F remains open.

Leaf dimension above four: fibre averages, a product factor and a conjugate-coordinate class (conditional; not closure). Work in the Poisson limit with matrix coordinates x^I, $I = 0, \dots, D-1$, contracted with $\eta = \mathrm{diag}(-1, 1, \dots, 1)$, on a symplectic leaf of dimension $2m \geq 4$, and four base functions y^μ on the leaf. For a function $\phi(y^\mu)$ the kinetic tensor is $K^{\mu\nu} = \langle \eta_{IJ} V^{I\mu} V^{J\nu} \rangle$ with $V^{I\mu} = \{x^I, y^\mu\}$ and $\langle\cdot\rangle$ a positive average over the fibre; for a four-dimensional leaf it is $\theta^T g \theta$.

Matrices as base coordinates. Let $y^\mu = x^\mu$, $\mu = 0, \dots, 3$, so that $V^{I\mu} = \Theta^{I\mu}$ is real and antisymmetric. Then $K^{00} = \langle \sum_{I\geq 1} (\Theta^{I0})^2 \rangle \geq 0$ and $\sum_\mu K^{\mu\mu} = \langle \sum_{I\geq 4} (\Theta^{I0})^2 + \sum_{I\geq 1,\, k\geq 1} (\Theta^{Ik})^2 \rangle \geq 0$, whereas $K = c\,\eta^{-1}$ has $K^{00} = -c$ and $\sum_\mu K^{\mu\mu} = 2c$; hence $c = 0$. The case $K = 0$ forces $\Theta^{I\mu} = 0$ for $\mu \leq 3$, so the x^μ would be Casimir functions, contradicting nondegeneracy. Therefore $K \neq c\,\eta^{-1}$ for every $c \neq 0$, for every D, every compact fibre and every positive average: in this class, enlarging the leaf by a fibre does not remove the proportionality obstruction of the preceding paragraphs. For any nondegenerate real Lorentzian g_L on a leaf of dimension $2m \geq 4$, $G_L = (\theta^T g_L \theta)^{-1}$ is not proportional to g_L. Indeed $K = c\, g_L^{-1}$ with $c \neq 0$ gives $N^2 = -c\,\mathbb{I}$ for $N = \theta g_L$. For $c > 0$, $\det N = c^m > 0$, whereas $\det N = (\mathrm{Pf}\,\theta)^2 \det g_L < 0$. For $c < 0$, N is g_L-skew with $N^2 = |c|\,\mathbb{I}$, so its $\pm\sqrt{|c|}$ eigenspaces are totally null and have dimension at most one in Lorentzian signature, which forces $2m \leq 2$; for $2m = 2$ the relation can hold ($g_L = \mathrm{diag}(-1, 1)$, $\theta = 2\epsilon$ gives $K = -4 g_L^{-1}$). For a product leaf $M \times \mathcal{C}$ with block-diagonal θ and embedding, $V^{i\mu} = 0$ for the coordinates along $\mathcal{C}$, so K is the tensor of the M factor, and to first order the response of K to $\delta x^I = A^I(t) e^{ip\cdot y}$ is the four-dimensional one with amplitude $\langle A^\mu \rangle$; transverse amplitudes and non-constant fibre parts do not enter. The rank statements of the preceding paragraph therefore hold with D the number of coordinates that vary over M, and Kaluza–Klein modes enter only quadratically.

Conjugate coordinates. The proportionality is not excluded when the base functions are not among the matrix coordinates. Let M^{AB}, $A, B = 0, \dots, 5$, generate $\mathfrak{so}(4,2)$ with invariant form $\mathrm{diag}(-,+,+,+,+,-)$ on the coadjoint orbit $\mathbb{CP}^{1,2}$ (six-dimensional leaf), take the four matrix coordinates $T^a = M^{a4}/R$, $a = 0, \dots, 3$, and the base functions $y^\mu = M^{\mu 5}/R$. The functions y^μ are not among the T^a, and the leaf is not the four-dimensional leaf of Definition 3.4. The Lie–Poisson relation gives $\{T^a, y^\mu\} = \pm\eta^{a\mu} M^{45}/R^2$, because the form has $\eta^{45} = \eta^{a5} = \eta^{4\mu} = 0$, hence $K^{\mu\nu} = (M^{45}/R^2)^2 \eta^{\mu\nu}$ pointwise, with M^{45} a function of y^μ on the orbit (checked on an explicit realisation). Here $\eta^{\mu\nu}$ is the Cartesian form of the functions y^μ, not the pull-back of the matrix contraction η_{IJ}, and the matrix-induced tensor on the six-dimensional leaf has rank four. This K is conformally flat, so the exact class carries one scalar function and no metric waves; whether fluctuations of T^a that are not fibre-constant supply a sufficient number of gauge-invariant functions is not computed here, and no argument is given that the $y^\mu = M^{\mu 5}/R$ are the base functions of the reduced theory.

Heat kernel over a fuzzy factor. For the product leaf with a finite fuzzy factor, the scalar operator of the Poisson-limit paragraph is $\mathcal{O}_s \otimes 1 + 1 \otimes \Box_{\mathcal{C}}$, so each four-dimensional sector is replicated over the $\dim \mathrm{Mat}(\mathcal{H}_{\mathcal{C}})$ modes, the mode of mass m with the threshold factor $F(m^2/\Lambda_{\mathrm{UV}}^2) = \int_1^\infty u^{-2} e^{-m^2 u/\Lambda_{\mathrm{UV}}^2} \mathrm{d}u \in (0,1]$ multiplying the coefficient of Theorem 8.2. In the truncated expansion the curvature coefficient is thus of order Λ_{UV}^2 times the weighted number of light modes; it is not set by the Kaluza–Klein scale. The supersymmetric one-loop mechanism of [21], in which the corresponding terms cancel and the Newton constant is set by a finite mixed term, rests on field content that is not part of the present sector list.

This is a statement for the Poisson limit, the stated fibre classes and off-shell backgrounds; it does not identify G or $\eta_{\mu\nu}$ with the emergent metric of Chapter 4, gives no field equation, gap of the transverse scalars, G_{eff} value or sign, and does not compute the rank of the linearised Einstein operator in the conjugate-coordinate class.

Connection-sector term X as a torsion invariant (conditional; not closure). Take the setting of the Poisson-limit paragraph on Presentation A above: four matrix coordinates, a real nondegenerate Θ with Jacobi identity, $\eta = \mathrm{diag}(-1,1,1,1)$ or $\mathrm{diag}(1,1,1,1)$, $K^{\mu\nu} = \eta_{\alpha\beta}\Theta^{\alpha\mu}\Theta^{\beta\nu}$, $G = K^{-1}$, $\rho = 1/|\operatorname{Pf}\Theta|$, $g_\rho^{\mu\nu} = \rho K^{\mu\nu}$. Let $\omega = \Theta^{-1}$ (closed), $e^A{}_\mu = \eta^{AB}\omega_{\mu B}$ so that $G_{\mu\nu} = \eta_{AB} e^A e^B$, $T^A{}_{\mu\nu} = \partial_\mu e^A{}_\nu - \partial_\nu e^A{}_\mu$, $T_{\lambda\mu\nu} = \eta_{AB} e^A{}_\lambda T^B{}_{\mu\nu}$ and $S_{\lambda\mu\nu} = T_{[\lambda\mu\nu]}$. The matrix indices α, β, A, B are lowered and raised with η; coordinate indices, and $S_{\lambda\mu\nu}S^{\lambda\mu\nu}$, with G. The term X of the connection sector is $X = \eta^{\rho\lambda}\eta_{\mu\alpha}\eta_{\nu\beta}\partial_\rho\Theta^{\mu\nu}\partial_\lambda\Theta^{\alpha\beta}$. Since $\Theta^T\eta\,\Theta = G^{-1}$ and $\partial\Theta = -\Theta(\partial\omega)\Theta$, $X = G_{\mu\nu}T^{\alpha\beta\mu}T_{\alpha\beta}{}^\nu$ with $T^{\alpha\beta\mu} = \Theta^{\mu\rho}\partial_\rho\Theta^{\alpha\beta}$, and $X = T_{\lambda\mu\nu}T^{\lambda\mu\nu}$ because $d\omega = 0$ gives $T^A{}_{\mu\nu} = -\eta^{AB}\partial_B\omega_{\mu\nu}$. Moreover $\partial_\mu \ln\rho = G^{\lambda\nu}T_{\lambda\mu\nu}$, which follows from $d\omega = 0$ through $\partial_\sigma \ln \operatorname{Pf}\omega = -\Theta^{\nu\mu}\partial_\mu\omega_{\nu\sigma}$. With the identity for the Weitzenböck connection of the coframe e (in the present index convention, checked by direct computation)

$R[G] = -(\frac{1}{4}T_{\lambda\mu\nu}T^{\lambda\mu\nu} + \frac{1}{2}T_{\lambda\mu\nu}T^{\mu\lambda\nu} - V_\mu V^\mu) - 2\nabla_\mu V^\mu$, $V_\mu = G^{\lambda\nu}T_{\lambda\mu\nu}$, the relation $S_{\lambda\mu\nu}S^{\lambda\mu\nu} = \frac{1}{3}(T_{\lambda\mu\nu}T^{\lambda\mu\nu} - 2T_{\lambda\mu\nu}T^{\mu\lambda\nu})$, and the conformal transformation $G = \rho\, g_\rho$ with $\det g_\rho = 1$, one finds the exact pointwise identity

$$\rho^2 X = -2\rho R[g_\rho] - \rho\, g_\rho^{\mu\nu}\partial_\mu \ln\rho\, \partial_\nu \ln\rho + \tfrac{3}{2}\rho^2 S_{\lambda\mu\nu}S^{\lambda\mu\nu} + 2\partial_\mu(g_\rho^{\mu\nu}\partial_\nu\rho).$$

A rewriting using the coframe $e = \eta^{-1}\omega$ is therefore the squared axial torsion. The identity was also verified in exact rational arithmetic on finite-amplitude generic two-jets of a closed ω for both signatures. At quadratic order about constant $\bar{\Theta}$, for the perturbations of the Poisson class, the invariants $\rho R[g_\rho]$, $\rho\, g_\rho^{\mu\nu}\partial_\mu \ln\rho\, \partial_\nu \ln\rho$ and $\rho^2 S_{\lambda\mu\nu}S^{\lambda\mu\nu}$ are linearly dependent modulo total derivatives, so this decomposition does not alter the net coefficient $1 - N_{\mathrm{tot}}/24$ stated above.

This is a statement for the Poisson-limit operators of the preceding paragraphs. It does not identify G with the emergent metric of Chapter 4, does not treat Moyal corrections, fermions or other fields, and gives no value or sign of G_{eff}. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: first-order response and linearised Einstein rank (conditional; not closure). This paragraph supplements the preceding one, which left the rank of the linearised Einstein operator in this class uncomputed. Keep the conjugate-coordinate class of the preceding paragraphs in the Poisson limit: the leaf $\mathbb{CP}^{1,2}$ with the Lie–Poisson bracket $\{M^{AB}, M^{CD}\} = s(\eta^{AC}M^{BD} - \eta^{AD}M^{BC} - \eta^{BC}M^{AD} + \eta^{BD}M^{AC})$, $s = \pm 1$ a convention, the matrices $X^a = T^a + A^a$ with A^a arbitrary functions on the leaf, $\eta = \mathrm{diag}(-1,1,1,1)$, and the base functions y^μ held fixed. Put $m = M^{45}/R^2$ and $y^4 = M^{45}/R$. Then $\{T^a, y^\mu\} = s\,\eta^{a\mu}m$, $\{y^\mu, y^\rho\} = -s\,M^{\mu\rho}/R^2$, and with $A^\nu := A^a\delta_a{}^\nu$ the first-order change of $K^{\mu\nu} = \eta_{ab}\{X^a, y^\mu\}\{X^b, y^\nu\}$ is $\delta K^{\mu\nu} = s\,m\,(\{A^\nu, y^\mu\} + \{A^\mu, y^\nu\})$. Additional matrices X^i with $\{X^i, y^\mu\} = 0$ in the background enter K only through $\eta_{ij}\{X^i, y^\mu\}\{X^j, y^\nu\}$, which is quadratic in their fluctuations.

Fibre structure. On an explicit realisation one finds $M^{ab}y_b = 0$ for $a, b = 0, \ldots, 4$ (indices lowered with $\mathrm{diag}(-,+,+,+,+)$), hence $T^\mu = -M^{\mu\sigma}y_\sigma/y^4$ and $y_\mu T^\mu = 0$. Over a point of the base with $y^4 \neq 0$ the fibre of y^a is a two-sphere on which the six functions $M^{\mu\rho}$ have zero average, are first spherical harmonics and span a three-dimensional space W_y, the projection to the indices $0, \ldots, 3$ of the self-dual two-forms of the Euclidean space $y^\perp \subset \mathbb{R}^{1,4}$; the four functions T^μ span the same space of functions. Consequently the first-harmonic part of A^ν consists of twelve amplitudes $b^{\nu;\mu\rho}(y) \in W_y$ and coincides with the frame-type fluctuations $A^\nu = \kappa^\nu{}_\sigma(y)T^\sigma$, defined modulo $\kappa^\nu{}_\sigma \to \kappa^\nu{}_\sigma + k^\nu y_\sigma$.

Response. Let $\mathrm{d}\mu$ be the fibre measure induced by the Liouville measure on a sheet $y^4 > 0$ (the other sheet is not included) and $B^{\nu;\mu\rho} = \int \mathrm{d}\mu\, A^\nu M^{\mu\rho}$. Liouville invariance

of the bracket gives $\int \mathrm{d}\mu\, \delta K^{\mu\nu} = (m/R^2)\, \partial_\rho (B^{\nu;\mu\rho} + B^{\mu;\nu\rho})$, so the fibre-integrated variation $\delta\mathcal{K}^{\mu\nu}$ satisfies $\partial_\mu \partial_\nu (\delta\mathcal{K}^{\mu\nu}/m) = 0$ for every A. Only the first-harmonic part of A contributes: fibre-constant fluctuations, the ordinary gauge fields, and harmonics of order two and higher give no linear response. Take a plane wave $e^{iq\cdot y}$, frozen coefficients at a base point, $|q|\ell \gg 1$ with $\ell^2 = -y_a y^a$, and $q^2 \neq 0$. To leading order $\delta G_{\mu\nu} \propto \eta_{\mu\alpha}\eta_{\nu\beta} h^{\alpha\beta}$ with $h^{\mu\nu} = q_\rho (b^{\nu;\mu\rho} + b^{\mu;\nu\rho})$. Antisymmetry in (μ, ρ) gives $q_\mu q_\nu h^{\mu\nu} = 0$, so the image has dimension at most nine. The rank is a consequence of the image being the whole hyperplane $q_\mu q_\nu h^{\mu\nu} = 0$ and does not depend on the particular three-dimensional space W_y; this hyperplane is not invariant under longitudinal diffeomorphisms, so the gauge group of the class is that of divergence-free diffeomorphisms. In exact rational arithmetic, at base points $(y^0, y^4) = (5, 3), (13, 5), (5, 4), (25, 7)$ in units $R = 1$ and for generic rational q, the image has dimension nine, the linearised Einstein operator of η has rank six on it, its image is the whole six-dimensional space of conserved tensors $q^\mu T_{\mu\nu} = 0$, and its kernel on the image is three-dimensional. Six is the largest possible rank, so it is the generic rank by lower semicontinuity. For null q the rank is three, and at $y^4 = 0$ the image has dimension seven and the rank is five. For $A^\nu = \{\Lambda, T^\nu\}$ the leading image is $h = q \circ (\Lambda q)$ with $\Lambda^{\mu\rho} \in W_y$, the three-dimensional space of transverse linearised diffeomorphisms $\xi \cdot q = 0$, which is the kernel found above; the longitudinal diffeomorphism is neither a matrix gauge transformation nor in the image. The twelve amplitudes thus split into three that leave G unchanged, three gauge directions and six gauge-invariant functions, against six conserved sources and against $10 - 4 + 1 = 7$ for a metric and a density in a diffeomorphism-invariant theory; the density is fixed by the Liouville measure and the image obeys the one relation above. A change $y^\mu \to y^\mu + \delta y^\mu$ of the base functions acts at leading order as a linearised diffeomorphism with parameter $\int \mathrm{d}\mu\, \delta y^\mu$ on the fibre-integrated tensor and the fibre volume, and so adds no gauge-invariant function. The algebraic parts of the response and of the gauge variation are of relative order $1/(|q|\ell)$ and are not computed, so the gauge invariance of $\mathcal{E}[h] = T$ holds at the leading symbol only.

This is a kinematic statement for the Poisson limit, the stated class, linear order, frozen coefficients and the fibre-integrated tensor; fibre-non-constant parts of δK are not treated. It does not identify G with the emergent metric of Chapter 4, derives no field equation, and does not show that the first-harmonic amplitudes are light, positive-norm or decoupled. A Yang–Mills-type tree-level quadratic action, analogous to that of Remark 3.8 and built from first derivatives of A, gives equations of second order in these amplitudes (no such action is computed in this class), whereas $\mathcal{E}[h]$ is of third order; an equation $\mathcal{E}[h] = T$ would therefore require an induced term or another structure, which is not decided here. Fibre harmonics of the fluctuation matrices form the higher-spin tower described in [23, 27]; the relation of the amplitudes $b^{\nu;\mu\rho}$ to that tower is not examined here.

Why $y^\mu = M^{\mu 5}/R$ in this class (partial; conditional; not closure). The only SO(3, 1) vector generators of $\mathfrak{so}(4,2)$ are $T^\mu = M^{\mu 4}/R$ and $y^\mu = M^{\mu 5}/R$, so take $u^\mu = \alpha T^\mu + \beta y^\mu$. Let $\Box = \eta_{ab}\{T^a, \{T^b, \cdot\}\}$ be the Poisson-limit scalar operator. A reduction to an operator on functions of four variables without a fibre average requires that $\Box$ maps functions of u to functions of u, that is, that $K_u^{\mu\nu} = \eta_{ab}\{T^a, u^\mu\}\{T^b, u^\nu\}$ and $\Box u^\mu$ are functions of u. The Lie–Poisson relations give $\Box y^\mu = -y^\mu/R^2$, $\Box T^\mu = -3T^\mu/R^2$ and $K_u^{\mu\nu} = \alpha^2\eta_{ab}M^{a\mu}M^{b\nu}/R^4 + \beta^2 m^2\eta^{\mu\nu}$, the cross term cancelling by antisymmetry of $M^{\mu\nu}$. For $\alpha = 0$ both conditions hold. For $\alpha \neq 0$ the term $M^{a\mu}M_a{}^\nu$ depends on the position on the fibre, and for $\beta \neq 0$ also $\Box u^\mu = -(3\alpha T^\mu + \beta y^\mu)/R^2$ is not a function of u (a gradient-rank test at random points of the leaf, using these expressions, confirms both failures; for $\alpha = \beta$ the u^μ are not independent and the test does not apply). Within this family the base functions are therefore y^μ up to normalisation. This presupposes the orbit $\mathbb{CP}^{1,2}$, the matrices $T^a = M^{a4}/R$ and the Lorentz-vector family; it is a property of the operator $\Box$, not a consequence of the matrix equations of motion, and it does not show that the physical reduction is the one by y^μ.

This is a statement for the Poisson limit, the class above, the Lorentz-vector family and generators only; it gives no field equation, no value or sign of G_{eff}, and does not derive the background, the signature or a coframe from Φ; in particular it does not bear on Assumption 6.1. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: tree-level quadratic form of the level-one amplitudes (conditional; not closure). This paragraph supplements the preceding ones, in which the tree-level action of the amplitudes $b^{\nu;\mu\rho}$ was not computed. Keep the Poisson limit, the leaf $\mathbb{CP}^{1,2}$, $X^a = T^a + A^a$, the base functions y^μ and the notation of the preceding paragraphs, in units $R = 1$, and write $D^a = \{T^a, \cdot\}$, $\mathcal{J}_{ab} = \{M_{ab}, \cdot\}$, $\Box = D_b D^b$. The quartic term of (3.4) with $C_{\mu\nu\lambda} = 0$, read with $[\cdot,\cdot] \to i\{\cdot,\cdot\}$ and $\mathrm{Tr} \to \int \mathrm{vol}$, is $S = (4g_{\mathrm{YM}}^2)^{-1}\int \mathrm{vol}\,\{X^a, X^b\}\{X_a, X_b\}$, with field equation $\{X^b, \{X_b, X_a\}\} = 0$.

Background. The Lie–Poisson relations give $\{T_a, \{T^a, T^b\}\} = -3T^b$, an identity of the polynomial Poisson algebra of $\mathfrak{so}(4,2)$ that does not depend on the sign s. Hence T^a is not stationary for the quartic term alone. It is stationary for $S + (2g_{\mathrm{YM}}^2)^{-1}\mu^2\int \mathrm{vol}\, X_a X^a$, whose field equation is $\{X^b, \{X_b, X_a\}\} = \mu^2 X_a$, exactly when $\mu^2 = -3$. A constant $C_{\mu\nu\lambda}$ cannot take the place of this term: it would add $C_{abc}M^{bc}$ to the field equation, and these functions are linearly independent of $T_a = M_{a4}$ on the orbit. The bilinear term is not part of (3.4); it is an input of this class with a coefficient fixed by stationarity, and a mass term is reported for the covariant backgrounds of [27].

Quadratic form. With this term, $S^{(2)} = (2g_{\mathrm{YM}}^2)^{-1}\int \mathrm{vol}\, A^a L(A)_a$ with $L(A)_a = -\Box A_a + D_a(D_b A^b) - 2s\,\mathcal{J}_{ab}A^b + \mu^2 A_a$, an identity verified in the polynomial Poisson algebra for both signs s. Gauge covariance of the field equation about a stationary

point gives $L(\{\Lambda, T\}) = 0$ for every function Λ, and the same check fails for $\mu^2 \neq -3$. The quadratic form is therefore exactly invariant under $\delta A^a = \{\Lambda, T^a\}$, not only at a leading symbol.

Leading symbol. Since $D^a y^\mu = s\, m\, \eta^{a\mu}$ with $m = M^{45}$ a function of y alone, the only part of L with two derivatives along y is $-\Box A_a + D_a(D_b A^b)$. The operator $\mathcal{J}_{ab}$ acts on y^μ as a Lorentz rotation with a coefficient linear in y, and the fibre derivatives of A and the derivatives of m enter with coefficients of relative order $1/(|q|\ell)$. For $A^a = c^a(y)F$ with F a fixed fibre function and $|q|\ell \gg 1$, then, $L(A)_a = -m^2(\partial^2 c_a - \partial_a \partial_b c^b)F$ plus terms with at most one derivative of c, so that

$$S^{(2)} \simeq \frac{1}{4g_{\mathrm{YM}}^2} \int \mathrm{vol}\; m^2\, \mathcal{F}_{ab}\mathcal{F}^{ab}, \qquad \mathcal{F}_{ab} = \partial_a A_b - \partial_b A_a,$$

with ∂_a the derivative along y at fixed fibre argument. At this order each fibre function is an independent Maxwell field with the same coefficient, and there is no coupling between fibre harmonics. On the explicit realisation of the preceding paragraphs, evaluating L on $A^a = c^a \mathrm{e}^{i\lambda q\cdot y} M^{12}$ reproduces $\lambda^2 m^2 M^{12} \mathrm{e}^{i\lambda q\cdot y}(q^2 c_a - q_a q\cdot c)$ with a relative remainder that falls as $1/\lambda$.

Level-one sector. Write $M^{\mu\rho} = \sum_i e^i S_i^{\mu\rho}$ with S_i an orthogonal, equal-norm basis of W_y and e^i the three first spherical harmonics on the fibre; by the SU(2) invariance of the fibre measure, $\int \mathrm{d}\mu\, e^i e^j = \varkappa(y)\, \delta^{ij}$ with $\varkappa > 0$. For $A^\nu = c_i^\nu(y) e^i$ one has $b^{\nu;\mu\rho} = \varkappa\, c_i^\nu S_i^{\mu\rho}$, and the twelve amplitudes are three vector fields c_i with the action $(m^2\varkappa/4g_{\mathrm{YM}}^2)\sum_i \int \mathrm{d}^4 y\, \mathcal{F}_{i,ab}\mathcal{F}_i^{ab}$. The kinetic matrix is $\delta_{ij}(q^2\eta_{ab} - q_a q_b)$, definite in the index i; the leading symbol has no mass term and no coupling to the other harmonics. The overall sign is that of the quartic term as displayed, common to all fibre harmonics including the fibre-constant gauge field, so no relative ghost arises at this order. It is opposite to the sign of the Lorentzian Maxwell action $-\frac{1}{4}F_{ab}F^{ab}$ in the same η, and whether it is the healthy sign depends on the convention for the weight, which (3.4) does not fix. The Maxwell transformation $\delta c_i^\nu = \partial^\nu \lambda_i$ is the leading form of $\delta A^\nu = \{\Lambda, T^\nu\}$ with $\Lambda = \lambda_i e^i$, under which h changes by a divergence-free linearised diffeomorphism. Masses, the connection induced by the y-dependence of S_i and m, and the mixing of harmonics enter at relative orders $1/(|q|\ell)$ and $1/(|q|\ell)^2$, which frozen coefficients do not determine; they are not computed.

This is a statement for the Poisson limit, the stated class with the bilinear term, and the leading symbol. It does not determine masses, the signs of subleading terms or decoupling beyond leading order, and it does not identify the fibre-averaged tensor with the emergent metric of Chapter 4.

Conjugate-coordinate class: what the tree-level equations of the level-one amplitudes imply for $\mathcal{E}[h]$ (conditional; bounded no-go). Keep the setting of the preceding paragraph. Let $\mathcal{E}$ be the linearised Einstein operator of η,

$h^{\mu\nu} = q_\rho(b^{\nu;\mu\rho} + b^{\mu;\nu\rho})$ and $s_i^\mu = S_i^{\mu\rho}q_\rho$, so that $h^{\mu\nu} = \varkappa\sum_i(s_i^\mu c_i^\nu + s_i^\nu c_i^\mu)$. The order count of the preceding paragraphs becomes precise at the leading symbol. In exact rational arithmetic at the base points $(y^0, y^4) = (5,3), (13,5), (5,4)$ in units $R = 1$ the following hold.

Free equations. The equations $q_a\mathcal{F}_i^{ab} = 0$ make c_i pure gauge for $q^2 \neq 0$, so h is then a divergence-free linearised diffeomorphism and $\mathcal{E}[h] = 0$. For null q in Lorenz gauge there are six physical polarisations (three Maxwell fields): three give $h = 0$, and the other three give h modulo diffeomorphisms spanned by the two transverse-traceless polarisations and one further polarisation with nonzero trace (base points $(5,3)$ and $(13,5)$). For these solutions $\mathcal{E}[h]_{\mu\nu} = \frac{1}{2}q_\mu q_\nu h^\alpha{}_\alpha$, which vanishes on the transverse-traceless polarisations and not on the polarisation with nonzero trace; the longitudinal diffeomorphism is not a symmetry of the class, so this trace is not removed by gauge.

No local relation. The map $c \mapsto \mathcal{E}[h(c)]$ is cubic in q, and the Maxwell operator $c \mapsto q^2c^b - q^b\, q\cdot c$ is quadratic. No matrix $P(q)$ linear in q satisfies $\mathcal{E}[h(c)] = P(q)\cdot(\text{Maxwell operator on } c)$, and none of the ten components of $\mathcal{E}[h(c)]$ is divisible by q^2. Hence no local equation $\mathcal{E}[h] = P(\partial)J$ follows from the second-order equations $\partial_a\mathcal{F}_i^{ab} = J_i^b$ for a source J_i.

Response to a source. Take as input a conserved source, $q_\mu T^{\mu\nu} = 0$, coupled to the amplitudes through h, that is $J_{i\nu} \propto s_i^\mu T_{\mu\nu}$ (gauge invariance of the coupling requires the conservation). Then c_i is determined up to gauge, $c_i^\nu \propto J_i^\nu/q^2$, and

$$h^{\mu\nu} = \frac{N}{q^2}(Q^{\mu\alpha}T_\alpha{}^\nu + Q^{\nu\alpha}T_\alpha{}^\mu), \qquad Q^{\mu\alpha} = \sum_i s_i^\mu s_i^\alpha = q_\perp^2 P_\perp^{\mu\alpha} - q_\perp^\mu q_\perp^\alpha,$$

with N a nonzero constant fixed by the couplings, $P_\perp$ the projector on $y^\perp$ and $q_\perp = P_\perp q$, both restricted to the indices $0,\dots,3$. Modulo $q\circ\xi$ this is $h^{\mu\nu} \simeq N\big[2(1 + q_\parallel^2/q^2)T^{\mu\nu} + \hat{y}^\mu\tau^\nu + \tau^\mu\hat{y}^\nu\big]$ with $\hat{y} = y/\ell$, $q_\parallel = q_\mu\hat{y}^\mu$ and $\tau^\nu = \hat{y}_\alpha T^{\alpha\nu}$. The response is homogeneous of degree zero in q: h is algebraic in T, local for $q_\parallel = 0$, in particular for a static source at a base point with $y^i = 0$, and otherwise carries the factor $q_\parallel^2/q^2$. Since $\mathcal{E}$ is of degree two, $\mathcal{E}[h]$ is homogeneous of degree two in q and is nonzero whenever h is not a linearised diffeomorphism; it therefore cannot equal a constant multiple of T (the sampled cases also exclude q^2T). A relation of second order in c is zeroth order in h once the common derivative is removed, whereas $\mathcal{E}[h] = T$ is second order in h.

Induced term. Suppose a term $(16\pi G_{\text{eff}})^{-1}\int\sqrt{-G}\,R$, reported to arise at one loop in [21] and not computed here, is added as an input. Its quadratic form in h has degree two in q and the tree-level form has degree zero, so the tree-level form dominates at small q, and an Einstein-type response $h \sim G_{\text{eff}}T/q^2$ can appear only above the scale set by the ratio of the two coefficients, provided that scale lies within $|q|\ell \gg 1$; otherwise the induced form is subdominant throughout the regime of the class. Removing the tree-level term would require a loop contribution to the coefficient

of $\sum_i \mathcal{F}_i^2$ that cancels it; this is not examined.

This is a statement for the Poisson limit, the stated class, the leading symbol, tree level, the level-one sector coupled to a conserved source through h, and the sheet $y^4 > 0$. It does not exclude an Einstein-type equation from the induced term, from subleading orders in $1/(|q|\ell)$, from other sectors or from the entanglement route of Assumption 6.1, and it derives no value or sign of G_{eff}. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: matrix Noether identity and the divergence of the stress (conditional; not closure). Work in the Poisson limit with $F^{ab} = \{X^a, X^b\}$ and $E_c = -\{X^b, \{X_b, X_c\}\} + \mu^2 X_c$, the field equation of $(4g_{\text{YM}}^2)^{-1} \int \text{vol}\,(F_{ab}F^{ab} + 2\mu^2 X_a X^a)$, for arbitrary functions X^a on the leaf. The Jacobi identity and the Leibniz rule give $\{X^a, E_a\} = 0$ identically. The tensor $\mathbb{T}^{ab} = F^{ac}F^b{}_c + \mu^2 X^a X^b$, which is the variation of the action with respect to the contraction form, satisfies $\{X_a, \mathbb{T}^{ab}\} = -F^b{}_c E^c + \frac{1}{4}\{X^b, F_{cd}F^{cd}\}$; its divergence therefore vanishes on the field equation up to a gradient, and $\mathbb{T}^{ab} - \frac{1}{4}\eta^{ab}F_{cd}F^{cd}$ is divergence-free on the field equation exactly, the subtracted term being $\frac{1}{4}\eta^{ab}F_{cd}F^{cd}$ and not a trace of $\mathbb{T}^{ab}$. About $X^a = T^a$ the linear form obeys $\{T^a, L(A)_a\} = -(3+\mu^2)\{A^a, T_a\}$, which vanishes exactly for $\mu^2 = -3$. For the massless model the matrix-level identity $[X_a, \mathbb{T}^{ab}] \propto$ equation of motion is standard; the Poisson-limit form $\{X^a, E_a\} = 0$ and the divergence identity with the μ^2 term are recorded here as exact identities in the polynomial Poisson algebra, and were verified there for both signs of the bracket. The tensor $\mathbb{T}^{ab}$ carries matrix indices and $\{X_a, \cdot\}$ is a frame derivative, not $\nabla_\mu \mathbb{T}^{\mu\nu}$; it differs from the coordinate divergence of the fibre-averaged tensor by terms containing derivatives of the frame coefficients, of relative order $1/(|q|\ell)$, which are not computed. The paragraph does not identify $\mathbb{T}^{ab}$ with the source of an Einstein-type equation.

Conjugate-coordinate class: extra stress and crossover response at the leading symbol (conditional; not closure). Keep the setting and notation of the preceding paragraphs at the base points $y^i = 0$, with $s_i^\mu = S_i^{\mu\rho}q_\rho$, $Q^{\mu\nu} = \sum_i s_i^\mu s_i^\nu$ and $G_{ij} = s_i \cdot s_j$ (four-dimensional contraction). Add as inputs a term $\frac{1}{2}w \int h^{\mu\nu}\mathcal{E}_{\mu\nu}[h]$, the quadratic form of an induced Einstein–Hilbert term with $w = -\zeta^2/(16\pi G_{\text{eff}})$ when $\delta g = \zeta h$, and a conserved source $T_{\mu\nu}$ coupled as $\int h^{\mu\nu}T_{\mu\nu}$; the normalisation of the tree-level term and of the coupling is absorbed in w and in Λ, the coefficient of $\frac{1}{2}c_i \cdot Mc_i$, $Mc^\nu = q^2 c^\nu - q^\nu q \cdot c$. Since $s_i \cdot q = 0$, $\{s_1, s_2, s_3, q\}$ is a basis for $q^2 \neq 0$; $c_i = D_{ij}s_j + \gamma_i q$ with γ_i the gauge part, $h = H(c)$ with H the linear map $c \mapsto h$, which depends only on the symmetric part D^S of D, the three antisymmetric components of D are the inert amplitudes, $c^T Mc = q^2 \text{tr}(DGD^T)$, and $G = q_\perp^2 \mathbf{1} - (y^0/\ell)^2 \mathbf{k}\mathbf{k}^T$ with $q_\perp^2 = q^2 + (q_\mu y^\mu)^2/\ell^2$ and $\mathbf{k} = \mathbf{q}/|\mathbf{q}|$; its eigenvalues are $g = q_\perp^2$ (twice) and $\xi = -\sigma q^2$, $\sigma = (y^4)^2/\ell^2$, along the spatial $\mathbf{q}$. The restriction to $y^i = 0$ only provides these closed

forms.

Extra stress. The amplitude equations are equivalent to $H^T X = -\Lambda M c$ with $X = w\mathcal{E}[h] - T$. In exact rational arithmetic at $(y^0, y^4) = (5,3), (13,5)$ and several rational q one finds that X is symmetric (equivalent to the three inert-amplitude equations), that $q^\mu X_{\mu\nu} = 0$, and that $X_{\mu\nu} = -\frac{1}{2}\Lambda \sum_i \tilde{s}^i_\mu (Mc_i)_\nu$ with $\tilde{s}^i = \sum_k G^{-1}_{ik} s_k$. The amplitude equations determine X up to $q_\mu q_\nu$, which H^T annihilates; $q^\mu X_{\mu\nu} = 0$ follows from the Bianchi identity of $\mathcal{E}$ and the conservation of T, which fixes that term, as the ambiguity of the anharmonicity tensor is fixed in [25, 26]. The closed form $X = -\frac{1}{2}\Lambda \sum_i \tilde{s}^i \otimes Mc_i = -\Lambda \mathcal{L}\mathcal{E}$ expresses the extra stress through the tree-level equations; it is not an independent derivation of its conservation, and the Einstein part $w\mathcal{E}$ is the supplied input.

Crossover. The operator $\mathcal{L}$ is defined on the six-dimensional span of the symmetrised products $s_i s_j$ and is invertible for $q^2 \neq 0$ and $\xi \neq 0$. With $\mathbb{L}[Z] = (\eta Q)Z + Z(Q\eta)$ one has $h_T = \mathbb{L}[Y]$, $Y = -2X/(\Lambda q^2)$, and eliminating the amplitudes gives exactly $(w + \Lambda\,\mathcal{L})\mathcal{E} = T$, $\mathcal{L} = \frac{1}{2}\mathbb{L}^{-1} \circ (2 - P\,\mathrm{tr})$, where P projects on $q^\perp$. The operator $\mathcal{L}$ has eigenvalues $1/(2g)$ (two modes), $1/(g+\xi)$ (two), $(g \pm \sqrt{g(g+8\xi)})/(8g\xi)$ (one each) and zero on the three inert modes. In the first channel $(1 + \varrho/(2q_\perp^2))\mathcal{E} = T/w$ with $\varrho = \Lambda/w$ the ratio of the tree-level to the induced coefficient, the form $(1 - m^2\Box^{-1})$ with $\Box \to -q_\perp^2$ and $m^2 = \varrho/2$; $q_\perp^2 = k^2 + \sigma\omega^2$ is positive definite in $(\omega, \mathbf{k})$, not the form q^2. The result is the supplied Einstein part $w\mathcal{E}$ plus a non-local, frame-dependent tree-level correction.

Roots. $\det(\Lambda A + wB) \propto \Lambda^3 (q^2)^9 (\Lambda + 2gw)^2 (\Lambda + (g+\xi)w)^2 (-\Lambda^2 + 2\Lambda g w + 8 g w^2 \xi)$, which reproduces the generalised eigenvalues of the tree-plus-induced form at the sampled points. Besides $q^2 = 0$ there are roots that depend on the direction of q through the frame: in the first channel $\omega^2 = -(k^2 + \varrho/2)/\sigma$, complex for every k if $\varrho > 0$ and for $k^2 > -\varrho/2$ if $\varrho < 0$; the helicity-zero roots are complex when $g + 8\xi < 0$. The relation is elliptic in ω rather than wave-like: for $\varrho > 0$ the first channel has no real-frequency root, and for $\varrho < 0$ real roots exist only for $k^2 < |\varrho|/2$, with $\omega^2 < 0$ and growth rate proportional to k above that. The anisotropy is that of the background direction $\hat{y} = y/\ell$, through $q_\perp^2 = q^2 + (q \cdot \hat{y})^2$, and G is indefinite for spacelike q; the anisotropy is $SO(3,1)$-covariant if y^μ is treated as a vector, and is not an explicit breaking of covariance. The sign of ϱ is that of Λ/w; the sign of the induced coefficient is not determined here. These are roots of the leading-symbol dispersion relation, not shown to be poles of a retarded response (constraints and gauge fixing are not treated). The extra modes reported in [26] are, as extracted, rotation-isotropic with a reduced velocity from the cosmic frame; they belong to the same family of non-boost-invariant effects, and no match is claimed, since the map between the present $q_\perp, \xi, \sigma$ and the notation there is not established.

This is a statement for the Poisson limit, the stated class with the bilinear term, the leading symbol, tree level plus the supplied induced form and conserved source,

and the base points with $y^i = 0$. It derives no value or sign of G_{eff} or of ϱ, no masses or subleading terms, no matrix-derived matter coupling, no new derivation of the conservation of the extra stress, and no identification of G with the emergent metric of Chapter 4; the comparison with the modified Einstein equation of [25, 26] is structural. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: the Θ-built heat kernel and the induced coefficient (conditional; bounded no-go). Keep the Poisson limit, the leaf $\mathbb{CP}^{1,2}$, $X^a = T^a + A^a$, the bilinear term with $\mu^2 = -3$ and the notation of the preceding paragraphs; $D^a = \{T^a, \cdot\}$, $J^{AB} = \{M^{AB}, \cdot\}$, $\Box = \eta_{ab} D^a D^b$. The regulator is the proper-time cutoff $s \geq \Lambda_{\text{UV}}^{-2}$ of the operator $\Box_X$ itself, applied to every sector; it is built from Poisson brackets and the matrix contraction only. Background gauge fixing $D_a A^a$ gives the ghost operator $-\Box$ (weight -2), the vector operator $H_V = -\Box - 2s J_{ab} \Sigma^{ab}$ (the same $J_{ab} A^b$ as in the quadratic form above, with μ^2 added), and, for n_{ex} further matrices with trivial background, $-\Box + \mu_i^2$ with μ_i^2 an additional input. Fermions are absent. In exact arithmetic in the polynomial Poisson algebra, for both signs s: $\Box = C_{\mathfrak{so}(4,1)} - C_{\mathfrak{so}(3,1)}$ with the generators J^{AB}, $A, B \leq 4$, and J^{ab}, $a, b \leq 3$; $[\Box, J^{ab}] = 0$ while $[\Box, D^a] \neq 0$; and on functions of $y^A = M^{A5}$, $A \leq 4$, one has $D^a = s\,(y^4 \partial^a - y^a \partial_4)$, a rotation field of the hyperboloid $y \cdot y = -\ell^2$. Fibre-constant functions therefore form an invariant subspace of $\Box$, on which $\Box = \rho^{-1} \partial_\mu (\rho\, m^2 \eta^{\mu\nu} \partial_\nu \cdot)$ with $m = y^4$, $\rho = 1/m$; for $G^{\mu\nu} = m^2 \eta^{\mu\nu}$ the Seeley–DeWitt density of this sector is $\sqrt{|G|}\,(R_G/6 - E) = (5m^2 - 3\ell^2)/(4m^6)$, which is not of definite sign (a formal evaluation for $\eta = \text{diag}(-1, 1, 1, 1)$; no Euclidean heat-kernel statement is made for this sector). Fibre harmonics of order ≥ 1 are not an invariant subspace of the perturbed operators, so this sector does not by itself give the response to the level-one amplitudes.

Hypoelliptic structure. At every sampled point of the orbit (random points, and the sequence $y^4 \to 0$) the fields D^a have rank four and D^a, J^{ab} together span the six-dimensional tangent space; the horizontal rank drops on $y^4 = 0$ (one singular value proportional to y^4), which is excluded. Hence, for the Euclidean contraction $\eta \to \delta$ (the operator $\Box_E = \sum_a (D^a)^2$, which is not the Casimir difference above), the operator is a sub-Laplacian of step two with homogeneous dimension $4 + 2 \cdot 2 = 8$, not an elliptic operator on a product; the ranks, the step and the dimension 8 depend only on the fields D^a, J^{ab}. The ghost and vector operators below are then built from $\Box_E$. Under equiregularity the diagonal of its heat kernel is $K(s) = c_0 s^{-4}(1 + O(s))$ with c_0 given by the nilpotent approximation [28], here a group with four horizontal generators and a two-dimensional centre. After Fourier transform along the centre (variable $\zeta \in \mathbb{R}^2$) the nilpotent operator is a four-dimensional charged Laplacian in a constant field $F_{ab}(\zeta) = \zeta_k S^{ab}{}_k$ of the two-plane strengths $f_1, f_2 \propto |\zeta|$, with Landau levels $f_1(2n_1+1) + f_2(2n_2+1)$ and the diagonal $\prod_j f_j/(4\pi \sinh s f_j)$ per unit horizontal volume

and per $\mathrm{d}^2\zeta/(2\pi)^2$. With ν the Liouville volume of the frame $(D^0,\dots,D^3,Z_1,Z_2)$ of the nilpotent algebra, $c_0 = \nu^{-1}(64\pi^4)^{-1}\int_0^{2\pi}\mathrm{d}\varphi\,\phi_1\phi_2\int_0^\infty \mathrm{d}u\,u^3/(\sinh u\phi_1\sinh u\phi_2)$, $f_j = |\zeta|\phi_j$; for $f_1 = f_2$ this is $3\zeta(3)/(64\pi^3\nu)$. With the Euclidean contraction $\eta \to \delta$ of the four matrix indices (the Poisson algebra kept) the ratio f_1/f_2 at 12 sampled points and 9 directions of ζ each lies between 0.006 and 0.84, never at 1; no claim is made over the whole leaf. For the Lorentzian contraction $F\eta$ has one real and one imaginary eigenvalue pair, the electric-type case, to which the Euclidean heat-kernel construction does not apply.

Consequences. (i) The statement that a fuzzy factor replicates the four-dimensional coefficient of Theorem 8.2 over its modes, with threshold weights, is the limit $sf_j \ll 1$ of the diagonal above, $\prod_j sf_j/\sinh sf_j = 1 - s^2(f_1^2+f_2^2)/6 + \dots$, valid when Λ_{UV}^2 exceeds the largest populated f_j. In the Poisson limit $|\zeta|$ is unbounded and the regime is the hypoelliptic one; the replication formula does not apply there, and the weighted mode count is replaced by $\nu^{-1}\int \mathrm{d}^2\zeta\,(2\pi)^{-2}\prod_j sf_j/\sinh sf_j$. (ii) In the same nilpotent limit H_V acquires the Zeeman term $\pm 2f_1, \pm 2f_2$, so its lowest eigenvalue at fixed ζ is $-|f_1 - f_2|$ (checked by diagonalisation in a truncated Fock space): zero modes of unbounded number for $f_1 = f_2$, and eigenvalues unbounded below for $f_1 \neq f_2$, neither regulated by the proper-time cutoff. The vector sector therefore needs a fibre truncation (a finite fuzzy fibre) or fermionic partners, and in the Euclidean continuation used here e^{-sH_V} is unbounded for every $s > 0$ at the leading symbol, so the proper-time trace of this sector diverges. (iii) For $sf_j \ll 1$, the vector, ghost and n_{ex} scalar sectors together give $2 + n_{\mathrm{ex}} + \frac{22-n_{\mathrm{ex}}}{6}s^2(f_1^2+f_2^2)$; the s^2 term has the structure of the quartic term $\{X^a,X^b\}^2$ and renormalises its coefficient logarithmically, not the Einstein–Hilbert coefficient. (iv) An Einstein–Hilbert coefficient would be the weight-two term $\propto s^{-3}$ of the diagonal and its second variation under A; no such computation is made here, and neither the sign nor the size of w, hence of $\varrho = \Lambda/w$, follows, with w and ϱ the induced and tree-level coefficients of the preceding paragraph.

This is a statement for the Poisson limit, the proper-time cutoff of $\Box_X$ as the regulator, one loop, the leading (nilpotent) symbol, equiregular points, the Euclidean contraction $\eta \to \delta$ with the Lie–Poisson algebra kept, background gauge and the bosonic content listed. It does not treat Moyal terms, fermions, the subleading symbol, the dependence on A, or the identification of G with the emergent metric of Chapter 4, and it does not bear on Assumption 6.1. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: where the flux eigenvalues of the Euclidean symbol coincide (conditional; bounded no-go). Keep the setting of the preceding paragraph: Poisson limit, leaf $\mathbb{CP}^{1,2}$, $D^a = \{T^a,\cdot\}$, the nilpotent (Landau) leading symbol with centre variable $\zeta \in \mathbb{R}^2$, the contraction $\eta \to \delta$ of the four matrix indices with the Lie–Poisson algebra $\mathfrak{so}(4,2)$ kept, and the sheet realised by the orbit. Write

$y^a = M^{a5}$, $y^4 = M^{45}$, $y \cdot y = -\ell^2$, and $Q^{ab} = (y^4)^2\eta^{ab} + y^a y^b$, the Gram matrix of the fields D^a in the ambient metric of the hyperboloid; Q is positive definite with $\det Q = \ell^2|y^4|^6$. At 300 random orbit points, with both centre directions, the matrix $Q^{-1/2}F_{ab}Q^{-1/2}$, $F_{ab} = \zeta_k S^{ab}{}_k$, is self-dual or anti-self-dual with respect to δ to a relative accuracy of 10^{-9}, with the sign equal to $\operatorname{sign} y^4$; this is the expected statement that the vertical curvature of the horizontal distribution over the constant-curvature base is self-dual for the ambient metric. Hence $F = Q^{1/2}\omega Q^{1/2}$ with ω self-dual, and the plane strengths of F are $f_{1,2} = |p| \pm |r|$, with p and r the δ-self-dual and anti-self-dual parts of F. Let $a \geq b$ be the eigenvalues of Q on the plane spanned by the time axis and $\vec{y}$, so that $ab = \ell^2(y^4)^2$ and the remaining eigenvalue is $(y^4)^2$ twice. For ζ such that ω points along the axis of that plane one finds $f_2/f_1 = \min(\ell, |y^4|)/\max(\ell, |y^4|)$; for ω orthogonal to it, $f_2/f_1 = \sqrt{b/a}$; for other directions the ratio lies between these values. The numerical range of f_2/f_1 at the sampled points lies inside this bracket to 10^{-13}.

Consequences. (i) The vector sector has no negative mode at fixed ζ only if $f_1 = f_2$, and a stable point must have this for every ζ, because the set of ζ with $f_1 = f_2$ is a linear subspace of the ζ-plane (a line, or the whole plane only if F is (anti-)self-dual). Conditional on the self-duality statement above, the closed forms give $f_1 = f_2$ for all ζ exactly on $\vec{y} = 0$, $|y^4| = \ell$, that is on the two-sphere fibres over the points $y^0 = -\sqrt{2}\,\ell$, $y^4 = \pm\ell$: codimension 4 in the six-dimensional leaf (the minimisation of the distance to this set found it, with $f_2/f_1 = 1$ to six digits at six points). On the rest of the leaf the lowest vector eigenvalue is $-|f_1 - f_2| < 0$ for generic ζ, and $|f_1 - f_2|/(f_1 + f_2) = (\ell - |y^4|)/(\ell + |y^4|)$ on the time axis grows linearly with the distance from the set. (ii) The same holds for the compact real form $\mathfrak{so}(6)$ with δ throughout: there $f_2/f_1 = |y^4|/\ell$ for every ζ and every fibre point (checked at 200 points, error $4 \cdot 10^{-16}$), equal to 1 only over the poles $\vec{y} = 0$. A compact Euclidean section with the contraction δ_{ab} therefore does not replace the instability by zero modes; self-duality holds identically only for the contraction by the position-dependent ambient metric Q^{-1}, which is not a constant contraction of the matrix indices. (iii) Constant additional fluxes $f_3, \ldots$ from transverse matrices change the lowest level to $\sum_k f_k - 2\max_k f_k$ (the Landau levels of the same form, $2\pi(\sum_i (2m_i + 1)f_i \pm 2f_k)$, of [29], eq. (35), and the no-tachyon condition of its Section 9, $f_1 \leq f_2 + \cdots + f_n$); because $f_1 - f_2$ grows linearly in $|\zeta|$, a ζ-independent transverse flux restores stability only for $|\zeta|$ below a bound set by that flux, that is on a finite fuzzy fibre, not in the Poisson limit. (iv) At the self-dual points the vector density $\prod_j f_j/(4\pi \sinh s f_j)$ times the Zeeman trace tends to $f^2/(2\pi^2)$ for $sf \gg 1$, independent of s, so the integral over ζ diverges there as well; the proper-time cutoff does not regulate it.

This is a statement for the Poisson limit, the leading symbol, equiregular points, the contraction $\eta \to \delta$ (or δ with $\mathfrak{so}(6)$), constant additional fluxes and bosonic content. The self-duality of $Q^{-1/2}FQ^{-1/2}$ is verified numerically at sampled points and is not

proved, so the classification of the set $f_1 = f_2$ is conditional on it. It does not treat a contraction by a position-dependent metric, Moyal terms, fermions, a finite fibre, or other backgrounds for the transverse matrices, and it determines neither the sign nor the size of w. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: the contraction by the ambient metric and the five-matrix presentation (conditional; bounded no-go and conditional route). Keep the Poisson limit, the leaf $\mathbb{CP}^{1,2}$ of $\mathfrak{so}(4,2)$ with $\eta_6 = \mathrm{diag}(-1,1,1,1,1,-1)$, the notation $y^a = M^{a5}$, $y^4 = M^{45}$, $Q^{ab} = (y^4)^2\eta^{ab} + y^a y^b$ of the preceding paragraph, and $T^a = M^{a4}$, $D^a = \{T^a, \cdot\}$. On the orbit $(\eta_6 M)^2 = -\ell^2\mathbf{1}$ (checked at 200 random points to 10^{-12}), whence $y \cdot y \equiv \eta_{AB}y^A y^B = -\ell^2$ $(A, B = 0, \ldots, 4)$ and $T^a y_a = 0$. The base metric induced from the ambient space is $g = \eta + y_\mu y_\nu/(y^4)^2$ with inverse $g^{-1} = \eta + yy/\ell^2$; it is positive definite, and the contraction of the fields D^a by Q^{-1} has base symbol exactly g^{-1} (checked numerically). The kinetic tensor of the four matrices T^a is $K = (y^4)^2\eta = (y^4)^2(g^{-1} - yy/\ell^2)$. The contraction by Q^{-1} equals $(y^4)^{-2}[\eta_{ab}D^aD^b + (y_aD^a)^2/\ell^2]$ and therefore replaces the Lorentzian tensor K by the Euclidean metric g of the hyperboloid; it is not a Wick rotation of η.

No supply in the four-matrix class. Hypothesis class: the Poisson limit; the four matrices $T^a = M^{a4}$; the quartic term plus one bilinear term, with Hessian $L(A)_a = -\Box A_a + D_a(D_bA^b) - 2s\mathcal{J}_{ab}A^b + \mu^2 A_a$; the background gauge function D_aA^a, the ghost operator $-\Box$, and a regulator that is a function of $\Box_X$; the matrix-index form constant, η_{ab}. In this class every operator is a polynomial with constant coefficients in the D^a and $J^{ab} = \{M^{ab}, \cdot\}$, that is, an element of the enveloping algebra of $\mathfrak{so}(4,1)'$, the Lie algebra generated by the D^a and J^{ab} (the subalgebra of $\mathfrak{so}(4,2)$ on the indices $0, \ldots, 4$; the generators M^{a5} and M^{45} are not in it). The operator $(y_aD^a)^2$ is not an element of that enveloping algebra: its coefficients are the non-constant functions y_a, which would have to be supplied by the missing generators. Hence in this class none of the action, the bilinear term, the background gauge, the ghosts and the $\Box_X$ regulator produces the term $(y_aD^a)^2$. The full Casimir $\frac{1}{2}\eta_{AC}\eta_{BD}J^{AB}J^{CD}$ of $\mathfrak{so}(4,2)$ has base symbol $-2\ell^2 g^{-1}$ (checked numerically), but it lies outside the enveloping algebra of $\mathfrak{so}(4,1)'$ and enters neither the Hessian nor the ghost operator. Extensions of the matrix set by y^a with a positive-norm contraction (eight matrices T^a, y^a; seven matrices T^a, y^i) have rank-six horizontal span, but the equation of motion then needs different bilinear coefficients for different matrices (checked by exact Poisson algebra). The eight-matrix set T^a, y^a with the y-block contracted by $-\eta$ is stationary for one coefficient $\mu^2 = -4$, with horizontal rank six, and is not examined further.

Five-matrix presentation. Take $X^A = M^{Ac_0}$, $A \neq c_0$, with the restriction of η_6 as contraction. Definition 3.4 $(D = 5)$ covers $c_0 \in \{0, 5\}$ up to relabelling; the forms for $c_0 \in \{1, \ldots, 4\}$, of signature $(3,2)$, lie outside it and are included as a comparison. The orbit relation gives $y \cdot y = \eta_{c_0c_0}\ell^2$ for $y^A = M^{Ac_0}$. At 25 random points for each

c_0 the horizontal rank is 4 and the centre rank 2; the induced contraction $h = c^T \eta c$ on the horizontal space is positive definite for $c_0 \in \{0, 5\}$ and has signature $(2, 2)$ for $c_0 \in \{1, \dots, 4\}$; a Lorentzian signature does not occur. For $X^A = M^{A5}$, $A = 0, \dots, 4$ (stabiliser $\mathfrak{so}(4, 1)$): $y_A D^A = 0$ on the orbit, the base symbol of $\eta_{AB} D^A D^B$ is $\ell^2 g^{-1}$ (checked to 10^{-12}), and at 300 random points and 91 directions of ζ, for both signs of the bracket, the vertical flux in an h-orthonormal frame is self-dual or anti-self-dual to 10^{-11}, so $f_1 = f_2$ identically. This constant contraction of the five matrices is consistent with the statement that self-duality holds identically only for the contraction by the position-dependent Q^{-1} of the four matrices T^a: its base symbol $\ell^2 g^{-1}$ is that of Q^{-1} up to the factor $(y^4)^2$. The bilinear coefficient required by stationarity is $\mu^2 = +4$ (exact, both signs of s), the Zeeman term of the gauge-fixed vector operator has the same modulus as in the four-matrix class, and in the polynomial Poisson algebra, exactly,

$$H_V = C_{\mathfrak{so}(4,2)} - C_G + C_\Sigma + \mu^2, \qquad \Box = C_{\mathfrak{so}(4,1)} - C_{\mathfrak{so}(4,2)},$$

where C_G is the Casimir of the total $\mathfrak{so}(4, 1)$ action on vector-valued functions and C_Σ that of the vector index. By the Weyl formulae the $\mathfrak{so}(4, 1)$ component $(k + 1, k)$ of the degree-k vector polynomials has $C_{\mathfrak{so}(4,2)} - C_G + C_\Sigma = 0$ for every k, so it is an eigenspace of H_V with eigenvalue μ^2 (verified directly for $k = 1$, dimension 35). On the 75-dimensional space of degree-one vector polynomials the formal eigenvalues of H_V are $-8, -4, -2, +2, +4$ with multiplicities $1, 5, 20, 14, 35$ (checked); polynomials are not normalisable, so the negative values have no spectral meaning. At the leading symbol the vector operator then has the lowest Landau level at zero for the four horizontal components, a zero-mode family with density $f^2/(2\pi^2)$ in ζ, independent of the proper time; the normal component has no Zeeman term, and the bosonic density per $\mathrm{d}^2\zeta/(2\pi)^2$ is $f^2(16\pi^2)^{-1}\big[8 + (3 + n_{\mathrm{ex}})/\sinh^2(sf)\big]$. The ζ-integral of the second term converges; that of the first diverges and is not regulated by the proper-time cutoff. The fibre of the leaf is compact, so ζ is expected to be bounded by the fibre harmonic number of the quantum space; the divergence is a feature of the Poisson limit.

Compact check at finite size. For the compact form $\mathfrak{so}(6)$ with δ throughout, on the N-th symmetric power of the fundamental representation, the exact Hessian of the quartic term plus $\frac{1}{2} m^2 \operatorname{Tr} X^2$ (finite matrices, $N = 1, \dots, 4$) satisfies $H + G^\dagger G = C_6 - C_5^{\mathrm{tot}} + C_5^{\mathrm{vec}} + m^2$ for $X^A = M^{A6}$, $m^2 = -4$. Its lowest eigenvalue is $-4 = m^2$ for $N \le 4$, with the family $(k + 1, k)$, $1 \le k \le N$, of total multiplicity $\sum_{k=1}^{N} (2k + 1)(2k + 3)(2k + 5)/3$ and 14 further modes at -2, so that the number of negative modes is $49, 154, 385, 814$ for $N = 1, \dots, 4$ and this compact background is unstable; for the four-matrix background $X^a = M^{a5}$, $m^2 = -3$, it is $-(N + 3)$, growing with N. The sign of m^2 is -4 on the sphere and $+4$ on the hyperboloid.

Consequences. (i) In the matrix sets examined (T^a; M^{Ac_0}; the extended sets listed above), the term $(y_a D^a)^2$ is supplied by the matrix data only if the matrices are the position functions $X^A = M^{A5}$; the price is that the horizontal metric induced by the contraction is then the Euclidean metric g, and the conformally flat Lorentzian K of the four-matrix class is not obtained, so the level-one and stress results of the four-matrix class are not obtained in this presentation. (ii) For the hyperboloid, the vector zero modes of the leading symbol are lifted by $+\mu^2$ only if the vanishing of the Casimir combination, which is exact on polynomial vector fields and on the compact form, continues to the normalisable sector; this continuation is not shown, and the negative formal eigenvalues $-8, -4, -2, +2$ of the same operator on degree-one polynomials, like the negative modes of the compact background, may persist on normalisable states. (iii) Neither the sign nor the size of w follows: the weight-two coefficient and its variation under A are not computed, and the passage from a Euclidean coefficient to the Lorentzian one is a Wick-rotation hypothesis tied to Problem E.

This is a statement for the Poisson limit, the leading symbol, equiregular points, the bosonic content and the single bilinear term; the self-duality of the flux is verified numerically and not proved, and the finite-size spectrum is for the compact form. It does not treat fermions, Moyal terms for the hyperboloid, or the Lorentzian emergent metric. It does not identify g with the emergent metric of Chapter 4, does not derive Lorentzian signature, and leaves Assumption 4.6 in place (Problem E). This is a scope statement and not a retraction of any earlier result. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: the Casimir of $\mathfrak{so}(4,2)$ and its combinations as proper-time regulators in the four-matrix class (conditional; bounded no-go and conditional route). Keep the Poisson limit, the leaf $\mathbb{CP}^{1,2}$ of $\mathfrak{so}(4,2)$, $T^a = M^{a4}$, $D^a = \{T^a, \cdot\}$, $J^{AB} = \{M^{AB}, \cdot\}$, $\Box = \eta_{ab} D^a D^b$, the vector operator H_V with $\mu^2 = -3$ and the notation of the preceding paragraphs. Write C_{42}, C_{41}, C_{31} for $\frac{1}{2}\eta_{AC}\eta_{BD} J^{AB} J^{CD}$ summed over the indices $0, \ldots, 5$, $0, \ldots, 4$, $0, \ldots, 3$, and $\sigma(\cdot)$ for the quadratic form $\sum w\,(\xi \cdot v)^2$ of the cotangent vector ξ of an operator $\sum w\, v^2$ of vector fields v (the principal symbol is its negative, so that the principal symbol of H_V is $\sigma(\Box)$ and that of Δ_κ below is $-\sigma(\Delta_\kappa)$).

Exact relations. In the polynomial Poisson algebra, for both signs of the bracket, $\Box = C_{41} - C_{31}$, $\mu^2 = -3$ and, with C_G the Casimir of the total $\mathfrak{so}(3,1)$ action on vector-valued functions (orbital part plus the vector index, the sign of the index action correlated with the sign of the bracket) and C_Σ that of the vector index,

$$H_V = C_G - C_{41} - C_\Sigma + \mu^2,$$

the unique solution of the linear fit $H_V = xC_G + yC_{31} + zC_{41} + wC_{42} + uC_\Sigma + \mu^2$

with $w = 0$. Hence C_{42} does not occur in H_V, and $C_{42} - C_{41} = -\sum_{A\le 4} \eta_{AA} (J^{A5})^2$. Since C_{42} is central in $U(\mathfrak{so}(4,2))$ and C_{41} commutes with $\mathfrak{so}(4,1)$, both commute with H_V, with $\Box$ and with the ghost operator $-\Box$; a function of C_{42} and C_{41} therefore commutes with every operator of the class and no ordering ambiguity arises.

Symbols. At 40 random orbit points the form $\sigma(C_{42})$ on the six-dimensional cotangent space has signature $(2,4)$, and so has $\sigma(C_{41})$. The pencil $\sigma(C_{41})v = \lambda\,\sigma(C_{42})v$ has the eigenvalues $\frac{1}{2}$ (four times) and 1 (twice) at 30 points (deviation $3 \cdot 10^{-13}$); on the four-dimensional eigenspace (the span of the dy^C, $y^C = M^{C5}$) $\sigma(C_{42})$ is negative definite and on the two-dimensional (vertical) eigenspace positive definite. The base symbol $-2\ell^2 g^{-1}$ of the preceding paragraph is the first of these; the fibre carries the opposite sign. Write $\sigma(C_{42}) = -\alpha q_b + \beta q_f$ in this splitting, so that $\sigma(C_{41}) = -\frac{1}{2}\alpha q_b + \beta q_f$. The operators $C_{41} - C_{42}$ (the five-matrix $\Box$, purely horizontal) and $2C_{41} - C_{42}$ (purely vertical) are the two degenerate members of the family

$$\Delta_\kappa = \kappa\, C_{42} - C_{41}, \qquad -\sigma(\Delta_\kappa) = \alpha(\kappa - \tfrac{1}{2}) q_b + \beta(1-\kappa) q_f,$$

which, as the principal symbol of Δ_κ, is positive definite exactly for $\frac{1}{2} < \kappa < 1$ (checked at 15 points for nine values of κ); the value $\kappa = \frac{3}{4}$ gives $-\sigma(\Delta_{3/4}) = \frac{1}{4}(\alpha q_b + \beta q_f)$, the absolute value of $\sigma(C_{42})$ in the splitting. The horizontal planes $\mathrm{span}\{D^a\}$ of the four-matrix class and $\mathrm{span}\{J^{A5}\}$ of the five-matrix class are different four-planes (their sum has dimension 6 at six points).

Consequences. (i) No function of C_{42} alone is an elliptic regulator: the level sets of an indefinite form in the cotangent fibre are non-compact, so $\int \mathrm{d}^6\xi\, \chi(\sigma(C_{42})/\Lambda_{\mathrm{UV}}^2)$ diverges for every χ of compact support, and a cutoff of C_{42} leaves the null cone $q_b = \beta q_f/\alpha$ unregulated; this is a statement about the leading symbol. (ii) Δ_κ, $\frac{1}{2} < \kappa < 1$, is built from the matrix data M^{AB} and the contraction η_{AB} only, commutes with H_V, $\Box$ and the ghost operator, and has a definite symbol on the whole leaf. For a cutoff function χ of compact support, the density $(2\pi)^{-6}\int \mathrm{d}^6\xi\, \chi(-\sigma(\Delta_\kappa)/\Lambda_{\mathrm{UV}}^2)\, F(\sigma(\Box))$ is finite at every point for every continuous F (at the level of the leading symbol); in particular the central variable ζ of the preceding paragraphs is confined to a ball $|\zeta| \lesssim \Lambda_{\mathrm{UV}}$, which regulates the ζ-integral left divergent there. It is a different regulator from a function of $\Box_X$, and it uses the generators J^{A5}, which are not in $U(\mathfrak{so}(4,1)')$; it is an additional element of the regularisation and not derived from the action. (iii) The volume of the window $\{-\sigma(\Delta_\kappa) \le \Lambda_{\mathrm{UV}}^2\}$ is proportional to $\Lambda_{\mathrm{UV}}^6[(\kappa - \frac{1}{2})^2(1-\kappa)]^{-1}$ (checked against $\det^{-1/2}$ at five values of κ), so the leading coefficient depends on κ and diverges at both ends of the interval, where the horizontal and the vertical directions respectively are unregulated; no value of κ is singled out by the algebra, and the quadratic coefficient, which is the one an Einstein–Hilbert term needs, is not independent of the choice of κ and χ without a further argument. (iv) For the Lorentzian η the form $\sigma(\Box)$ has signature $(3,1)$ with two null directions, and in the

window of $\Delta_{3/4}$ the generalised eigenvalues of $(\sigma(\Box), -\sigma(\Delta_{3/4}))$ are -2 (at eight points, the same value at each) and three positive values that grow with $(y^4)^2$ (so the negative bound is uniform in position and the positive range is not); the principal symbol $\sigma(\Box)$ of H_V is therefore bounded by $\mp$ a multiple of Λ_{UV}^2 on the window but is negative on part of it, so the symbol of e^{-sH_V} is bounded by $e^{2s\Lambda_{\mathrm{UV}}^2}$ there and the Euclidean proper-time integral at large s still diverges. The window removes the divergence at large ζ, not the negative direction: for the Euclidean contraction $\eta \to \delta$ of the preceding paragraphs the lowest Zeeman level $-|f_1 - f_2|$ is bounded in the window of $\Delta_{3/4}$ by $d_*\Lambda_{\mathrm{UV}}$ (in the normalisation of the forms above) with $1.7 \le d_* \le 5.6$ at six points, linear in the cutoff.

Compact check. For the compact form $\mathfrak{so}(6)$ with δ, $X^a = M^{a5}$, $a = 1, \dots, 4$, $m^2 = -3$, on the N-th symmetric power (the matrices $\mathrm{End}(\mathrm{Sym}^N \mathbb{C}^4) = \bigoplus_{k \le N}(k, k, 0)$ of $\mathfrak{so}(6)$, so that a cutoff on $C_{\mathfrak{so}(6)} = 2k^2 + 6k$ is the truncation of the matrix algebra), the gauge-fixed Hessian $H + G^\dagger G$, $Ga = \sum_a [X^a, a^a]$, has the spectrum

$$\lambda = c_5(a, b) - c_4(c', d') + 3 + m^2,$$

with $c_5(a, b) = a(a + 3) + b(b + 1)$, $c_4(c, d) = c(c + 2) + d^2$, $a = k$ the label of $\mathfrak{so}(6)$, $0 \le b \le a$, (c, d) the $\mathfrak{so}(4)$ content of (a, b) and (c', d') that of $(c, d) \otimes (1, 0)$; this was checked against the numerical spectrum for $N = 1, 2, 3$ (differences below 10^{-12}). The lowest eigenvalue in the window $k \le A$ is $-(A + 3)$, the number of negative modes is $38, 124, 344, 750$ for $A = 1, \dots, 4$, and their fraction of the $4\binom{A+3}{3}^2$ modes falls numerically to $2.5 \cdot 10^{-3}$ at $A = 48$ (local exponents 3.9 and 5.7, approaching 4 and 6; not proved). The cutoff by the Casimir therefore bounds the negative direction by a quantity growing linearly in the cutoff and removes none of it.

Eight matrices. For T^a and $y^a = M^{a5}$, $a = 0, \dots, 3$, with the contraction η on T and $\lambda\eta$ on y, exact stationarity requires the bilinear coefficients $\lambda - 3$ for T and $3\lambda - 1$ for y, equal only at $\lambda = -1$ (common value -4). The contraction form on the tangent space has signature $(3, 3)$ at $\lambda = -1$ and $-100 \le \lambda \le -0.01$ (thirty points each), $(5, 1)$ at $\lambda = 0.01, 0.1, 10$ and is degenerate of rank 4 at $\lambda = 1$; the horizontal span has rank 6 throughout, so there is no centre variable. A $(5, 1)$ induced form with no centre variable thus occurs at the sampled values $\lambda = 0.01, 0.1, 10$, and requires two independent bilinear coefficients; the Hessian of that set is not examined.

This is a statement for the Poisson limit, one loop, the leading symbol, the four-matrix class with the bilinear term, the regulators that are functions of C_{42} and C_{41}, and the sampled points; the signatures and the pencil are numerical, the relations in the first paragraph exact. It does not treat fermions, the Moyal terms, a Lorentzian $i\epsilon$ integral, the identification of the window with any admissibility condition of the fibre, or the subleading symbol, it does not compute the weight-two coefficient or its variation under A, and neither the sign nor the size of w follows. It identifies

no induced metric with the emergent metric of Chapter 4 and, since a Euclidean coefficient is not a Lorentzian one, leaves Assumption 4.6 in place (Problem E). This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: fibre admissibility and the Lorentzian $i\epsilon$ window (conditional; bounded no-go). Keep the setting and notation of the preceding paragraph (Poisson limit, leaf $\mathbb{CP}^{1,2}$, $T^a = M^{a4}$, $\Box = \eta_{ab}D^aD^b$, H_V with $\mu^2 = -3$, C_{42}, C_{41}, C_{31}, $\sigma(\cdot)$, Δ_κ); write $S^2 = 2C_{41} - C_{42}$, whose symbol βq_f is purely vertical.

Fibre cutoffs. In the Poisson limit the fibre of the leaf is a compact two-sphere and $|\zeta|$ is unbounded; it is expected to be bounded by the fibre harmonic number of the quantum space. Two bounds on the harmonic number are in the literature. (a) $S^2 = 2s(s+1)$ on the sectors $\mathcal{C}^s$ with $s \leq n$, from the $(n+1)$-dimensional representation of the stabiliser $SU(2)_L$ on the lowest-weight state [24]; the classical counterpart is the radius of the fibre as a coadjoint orbit of $SU(2)_L$, the self-dual part of M^{ab} on the η-orthogonal complement of y: at 10 orbit points exactly one of the two radii vanishes and the other equals ℓ, independent of y. (b) A local bound $s \leq m = r^{-1}x^0$ from the uncertainty of the coherent states, which the same group states to correct earlier statements on the higher-spin cutoff [23]; it grows with the point and is not the area bound of (a), and the two are not reconciled here. Neither is a sharp cutoff of commuting matrix-built operators.

Admissible cone. For admissible modes, the principal series of $SO(4,1)$, one has $r^{-2}\Box_H > s^2 + s + 9/4$ with $\Box_H = -C_{42} + C_{41}$ in the convention of [24], equivalent, by (2.7) and (3.43) of that reference, to $-C_{42} > 9/2$ in that convention (its footnote 8 prints $C_2[\mathfrak{so}(4,2)] > 9/2$ and states the identification with the $SO(4,2)$ content of $\mathrm{End}(\mathcal{H}_n)$ as plausible, not investigated). With $\Box_H$ horizontal, S^2 vertical and $\sigma(C_{42}) = -\alpha q_b + \beta q_f$ of the preceding paragraph, the semiclassical content of this bound is the cone $-\sigma(C_{42}) \geq 0$, that is $\alpha q_b \geq \beta q_f$; the Casimirs here and there differ by a sign at the symbol level. Conditional on this identification, a cutoff of C_{42} restricted to the cone together with a bound $\beta q_f \leq Q$ is a compact region, so a function of the central operator C_{42} regularises once the fibre is bounded. The cone restricts to the admissible sector and is not what bounds the window: the bound $\beta q_f \leq Q$ is a cutoff of $S^2 = 2C_{41} - C_{42}$, so this relaxes the hypothesis of the preceding paragraph and does not contradict it, whose consequence (i) for functions of C_{42} alone stands. The Lorentzian moments, the flux and the exact-sector results below are for the Gaussian window of Δ_κ, not for this fibre-bounded window. For fixed Q and $\Lambda^2_{\mathrm{UV}} \to \infty$ the window $\{-\sigma(\Delta_\kappa) \leq \Lambda^2_{\mathrm{UV}}\}$ has its vertical term bounded by $\beta(1-\kappa)Q$, so κ only rescales Λ^2_{UV} by $2\kappa - 1$ at leading order and the divergence at $\kappa \to 1$ is absent. The cutoff function and the base scale remain external: the quantisation map is $SO(4,2)$-equivariant and so acts on each irreducible component by a scalar, but this multiplier is not computed, and neither χ nor the base scale is fixed. The window

uses M^{AB}, which are background data and not dynamical matrices, so the regulated trace is not shown to be a diffeomorphism-covariant functional of X^a.

Lorentzian trace in a window. Define the one-loop sum as $\sum \ln(\lambda - i\epsilon) = -\int_0^\infty \frac{\mathrm{d}s}{s} \operatorname{Tr}\left[\chi\, e^{-is(H_V - i\epsilon)}\right]$ up to a scale-dependent constant, with χ a function of the commuting operator Δ_κ; the contour is the proper time of H_V with the factor $e^{-\epsilon s}$, and uses no metric. In a window of compact support (or a Gaussian) the number of modes per unit volume is finite, so the trace is finite for every s, and $\sum \ln(\lambda - i\epsilon)$ is a finite product over the window whether or not the Schwinger integral converges. For $\chi = e^{-\sigma_W/\Lambda_{\mathrm{UV}}^2}$, $\sigma_W = -\sigma(\Delta_\kappa)$, the leading-symbol density is the Gaussian $(4\pi)^{-3}\Lambda_{\mathrm{UV}}^6 \det(\sigma_W)^{-1/2} \prod_{k=1}^4 (1 + is\Lambda_{\mathrm{UV}}^2\lambda_k)^{-1/2}$, with λ_k the nonzero generalised eigenvalues of $(\sigma(\Box), \sigma_W)$; the kernel of $\sigma(\Box)$ is not the vertical plane (checked at eight points). The mass-type moment is $\langle(\sigma - i0)^{-1}\rangle = i\int_0^\infty \mathrm{d}s \prod_k (1 + is\lambda_k)^{-1/2}$, whose real part is the principal value. For $\lambda = (-c, a, a, a)$ the real part is $2/(a + c)$ (closed form, checked), and for 4000 samples with $c \in [0.1, 10]$, $a_k \in [10^{-3}, 10^3]$ it is positive. At eight orbit points ($\kappa = \frac{3}{4}$, $(y^4)^2$ from 0.08 to 144) it is 1.1 to 2.3 times the value for the Euclidean contraction $\eta \to \delta$, with the same sign, and the imaginary part is positive.

Flux. At the same points the Lorentzian centre flux $F_{ab}(\zeta)$ has ηF with one real pair $\pm a$ and one imaginary pair $\pm ib$, $a, b > 0$ at all sampled ζ, and $F_{ab}F^{ab}$ takes both signs (negative for $(y^4)^2 \lesssim 0.4$, positive above). The vector block therefore has Zeeman eigenvalues $\pm 2b$ (phases in $e^{-is(H_V - i\epsilon)}$) and $\pm 2ia$ (a factor $e^{\pm 2sa}$). The Schwinger integrand of the vector sector then grows for the electric pair, bounded by e^{2sa} with a linear in the cutoff, while $\ln\det(\sigma + Z) = \ln(\sigma^2 - 4b^2) + \ln(\sigma^2 + 4a^2)$ has integrable logarithmic singularities at $\sigma = \pm 2b$. The exponential growth of the Euclidean contraction is absent order by order in the flux (each order is a polynomial in s times $e^{-is\sigma}$) and for the magnetic pair, at the leading symbol and at the sampled points, and is present for the electric pair as a property of the integral representation and not of the determinant. The order-F^2 term is proportional to $F_{ab}F^{ab}$, so its sign is not definite on the leaf.

Invariant sector, exactly. On fibre-constant functions $\Box = -(\Delta_{H^4} - \Delta_{H^3})$ with $H^4 = \mathbb{R}_\tau \times H^3$, metric $\mathrm{d}\tau^2 + \cosh^2\tau\, \mathrm{d}H_3^2$, $y^4 = \ell \sinh\tau$, and the diagonal of $e^{z\Delta_{H^4} + w\Delta_{H^3}}$ is the H^3 integral of the product of the two heat kernels. Its expansion gives $D_0 = (4\pi)^{-2} z^{-1/2} (z + w\cosh^2\tau)^{-3/2}$, the Gaussian symbol formula above, and, for the pure $\Box$ term (window removed), the Seeley–DeWitt coefficient $(5m^2 - 3\ell^2)/(4m^2)$, $m = y^4$, whose product with $\sqrt{|G|} = m^{-4}$ is the density $(5m^2 - 3\ell^2)/(4m^6)$ of the preceding paragraphs, obtained by an independent route. With the window the weight-two part of $\sum \ln(\lambda - i\epsilon)$ is $-(4\pi)^{-2}\Lambda_{\mathrm{UV}}^2\, Q(a)$, $a = \sinh\tau$, plus -2 times a scale-dependent logarithm proportional to the number of modes, which is the $R/6$ curvature term of the auxiliary H^4 window geometry; $\operatorname{Re} Q$ is indefinite, and with the subtraction at $x = 1$ it changes sign at $a \simeq 0.62$ (not at $a^2 = \frac{3}{5}$, the zero of the coefficient above), a location

that moves with the subtraction point and has no invariant meaning; $\operatorname{Im} Q \to \pi$. This is the Lorentzian density of one scalar sector at the background $A = 0$, with a scale-dependent additive constant; it is not the Einstein–Hilbert coefficient.

Consequences. (i) The coefficient carrying w is the weight-two term, a local Poisson-algebraic coefficient multiplied by window moments; the signature enters through the moments and the window, and in the potential channel the Lorentzian moment is finite and positive in the class sampled. (ii) The mode-count term is proportional to the difference between the numbers of bosonic and fermionic modes in the window, so its logarithmic scale dependence cancels only if that difference vanishes at this order. (iii) The weight-two coefficient as a quadratic form in A is not computed, the vector, ghost and fibre-harmonic sectors are not computed, and neither the sign nor the size of w, hence no sign of an attractive coupling, follows.

This is a statement for the Poisson limit, one loop, the leading symbol and the invariant sector, the four-matrix class with $\mu^2 = -3$, Gaussian windows and the sampled points; the admissible cone is conditional on the identification stated, the Lorentzian moments are not computed in the fibre-bounded window, and the positivity of the moment is numerical. It does not treat fermions, Moyal terms, the dependence on A, or a window built from dynamical matrices only; it identifies no induced metric with the emergent metric of Chapter 4 and, since a Euclidean coefficient is not a Lorentzian one, leaves Assumption 4.6 in place (Problem E). This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: Lorentzian moments in the fibre-bounded cone window, and the vector, ghost and spin terms of the invariant sector (conditional; bounded no-go). Keep the setting and notation of the two preceding paragraphs (Poisson limit, leaf $\mathbb{CP}^{1,2}$, $T^a = M^{a4}$, $\Box$, H_V with $\mu^2 = -3$, C_{42}, C_{41}, C_{31}, $\sigma(\cdot)$, $S^2 = 2C_{41} - C_{42}$ with symbol βq_f, the base symbol αq_b, $a = \sinh\tau = y^4/\ell$, and the function $Q(a)$ of the exact invariant sector). At a point of the leaf take coordinates $(u, \zeta) \in \mathbb{R}^4 \times \mathbb{R}^2$ on the cotangent space with $|u|^2 = \alpha q_b$ and $|\zeta|^2 = \beta q_f$, so that $-\sigma(C_{42}) = |u|^2 - |\zeta|^2$, and write K for the form $\sigma(\Box)$ (rank four, signature $(3,1)$) in these coordinates, with blocks K_{uu}, $\mathsf{K}_{u\zeta}$, $\mathsf{K}_{\zeta\zeta}$.

Gaussian windows. Let $\lambda_0 = -c$ and $\lambda_1, \lambda_2, \lambda_3$ be positive and $\mathcal{M} = \mathrm{i}\int_0^\infty \mathrm{d}s \prod_{k=0}^3 (1+\mathrm{i}s\lambda_k)^{-1/2}$, the Gaussian average of $(\sigma - \mathrm{i}0)^{-1}$. Rotating $s = -\mathrm{i}u$ (the fourth quadrant contains no singularity) gives $\mathcal{M} = \int_0^\infty \mathrm{d}u\,(1 - cu - \mathrm{i}0)^{-1/2} \prod_{k\geq 1}(1+\lambda_k u)^{-1/2}$, whose first factor is real for $u < 1/c$ and equals $\mathrm{i}(cu-1)^{-1/2}$ for $u > 1/c$. With $u = (1 \mp v^2)/c$,

$$\operatorname{Re}\mathcal{M} = \frac{2}{c}\int_0^1 \mathrm{d}v \prod_{k\geq 1}\Big(1+\frac{\lambda_k(1-v^2)}{c}\Big)^{-1/2} > 0, \qquad \operatorname{Im}\mathcal{M} = \frac{2}{c}\int_0^\infty \mathrm{d}v \prod_{k\geq 1}\Big(1+\frac{\lambda_k(1+v^2)}{c}\Big)^{-1/2} >$$

and $\operatorname{Re}\mathcal{M} = 2/(\lambda + c)$ for $\lambda_k = \lambda$. The pencil of $\sigma(\Box)$ with any positive definite

window form has one negative, three positive and two zero generalised eigenvalues, so the principal-value mass moment of every Gaussian window of the class is positive; this proves, for Gaussian windows, the positivity that the preceding paragraph found by sampling.

Geometry of the cone window. At 60 random orbit points (agreement 10^{-9} or better, not proved): the eigenvalues of K_{uu} are $-\frac{1}{2}$ and $\frac{1}{2}\sinh^2\tau$ (three times); the Schur complement of K_{uu} in K vanishes, so $\sigma = Q_L(u + w\zeta)$ with $Q_L = \mathsf{K}_{uu}$ and $w = \mathsf{K}_{uu}^{-1}\mathsf{K}_{u\zeta}$; w is conformal, $|w\zeta| = \coth\tau\,|\zeta|$, and its image is orthogonal to the time eigenvector of Q_L. In the variable $\boldsymbol{x} = u + w\zeta$ the removed region $|u| < |\zeta|$ is therefore the ball of radius $r = |\zeta|$ about the spacelike point $p = w\zeta$, at distance $r\coth\tau = r\sqrt{1 + c/\lambda}$ from the apex $\boldsymbol{x} = 0$ with $c = \frac{1}{2}$, $\lambda = \frac{1}{2}\sinh^2\tau$: the cone $|u| = |\zeta|$ touches the null cone of Q_L.

Mass moment in the fibre-bounded cone window. Let $\chi \geq 0$ be bounded and integrable on $[0, \infty)$ and take the window function $\chi(|u|^2 - |\zeta|^2)$ on $|\zeta|^2 \leq Q_f$ and $|u| \geq |\zeta|$. For a ball of radius R about p with $|p| < R$, polar coordinates about the apex give $\mathrm{PV}\int_{|\boldsymbol{x}-p|<R} \mathrm{d}^4\boldsymbol{x}\, Q_L^{-1} = \frac{1}{2}(R^2 - |p|^2)I_0 + p^{\mathsf{T}}\mathsf{N}p$, where $I_0 = \mathrm{PV}\oint \mathrm{d}\Omega\, Q_L^{-1} = 8\pi^2/\cosh^2\tau$ and $\mathsf{N} = \mathrm{PV}\oint \mathrm{d}\Omega\, nn^{\mathsf{T}}Q_L^{-1}$, because the remaining term of the exit radius is odd in the direction n and Q_L is even (checked against a direct quadrature to 10^{-4}). For spacelike p with $|p| \geq R$, quadrature indicates that the same expression continues to hold up to the tangency distance $R\coth\tau$ (agreement to 10^{-3} in the sampled cases) and departs from it beyond; this is a numerical observation and is not proved here. Under this observation every shell $r \leq |u| \leq \sqrt{r^2 + \Lambda^2}$ has principal-value moment $\frac{1}{2}\Lambda^2 I_0$, independent of p, and so

$$\mathrm{Re}\,\mathcal{N}_W = \tfrac{1}{2}\,\varrho\,\pi Q_f\, I_0 \int_0^\infty \chi(\lambda)\,\mathrm{d}\lambda > 0, \qquad \mathcal{N}_W = \int_W \varrho\,\mathrm{d}^4u\,\mathrm{d}^2\zeta\;\chi\,(\sigma - \mathrm{i}0)^{-1},$$

with $\varrho > 0$ the Jacobian of the coordinates (u, ζ); the sign of the real part is conditional on the numerical observation, and the real part depends on the cutoff function only through $\int\chi$ and on the fibre bound only through the area πQ_f. The imaginary part is $\pi\int_W \varrho\,\chi\,\delta(\sigma) \geq 0$. Finiteness of $\mathcal{N}_W$ is likewise rested on the shell formula: the apex singularity is of order $|\boldsymbol{x}|^{-2}$ in four dimensions and integrable, but the window function has a jump at the cone edge when $\chi(0) \neq 0$, and the edge touches the null cone of Q_L at one point, so finiteness there is not shown independently.

Zeeman field and electric growth. The first-order term of H_V is exactly the Zeeman matrix of the field $F_{ab}(\xi) = \xi(\mathcal{J}_{ab})$, linear in the covector ξ; let ηF have eigenvalues $\pm f_e$ (electric pair, factor $\mathrm{e}^{\pm 2sf_e}$ in $\mathrm{e}^{-\mathrm{i}sH_V}$) and $\pm\mathrm{i}f_m$ (magnetic pair, phases), so $f_e^2 - f_m^2 = -F_{ab}F^{ab}/2 = -\sigma(C_{31})(\xi)$. At 59 of the 60 points (one point is numerically singular in the whitening, a fibre Gram eigenvalue vanishing in double precision), to 10^{-9} except for the maximum over the base plane, which is obtained from 3000 sampled directions to $6 \cdot 10^{-5}$: on the base plane $f_m = 0$ and $\max f_e = |u|\cosh\tau/\sqrt{2}$; on the

vertical plane of the pencil $f_e = |\zeta| \sinh\tau/\sqrt{2}$, $f_m = |\zeta|/\sqrt{2}$ in every direction; on the kernel plane of K (the centre variable of the nilpotent model) $f_e^2 = |\zeta|^2/(2(1+2\sinh^2\tau))$ and $f_m^2 = |\zeta|^2 \sinh^2\tau/(2(1+2\sinh^2\tau))$, so that on this plane $F_{ab}F^{ab}/2$ is negative for $|y^4| < \ell$, positive for $|y^4| > \ell$ and changes sign exactly at $|y^4| = \ell$; this exact threshold replaces the sample-dependent value of the preceding paragraph on that plane (on the vertical plane the signs are opposite). The fibre-driven electric rate $2f_e$ is $2s_f \sinh\tau$ with $\beta q_f = 2s_f^2 \le 2s_{\max}^2$, whereas the base-driven rate grows linearly with $|u|$; the electric rate is thus bounded by a quantity linear in the base cutoff, whose fibre part is bounded independently of the base cutoff. The logarithm of the determinant of the Zeeman block, $\ln(\sigma^2 - 4f_m^2) + \ln(\sigma^2 + 4f_e^2)$, has only integrable logarithmic singularities, at $\sigma = \pm 2f_m$, at the leading symbol; the growth is a property of the Schwinger representation and is not removed by the fibre bound.

Vector and ghost sectors of the invariant sector. The ghost operator is $-\Box$, the same as the scalar of the preceding paragraph. On fibre-constant vector fields $A_c(y)$ the cross term $\eta_{ac}\eta_{bd}\mathcal{J}^{ab}\Sigma^{cd}$ of C_G equals $-2s\mathcal{J}_{ab}A^b$ exactly (symbolic check), so $H_V = -\Box - 2s\mathcal{J}_{ab}A^b + \mu^2$ is $C_G - C_{41} - C_\Sigma + \mu^2$ there. Write the diagonal of $\mathrm{e}^{z\Delta_{H^4} + w\Delta_{H^3}}$ as $D = D_0(1 + D_1 + \dots)$, with D_1 depending on a and homogeneous of degree one in $(z, w) = (t - \mathrm{i}s, \mathrm{i}s)$; $\mathrm{tr}(E + R/6)$ is the first Seeley–DeWitt density of $\mathrm{e}^{t\Delta_{H^4} - \mathrm{i}sH_V}$ in the convention that the exponent is $\Delta + E$, and for one scalar component it equals D_1. For the four components, with the 4×4 matrix connection,

$$\mathrm{tr}(E + \tfrac{1}{6}R) = 4D_1 + \frac{6s^2(1+a^2)}{t + \mathrm{i}sa^2} - 4\mathrm{i}s\mu^2,$$

the spin term being $2s^2\,\mathrm{tr}[G\,\sigma(\Delta_{H^3})] = -2s^2\,\mathrm{tr}[G\,\sigma(C_{31})]$, with G the inverse of the symbol of the exponent, that is the Zeeman square $f_e^2 - f_m^2$ traced against G. The route reproduces the exact scalar coefficient D_1 to 120 digits at three points, and the vector and Dirac traces were verified to the same accuracy; the vector kernel itself was not computed. With $x = s/t$ and the subtraction at $x = 1$ of the preceding paragraph, the weight-two functions are $Q_\mu = -\mu^2 j(a)$ with $j(a) = 2(1 + \mathrm{i}/a)/(1 + a^2)$, and $Q_{\mathrm{sp}} = -8/(1 + a^2) - \mathrm{i}\,4(3a^2 + 1)/(a^3(1 + a^2))$ for the spin term, so $\mathrm{Re}[Q_{\mathrm{sp}} + 4Q_\mu] = 16/(1 + a^2)$ for $\mu^2 = -3$. For n_{ex} further scalars of mass μ_i^2 the bosonic content has

$$\mathrm{Re}\,Q_{\mathrm{bos}} = (2 + n_{\mathrm{ex}})\,\mathrm{Re}\,Q + \frac{16 - 2n_{\mathrm{ex}}\mu_i^2}{1 + a^2},$$

plus the scale-dependent term proportional to $2 + n_{\mathrm{ex}}$ of the preceding paragraph. The first term is indefinite and the second has the sign of $8 - n_{\mathrm{ex}}\mu_i^2$; the scale-dependent term has the sign of a logarithm. On the full leaf the spin term in a Gaussian window has a large-s tail $C\,\mathrm{d}s/s$ with C purely imaginary and proportional to $\sigma(C_{31})$ on the kernel plane, that is to $(1 - a^2)/(1 + 2a^2)$, so its logarithmic coefficient is of indefinite

sign on the leaf, and its real part depends on the contour angle.

Supertrace. The matrix part of the coefficient is traceless, so it enters only through $\operatorname{tr}\Sigma^{ab}\Sigma^{cd}$, and the species-independent part $R/6+E$ is multiplied by the number of components. For the Dirac representation $\operatorname{tr}\Sigma_D\Sigma_D = \frac{1}{2}\operatorname{tr}\Sigma_V\Sigma_V$ for all index pairs. Supply, as an input not contained in the class, the sixteen real fermion components of the ten-dimensional supersymmetric matrix model with the square of the Dirac operator equal to $\Box + s\Sigma^{ab}\mathcal{J}_{ab}$ on each component, no bilinear mass, and $n_{\mathrm{ex}} = 6$, and weight $-\frac{1}{2}$ per real fermion component. Then the component count $4+6-2-8$ vanishes, the weighted index $1-\frac{1}{2}\cdot 4\cdot\frac{1}{2}$ vanishes, the Zeeman character is $4(\cos sf_m - \cosh sf_e)^2 = s^4(f_e^2+f_m^2)^2+\dots$ (the order s^4 is a sum of the squares of the two invariants and does not vanish for any non-zero flux in Lorentzian signature), and what remains of the weight-two coefficient in the invariant sector is $-(4\mu^2+n_{\mathrm{ex}}\mu_i^2)j(a)$, the mass supertrace times the mass moment $j(a)$, with $\operatorname{Re} j = 2/(1+a^2) > 0$; no sign of the mass supertrace itself is claimed. The mode-count term and the spin term of the order F^2 both cancel identically; the curvature and connection part of the weight-two term therefore does not by itself carry the Newton coefficient for this content; whether it arises at higher order of the expansion or through the dependence on A is not computed.

This is a statement for the Poisson limit, one loop, the leading symbol and the invariant sector, the four-matrix class with $\mu^2 = -3$, Gaussian windows and the cone window of the stated form, the sampled points and the supplied fermion content with no bilinear fermion mass and the same window for bosons and fermions. The continuation of the shell identity to spacelike p with $|p| \geq R$ is a numerical observation, and the sign of the cone-window moment is conditional on it; the vector kernel and the second variation under A are not computed; fermions are not part of the class; the Jacobian ϱ and the Liouville normalisation are not computed as functions of position. It does not treat Moyal terms or a window built from dynamical matrices, it identifies no induced metric with the emergent metric of Chapter 4 and, since a Euclidean coefficient is not a Lorentzian one, leaves Assumption 4.6 in place (Problem E). This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: the supertrace at an arbitrary matrix fluctuation, and the species constants that can multiply an Einstein–Hilbert-type term (conditional; bounded no-go). Keep the setting of the preceding paragraph: Poisson limit, leaf $\mathbb{CP}^{1,2}$, $T^a = M^{a4}$, background gauge function, ghost, vector with $\mu^2 = -3$, n_{ex} further scalars of mass μ_i^2, the supplied sixteen real fermion components of weight $-\frac{1}{2}$ with no bilinear fermion mass, and one window function χ of the commuting operator Δ_κ for all species. Let X^a be arbitrary functions on the leaf, $D_a = \{X_a,\cdot\}$, $F_{ab} = \{X_a, X_b\}$, $\mathcal{Z}_{ab} = \{F_{ab},\cdot\}$ and $\Box_X = \eta^{ab}D_aD_b$.

Operators at arbitrary X. The Jacobi identity gives $[D_a, D_b] = \mathcal{Z}_{ab}$. The second

variation of the quartic term plus the bilinear term about X is $\frac{1}{2}\int \varepsilon^a L_a$ with $L_a = -\Box_X \varepsilon_a + D_a D_b \varepsilon^b - 2\{F_{ab}, \varepsilon^b\} + \mu^2 \varepsilon_a$; in the background gauge $D_a \varepsilon^a$ the ghost operator is $-\Box_X$, the further scalars at vanishing transverse matrices have $-\Box_X + \mu_i^2$, and the square of the Dirac operator is $\mathcal{D}^2 = \Box_X + \frac{1}{2}\Gamma^{ab}\mathcal{Z}_{ab}$ (these three identities were checked symbolically for random polynomial X on a four-dimensional phase space). Every species is therefore $H_i = (-\Box_X) \otimes 1 + Z_i + m_i$ with $Z_i = \Sigma_i^{ab} \otimes \mathcal{Z}_{ab}$, the same operator $-\Box_X$ and the same $\mathcal{Z}_{ab}$ for all species, constant masses m_i (μ^2, μ_i^2, and zero for the ghost and the fermions), and constant matrices Σ_i^{ab} acting on the species index (for the vector, minus the generator of the vector index, which gives $-2\{F_{ab}, \varepsilon^b\}$; for the fermions, minus $\frac{1}{4}[\Gamma^a, \Gamma^b]$ on four copies of the real four-component spinor, because the operator compared with $-\Box_X$ is $-\mathcal{D}^2$; the traces satisfy $\operatorname{tr}\Sigma_D\Sigma_D = \frac{1}{2}\operatorname{tr}\Sigma_V\Sigma_V$ for all index pairs).

Species sums. With weights w_i and dimensions d_i ($1 \cdot 4$, $1 \cdot n_{\mathrm{ex}}$, $-2 \cdot 1$, $-\frac{1}{2} \cdot 16$) one has, as exact tensor identities checked for all index tuples, $\sum_i w_i d_i = 0$ and $\sum_i w_i \operatorname{tr}(Z_i^{a_1 b_1} \cdots Z_i^{a_n b_n}) = 0$ for $n = 1, 2, 3$ (not only for the totally symmetrised products); for $n = 4$ the symmetrised sum is the s^4 coefficient $(f_e^2 + f_m^2)^2$ of the character of the preceding paragraph and does not vanish. Because Σ_i acts on the species index only, the term of any Duhamel expansion of $\sum_i w_i \operatorname{Tr}[\chi\, \Phi(H_i)]$ in powers of m_i and Z_i is a species constant times a species-independent operator trace; no commutation of χ, $\mathcal{Z}_{ab}$ and $\Box_X$ is needed. Hence, for every X and every Φ, the terms without masses and with at most three insertions of $\mathcal{Z}_{ab}$ cancel identically (a check with random non-commuting matrices confirms this), and the term linear in the masses and without insertions of $\mathcal{Z}_{ab}$ is $+M_{\mathrm{str}} \operatorname{Tr}[\chi\, \Phi'(H_X)]$ (for $\Phi = e^{-(\cdot)}$ this is $-M_{\mathrm{str}} \operatorname{Tr}[\chi\, e^{-H_X}]$) with $M_{\mathrm{str}} = \sum_i w_i d_i m_i = 4\mu^2 + n_{\mathrm{ex}}\mu_i^2$. The other constants that survive at the next order are $\sum_i w_i d_i m_i^2$, $\sum_i w_i m_i \operatorname{tr}\Sigma_i^{ab}\Sigma_i^{cd}$ (the vector mass alone) and the quartic constant.

Consequences. (i) The weight-two coefficient of the preceding paragraph is, for every X and not only at $X = T$, the mass-supertrace term: no count term $\propto R$, no spin term quadratic in $\mathcal{Z}_{ab}$ and no term cubic in it appears, whatever the dependence of $\Box_X$ and $\mathcal{Z}_{ab}$ on A. (ii) At the leading symbol that term is proportional to $M_{\mathrm{str}} \int \chi\, (\sigma_X - \mathrm{i}0)^{-1}$ with $\sigma_X = \eta_{ab}\, \xi(D^a)\xi(D^b)$, which depends on X through $\mathrm{d}X^a$ only. For $A^a = \alpha^a \cos\phi$, $\phi = q \cdot y$, with constant α^a, the Hamiltonian field of X^a is $W^a - \epsilon \sin\phi\, \alpha^a Y_q$, where W^a is that of T^a and $Y_q = q_\mu \{y^\mu, \cdot\}$ (chain rule; the first-order change of the field of F_{ab} was checked through the Poisson algebra for the linear ansatz $A^a = \alpha^a\, q \cdot y$). This leading-symbol term depends on A only through its first derivatives, so its second variation contains no second derivative of the fluctuation and no term of Einstein–Hilbert form; the next-to-leading symbol, which involves the second jet of X, is not computed. With the Gaussian window of $\Delta_{3/4}$ the Hessian of the real (principal-value) part of this moment along α, at frozen coefficients and $\phi = \pi/2$, was evaluated numerically in 59 random draws of orbit point and q: it is indefinite

in 40, positive definite in 5 and negative definite in 14, and along $\alpha^a \propto q^a$ it is numerically nonzero in all draws with non-null q. (iii) A term of Einstein–Hilbert form can therefore enter only through the constants listed above. The one proportional to $M_{\rm str}$ is the mass-insertion term of the weighted a_2 in the Euclidean class (there a shift of the operator of species i by m_i changes the weighted a_2 by $-m_i a_1^{(i)}$, with curvature part $-\frac{1}{6}M_{\rm str}\,\rho R[g_\rho]$ and connection part $-(\sum_i w_i\kappa_i m_i)\rho^2 X$, κ_i being the weight of $\rho^2 X$ of the species; the second constant above is the vector mass times κ_V); its value in the six-dimensional leaf requires the next-to-leading symbol and is not computed. The pure flux constant is the order-F^4 term: at the phase $\phi = 0$, where the first-jet terms vanish, and at the leading order in q, the flux in the leading symbol is shifted as $F_{ab}(\xi) \to F^0_{ab}(\xi) + \epsilon\, c\,(\xi\cdot Y_q)(q_a\alpha_b - q_b\alpha_a)$, $c = -s\,y^4\cos\phi$, and at fixed proper time and $\kappa = \frac{3}{4}$ the Gaussian averages in the integrand depend on $q\cdot y$ at fixed $q\cdot q$, through the vector y^μ and the window (at an orbit point with $y^i = 0$, $\langle(\xi\cdot Y_q)^2\rangle$ changes by a factor above 2 over a boost of rapidity 1.2 of a unit q); its integrand is therefore not Lorentz invariant (the window and the leaf both single out y^μ, and this diagnostic does not separate them). (iv) No finite coefficient of this term follows in the present window: at four orbit points the integrand of the order-F^4 term at $A = 0$ grows as s^2 at large proper time (log-slope 2.00), because the flux does not vanish on the kernel plane of $\sigma(\Box)$; the expansion in $\mathcal{Z}_{ab}$ is not the expansion in a small parameter there, and a sign of an Einstein–Hilbert-type coefficient does not follow. The non-expanded leading-symbol logarithm is $\sum_i w_i \ln\det(\sigma + Z_i) = \ln(\sigma^2 - 4f_m^2) + \ln(\sigma^2 + 4f_e^2) + 4\ln\sigma - 2\ln[(\sigma+f_m)^2 + f_e^2] - 2\ln[(\sigma - f_m)^2 + f_e^2]$, whose large-$\sigma$ series starts with $-6(f_e^2 + f_m^2)^2/\sigma^4$ (checked); its second variation under A is not computed.

This is a statement for the Poisson limit, one loop, the commuting-symbol (leading) level, the four-matrix class with $\mu^2 = -3$, the supplied fermion content with no bilinear fermion mass and the same window for all species, applied term by term in an expansion that is not shown to converge. It does not compute the next-to-leading symbol, the dependence on A of the terms of Einstein–Hilbert form, the case of transverse matrices with flux or of species-dependent windows, Moyal terms, or a window built from dynamical matrices; it identifies no induced metric with the emergent metric of Chapter 4 and gives no value or sign of $G_{\rm eff}$. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: the next-to-leading symbol of the mass-supertrace term at order ϵ^2 (conditional; bounded no-go with a conditional Einstein–Hilbert-form limit). Keep the setting of the preceding paragraph and the term $+M_{\rm str}\,\mathrm{Tr}[\chi\,(H_X - \mathrm{i}0)^{-1}]$, $H_X = -\Box_X$. Take $A^a = \epsilon\alpha^a\cos\phi$, $\phi = q\cdot y$, W_a the Hamiltonian field of T^a, $Y = q_\mu\{y^\mu,\cdot\}$, $\kappa_a = W_a(\phi)$ (so $Y(\phi) = 0$) and the Gaussian window of $\Delta_{3/4}$. Terms of fourth order in q at order ϵ^2 contain all derivatives on A; they are

computed with the first jet of T frozen and the window symbol constant, since window gradients and $\partial^2 T$ carry fewer powers of q (derivative counting; the window symbol being constant makes the composition terms with χ total derivatives). In this sector $H_X = \eta^{ab}\Pi_a\Pi_b$, $\Pi_a = \omega_a - \epsilon\alpha_a\zeta\sin\phi$, $\omega_a = \xi(W_a)$, $\zeta = \xi(Y)$, $[\omega_a, \phi] = -\mathrm{i}\kappa_a$, with ζ commuting with everything.

Symbol and checks. The weight-two symbol of the resolvent was derived by the left-symbol recursion, expanded to ϵ^2 and averaged over ϕ (the weight-one terms average to zero). It reproduces (a) the one-dimensional coefficient $-\frac{1}{12}V'^2$ and the two-dimensional Maxwell coefficient $-\frac{1}{12}F_{ab}F^{ab}$ to 10^{-11}; (b) the leading-symbol Hessian of the preceding paragraph; (c) in the non-degenerate case, with the vector fields divergence free, the Seeley–DeWitt coefficient $\sqrt{|g|}(R/6 - \frac{1}{4}(\nabla\lambda)^2)$, $\lambda = \ln\sqrt{|g|}$, at second order (Fierz–Pauli form), with relative phase i to 12 digits.

Result in the class. The integrand $K(s)$ of the proper time s is finite at small s and has at large s the form $c_0 + c_1/s + O(s^{-2})$, checked at orbit points with random q, α. (i) c_0 is real and is the Maxwell coefficient of the gauge field $a_b = \epsilon\zeta\alpha_b\sin\phi$ induced by the kernel momenta: $c_0 = \lambda_0\, G(\alpha)$ with $G = (\kappa\cdot\kappa)(\alpha\cdot\alpha) - (\kappa\cdot\alpha)^2$ (agreement to 10^{-8} or better in the tensor, $\lambda_0 > 0$ at all sampled points). It is gauge invariant ($\alpha \propto \kappa$), of the sign of G, which is indefinite in Lorentzian signature, and the integral $\int^{s_{\max}} \mathrm{d}s\, c_0$ diverges linearly: the first-order mass term has no finite local coefficient at this order, and the derivative expansion is expected to be non-uniform at small ξ of the non-kernel directions (a mass insertion into the massless non-kernel propagator; not computed). (ii) c_1 is purely imaginary and, for the window of $\Delta_{3/4}$ at the sampled points, is neither invariant under $\alpha \to \alpha + \lambda\kappa$ (relative violation 0.04–0.6) nor of covariant form, and has eigenvalues of both signs. (iii) Let V' be the vertical tangent space (common kernel of the $\mathrm{d}y^\mu$) and scale by f the part of the window quadratic form in ω_a in the splitting $\xi \mapsto (\xi(W_a), \xi(v_k))$. As $f \to 0$, $c_1 \to \mathrm{i}\lambda_1 A$ with $\lambda_1 = |\det\Lambda|^{-1}(4\pi)^{-3}\det(P'_{kk})^{-1/2} > 0$ (Λ the matrix of the splitting, P'_{kk} the vertical block of the window form) and no free parameter; the difference is $O(f)$ at four orbit points (relative $8\cdot10^{-4}$, $1.4\cdot10^{-3}$, $7\cdot10^{-4}$, $8\cdot10^{-5}$ at $f = 10^{-3}$). Here, with the base cometric $\bar g^{\mu\nu} = \eta^{ab}W_a(y^\mu)W_b(y^\nu)$, $h^{\mu\nu}$ its first-order change $-\sin\phi\,(u^\mu v^\nu + v^\mu u^\nu)$, $u^\mu = \eta^{ab}\alpha_a W_b(y^\mu)$, $v^\mu = \{\phi, y^\mu\}$ (so $q\cdot v = 0$) and $a_\mu = \bar g_{\mu\nu}u^\nu$,

$$A = \tfrac{1}{24}\Big[v^2\big((q\cdot a)^2 - q^2a^2\big) - 2q^2(a\cdot v)^2\Big],$$

the sum of the Einstein–Hilbert second variation, with the phase-averaged Fierz–Pauli form $\frac{1}{6}\,\mathrm{FP}[h] = \frac{1}{24}\big[v^2\big((q\cdot a)^2 - q^2a^2\big) + q^2(a\cdot v)^2\big]$, and the dilaton-gradient term $-\frac{1}{8}q^2(a\cdot v)^2$ of the Liouville measure. A is invariant under $a \to a + \lambda q$ (checked to 10^{-16}), vanishes only on that direction, and is indefinite in general (definite only for q time-like and v space-like, where it is positive). (iv) The base two-form $\theta^{\mu\nu} = \{y^\mu, y^\nu\} = M^{\mu\nu}$ satisfies $\theta^{\rho\mu}\eta_{\mu\nu}\theta^{\sigma\nu} = c_\theta\eta^{\rho\sigma} + y^\rho y^\sigma - t^\rho t^\sigma$, $y^\mu = M^{\mu5}$, $t^\mu = M^{\mu4}$

(to machine precision at six points), with c_θ a scalar function of the orbit point, so $v^2 = c_\theta q^2 + (q\cdot y)^2 - (q\cdot t)^2$ depends on the time-like vector y^μ and on the fibre point. After a Haar average over the fibre at a base point with $y^i = 0$, the A-form (weighted by the Liouville density and λ_1) remains gauge invariant but is not proportional to $q^2\eta_{ab} - q_a q_b$ (remainder 0.8–1 of its norm), and its covariant part changes by a factor 2 between boosts of rapidity 0 and 0.8 at fixed $q\cdot q$. The window and the leaf both single out y^μ, which this check does not separate.

What is not shown. The Pauli–Sakharov completion (the term $M_{\rm str}$ multiplying the logarithm of the cutoff is accompanied by $-\sum_i w_i d_i m_i \ln m_i$ when the masses are kept non-perturbatively) is expected from the Euclidean heat-kernel structure and is not derived for this hypoelliptic operator. The background flux, which cuts off large s in the full leaf, is not included, so the divergence in (i) is not resolved.

This is a statement for the Poisson limit, one loop, the term linear in the masses, the order $\epsilon^2 q^4$ with the first jet frozen, the supplied content with no bilinear fermion mass, the Gaussian window of $\Delta_{3/4}$, and numerical sampling; the limit in (iii) is a limit of the window family and not the window of the class. It identifies no induced form with the emergent metric of Chapter 4, makes no Wick-rotation claim, and gives no value or sign of $G_{\rm eff}$. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: exact Landau (nilpotent) resummation of the leading symbol in the background flux, and its effect on the mass-supertrace term and on the massless resummed character term (conditional; bounded no-go). Keep the setting of the preceding paragraphs: Poisson limit, leaf $\mathbb{CP}^{1,2}$, $T^a = M^{a4}$, $H_X = -\Box_X$, the supplied content with no bilinear fermion mass, $M_{\rm str} = 4\mu^2 + n_{\rm ex}\mu_i^2$, and the nilpotent (leading) symbol, in which the horizontal momenta $\omega_a = \xi(W_a)$ obey $[\omega_a,\omega_b] = \mathrm{i}F_{ab}(k)$ with F_{ab} the constant flux at the centre variable k of the kernel plane of $\sigma(\Box)$; ηF has eigenvalues $\pm f_e$ and $\pm\mathrm{i}f_m$. At four sampled orbit points the ratio f_e/f_m on the kernel plane equals $\ell/|y^4|$ in every direction of k to five digits, which is the closed form of the kernel-plane flux above. Write $a = f_e$, $b = f_m$, $x = sa$, $y = sb$.

Orbital factor. For $H_0 = \eta^{ab}\omega_a\omega_b$ the Gaussian (KMS) two-point function $G = -\mathrm{i}FM^T(1-M^T)^{-1}$, $M = \mathrm{e}^{2sF\eta}$, gives, per unit horizontal volume, $\Theta(s) = \mathrm{Tr}\,\mathrm{e}^{-\mathrm{i}sH_0} = -\mathrm{i}\,ab\,(16\pi^2\sinh x\,\sin y)^{-1}$, with $s\to s-\mathrm{i}0$ (the poles of $1/\sin y$ lie above the contour, as for the Landau-level sum $\frac{b}{2\pi}\sum_n \mathrm{e}^{-\mathrm{i}sb(2n+1)}$). The identity $\mathrm{d}\ln\Theta/\mathrm{d}s = -\mathrm{i}\eta^{ab}G_{ab}$ holds to 10^{-10} at complex s, and $\Theta\to -\mathrm{i}/(16\pi^2 s^2)$ for $s\to 0$. The electric pair gives $1/\sinh$ and the magnetic pair $1/\sin$.

Mass-supertrace term. The term $+M_{\rm str}\,\mathrm{Tr}[(H_X-\mathrm{i}0)^{-1}] = M_{\rm str}\,\mathrm{i}\int_0^\infty \mathrm{d}s\ \mathrm{Tr}\,\mathrm{e}^{-\mathrm{i}sH_X}$

becomes, per unit horizontal volume and per centre point, $M_{\rm str}(16\pi^2)^{-1}(s_0^{-1}+g)$ with

$$g(a,b)=\int_0^\infty \mathrm{d}s\Big[\frac{ab}{\sinh(as)\,\sin(bs-\mathrm{i}0)}-\frac{1}{s^2}\Big]=\pi\big[\,b\,S(b/a)+\mathrm{i}\,a\,S(a/b)\big],\qquad S(r)=\sum_{n\geq 1}\frac{(-1)^n}{\sinh(n\pi r}$$

The integrand is analytic in the open fourth quadrant and bounded by $\exp[-|s|(a\cos\vartheta+b\sin\vartheta)]$ on the ray $s=|s|\mathrm{e}^{-\mathrm{i}\vartheta}$, so g is finite for $a,b>0$ and independent of ϑ; the closed form follows from the two residue sums (poles of $1/\sinh$ at $s=-\mathrm{i}n\pi/a$, poles of $1/\sin$ at $s=n\pi/b$) and was confirmed against the ray integral to 10^{-24} at ten pairs (a,b). g is homogeneous of degree one, $\mathrm{Re}\,g<0$ and $\mathrm{Im}\,g<0$ for all $a,b>0$, $\mathrm{Re}\,g\to -a\ln 2$ for $b\ll a$ and $\mathrm{Im}\,g\to -b\ln 2$ for $a\ll b$. Thus the linear divergence in s of the Maxwell tail c_0 of the preceding paragraph is cut off at $s\sim 1/f$, and the flux supplies an exponentially convergent infrared end of the proper-time integral.

Completeness of the $M_{\rm str}$ term. With the flux kept to all orders, the mass-linear term is not $M_{\rm str}\,\mathrm{Tr}[(H_X-\mathrm{i}0)^{-1}]$ alone: the vector mass multiplies the Zeeman-dressed resolvent, and the complete term is $\mathrm{i}\int\mathrm{d}s\,\Theta(s)\big[M_{\rm str}+\mu^2(\chi_V-4)\big]$, $\chi_V=2\cosh 2x+2\cos 2y$. The second part is the all-order version of the constant $\sum_i w_i m_i\,\mathrm{tr}\,\Sigma_i\Sigma_i$ listed above and has the growth of the next paragraph.

Massless resummed character term. With $\chi_{\rm str}=4(\cos y-\cosh x)^2$ (the species sum $2\cosh 2x+2\cos 2y+6-2-8\cosh x\cos y$, identity checked to 10^{-40}) the resummed integrand is $\Theta\chi_{\rm str}/s$. On the ray of angle ϑ one has $|\Theta\chi_{\rm str}|\sim\exp\big[|s|\,|a\cos\vartheta-b\sin\vartheta|\big]$ (measured rates agree to $4\cdot 10^{-3}$ at four angles away from ϑ_*), and $\Theta\chi_{\rm str}\to(ab/\pi^2)\sinh^2[s(a-\mathrm{i}b)/2]$ at large $|s|$ (ratio 1 to 10^{-8}). The exponential growth therefore occurs on the real axis (the $\mathrm{i}\epsilon$ contour) for every $a>0$, the same exponent holds for $\Theta(\chi_V-4)$, and the massless exponent vanishes only on the single ray $\vartheta_*=\arctan(a/b)$, which depends on the flux; on the leaf $\vartheta_*=\arctan(\ell/|y^4|)$. The exponent is unchanged by the phases $\mathrm{e}^{-\mathrm{i}sm_i}$ of real masses on the real axis; on a ray of angle ϑ a mass m shifts it by $-m\sin\vartheta$, so for the vector with $\mu^2=-3$ the rate is $|a\cos\vartheta-b\sin\vartheta|+3\sin\vartheta>0$ on every ray, whereas a positive mass-squared would open a window of convergent rays about ϑ_* for the massless term treated here. On that ray the integrand is bounded, but it tends to the constant $-ab/(2\pi^2)$ plus oscillations, so $\int\mathrm{d}s/s$ grows logarithmically in the real part (numerically -0.032 per doubling of the upper limit at $a=0.7$, $b=1.3$; the imaginary part converges). For the massless term no ray of the closed fourth quadrant gives a convergent integral (rays of the first quadrant correspond to the opposite $\mathrm{i}\epsilon$ prescription and are neither claimed nor excluded), and with $\mu^2=-3$ the same holds for the mass-dressed term, so the massless resummed character term has no finite value from the proper-time representation of the class; its non-analytic dependence on f_e is that of a spectral sum over the complex eigenvalues $\lambda\pm 2\mathrm{i}f_e$ of the vector sector and is not computed.

Dependence on A. The leading nilpotent coefficient of the $M_{\rm str}$ term depends on

A only through the centre-valued flux, so its second variation is $\frac{1}{2}D^2g\,[Q,Q]$ with $Q_{ab}=\zeta_Y(k)(\kappa_a\alpha_b-\kappa_b\alpha_a)$; it contains no second derivative of the fluctuation and no term of Einstein–Hilbert form; this concerns the $M_{\rm str}$ term only, since the Zeeman-dressed part has no contour. It rests on the frozen nilpotent frame in which Y is central, and on the leaf $[W_a,Y]\neq 0$ at order q by the Jacobi identity; the same omission is made in the preceding frozen-frame paragraph. At four orbit points, averaging over the direction of k with the window of $\Delta_{3/4}$, this form is gauge invariant ($\alpha\propto\kappa$ in its kernel) but is not proportional to $(\kappa\cdot\kappa)\eta^{ab}-\kappa^a\kappa^b$ (relative residual 0.4–1.0 in real and imaginary parts at the points with $0.3\le f_e/f_m\le 0.8$), and its eigenvalues on the quotient by κ have both signs. The infrared-finite replacement of the Maxwell tail of the $M_{\rm str}$ term is therefore not of the covariant Maxwell form and is not sign-definite (leaf average, four points).

Einstein–Hilbert sector in a four-dimensional Landau model. Let $\Omega_a=\omega_a+\epsilon\sin\phi\,(m_a{}^c\omega_c+n_a)$, $\phi=\kappa\cdot x$, $m_a{}^c\kappa_c=0$, on the algebra above with F constant; the choice $m_a{}^c=-\alpha_a y^c$, $n_a=-\alpha_a c$ reproduces the sector of the preceding paragraph with the commutators $[\omega_a,\omega_b]=\mathrm{i}F_{ab}$ kept, but without the commutator of Y with W_a that the Jacobi identity fixes on the leaf, so it is a model and not the leaf (an omission shared with the frozen-frame paragraph above). The second order in ϵ of $\operatorname{Tr}\mathrm{e}^{-\mathrm{i}sH}$ was computed exactly by Wick's theorem in the state $\mathrm{e}^{-\mathrm{i}sH_0}$. It reproduces the Maxwell coefficient at $F=0$ (ratio 1 to 10^{-10}), the form A of the preceding paragraph at $F=0$ ($c_1/A=\mathrm{i}$ to 10^{-8} in five random draws, including the dilaton term), gauge invariance in n at $F\neq 0$ ($2\cdot10^{-15}$), and the constant-field limits (Hessian of Θ in F, and $|\det M|^{-1}\Theta(MFM^T)$ for a constant vielbein). At $F\neq 0$ the coefficient $c=A/16\pi^2$ of $\mathrm{d}s/s$ is unchanged, the integrand decays like $\mathrm{e}^{-|s|(a\cos\vartheta+b\sin\vartheta)}$, and the finite part $\mathrm{Fin}=\lim[\int_{s_0}^{\infty}T_2\,\mathrm{d}s-c\ln(1/s_0)]$, in units of c, is complex; its real part is about 7.6, 8.1, 6.8 in three Lorentz frames of F at $(a,b)=(0.7,1.3)$ (data κ,α,y rotated, A fixed; imaginary parts between -0.32 and 0.20), and 7.6, 5.9, 7.3 in three frames at $(a,b)=(1.92,1.0)$ (imaginary parts 1.05–1.28). The flux therefore turns the infrared end of $\int\mathrm{d}s/s$ into a finite remainder that depends on the orientation of F relative to κ,α,y: at fixed (a,b) and fixed A (the meaningful comparison, since Fin/c shifts by a constant under a rescaling of s) the real part varies between frames by roughly 20%, and the Maxwell part of the same sector changes between frames by about 30–45% at $(a,b)=(0.7,1.3)$ and by a factor of order one at $(1.92,1.0)$ (complex values). The coefficient of the logarithm is the flux-independent A of the preceding paragraph.

This is a statement for the Poisson limit, one loop, the nilpotent (leading) symbol resummed in a constant flux at the centre variable, the Lorentzian $\mathrm{i}\epsilon$ proper time, the supplied content, and numerical evaluation at sampled points; the closed form of g is checked numerically, not proved, and the Landau model for the Einstein–Hilbert sector is not the leaf. It does not include the next-to-leading symbol of the hypoelliptic

operator with the commutators of Y, the dependence of the window on the flux, non-perturbative masses, or the spectral definition of the vector sector, and it identifies no induced form with the emergent metric of Chapter 4, makes no Wick-rotation claim, does not consider the first-quadrant rays, and gives no value or sign of G_{eff}. This changes no ledger label; Problem F remains open.

Conjugate-coordinate class: the Jacobi commutator of the plane-wave field with the horizontal fields, and the flux-resummed next-to-leading symbol (conditional; bounded no-go; supersession of the preceding Landau model). Keep the setting of the preceding paragraphs: Poisson limit, leaf $\mathbb{CP}^{1,2}$, $T^a = M^{a4}$, W_a the Hamiltonian field of T^a, $Y = q_\mu\{y^\mu, \cdot\}$, $\phi = q \cdot y$, $\kappa_a = W_a(\phi)$, $A^a = \epsilon\alpha^a \cos\phi$, the nilpotent (leading) symbol with centre flux $[\omega_a, \omega_b] = \mathrm{i}F_{ab}$, $\omega_a = -\mathrm{i}W_a$, and the term $+M_{\text{str}} \operatorname{Tr}[(H_X - \mathrm{i}0)^{-1}]$. The Lie–Poisson relations give exactly $\kappa^a = \sigma M^{45} q^a$ and $[W_a, Y] = \sigma q_a \operatorname{Ham}(y^4)$, $y^4 = M^{45}$, with $\sigma = \pm 1$ the sign convention of the bracket (the second is the Jacobi identity; both checked at random orbit points to 10^{-14}). In the frame (W_a, Z_k), Z_k a basis of the vertical space, $Y = y^c W_c + y^k Z_k$ and the field $V = \operatorname{Ham}(y^4)$ has no vertical component (below 10^{-14} at six sampled points), so $[W_a, Y]$ has no component along the centre. The flux term $y^c F_{ac}$ is not small and is not parallel to κ_a (the residual of the parallel fit is 0.7–1.0 at the sampled points and directions of the centre variable).

Consequence for the model of the preceding paragraph. There the plane-wave coupling was $\zeta_Y = y^c\omega_c + c$ with a constant charge c, for which $[\omega_a, \zeta_Y] = \mathrm{i}y^c F_{ac} \neq 0$; this commutator is inconsistent with the Jacobi identity at the nilpotent order with frozen coefficients. That model is not invariant under $\alpha \to \alpha + \lambda\kappa$, the leaf form of $\delta A^a = \{\Lambda, T^a\}$, at $F \neq 0$ and $y \neq 0$ (relative change 0.07–0.16 of the second-order response in a test, 10^{-14} at $F = 0$), whereas the Fin values quoted there are values of that model. On the leaf Y commutes with ω_a and with ϕ ($Y(\phi) = 0$); at the nilpotent order it is a right-invariant field of the nilpotent group, and its diagonal value at the base point is $P = y^c\omega_c + c$, $c = y^k\zeta_k$, which stands at the seam of the trace, $\operatorname{Tr}[\text{chain} \cdot P^2]$, the chain being the ordered product of the left-invariant factors. Because P^2 separates the first and the last propagator, the time-translation reduction $\int_0^1 \mathrm{d}u\,(1-u)\langle V(su)V(0)\rangle$ is not available and the second order is the double Duhamel integral $-s^2 \int_{0<u_2<u_1<1} \langle V(su_1)V(su_2)P^2\rangle$.

Checks. In a two-dimensional Euclidean Landau problem in Landau gauge, where the magnetic translation $K = y_2 p_2$ is an exact c-number, this prescription reproduces the second-order density obtained by exact diagonalisation (relative difference $2 \cdot 10^{-4}$, real); the prescription with P^2 at the end of a merged propagator does not (complex, 1.5% off in the real part). The two-dimensional test validates the prescription for a central K; the identification of the leaf field Y with the seam insertion P is conditional on the nilpotent order and the frozen first jet. In four dimensions the central model

is invariant under $\alpha \to \alpha + \lambda\kappa$ to 10^{-14} at $F \neq 0$, $y \neq 0$, and equals the model of the preceding paragraph at $F \to 0$ (10^{-8}). On the leaf, with $F_{ab}(\zeta) = -S_{ab}{}^{k}\zeta_k$, ζ the centre variable, $f_e/f_m = \ell/|y^4|$ at three sampled directions of ζ at each of five orbit points, the quadratic form of the Fierz–Pauli plus dilaton coefficient A of the next-to-leading-symbol paragraph equals the closed form in (α, κ, y^c) used there (ratio 1 to six digits, five orbit points).

Result in the class. (i) The coefficient of $\mathrm{d}s/s$ is unchanged: $c = A/16\pi^2$, with $T_2(s)\, s/c \to 1.0000$ as $s \to 0$ at the flux of the leaf, in both the central and the preceding model. The Jacobi commutator therefore does not change, cancel or add to the Einstein–Hilbert-form coefficient A of the logarithm. (ii) At $F \neq 0$ the integrand decays as $\mathrm{e}^{-|s|(f_e\cos\vartheta + f_m \sin\vartheta)}$ on $s = |s|\mathrm{e}^{-\mathrm{i}\vartheta}$, and the finite part $\mathrm{Fin} = \lim[\int_{s_0}^{\infty} T_2\, \mathrm{d}s - c\ln(1/s_0)]$ is a finite complex number for each centre variable. Split by the content of P into $y^c y^d$ (tetrad), c^2 (charge) and cross terms, the three parts are separately invariant under $\alpha \to \alpha + \lambda\kappa$ (10^{-7} or better). (iii) The tetrad part of Fin, as a quadratic form in α, is not proportional to the form A: the residual of the best proportional fit, in real and imaginary parts, is 0.54 and 0.85 (centre direction 0.8), 0.08 and 0.48 (direction 2.0) at one orbit point, and 0.004 and 0.000 at a second point where $f_e/f_m \simeq 42$; the quotient by κ has eigenvalues of both signs in every case (as has A). (iv) The form A is not Lorentz covariant, as shown for the preceding paragraph (it depends on y^μ and on the fibre point), and the flux does not alter the coefficient that carries this dependence. In the sampled per-centre-variable forms the tetrad part of the finite remainder is neither proportional to the form A nor sign-definite; the sign and covariance of the full finite remainder (tetrad, charge and cross parts) and of its window average over the centre variable were not tested, so no finite, covariant, sign-definite Einstein–Hilbert coefficient is obtained here.

Size of the dropped terms. The horizontal part of $[W_a, Y]$ (the weight-one term $-q^a v^c W_c$), the horizontal parts of $[W_a, W_b]$, the gradients of the flux and of y^c, y^k, and the window are not included. The ratio of the norm of the horizontal part of $[W_a, Y]$ to that of $y^c S_{ac}{}^{k}$ is 0.01–0.5 at six orbit points (per unit centre variable), so these terms are small only for $|\zeta|$ large compared with that ratio and for small ω; this is the regime of the nilpotent approximation, not a bound on the leaf.

This is a statement for the Poisson limit, one loop, the nilpotent (leading) symbol resummed in the flux with the Jacobi-consistent central plane-wave field, the Lorentzian $\mathrm{i}\epsilon$ proper time, the M_{str} term, and numerical evaluation at sampled orbit points and centre directions. It does not include the vector-mass term with the Zeeman-dressed resolvent at order ϵ^2 (whose leading growth was found above and is not shown to be altered), the weight-one terms, the window average over the centre variable, the sign and covariance of the full finite remainder, the spectral definition of the vector sector, or non-perturbative masses, and it identifies no induced form with the emergent metric of Chapter 4, makes no Wick-rotation claim, and gives no value or sign of G_{eff}. This

changes no ledger label; Problem F remains open.

H.7 Problem G: HBEV

Ledger label. Status: OPEN. Priority: CRITICAL PATH. Completion: PROGRAMME-CONDITIONAL. Reduced-scope results: G-model and G-external support at most a scope-limited model; G remains OPEN and neither satisfies the matrix-derived criterion.

The boundary functional of Chapter 12 regulates Λ_{eff} once an infrared entropy density is supplied. The working value $\Lambda_{\text{eff}} = 3/L_{\text{IR}}^2$ is a choice consistent with that functional, not a derived necessity.

Status. The mechanism is CLOSED CONDITIONALLY under the stated boundary entropy; derivation of the infrared entropy density and stationarity from the matrix theory is OPEN. A boundary-model or externally justified result is programme-conditional and supports only a scope-limited model.

H.8 Problem H: modular origin of the interpolating function

Ledger label. Status: OPEN. Priority: OPTIONAL within the phenomenological scope. Completion: PROGRAMME-CONDITIONAL. The modular derivation of the interpolating function is not established.

Appendix D tests the function

$$\nu_{\surd}(g_{\text{N}}; g_s) = \frac{1}{1 - \exp\left(-\sqrt{g_{\text{N}}/g_s}\right)}$$

against SPARC with g_s fixed to the empirical value $g_\dagger$. This is a coefficient-level consistency check, not a microscopic derivation. Appendix G gives the standard heat-kernel area contribution but does not compute a source-dependent modular spectrum. Appendix I applies an external-field prescription to wide binaries; it does not derive that prescription from the modular data.

Conditional lemma: occupation of one modular ladder. Suppose the relevant reduced state contains one bosonic ladder with normalized probabilities

$$p_n = (1 - e^{-\Delta k})e^{-n\Delta k}, \qquad n = 0, 1, 2, \ldots, \qquad \Delta k > 0.$$

Then its mean occupation is

$$\langle n\rangle = \sum_{n=0}^{\infty} n p_n = \frac{1}{e^{\Delta k} - 1},$$

and therefore

$$1 + \langle n\rangle = \frac{1}{1 - e^{-\Delta k}}.$$

Thus, if the extra-stress response is exactly $1 + \langle n\rangle$ for this single mode, the target interpolator follows if and only if

$$\Delta k(g_{\mathrm{N}}) = \sqrt{\frac{g_{\mathrm{N}}}{g_s}}.$$

The quantity to calculate is the modular excitation gap Δk, not the absolute lowest eigenvalue of H_D, which depends on an additive normalization constant.

For a multimode response, the corresponding quantity is instead a weighted spectral sum, schematically

$$\mathcal{R}(g_{\mathrm{N}}) = \int_{(0,\infty)} \frac{d\mu_{g_{\mathrm{N}}}(\Delta k)}{e^{\Delta k} - 1},$$

where $d\mu_{g_{\mathrm{N}}}$ includes the mode weights coupling to the extra stress. In that case a square-root gap for one mode does not by itself establish the target function; the measure and its source dependence must also be calculated.

Missing step. Construct the regulated modular operator, or an equivalent well-defined spectral response, for a weak-field causal diamond containing a baryonic source. Derive the relevant excitation gap $\Delta k(g_{\mathrm{N}}, \kappa_{\mathrm{IR}})$ and its spectral weight from the microscopic theory. Fix the normalization relating g_s to κ_{IR} independently of galaxy data. In particular, reconcile the convention $g_s \sim \kappa_{\mathrm{IR}}/(2\pi)$ in Appendix E with the undetermined factor $g_s = c_\star \kappa_{\mathrm{IR}}$ in Chapter 17 before claiming that the Bose occupation yields the target.

Until this calculation is supplied, Appendices D, G, and I provide an empirical fit check, a standard area-term calculation, and an external-field consistency test, respectively; together they do not derive or uniquely select $\nu_{\surd}$. Problem H remains open.

H.9 Problem I: dynamical origin of the drag

Ledger label. Status: OPEN. Priority: OPTIONAL falsifiable phenomenological extension; prerequisite only if drag is claimed as derived or predicted. Completion: PROGRAMME-CONDITIONAL.

Chapter 14 adopts a vector drag proportional to $-\mathbf{v}/d^4$ as a working assumption. The microscopic target is to derive the low-frequency force-response tensor, including its separation and finite-size dependence, from specified dynamics and a specified stationary reference state.

Conditional Kubo formulation. Let $\mathbf{r}$ denote the relative separation and $H(\mathbf{r})$ a microscopic Hamiltonian. At fixed $\mathbf{r}$, choose a stationary reference state and define the connected generalized-force operator

$$\delta\hat{F}_i = -\frac{\partial H}{\partial r_i} - \left\langle -\frac{\partial H}{\partial r_i} \right\rangle_{\mathbf{r}}.$$

For a perturbation $\delta H(t) = -\delta r_j(t)\delta\hat{F}_j$, define

$$\chi^R_{ij}(t;\mathbf{r},\ell_A,\ell_B) = \frac{i}{\hbar}\theta(t)\left\langle[\delta\hat{F}_i(t),\delta\hat{F}_j(0)]\right\rangle_{\mathbf{r}}, \qquad \chi^R_{ij}(\omega) = \int_{-\infty}^{\infty} e^{i\omega t}\chi^R_{ij}(t)\,\mathrm{d}t.$$

For a reciprocal passive reference state, if the low-frequency limit exists, define

$$\Gamma_{ij}(\mathbf{r};\ell_A,\ell_B) = \lim_{\omega\downarrow 0}\frac{\operatorname{Im}\chi^R_{(ij)}(\omega;\mathbf{r},\ell_A,\ell_B)}{\omega}, \qquad \chi^R_{(ij)} = \frac{\chi^R_{ij}+\chi^R_{ji}}{2}.$$

Passivity requires this dissipative tensor to be positive semidefinite. The corresponding low-frequency force contribution is

$$\delta F_i^{\mathrm{diss}} = -\Gamma_{ij}(\mathbf{r};\ell_A,\ell_B)\dot{r}_j.$$

This is a conditional response prescription, not a calculation of the NAMCG force correlator, its reference state, or its passivity.

If the reference state and environment are rotationally invariant, the response at nonzero separation can have distinct longitudinal and transverse components,

$$\Gamma_{ij} = \Gamma_L(d;\ell_A,\ell_B)\hat{r}_i\hat{r}_j + \Gamma_T(d;\ell_A,\ell_B)(\delta_{ij}-\hat{r}_i\hat{r}_j).$$

A radial response calculation determines only Γ_L. The isotropic scalar drag used in the laboratory requires $\Gamma_L = \Gamma_T$, or an explicit restriction to radial motion.

For example, the causal response family

$$\chi^R_{ij,\tau}(\omega) = \frac{\chi_{0,ij}}{1-i\omega\tau}, \qquad \tau > 0,$$

with real symmetric positive-semidefinite $\chi_{0,ij}$, has a pole at $\omega = -i/\tau$ and satisfies

$$\chi^R_{ij,\tau}(0) = \chi_{0,ij}, \qquad \lim_{\omega\downarrow 0}\frac{\operatorname{Im}\chi^R_{ij,\tau}(\omega)}{\omega} = \tau\chi_{0,ij}.$$

Thus a fixed static susceptibility does not determine the low-frequency dissipative coefficient. This example is not a NAMCG derivation.

Positivity of mutual information establishes $I(A:B) \geq 0$ for each state. It does not establish an ordering between adiabatic and non-adiabatic mutual informations, or the sign or separation dependence of a dissipative response.

Static tail, distinct from the response. Chapters 6 and 14 and Appendix E use the same large-separation scaling,

$$I(A:B) \sim c_{p_*} \left(\frac{\ell_A \ell_B}{d^2} \right)^{p_*}, \qquad d \gg \ell_A, \ell_B,$$

which is dimensionless. It is established in the literature for Rényi mutual information of regions in a conformal vacuum, with the von Neumann limit shown for free-scalar spheres [3]; its leading power for NAMCG cores is an assumption. Neither this static tail nor a coincidence of its power with the assumed drag exponent determines the retarded force response, its tensor structure, or its coefficient.

Missing step. Specify the microscopic dynamics, stationary reference state, and force operators for the finite cores. Compute $\chi^R_{ij}(\omega; \mathbf{r}, \ell_A, \ell_B)$, justify the passivity and low-frequency assumptions, and derive the longitudinal and transverse responses. Determine whether they reduce to the isotropic d^{-4} laboratory law and calculate its coefficient before estimating η at tidal or LIGO-band scales. Problem I remains open.

Status. Optional, not part of the matrix-derived completion criterion unless drag is claimed to be derived or predicted. Its absence does not kill the framework within a scope that treats the drag law as phenomenological.

H.10 Problem J: spherical ECG solutions

Ledger label. Status: OPEN. Priority: OPTIONAL within the stated ECG model. Completion: PROGRAMME-CONDITIONAL. The reduced results do not derive singularity resolution from the matrix theory.

For the explicit invariant $\mathcal{P}$ in (9.7), consider the reduced action per unit time and angular volume

$$L[N, f] = N r^2 \big(R - 2\Lambda_{\text{eff}} + \kappa \mathcal{P}\big), \qquad \kappa = 16\pi G_{\text{eff}} \beta_{\text{ECG}}.$$

This reduction assumes that the static, spherically symmetric variations capture the corresponding field equations; an appropriate symmetric-criticality argument is not

established here. The reduced Euler–Lagrange operators, including all higher-derivative terms, are

$$\mathcal{E}_q = \frac{\partial L}{\partial q} - \frac{\mathrm{d}}{\mathrm{d}r}\frac{\partial L}{\partial q'} + \frac{\mathrm{d}^2}{\mathrm{d}r^2}\frac{\partial L}{\partial q''} - \cdots, \qquad q = N, f.$$

Variations are taken with the fields and the derivatives required by the order of L fixed at the endpoints (or with compact support); otherwise the associated boundary terms must be retained. For the specified $\mathcal{P}$, these reduced equations have been checked symbolically:

$$\mathcal{E}_f|_{N=1} \equiv 0, \qquad \mathcal{E}_N|_{N=1} = \frac{\mathrm{d}}{\mathrm{d}r}\left[2r(1-f) + 3\kappa\mathcal{B}[f] - \frac{2}{3}\Lambda_{\mathrm{eff}} r^3\right].$$

Thus $N = 1$ is a consistent vacuum branch of the reduced equations, and, with $\alpha = -\kappa$, the lapse equation has first integral

$$\mathcal{F}[f] - \frac{2}{3}\Lambda_{\mathrm{eff}} r^3 = 4M, \qquad \mathcal{F}[f] = 2r(1-f) - 3\alpha\,\mathcal{B}[f],$$

where α is the reduced ECG coupling and

$$\mathcal{B}[f] = \frac{4}{3}(f')^3 - \frac{8}{r}\,ff''\left(\frac{rf'}{2} + 1 - f\right) + \frac{4}{r}(f')^2 - \frac{8}{r^2}ff'(f-1).$$

For the rational profile with $a = r_0^3$ and $\Lambda_{\mathrm{eff}} = 0$,

$$f_{\mathrm{rat}}(r) = 1 - \frac{2Mr^2}{r^3 + a} = 1 - \frac{2M}{r} + \frac{2Ma}{r^4} + O(r^{-7}),$$

substitution gives

$$\mathcal{F}[f_{\mathrm{rat}}] - 4M = -\frac{4Ma}{r^3} - \frac{432\alpha M^2}{r^5} + O(r^{-6}).$$

The r^{-3} term cannot be cancelled by the cubic contribution, which begins at r^{-5}. Hence, for the displayed equation, this profile is not an exact vacuum solution when $Mr_0 \neq 0$.

Moreover, a smooth regular center has $f(r) = 1 + O(r^2)$, which implies $\mathcal{B}[f] \to 0$ and $\mathcal{F}(0) - \frac{2}{3}\Lambda_{\mathrm{eff}} 0^3 = 0$. Consequently the integration constant vanishes: this displayed equation has no smooth regular-centre vacuum solution with a nonzero integration constant, for any Λ_{eff}. For $\Lambda_{\mathrm{eff}} = 0$, the asymptotic falloff $f = 1 - 2M/r + o(r^{-1})$ identifies that constant as $4M$ and M as the ADM mass. For nonzero Λ_{eff}, its interpretation is not automatically the ADM mass.

Maximally symmetric branch. For $f = 1 - kr^2$, the first integral gives

$$\frac{\mathcal{F}[f]}{r^3} = 2k - 16\alpha k^3, \qquad k - 8\alpha k^3 = \frac{\Lambda_{\text{eff}}}{3}.$$

This is a conditional background-curvature relation, not a stability claim.

Large-radius tail. For the displayed equation, write $f = 1 - 2M/r + \delta(r)$. Then $2r(1-f) = 4M - 2r\delta$ and $\mathcal{B}[1-2M/r] = 144M^2/r^5 + O(r^{-6})$. The leading correction and the next asymptotic order give

$$f(r) = 1 - \frac{2M}{r} - \frac{216\,\alpha M^2}{r^6} + \frac{368\,\alpha M^3}{r^7} + \cdots .$$

The rational profile instead has a $+2Ma/r^4$ term. The tail expansion is an asymptotic check of the displayed first integral; it does not determine the model parameter r_0 or establish a match to an independently published expansion. In the stated action normalization $\alpha = -16\pi G_{\text{eff}}\beta_{\text{ECG}}$. If $G_{\text{eff}} > 0$, then $\alpha > 0$ iff $\beta_{\text{ECG}} < 0$, which fixes the sign of the leading correction. For $M > 0$, its magnitude defines the characteristic scale $r_* \sim |\alpha M|^{1/5}$; it is not a core scale.

Status. Conditional on the stated symmetric reduction, the reduced-action identities, first integral, regular-centre constraint, maximally symmetric curvature relation, and asymptotic tail above have been checked for the specified $\mathcal{P}$. The cited Einstein-like linearized spectrum on maximally symmetric backgrounds is a separate result; no nonlinear, sourced, or non-maximally-symmetric stability claim follows. Problem J remains open because the reduction's relation to the full field equations and the cubic truncation and its matching to the matrix theory are not established. A regular massive core still requires a specified source, different boundary conditions, or modified field equations; the entanglement stress for such a source is not specified.

H.11 Problem N: control of the effective expansion

Ledger label. Status: OPEN. Priority: CRITICAL PATH for the matrix-derived label. Completion: PROGRAMME-CONDITIONAL.

Assumption 9.2 treats the four-derivative terms as expansion vertices at order $(p/\Lambda_{\text{UV}})^2$ and not as propagating modes. It is stated in Chapter 9 and justified there only by the composite character of the metric. The ghost mass of Stelle gravity is of order $1/\sqrt{\alpha G}$, and its relation to Λ_{UV} depends on the physical G_{eff} and on α, which are part of Problem F.

Missing step. Derive from the matrix path integral that the graviton two-point function has no propagating Ostrogradsky ghost for $p < \Lambda_{\rm UV}$, for the values of α and $G_{\rm eff}$ computed in Problem F (here α is the Weyl-squared coupling of (9.5), whose $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$ coefficient is $-\alpha/2$; with the continuation convention stated below, $c_W = -\alpha/2$ in (9.3); it is not any other coupling denoted α in this book), with the region of those values in which it holds stated. This is the requirement of item 6 of Chapter 1 and no stronger.

Induced Weyl term and the extra spin-two pole (conditional; not closure). Work in the truncated identity of Theorem 8.2, expanded about a flat background with the vacuum term cancelled or neglected so that flat space is a solution, with every included sector assigned the common limits $\Lambda_{\rm UV}, \mu$, $\mu \ll \Lambda_{\rm UV}$, with $S_E = -iS_L$ as in Appendix A, and with curvature invariants continued without change of sign (an added convention). Rewrite the traced $a_{2,s}$ of (A.7) in the basis $\{C^2, R^2, E_{\rm GB}\}$, and let $\mathcal{W}_C := \sum_s w_s c_{C,s}$ and $\mathcal{W}_R := \sum_s w_s c_{R,s}$, where $c_{C,s}$ and $c_{R,s}$ are the coefficients of $C_{\mu\nu\rho\sigma}C^{\mu\nu\rho\sigma}$ and R^2. For a real scalar sector with $E = E_0 + QR$, $c_{C,s} = \operatorname{tr}\mathbb{I}_s/120$ and $c_{R,s} = \frac{1}{2}\operatorname{tr}(\frac{1}{6}\mathbb{I}_s - Q)^2$; for other sectors $c_{C,s}$ also depends on E and on the bundle curvature and is not computed here. The logarithmic term of (8.8) gives, in (9.3),

$$c_W = \frac{\mathcal{W}_C L}{32\pi^2}, \qquad c_R = \frac{\mathcal{W}_R L}{32\pi^2}, \qquad L := \ln\frac{\Lambda_{\rm UV}^2}{\mu^2},$$

that is, $\alpha = -2c_W$ and $\beta = 2c_R$ in (9.5); the Gauss–Bonnet term does not contribute at quadratic order on a flat background. With $g_{\mu\nu} = \eta_{\mu\nu} + h_{\mu\nu}$ and $k^2 = \eta^{\mu\nu}k_\mu k_\nu$, the spin-two part of the quadratic form of $R/(16\pi G_{\rm eff}) + c_W C^2 + c_R R^2$ is proportional to $k^2(1 - 32\pi G_{\rm eff} c_W k^2)$ and its spin-zero part to $k^2(1 + 96\pi G_{\rm eff} c_R k^2)$ (a symbolic check, not a proof). There is therefore a spin-two pole at $k^2 = -m_2^2$ with residue opposite in sign to that of the massless graviton, and a spin-zero pole at $k^2 = -m_0^2$ with residue of the graviton's sign, where

$$m_2^2 = -\frac{1}{32\pi G_{\rm eff} c_W} = -\frac{\Upsilon\Lambda_{\rm UV}^2}{L}, \qquad m_0^2 = \frac{1}{96\pi G_{\rm eff} c_R}, \qquad \Upsilon := \frac{\pi}{G_{\rm eff}\mathcal{W}_C\Lambda_{\rm UV}^2}.$$

If $G_{\rm eff}\mathcal{W}_C > 0$, the extra spin-two pole of this fixed-L form is a tachyonic ghost, with $|m_2| < \Lambda_{\rm UV}$ exactly when $L > \Upsilon$. If $1/G_{\rm eff}$ is taken from the universal a_1 term alone, $1/G_{\rm eff} = \mathcal{A}_1(\Lambda_{\rm UV}^2 - \mu^2)/(2\pi)$ with $\mathcal{A}_1 := \sum_s w_s \mathcal{A}_{1,R,s}$, and $\Upsilon = \mathcal{A}_1/(2\mathcal{W}_C)$ up to $O(\mu^2/\Lambda_{\rm UV}^2)$, independent of $\Lambda_{\rm UV}$. For sectors of minimally coupled real scalars, $\mathcal{A}_1 = \mathcal{N}_0/6$ and $\mathcal{W}_C = \mathcal{N}_0/120$, so $\Upsilon = 10$ for every $\mathcal{N}_0 > 0$, $|m_2| < \Lambda_{\rm UV}$ in this fixed-L form once $\Lambda_{\rm UV}/\mu > e^5$, and $m_0^2 = 2\Lambda_{\rm UV}^2/L$. Since $|m_2|^2/\mu^2 = \Upsilon e^L/L$, which is much larger than one for $\mu \ll \Lambda_{\rm UV}$ unless Υ is exponentially small, this pole lies at momenta where the fixed-L form does not apply (see below); it is a property of

that form, not a one-loop prediction. For $|k| \lesssim \mu$, where the fixed-L form applies, the Weyl correction relative to the Einstein term is $32\pi G_{\rm eff} c_W k^2 = (L/\Upsilon)\, k^2/\Lambda_{\rm UV}^2$.

The fixed-L form is not the one-loop result at momenta $\mu \ll |k| \ll \Lambda_{\rm UV}$. At second order in curvature the one-loop action is nonlocal, with momentum-dependent form factors [30]; assuming that the Weyl form factor decays when $sk^2 \gg 1$, its proper-time integral is cut off at $s \sim k^{-2}$ rather than at μ^{-2}, and for massless sectors L is replaced by $\ln(\Lambda_{\rm UV}^2/k^2) + \delta_W$, with a constant δ_W fixed by the non-logarithmic part of the form factor and by the finite Weyl counterterm. For the analogous second-order form factor $\int_0^1 {\rm d}\tau\, e^{-\tau(1-\tau)sk^2}$ of a potential insertion, a direct numerical evaluation gives $\ln(\Lambda_{\rm UV}^2/k^2) + 2 - \gamma_{\rm E}$, with $\gamma_{\rm E}$ Euler's constant, independent of μ for $k \gg \mu$; this illustrates the mechanism and does not compute δ_W. With $x := k^2/\Lambda_{\rm UV}^2$ and the principal logarithm, the spin-two factor becomes

$$F(x) = 1 - \frac{x}{\Upsilon}\Big(\ln\frac{1}{x} + \delta_W\Big),$$

whose zeros on the principal sheet are $x = -\Upsilon/W_j(-\Upsilon e^{-\delta_W})$, $j \in \{0, -1\}$, with W_j the branches of the Lambert function; μ no longer enters. At $\delta_W = 0$ the zeros cross $|x| = 1$ at $x = \pm i$ when $\Upsilon = \pi/2$. A numerical scan with an argument-principle count (a check, not a proof) finds that F has no zero in the disc $|k^2| < \Lambda_{\rm UV}^2$ exactly when $\Upsilon > \Upsilon_*(\delta_W)$, with $\Upsilon_*(0) = \pi/2$ and $\Upsilon_*(\delta_W) = \delta_W$ for $\delta_W \geq 1$, where the crossing is at $x = 1$. For $\Upsilon = 10$ the condition is $-9.58 < \delta_W < 10$; at $\delta_W = 0$ the zeros are a complex-conjugate pair with $|k^2| \approx 3.9\Lambda_{\rm UV}^2$.

These statements are conditional on the truncated identity, the continuation and common limits stated, massless sectors (massive sectors require the threshold functions of Problem F), and the use of the leading-logarithmic form factor up to $|k^2| \sim \Lambda_{\rm UV}^2$, where the derivative expansion is not controlled. They do not compute $\mathcal{W}_C$, $\mathcal{A}_1$, $G_{\rm eff}$ or δ_W for the locked block, do not derive the induced coefficients from the matrix path integral, and do not decide whether complex-conjugate zeros beyond $\Lambda_{\rm UV}$ are propagating ghosts. Within the leading-logarithmic model F, absence of a zero in $|k^2| < \Lambda_{\rm UV}^2$ is a numerical condition on Υ and δ_W, which are outputs of Problem F, and is not implied by the metric being composite. Problems F and N remain open.

Status. Remains Assumption 9.2.

H.12 What this appendix does not do

It does not produce an unsuppressed laboratory coupling between a controllable external field and $T_{\mu\nu}^{\rm ent}$ or $G_{\rm eff}$. The observational window therefore remains empty. It does not alter the chain of Chapter 16 or the agenda of Chapter 17.

Appendix I

Gaia DR3 Wide Binaries: A Coefficient-Level Consistency Check

This appendix records a comparison of Gaia DR3 wide-binary kinematics with the external-field-corrected form of the coefficient-level interpolator already examined on SPARC in Appendix D. It is not a test of Theorems 7.3–12.4, not a derivation of $\nu_{\surd}$, and not evidence that Problem H is solved.

I.1 What is being compared

The relevant acceleration scale for a solar-neighbourhood binary is not the internal Newtonian acceleration alone. The Galactic external field at the Sun satisfies

$$g_{\rm gal} \approx (1.5\text{–}2.2) \times 10^{-10}\,\mathrm{m\,s^{-2}} > g_\dagger .$$

Under the external-field saturation already noted for the coefficient layer (Assumption 13.3), the isolated deep-regime boost is suppressed. The working prediction used here is therefore the residual enhancement obtained by evaluating $\nu_{\surd}$ on the quadrature sum of the internal and external fields (or the equivalent AQUAL/QUMOND external-field prescription). Asymptotically this residual is of order 20%, not the factor ~ 4 of the isolated map.

The comparison is performed on the public El-Badry et al. Gaia DR3 wide-binary catalogue after the standard quality cuts (RUWE$<$ 1.3, fractional parallax error < 0.06, projected separations 2–30 kAU). Projection, unresolved companions, and chance alignments are modelled by the same mixture techniques employed in the controlled literature analyses; they are not optional.

I.2 What the data show

Controlled analyses that include the Galactic external field, realistic triple contamination, and the quality cuts above find relative velocities consistent with Newtonian gravity. The residual MOND/RAR-like enhancement is excluded at high significance (Banik et al. 2024; Pittordis, Sutherland & Shepherd 2025; quality-framework study 2026). Apparent boosts reported in less filtered samples diminish once these systematics are treated.

Consequently the external-field-corrected $\nu_{\surd}$ (the coefficient-level map of Appendix D) is also consistent with the data only to the extent that its residual boost remains inside the observational uncertainties. It does not outperform Newtonian gravity on this sample, nor does it produce a statistically preferred rising velocity ratio of the size generated by the isolated interpolator.

I.3 What this is not

- It is not a derivation of $\nu_{\surd}$ from $I(A : B)$ or from the modular spectrum. Problem H remains open.
- It is not a constraint on $m_T = 0$, on the heat-kernel map for G_{eff}, or on the conservation identity. A null result under the external-field-corrected prediction is the outcome expected once Assumption 13.3 is respected.
- It is not evidence that an isolated reading of the RAR interpolator applies to solar-neighbourhood binaries. That reading is already ruled out by $g_{\text{gal}} > g_{\dagger}$.
- It is not a laboratory handle on the entanglement stress.

I.4 Status

The Gaia DR3 wide-binary sample, analysed with the external field included, is compatible with Newtonian gravity and with the modest residual enhancement allowed by the coefficient-level NAMCG/RAR map under external-field saturation. It supplies no new support for the framework beyond the SPARC consistency check already recorded, and it does not alter the open-problem ledger of Appendix H.

Appendix J

Infrared Limits That Follow from the Proved Chain

This appendix collects the infrared limits that follow directly from the theorems and structural results already established in Chapters 3–12. It does not treat open problems as closed, and it does not claim additional recovery of general relativity beyond those limits.

J.1 Limits that are obtained

Under the named assumptions of the monograph the following hold.

1. **High-acceleration Einstein equation.** When the reversible entanglement stress is negligible compared with ordinary matter stress, the infrared equation reduces to
$$G_{\mu\nu} + \Lambda_{\text{eff}} g_{\mu\nu} = 8\pi G_{\text{eff}} T^{\text{matter}}_{\mu\nu}. \tag{J.1}$$
This is the content of the high-acceleration limit of the equation already written after Theorem 11.1.

2. **Massless tensor polarizations.** The entanglement-equilibrium condition of Chapters 6–7 forces $m_T = 0$, so the infrared spectrum contains the two transverse-traceless modes of a massless spin-2 field. This is Theorem 7.3; it is not re-derived here.

3. **Conservation.** The total stress (matter + tensor + entanglement) is divergenceless,
$$\nabla^\mu (T^{\text{matter}}_{\mu\nu} + T^{\text{tensor}}_{\mu\nu} + T^{\text{ent}}_{\mu\nu}) = 0, \tag{J.2}$$
by Theorem 11.1. This identity constrains the total stress; it does not by itself

derive any reduced drag used in later chapters, whose microscopic origin remains Problem I.

4. **Structural form of G_{eff}.** The conditional, truncated heat-kernel identity is recorded in Theorem 8.2; the physical coefficient calculation remains open in Problem F.

5. **Regulation of the quartic term.** The holographic boundary-entropy vacuum of Chapter 12 supplies a consistent mechanism that can cancel the leading Λ_{UV}^4 contribution against a boundary term. The specific working value $\Lambda_{\text{eff}} = 3/L_{\text{IR}}^2$ is a choice compatible with that mechanism, not a derived necessity.

J.2 Limits that are not obtained

The following are frequently listed as recoveries but do not follow from the proved chain.

- Entanglement equilibrium from the microscopic path integral (Problem B). The fixed-algebra first law does not derive $\delta S_D = 0$ for the non-conformal tensor block.

- Matrix-derived local Lorentz symmetry and a full field-content account, including the independent connection and ghost operators (Problem D). Imposing local Lorentz symmetry gives only a scope-limited Presentation-B model.

- The infrared boundary entropy density and stationarity derived from the matrix theory (Problem G). The conditional mechanism and an assumed boundary-model scaling do not derive these inputs.

- Exact $\gamma - 1 = 0$ in the solar system. The conformal mode ψ is not identical to the radial Higgs; its mass matrix has not been diagonalized (Problem C). Cassini therefore remains a constraint on any light eigenmode that may survive.

- A first-principles drag exponent or numerical rate. The static conformal tail is conditional on its stated assumptions and does not determine a drag law. The assumed d^{-4} drag is a phenomenological input of Chapter 14 (Problem I). Its microscopic origin and compatibility with a covariantly conserved stress remain unresolved.

- Singularity resolution from the matrix theory. The regular-core profile is a phenomenological model. For $\Lambda_{\text{eff}} = 0$ and $Mr_0 \neq 0$ it is not an exact vacuum solution of the displayed first-integral equation in Appendix H (Problem J), conditional on the stated reduction. That equation also excludes a nonzero

integration constant for a smooth regular centre at any Λ_{eff}. Neither result derives the profile from the microscopic action.

- Lorentzian signature from the matrix algebra (Problem E). It remains under the assumption stated in Chapter 3.
- An interpolating function for galactic or binary kinematics derived from $I(A : B)$ (Problem H). Coefficient-level compatibility with the published RAR map is recorded in Appendix D; it is not a derivation.

J.3 Status

The infrared theory obtained from the proved chain is Einstein gravity with a structural G_{eff}, a massless spin-2 sector, a divergenceless entanglement stress, and a consistent regulation of the leading vacuum-energy term. Every stronger recovery claim requires the closure of at least one open problem listed in Appendix H; a completed matrix-derived infrared derivation requires all the critical-path problems listed in Chapter 17 to be DERIVED, as specified there. This appendix adds no further recovery.

Appendix K

Heat-Kernel Response to an External Field Strength

This appendix records the leading response of the Seeley–DeWitt coefficient a_1 to a laboratory field strength and the precision constraints that any residual conformal mixing must satisfy. It does not recompute a_1 on the locked block, does not derive a kinetic mixing from the entanglement functional, and does not open an observationally allowed laboratory window.

K.1 Vanishing of the linear term in a_1

On a Laplace-type operator $\Delta = -D^\mu D_\mu + E$ the first Seeley–DeWitt coefficient, in the convention of (A.6), is

$$a_1 = \mathrm{tr}\Big(\tfrac{1}{6}R - E\Big) = \tfrac{1}{6}R\,\dim - \mathrm{tr}(E). \tag{K.1}$$

If a controllable field strength enters the endomorphism only through an antisymmetric generator,

$$E = E_0 + \xi F_{\alpha\beta} S^{\alpha\beta}, \tag{K.2}$$

the trace of $S^{\alpha\beta}$ vanishes on every representation that appears in the locked 4×4 block and in the diffeomorphism and local-Lorentz ghost sectors. Consequently

$$\delta a_1 = 0. \tag{K.3}$$

This is a standard algebraic fact. It is not a recomputation of a_1 on the NAMCG fluctuation operator (Problem F remains open), nor does it cancel the higher-order suppression by Λ_{UV} already identified for laboratory fields.

K.2 Quadratic appearance in a_2

The first non-vanishing correction proportional to $F_{\mu\nu}F^{\mu\nu}$ appears in a_2 and is quadratic in the laboratory field. Variation of the resulting effective action with respect to the metric produces a contribution proportional to $g_{\mu\nu}$. For the quadratic term, if its metric variation has this form, the coupling is confined to the conformal mode and generates no transverse-traceless response. This statement does not cover a term linear in $F_{\alpha\beta}$ proportional to $\mathrm{tr}(S^{\alpha\beta}E_0)$, which is not excluded by $\mathrm{tr}\, S^{\alpha\beta} = 0$ alone, and the proportionality to $g_{\mu\nu}$ is not derived here.

K.3 Precision constraints on conformal mixing

Any additional kinetic mixing between the conformal mode ψ and a laboratory scalar that might be introduced by hand is constrained by existing data:

- Cassini: $|\gamma - 1| \lesssim 2 \times 10^{-5}$;
- torsion-balance equivalence-principle tests: $\sin^2\theta \lesssim 10^{-13}$.

A mixing angle large enough to produce a macroscopic laboratory acceleration violates these bounds. No calculation within the present framework cancels the parametric suppression $F/\Lambda_{\mathrm{UV}}^2$ while leaving a residual that survives the bounds. The observational window therefore remains empty.

K.4 What is not shown

- A completed evaluation of a_1 (or the relevant part of a_2) on the locked 4×4 block with ghosts.
- A derivation of a kinetic mixing term from $U_{\mathrm{ent}}[I(A:B)]$ or from the modular Hamiltonian.
- A dynamical cancellation of quadratic ultraviolet divergences that leaves a finite infrared residual.
- Any projection of galactic rotation-curve discrepancies onto solar-system scales, or any fit to pulsar-timing-array correlation data. Those exercises do not follow from the heat-kernel expansion and are omitted.

K.5 Status

The linear coupling of a laboratory field strength to a_1 vanishes for antisymmetric generators, and any residual conformal mixing is limited by existing precision tests

to a level far below laboratory interest. This appendix adds no new handle on the entanglement stress or on G_{eff} and does not alter the open-problem ledger of Appendix H.

Appendix L

A Phenomenological Redshift-Dependent Multiplier (Outside the Proved Chain)

This appendix records a purely phenomenological modification that lies outside the NAMCG derivation. It is not obtained from the entanglement stress, the heat-kernel map, or any theorem of Chapters 3–12, and it does not close any open problem.

L.1 The inserted multiplier

One may choose, by hand, a redshift-dependent factor

$$\beta(z) = \beta_{\mathrm{ref}}\Big(\frac{1+z_{\mathrm{ref}}}{1+z}\Big)^{\gamma} \tag{L.1}$$

and insert it into the standard linear growth factor according to

$$D(z) = D_{\Lambda\mathrm{CDM}}(z)\big[1+\beta(z)\ln(1+z)\big]. \tag{L.2}$$

The parameters β_{ref}, z_{ref} and γ are free. No equation in the monograph fixes them.

L.2 Illustrative high-redshift comparison

With the particular choice $\beta_{\mathrm{ref}} = 0.5$, $z_{\mathrm{ref}} = 11$, $\gamma = 0.8$, the modified growth can be fed into a schematic Sheth–Tormen mass function and a fixed mass-to-light conversion. The resulting curves can be plotted against a small set of representative high-redshift UV luminosity points. Agreement or disagreement with those points is adjustable by changing the three free parameters and does not test the infrared equation, the

conservation identity, or the value of G_{eff}.

L.3 What the illustration is not

- It is not a derivation of an epoch-dependent coupling from the matrix condensate or from mutual information.
- It is not a solution of the continuity or Euler equation sourced by $T^{\text{ent}}_{\mu\nu}$.
- It is not a statistical fit to the public JADES or CEERS catalogs.
- It does not constrain, or supply evidence for, any load-bearing theorem.
- It does not open a laboratory handle on the entanglement stress.

L.4 Status

The construction is an external multiplier that can be tuned to high-redshift galaxy counts. It stands outside the proved chain and is recorded only as an illustration of how additional free functions may be added once one leaves the framework. It is omitted from the claim list of Chapter 16 and from the open-problem ledger of Appendix H.

Conclusion

The monograph constructs a single derivation chain from a non-Abelian matrix order parameter to an infrared equation that is Einstein gravity plus a divergenceless entanglement stress, with a structural heat-kernel map for G_{eff} and a massless spin-2 sector forced by entanglement equilibrium. That chain is complete only under the named assumptions of Chapters 3–12. Several critical-path derivations remain open.

What has been obtained

Under those assumptions one has:

- a composite metric defined from the condensate,
- $m_T = 0$ from the diamond Ward identity,
- the structural form of $1/G_{\text{eff}}$,
- covariant conservation of the total stress including $T^{\text{ent}}_{\mu\nu}$,
- a consistent regulation of the leading vacuum-energy term via the holographic boundary functional.

Coefficient-level compatibility of the published RAR interpolator with the SPARC sample is recorded in Appendix D. It is not a derivation of that interpolator.

What has not been obtained

- entanglement equilibrium from the microscopic path integral (Problem B);
- a matrix-derived local Lorentz symmetry and specified full field content (Problem D1);
- the infrared boundary entropy and stationarity derived from the matrix theory (Problem G);

- a derivation of $\nu_{\surd}$ (or any other interpolator) from mutual information or the modular spectrum (Problem H);
- scheme-fixed values of the heat-kernel coefficients A, B, C (Problem F);
- diagonalization of the conformal-mode mass matrix (Problem C);
- the dictionary map into the modular Hamiltonian (Problem A);
- an unsuppressed, observationally allowed laboratory coupling to the entanglement stress;
- empirical support from Gaia DR3 wide binaries once the Galactic external field is included (Appendix I);
- any redshift-dependent growth multiplier; such constructions lie outside the proved chain.

The framework therefore does not yet resolve the dark-matter problem, nor does it supersede existing phenomenological fits on galactic or binary scales.

Next steps

Progress toward a completed matrix-derived infrared derivation requires deriving, not retaining, the critical-path steps:

1. Problem A: the dictionary of Assumption 3.6 derived from the microscopic definition;
2. Problem B: entanglement equilibrium from the microscopic path integral, including non-conformal tensor variations;
3. Problem C: the scalar operator with a coupling test;
4. Problem D1: local Lorentz symmetry and a non-degenerate frame derived from the matrix algebra, with the ω, A_μ, B and ghost operators;
5. Problem E: Lorentzian signature derived, not retained;
6. Problem F: A, B, C and G_{eff} from specified operators, with G_{eff} not an input;
7. Problem G: infrared entropy density and stationarity derived from the matrix theory;
8. Problem N: the effective expansion controlled, with no propagating Ostrogradsky ghosts for $p < \Lambda_{\text{UV}}$.

Alternate scope. Specifying the operator content of an explicitly scope-limited Presentation-B model (imposed local Lorentz symmetry, externally justified boundary datum, or an input G_{eff}) can make that model calculable but does not derive the critical-path steps and leaves the matrix-derived completion open. Problems H, I, and J are optional within their stated scopes; Problem I is a prerequisite only if drag is claimed to be derived or predicted.

Until all the critical-path problems are derived, the infrared equation remains exactly as written in the main text and no stronger claim is licensed. The open-problem ledger of Appendix H is the working list.

Bibliography

[1] Pablo Bueno and Pablo A. Cano. Einsteinian cubic gravity. *Physical Review D*, 94:104005, 2016.

[2] Philipp Dorau and Albert Much. From quantum relative entropy to the semiclassical Einstein equations. *Physical Review Letters*, 136(9):091602, 2026. doi: 10.1103/lmq8-nsty.

[3] John Cardy. Some results on mutual information of disjoint regions in higher dimensions. *Journal of Physics A*, 46:285402, 2013. doi: 10.1088/1751-8113/46/28/285402.

[4] Stacy S. McGaugh, Federico Lelli, and James M. Schombert. The radial acceleration relation in rotationally supported galaxies. *Physical Review Letters*, 117: 201101, 2016.

[5] B. Bertotti, L. Iess, and P. Tortora. A test of general relativity using radio links with the Cassini spacecraft. *Nature*, 425:374, 2003.

[6] Federico Lelli, Stacy S. McGaugh, James M. Schombert, and Marcel S. Pawlowski. One law to rule them all: The radial acceleration relation of galaxies. *The Astrophysical Journal*, 836:152, 2017.

[7] M. Milgrom. A modification of the Newtonian dynamics as a possible alternative to the hidden mass hypothesis. *The Astrophysical Journal*, 270:365, 1983. doi: 10.1086/161130.

[8] Federico Lelli, Stacy S. McGaugh, and James M. Schombert. Testing Verlinde's emergent gravity with the radial acceleration relation. *Monthly Notices of the Royal Astronomical Society: Letters*, 468:L68, 2017.

[9] Markus B. Fröb. Modular Hamiltonian for de Sitter diamonds. *Journal of High Energy Physics*, 12:074, 2023. doi: 10.1007/JHEP12(2023)074.

[10] Sergey N. Solodukhin. Entanglement entropy of black holes. *Living Reviews in Relativity*, 14(1):8, 2011. doi: 10.12942/lrr-2011-8.

[11] Leonard Susskind and John Uglum. Black hole entropy in canonical quantum gravity and superstring theory. *Physical Review D*, 50:2700–2711, 1994. doi: 10.1103/PhysRevD.50.2700.

[12] Ted Jacobson. Entanglement equilibrium and the Einstein equation. *Physical Review Letters*, 116:201101, 2016.

[13] N. Ishibashi, H. Kawai, Y. Kitazawa, and A. Tsuchiya. A large-n reduced model as superstring. *Nuclear Physics B*, 498:467, 1997.

[14] Horacio Casini and Marina Huerta. Entanglement entropy in free quantum field theory. *J. Phys. A*, 42(50), 2009. doi: 10.1088/1751-8113/42/50/504007.

[15] Dmitri V. Fursaev and Gennaro Miele. Cones, spins and heat kernels. *Nucl. Phys. B*, 484:697–723, 1997. doi: 10.1016/S0550-3213(96)00631-1.

[16] Alessandro Manta, Harold C. Steinacker, and Tung Tran. hs-extended gravity from the IKKT matrix model, 2024.

[17] Daniel Kabat. Black hole entropy and entropy of entanglement. *Nuclear Physics B*, 453(1–2):281–299, 1995. doi: 10.1016/0550-3213(95)00443-V.

[18] William Donnelly and Aron C. Wall. Geometric entropy and edge modes of the electromagnetic field. *Physical Review D*, 94(10):104053, 2016. doi: 10.1103/PhysRevD.94.104053.

[19] Christopher P. Herzog and Tatsuma Nishioka. The edge of entanglement: Getting the boundary right for non-minimally coupled scalar fields. *Journal of High Energy Physics*, 2016(12):138, 2016. doi: 10.1007/JHEP12(2016)138.

[20] Aitor Lewkowycz and Juan Maldacena. Generalized gravitational entropy. *Journal of High Energy Physics*, 2013(8):090, 2013. doi: 10.1007/JHEP08(2013)090.

[21] Harold C. Steinacker. One-loop effective action and emergent gravity on quantum spaces in the IKKT matrix model. *Journal of High Energy Physics*, 05:129, 2023.

[22] Harold Steinacker. Emergent gravity on covariant quantum spaces in the IKKT model, 2016.

[23] Alessandro Manta and Harold C. Steinacker. Minimal covariant quantum space-time, 2025.

[24] Harold C. Steinacker. Higher-spin kinematics & no ghosts on quantum space-time in Yang–Mills matrix models, 2019.

[25] Kaushlendra Kumar and Harold C. Steinacker. Modified einstein equations from the 1-loop effective action of the IKKT model, 2023.

[26] Harold C. Steinacker. Modified gravity at large scales on quantum spacetime in the IKKT model, 2026.

[27] Marcus Sperling and Harold C. Steinacker. Covariant cosmological quantum space-time, higher-spin and gravity in the IKKT matrix model. *Journal of High Energy Physics*, 2019(7):010, 2019. doi: 10.1007/JHEP07(2019)010.

[28] Gérard Ben Arous. Développement asymptotique du noyau de la chaleur hypoelliptique sur la diagonale. *Annales de l'Institut Fourier*, 39(1):73–99, 1989. doi: 10.5802/aif.1158.

[29] Jan Troost. Constant field strengths on T^{2n}. *Nucl. Phys. B*, 568:180–194, 2000. doi: 10.1016/S0550-3213(99)00721-X.

[30] A. O. Barvinsky and G. A. Vilkovisky. Covariant perturbation theory (II). second order in the curvature. general algorithms. *Nuclear Physics B*, 333:471–511, 1990. doi: 10.1016/0550-3213(90)90047-H.

[31] Ted Jacobson. Thermodynamics of spacetime: The Einstein equation of state. *Physical Review Letters*, 75:1260, 1995.

[32] Bryce S. DeWitt. *Dynamical Theory of Groups and Fields*. Gordon and Breach, 1965.

[33] T. Padmanabhan. General relativity from a thermodynamic perspective. *General Relativity and Gravitation*, 46:1673, 2014.

[34] M. Ostrogradsky. Mémoires sur les équations différentielles, relatives au problème des isopérimètres. *Mem. Acad. St. Petersbourg*, 6:385, 1850.

[35] Federico Lelli, Stacy S. McGaugh, and James M. Schombert. SPARC: Mass models for 175 disk galaxies with Spitzer photometry and accurate rotation curves. *The Astronomical Journal*, 152:157, 2016.

[36] A. D. Sakharov. Vacuum quantum fluctuations in curved space and the theory of gravitation. *Doklady Akademii Nauk SSSR*, 177:70, 1967.

[37] W. G. Unruh. Experimental black-hole evaporation? *Physical Review Letters*, 46: 1351, 1981.

[38] Nathan Seiberg and Edward Witten. String theory and noncommutative geometry. *Journal of High Energy Physics*, 09:032, 1999.

[39] Harold Steinacker. Emergent geometry and gravity from matrix models: An introduction. *Classical and Quantum Gravity*, 27:133001, 2010.

[40] R. T. Seeley. Complex powers of an elliptic operator. *Proceedings of Symposia in Pure Mathematics*, 10:288, 1967.

[41] D. V. Vassilevich. Heat kernel expansion: User's manual. *Physics Reports*, 388: 279, 2003.

[42] K. S. Stelle. Renormalization of higher-derivative quantum gravity. *Physical Review D*, 16:953, 1977.

[43] Robie A. Hennigar, Mohammad Bagher Jahani Poshteh, and Robert B. Mann. Shadows, signals, and stability in einsteinian cubic gravity. *Physical Review D*, 97:064041, 2018.

[44] Erik Verlinde. On the origin of gravity and the laws of Newton. *Journal of High Energy Physics*, 04:029, 2011.

[45] Erik P. Verlinde. Emergent gravity and the dark universe. *SciPost Physics*, 2: 016, 2017.

[46] Kyu-Hyun Chae, Federico Lelli, Harry Desmond, Stacy S. McGaugh, Pengfei Li, and James M. Schombert. Testing the strong equivalence principle: Detection of the external field effect in rotationally supported galaxies. *The Astrophysical Journal*, 904:51, 2020.

[47] Douglas Clowe, Maruša Bradač, Anthony H. Gonzalez, Maxim Markevitch, Scott W. Randall, Christine Jones, and Dennis Zaritsky. A direct empirical proof of the existence of dark matter. *The Astrophysical Journal Letters*, 648: L109, 2006.

[48] B. P. Abbott et al. GW170817: Observation of gravitational waves from a binary neutron star inspiral. *Physical Review Letters*, 119:161101, 2017.

[49] B. P. Abbott et al. Gravitational waves and gamma-rays from a binary neutron star merger: GW170817 and GRB 170817a. *The Astrophysical Journal Letters*, 848:L13, 2017.

[50] Benoit Famaey and Stacy S. McGaugh. Modified Newtonian dynamics (MOND): Observational phenomenology and relativistic extensions. *Living Reviews in Relativity*, 15:10, 2012.

[51] Robert M. Wald. Black hole entropy is the Noether charge. *Physical Review D*, 48:R3427, 1993.

[52] J. J. Bisognano and E. H. Wichmann. On the duality condition for quantum fields. *Journal of Mathematical Physics*, 17:303, 1976.

[53] Kyu-Hyun Chae. Breakdown of the Newton-Einstein standard gravity at low acceleration in internal dynamics of wide binary stars. *The Astrophysical Journal*, 952:128, 2023. doi: 10.3847/1538-4357/ace101.

[54] Gaia Collaboration, A. Vallenari, A. G. A. Brown, et al. Gaia data release 3: Summary of the content and survey properties. *Astronomy & Astrophysics*, 674: A1, 2023. doi: 10.1051/0004-6361/202243940.

[55] Lily Whitler, Daniel P. Stark, Michael W. Topping, et al. The $z \gtrsim 9$ galaxy UV luminosity function from the JWST advanced deep extragalactic survey: Insights into early galaxy evolution and reionization. *The Astrophysical Journal*, 992:63, 2025.