
Turbot in IV Diagnostics − BTS−ISIS, age 1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − BTS−ISIS, age 2
a) Observed and fitted values time series
Observed
Fitted
values
0.2
0.5
2.0
5.0
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
0.2
0.5
2.0
5.0
0.2
0.5
1.0
2.0
5.0
10.0
Fitted values
c) Standardised residuals over time
Standardised Residuals
−2
−1
0
1
2
2000
2005
2010
2015
Year
d) Tukey−Anscombe plot
Standardised Residuals
−2
−1
0
1
2
0.2
0.5
1.0
2.0
5.0
10.0
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−2
−1
0
1
2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
12
Lag (yrs)


Turbot in IV Diagnostics − BTS−ISIS, age 3
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − BTS−ISIS, age 4
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − BTS−ISIS, age 5
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − BTS−ISIS, age 6
a) Observed and fitted values time series
Observed
Fitted
values
2
5
20
50
200
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
2
5
10
20
50
100
200
500
c) Standardised residuals over time
Standardised Residuals
1.5
0.5
−0.5
−1.5
Year
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
−2
−1
0
1
2
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.4
0.2
0.0
−0.4
Lag (yrs)
2
4
6
8
10
12


Turbot in IV Diagnostics − BTS−ISIS, age 7
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − catch, age 1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e+00
1e−01
5e−03
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
5e−03
5e−02
5e−01
5e+00
c) Standardised residuals over time
Standardised Residuals
3
2
1
0
−1
−2
−3
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
−2
−1
0
1
2
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.4
0.2
0.0
−0.4
Lag (yrs)
2
4
6
8
10
12


Turbot in IV Diagnostics − catch, age 2
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
0.2
0.5
2.0
5.0
20.0
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
0.2
0.5
1.0
2.0
5.0
10.0
20.0
c) Standardised residuals over time
Standardised Residuals
−2
−1
0
1
2
Year
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
12
Lag (yrs)


Turbot in IV Diagnostics − catch, age 3
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
0.5
1.0
2.0
5.0
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − catch, age 4
a) Observed and fitted values time series
Observed
Fitted
values
100
500
2000
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
100
500
2000
100
200
500
1000
2000
5000
Fitted values
c) Standardised residuals over time
Standardised Residuals
−2
−1
0
1
2
2000
2005
2010
2015
Year
d) Tukey−Anscombe plot
Standardised Residuals
−2
−1
0
1
2
100
200
500
1000
2000
5000
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−2
−1
0
1
2
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
12
Lag (yrs)


Turbot in IV Diagnostics − catch, age 5
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − catch, age 6
a) Observed and fitted values time series
Observed
Fitted
values
1000
200
50
10
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
1000
200
50
10
10
20
50
100
200
500
1000
Fitted values
c) Standardised residuals over time
Standardised Residuals
3
2
1
0
−1
−3
2000
2005
2010
2015
Year
d) Tukey−Anscombe plot
Standardised Residuals
3
2
1
0
−1
−3
10
20
50
100
200
500
1000
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
3
2
1
0
−1
−3
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.4
0.2
0.0
−0.4
2
4
6
8
10
12
Lag (yrs)


Turbot in IV Diagnostics − catch, age 7
a) Observed and fitted values time series
Observed
Fitted
values
5
20
50
200
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
5
20
50
200
5
10
20
50
100
200
500
Fitted values
c) Standardised residuals over time
Standardised Residuals
−1
0
1
2000
2005
2010
2015
Year
d) Tukey−Anscombe plot
Standardised Residuals
−1
0
1
5
10
20
50
100
200
500
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−1
0
1
−2
−1
0
1
2
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
12
Lag (yrs)


Turbot in IV Diagnostics − catch, age 8
a) Observed and fitted values time series
Observed
Fitted
values
2
5
20
50
200
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
2
5
10
20
50
100
200
500
c) Standardised residuals over time
Standardised Residuals
−1
0
1
Year
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
−2
−1
0
1
2
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
Lag (yrs)
4
6
8
10
12


Turbot in IV Diagnostics − catch, age 9
a) Observed and fitted values time series
Observed
Fitted
values
0.5
2.0
10.0
50.0
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
0.5
1.0
2.0
5.0
20.0
50.0
200.0
Fitted values
c) Standardised residuals over time
Standardised Residuals
−3
−2
−1
0
1
2
3
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
12
Lag (yrs)


Turbot in IV Diagnostics − catch, age 10
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − NL_LPUE, age −1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − SNS, age 1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − SNS, age 2
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV Diagnostics − SNS, age 3
a) Observed and fitted values time series
Observed
Fitted
values
2
5
20
50
200
2004
2006
2008
2010
2012
2014
2016
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
2
5
10
20
50
100
200
c) Standardised residuals over time
Standardised Residuals
−1.5
−0.5
0.5
1.5
Year
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
−1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
Lag (yrs)
2
4
6
8
10


Turbot in IV Diagnostics − SNS, age 4
a) Observed and fitted values time series
Observed
Fitted
values
0.5
2.0
10.0
50.0
2004
2006
2008
2010
2012
2014
2016
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
0.5
1.0
2.0
5.0
10.0
20.0
50.0
c) Standardised residuals over time
Standardised Residuals
−2
−1
0
1
2
Year
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
1:1 line
Quantiles of the Normal Distribution
−1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
Lag (yrs)


Turbot in IV Diagnostics − SNS, age 5
a) Observed and fitted values time series
Observed
Fitted
values
50
10
5
2
1
2004
2006
2008
2010
2012
2014
2016
Year
b) Observed vs fitted values
1:1 line
Observed values
50
10
5
2
1
1
2
5
10
20
50
100
Fitted values
c) Standardised residuals over time
Standardised Residuals
1.0
0.5
0.0
−1.0
2004
2006
2008
2010
2012
2014
2016
Year
d) Tukey−Anscombe plot
Standardised Residuals
1.0
0.5
0.0
−1.0
1
2
5
10
20
50
100
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
1.0
0.5
0.0
−1.0
−1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.4
0.2
0.0
−0.4
2
4
6
8
10
Lag (yrs)


Turbot in IV Diagnostics − SNS, age 6
a) Observed and fitted values time series
Observed
Fitted
values
0.5
2.0
5.0
20.0
2004
2006
2008
2010
2012
2014
2016
Year
b) Observed vs fitted values
1:1 line
Observed values
0.5
1.0
2.0
5.0
10.0
20.0
50.0
Fitted values
c) Standardised residuals over time
Standardised Residuals
−1.5
−0.5
0.5
1.5
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
−1.5
−1.0
−0.5
0.0
0.5
1.0
1.5
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
−0.4
0.0
0.2
0.4
2
4
6
8
10
Lag (yrs)


Turbot in IV
itrans_rho.1
transfIRARdist.2
transfIRARdist.1
logSdLogObs.14
logSdLogObs.13
logSdLogObs.12
logSdLogObs.11
logSdLogObs.10
logSdLogObs.9
logSdLogObs.8
logSdLogObs.7
logSdLogObs.6
logSdLogObs.5
logSdLogObs.4
logSdLogObs.3
logSdLogObs.2
logSdLogObs.1
logSdLogN.2
logSdLogN.1
logSdLogFsta.5
logSdLogFsta.4
logSdLogFsta.3
logSdLogFsta.2
logSdLogFsta.1
logFpar.8
logFpar.7
logFpar.6
logFpar.5
logFpar.4
logFpar.3
logFpar.2
logFpar.1
logFpar.1
logFpar.2
logFpar.3
logFpar.4
logFpar.5
logFpar.6
logFpar.7
logFpar.8
logSdLogFsta.1
logSdLogFsta.2
logSdLogFsta.3
logSdLogFsta.4
logSdLogFsta.5
logSdLogN.1
logSdLogN.2
logSdLogObs.1
logSdLogObs.2
logSdLogObs.3
logSdLogObs.4
logSdLogObs.5
logSdLogObs.6
logSdLogObs.7
logSdLogObs.8
logSdLogObs.9
logSdLogObs.10
logSdLogObs.11
logSdLogObs.12
logSdLogObs.13
logSdLogObs.14
transfIRARdist.1
transfIRARdist.2
itrans_rho.1
1.0
0.5
0.0
−0.5
−1.0


Observation variance vs uncertainty
catch 2
catch 1
NL_LPUE 1
catch 6
SNS 4
BTS−ISIS 4
BTS−ISIS 5
0.5
0.4
0.3
0.2
CV of estimate
0.1
0.2
0.5
1.0
Observation variance


Survey catchability parameters
SNS
BTS−ISIS
Catchability
0.005
0.010
0.015
0.020
1e−04
2e−04
3e−04
1
2
3
4
5
6
7
Age


Observation variances by data source
Observation Variance
0.0
0.2
0.4
0.6
0.8
catch
SNS
BTS−ISIS
NL_LPUE
NL_LPUE 1
catch 3
catch 4
catch 2
catch 5
catch 6
BTS−ISIS 1
BTS−ISIS 2
BTS−ISIS 3
BTS−ISIS 4
SNS 1
SNS 2
catch 1
SNS 3
catch 7
catch 8
catch 9
catch 10
SNS 5
SNS 6
BTS−ISIS 5
SNS 4
BTS−ISIS 6
BTS−ISIS 7


Selectivity of the Fishery by Pentad
2000's
2005's
2010's
2015's
F/Fbar
Age
0.0
0.5
1.0
1.5
2.0
2
4
6
8
10


Turbot in IV
Spawning stock biomass
SSB
30000
20000
10000
0
Fishing mortality
Fbar
0.5
0.4
0.3
0.2
0.1
0.0
Recruitment
Rec
20000
15000
10000
5000
0
2000
2005
2010
2015
Year


Residuals by year Catch
age
year
10
8
6
4
2
2000
2005
2010
2015


Residuals by survey
BTS−ISIS
SNS
2000
2005
2010
2015
7
6
5
4
3
2
1
age
year

SNS




Spawning stock biomass
SSB
25000
20000
15000
10000
5000
0
Fishing mortality
Fbar
0.4
0.3
0.2
0.1
0.0
Recruitment
Rec
10000
5000
0
2000
2005
2010
2015
Year
2016
2015
2014
2013
2012
2011
2010
2009


logSdLogN
logSdLogObs
logFpar
logSdLogFsta
Parameter value
Assessment year
2016 2015 2014 2013 2012 2011 2010 2009
0.0
0.1
0.2
0.3
0.4
0.2
0.4
0.6
0.8
1.0
0.000
0.005
0.010
0.015
0.020


logSdLogN
logSdLogObs
logFpar
logSdLogFsta
Parameter value
Assessment year
2016 2015 2014 2013 2012 2011 2010 2009
0.0
0.1
0.2
0.3
0.4
0.2
0.4
0.6
0.8
1.0
0.000
0.005
0.010
0.015
0.020


Retrospective pattern in F at age
2013
2014
2015
2010
2011
2012
2007
2008
2009
F
Ages
2008
2009
2010
2011
2012
2013
2014
2015


AIC
1000
900
800
700
600
500
Mohns rho (7−year peel)
−20
−25
−30
−35
−40
−45