
Turbot in IV pgSurvey 4 Diagnostics − BTS−ISIS, age 1
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV pgSurvey 4 Diagnostics − BTS−ISIS, age 2
a) Observed and fitted values time series
Observed
Fitted
values
0.2
0.5
2.0
5.0
1990
1995
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV pgSurvey 4 Diagnostics − BTS−ISIS, age 3
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV pgSurvey 4 Diagnostics − BTS−ISIS, age 4
a) Observed and fitted values time series
Observed
Fitted
values
Year
b) Observed vs fitted values
1:1 line
Observed values
Fitted values
c) Standardised residuals over time
Standardised Residuals
Year
d) Tukey−Anscombe plot
Standardised Residuals
Fitted values
e) Normal Q−Q plot
qqline
1:1 line
Standardised Residuals
Quantiles of the Normal Distribution
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
Lag (yrs)


Turbot in IV pgSurvey 4 Diagnostics − catch, age 1
a) Observed and fitted values time series
Observed
Fitted
values [10^3]
5e−03
5e−02
5e−01
5e+00
1980
1985
1990
1995
2000
2005
2010
2015
Year
b) Observed vs fitted values
1:1 line
Observed values [10^3]
Fitted values [10^3]
c) Standardised residuals over time
Standardised Residuals
−1
0
1
d) Tukey−Anscombe plot
e) Normal Q−Q plot
qqline
Quantiles of the Normal Distribution
−2
−1
0
1
2
f) Autocorrelation of Residuals
95% Conf. Int.
ACF
0.3
0.1
−0.1
−0.3
2
4
6
8
10
12
14
Lag (yrs)


a) Observed and fitted values time series
Observed
Fitted
values [10^3]
20.0
5.0
1.0
0.2
1980
1985
1990
1995
2000
2005
2010
2015
Year


b) Observed vs fitted values
1:1 line
Observed values [10^3]
20.0
5.0
1.0
0.2
0.2
0.5
1.0
2.0
5.0
10.0
20.0
Fitted values [10^3]

Turbot in IV pgSurvey 4




Survey catchability parameters
SNS
BTS−ISIS
Catchability
0.025
0.020
0.015
0.010
0.005
0.00035
0.00030
0.00025
0.00020
0.00015
0.00010
1.0
1.5
2.0
2.5
3.0
3.5
4.0
Age
Observation variances by data source
Observation Variance
0.0
0.2
0.4
0.6
0.8
catch
SNS
BTS−ISIS
NL_LPUE
IBTS_Q1
catch 3
catch 4
LPUE 1
catch 5
catch 6
catch 2
S−ISIS 1
S−ISIS 2
S−ISIS 3
SNS 1
SNS 2
S−ISIS 4
catch 7
catch 8
catch 9
catch 10
TS_Q1 1
catch 1
SNS 3
SNS 4


Selectivity of the Fishery by Pentad
1980's
1985's
1990's
1995's
2000's
2005's
2010's
2015's
F/Fbar
Age
0.0
0.5
1.0
1.5
2.0
2
4
6
8
10


Turbot in IV pgSurvey 4
Spawning stock biomass
SSB
30000
20000
10000
0
Fishing mortality
Fbar
1.2
1.0
0.8
0.6
0.4
0.2
0.0
Recruitment
Rec
15000
10000
5000
0
1980
1990
2000
2010
Year


Residuals by year Catch
age
year
10
8
6
4
2
1980
1990
2000
2010


Residuals by survey
BTS−ISIS
SNS
1990
1995
2000
2005
2010
2015
4.0
3.5
3.0
2.5
2.0
1.5
1.0
age
year

SNS




Spawning stock biomass
SSB
25000
20000
15000
10000
5000
0
Fishing mortality
Fbar
0.8
0.6
0.4
0.2
0.0
Recruitment
Rec
12000
10000
8000
6000
4000
2000
0
1980
1990
2000
2010
Year
2016
2015
2014
2013
2012
2011
2010
2009


logSdLogN
logSdLogObs
logFpar
logSdLogFsta
Parameter value
Assessment year
2016 2015 2014 2013 2012 2011 2010 2009
0.15 0.20 0.25 0.30 0.35
0.0 0.2 0.4 0.6 0.8 1.0
0.000 0.005 0.010 0.015 0.020 0.025
0.20 0.25 0.30 0.35 0.40 0.45


logSdLogN
logSdLogObs
logFpar
logSdLogFsta
Parameter value
Assessment year
2016 2015 2014 2013 2012 2011 2010 2009
0.15 0.20 0.25 0.30 0.35
0.0 0.2 0.4 0.6 0.8 1.0
0.000 0.005 0.010 0.015 0.020 0.025
0.20 0.25 0.30 0.35 0.40 0.45


Retrospective pattern in F at age
2015
2016
2011
2012
2013
2014
2007
2008
2009
2010
F
Ages
2009
2010
2011
2012
2013
2014
2015
2016


AIC
1000
900
800
700
600
500
Mohns rho (7−year peel)
−25
−30
−35
−40
−45
−50