

Skolemization Beyond Intuitionistic Logic: The Role of Quantifier Shifts


Raheleh Jalali

(joint work with Matthias Baaz, Mariami Gamsakhurdia, and Rosalie Iemhoff)

TABLEAUX 2025

28 September, 2025





- 1 Motivation
- 2 Logic of Quantifier Shifts, QFS
- 3 QFS admits Skolemization

Skolemization

A general practice: Suppose in the process of proving a theorem, you get
“for each x there is a y such that $A(x, y)$ ”.

Then, it is convenient to introduce a fresh function f that picks a y for each x , such that $A(x, f(x))$ holds for each x .

Skolemization

A general practice: Suppose in the process of proving a theorem, you get
“for each x there is a y such that $A(x, y)$ ”.

Then, it is convenient to introduce a fresh function f that picks a y for each x , such that $A(x, f(x))$ holds for each x .

Skolemization is defined either in the context of derivability or satisfiability:

- (model-theoretic view): replacing **existential** quantifiers by new function symbols (then, a satisfiability-equivalent formula is obtained).

Skolemization

A general practice: Suppose in the process of proving a theorem, you get
“for each x there is a y such that $A(x, y)$ ”.

Then, it is convenient to introduce a fresh function f that picks a y for each x , such that $A(x, f(x))$ holds for each x .

Skolemization is defined either in the context of derivability or satisfiability:

- (model-theoretic view): replacing **existential** quantifiers by new function symbols (then, a satisfiability-equivalent formula is obtained).
- (proof-theoretic view): replacing **universal** quantifiers by new function symbols (then, a validity-equivalent formula is obtained).

Skolemization

A general practice: Suppose in the process of proving a theorem, you get
“for each x there is a y such that $A(x, y)$ ”.

Then, it is convenient to introduce a fresh function f that picks a y for each x , such that $A(x, f(x))$ holds for each x .

Skolemization is defined either in the context of derivability or satisfiability:

- (model-theoretic view): replacing **existential** quantifiers by new function symbols (then, a satisfiability-equivalent formula is obtained).
- (proof-theoretic view): replacing **universal** quantifiers by new function symbols (then, a validity-equivalent formula is obtained).

As we are considering Skolemization for any logic in general, we will choose the proof theoretic view.

Skolemization in Classical logic

Skolemization

Skolemization (proof-theoretic view): a method to remove *strong* quantifiers (i.e., **positive occurrences of the universal quantifier** and **negative occurrences of the existential quantifier**) from a first-order formula φ , and replace them with *fresh* function symbols.



Skolemization in Classical logic

Skolemization

Skolemization (proof-theoretic view): a method to remove *strong* quantifiers (i.e., **positive occurrences of the universal quantifier** and **negative occurrences of the existential quantifier**) from a first-order formula φ , and replace them with *fresh* function symbols.



The result is called Skolemization of φ and is denoted by φ^S .

Example

$$(\forall x \exists y \forall z \varphi(x, y, z))^S = \exists y \varphi(c, y, f(y)).$$

- This idea goes back to Skolem and it works well in the classical predicate logic, CQC. Informally, Skolemization is blind with respect to the place of the strong quantifier; removes it and replaces it with a fresh function.

- This idea goes back to Skolem and it works well in the classical predicate logic, CQC. Informally, Skolemization is blind with respect to the place of the strong quantifier; removes it and replaces it with a fresh function.
- First impression: Skolemization may not be done for intuitionistic predicate logic, IQC, because we know that quantifiers and connectives do not commute freely.

- This idea goes back to Skolem and it works well in the classical predicate logic, CQC. Informally, Skolemization is blind with respect to the place of the strong quantifier; removes it and replaces it with a fresh function.
- First impression: Skolemization may not be done for intuitionistic predicate logic, IQC, because we know that quantifiers and connectives do not commute freely.
- Indeed! In the context of intuitionistic logic, as well as many intermediate logics, Skolemization is a non-trivial affair.

Definition

Skolemization is *sound* and *complete* for a logic L , (\Rightarrow and \Leftarrow , resp.) when for any formula φ we have

$$L \vdash \varphi \Leftrightarrow L \vdash \varphi^S.$$

Sometimes we also say L *admits* Skolemization.

Definition

Skolemization is *sound* and *complete* for a logic L , (\Rightarrow and \Leftarrow , resp.) when for any formula φ we have

$$L \vdash \varphi \Leftrightarrow L \vdash \varphi^S.$$

Sometimes we also say L *admits* Skolemization.

- (Mints '66) For prenex formulas Skolemization is sound and complete in the setting of IQC.
- (Baaz, Iemhoff '10, '16, '21) Alternative methods of Skolemization in intermediate logics (in certain conservative extensions of IQC).
- (Baaz, Metcalfe, Cintula '08, '15) Skolemization for substructural and fuzzy logics.

A natural question

What if we strengthen IQC in such a way that **quantifiers and connectives commute** freely? (and as a consequence for each formula there is a provably equivalent formula in ***prenex normal form***)

Does the new logic have Skolemization? If not, for which class of formulas does the Skolemization hold?

The motivation of this research!

Outline

- 1 Motivation
- 2 Logic of Quantifier Shifts, QFS
- 3 QFS admits Skolemization

Quantifier shift principles

Consider the commuting of each quantifier \forall, \exists with each connective $\wedge, \vee, \rightarrow$. The following principles, that we call the *quantifier shifts*, are the only ones not valid in IQC:

Quantifier shift principles

Consider the commuting of each quantifier \forall, \exists with each connective $\wedge, \vee, \rightarrow$. The following principles, that we call the *quantifier shifts*, are the only ones not valid in IQC:

Definition

Let $A(x)$ and B be formulas in the first-order language and the variable x is not free in B .

- (Constant Domain) $\forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$ (CD)

Quantifier shift principles

Consider the commuting of each quantifier \forall, \exists with each connective $\wedge, \vee, \rightarrow$. The following principles, that we call the *quantifier shifts*, are the only ones not valid in IQC:

Definition

Let $A(x)$ and B be formulas in the first-order language and the variable x is not free in B .

- (Constant Domain) $\forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$ (CD)
- (Quantifier Switch) $(\forall xA(x) \rightarrow B) \rightarrow \exists x(A(x) \rightarrow B)$ (SW)

Quantifier shift principles

Consider the commuting of each quantifier \forall, \exists with each connective $\wedge, \vee, \rightarrow$. The following principles, that we call the *quantifier shifts*, are the only ones not valid in IQC:

Definition

Let $A(x)$ and B be formulas in the first-order language and the variable x is not free in B .

- (Constant Domain) $\forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$ (CD)
- (Quantifier Switch) $(\forall xA(x) \rightarrow B) \rightarrow \exists x(A(x) \rightarrow B)$ (SW)
- (Existential Distribution) $(B \rightarrow \exists xA(x)) \rightarrow \exists x(B \rightarrow A(x))$ (ED)

Quantifier shift principles

Consider the commuting of each quantifier \forall, \exists with each connective $\wedge, \vee, \rightarrow$. The following principles, that we call the *quantifier shifts*, are the only ones not valid in IQC:

Definition

Let $A(x)$ and B be formulas in the first-order language and the variable x is not free in B .

- (Constant Domain) $\forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$ (CD)
- (Quantifier Switch) $(\forall xA(x) \rightarrow B) \rightarrow \exists x(A(x) \rightarrow B)$ (SW)
- (Existential Distribution) $(B \rightarrow \exists xA(x)) \rightarrow \exists x(B \rightarrow A(x))$ (ED)

None of these formulas are provable in IQC. However, in CQC, both these formulas and their converses are provable.

Quantifier shift principles

Consider the commuting of each quantifier \forall, \exists with each connective $\wedge, \vee, \rightarrow$. The following principles, that we call the *quantifier shifts*, are the only ones not valid in IQC:

Definition

Let $A(x)$ and B be formulas in the first-order language and the variable x is not free in B .

- (Constant Domain) $\forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$ (CD)
- (Quantifier Switch) $(\forall xA(x) \rightarrow B) \rightarrow \exists x(A(x) \rightarrow B)$ (SW)
- (Existential Distribution) $(B \rightarrow \exists xA(x)) \rightarrow \exists x(B \rightarrow A(x))$ (ED)

None of these formulas are provable in IQC. However, in CQC, both these formulas and their converses are provable.

Denote the logic $\text{IQC} + \{\text{CD}, \text{SW}, \text{ED}\}$ by QFS, which we call the *logic of quantifier shifts*.

Main Result

We provide a characterization of all intermediate logics that satisfy the soundness and completeness of Skolemization.

Theorem

An intermediate logic admits Skolemization if and only if it contains all quantifier shift principles.

Definition

- A *Kripke frame* for IQC is a triple (W, \leq, D) , where $W \neq \emptyset$ is a set of worlds, \leq is a binary **reflexive** and **transitive** relation over W , and D is a function assigning to each $w \in W$ a non-empty set $D(w)$, called the *domain* of w , such that if $w \leq w'$ then $D(w) \subseteq D(w')$.
- A *Kripke model* for IQC is a quadruple (W, \leq, D, V) where (W, \leq, D) is a Kripke frame and V is a valuation function in its usual sense.

Definition

A formula A is defined to be *valid* in a frame F , denoted by $F \models A$, and valid in a model M , denoted by $M \models A$, as usual:

- ▶ $M, v \Vdash \top$, $M, v \not\Vdash \perp$
- ▶ $M, v \Vdash p$ iff $v \in V(p)$, for a propositional variable p ,
- ▶ $M, v \Vdash A \wedge B$ iff $M, v \Vdash A$ and $M, v \Vdash B$
- ▶ $M, v \Vdash A \vee B$ iff $M, v \Vdash A$ or $M, v \Vdash B$
- ▶ $M, v \Vdash A \rightarrow B$ iff $\forall w \succcurlyeq v$ if $M, w \Vdash A$ then $M, w \Vdash B$
- ▶ $M, v \Vdash \exists x A(x)$ iff $\exists d \in D(v)$ such that $M, v \Vdash A(d)$
- ▶ $M, v \Vdash \forall x A(x)$ iff $\forall w \succcurlyeq v$ and $\forall d \in D(w)$ we have $M, w \Vdash A(d)$

Definition

A formula A is defined to be *valid* in a frame F , denoted by $F \models A$, and valid in a model M , denoted by $M \models A$, as usual:

- ▶ $M, v \Vdash \top$, $M, v \not\Vdash \perp$
- ▶ $M, v \Vdash p$ iff $v \in V(p)$, for a propositional variable p ,
- ▶ $M, v \Vdash A \wedge B$ iff $M, v \Vdash A$ and $M, v \Vdash B$
- ▶ $M, v \Vdash A \vee B$ iff $M, v \Vdash A$ or $M, v \Vdash B$
- ▶ $M, v \Vdash A \rightarrow B$ iff $\forall w \geq v$ if $M, w \Vdash A$ then $M, w \Vdash B$
- ▶ $M, v \Vdash \exists x A(x)$ iff $\exists d \in D(v)$ such that $M, v \Vdash A(d)$
- ▶ $M, v \Vdash \forall x A(x)$ iff $\forall w \geq v$ and $\forall d \in D(w)$ we have $M, w \Vdash A(d)$

A Kripke frame is *linear* when for any $w, w' \in W$ either $w \leq w'$ or $w' \leq w$.

Definition

A formula A is defined to be *valid* in a frame F , denoted by $F \models A$, and valid in a model M , denoted by $M \models A$, as usual:

- ▶ $M, v \Vdash \top$, $M, v \nVdash \perp$
- ▶ $M, v \Vdash p$ iff $v \in V(p)$, for a propositional variable p ,
- ▶ $M, v \Vdash A \wedge B$ iff $M, v \Vdash A$ and $M, v \Vdash B$
- ▶ $M, v \Vdash A \vee B$ iff $M, v \Vdash A$ or $M, v \Vdash B$
- ▶ $M, v \Vdash A \rightarrow B$ iff $\forall w \geq v$ if $M, w \Vdash A$ then $M, w \Vdash B$
- ▶ $M, v \Vdash \exists x A(x)$ iff $\exists d \in D(v)$ such that $M, v \Vdash A(d)$
- ▶ $M, v \Vdash \forall x A(x)$ iff $\forall w \geq v$ and $\forall d \in D(w)$ we have $M, w \Vdash A(d)$

A Kripke frame is *linear* when for any $w, w' \in W$ either $w \leq w'$ or $w' \leq w$.
A Kripke frame is *constant domain* if for any $w, w' \in W$, $D(w) = D(w')$.

Definition

A formula A is defined to be *valid* in a frame F , denoted by $F \models A$, and valid in a model M , denoted by $M \models A$, as usual:

- ▶ $M, v \Vdash \top$, $M, v \nVdash \perp$
- ▶ $M, v \Vdash p$ iff $v \in V(p)$, for a propositional variable p ,
- ▶ $M, v \Vdash A \wedge B$ iff $M, v \Vdash A$ and $M, v \Vdash B$
- ▶ $M, v \Vdash A \vee B$ iff $M, v \Vdash A$ or $M, v \Vdash B$
- ▶ $M, v \Vdash A \rightarrow B$ iff $\forall w \geq v$ if $M, w \Vdash A$ then $M, w \Vdash B$
- ▶ $M, v \Vdash \exists x A(x)$ iff $\exists d \in D(v)$ such that $M, v \Vdash A(d)$
- ▶ $M, v \Vdash \forall x A(x)$ iff $\forall w \geq v$ and $\forall d \in D(w)$ we have $M, w \Vdash A(d)$

A Kripke frame is *linear* when for any $w, w' \in W$ either $w \leq w'$ or $w' \leq w$.
A Kripke frame is *constant domain* if for any $w, w' \in W$, $D(w) = D(w')$.
For any $w \in W$ define $\leq [w] := \{v \in W \mid v \geq w\}$.

Example of failure of Skolemization in IQC

Example

The axiom $\forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$ is not provable in IQC (as illustrated below) but its Skolemization $\forall x(A(x) \vee B) \rightarrow A(c) \vee B$ is.

$$D_v = \{c, d\} \quad v \models A(c), B \quad v \not\models A(d)$$



$$D_w = \{c\} \quad w \models A(c) \quad w \not\models B$$

Because

$$w \models \forall x(A(x) \vee B) \quad \text{but} \quad w \not\models \forall xA(x) \vee B$$

Class of frames

As the first step of studying the logic of quantifier shifts, let us investigate the semantics.

Definition

Let $F = (W, \leq, D)$ be a frame. Define:

(WF) $\forall w \in W$ the set $\leq [w]$ is well-founded.

(cWF) $\forall w \in W$ the set $\leq [w]$ is conversely well-founded.

Class of frames

As the first step of studying the logic of quantifier shifts, let us investigate the semantics.

Definition


Let $F = (W, \leq, D)$ be a frame. Define:

(WF) $\forall w \in W$ the set $\leq [w]$ is well-founded.

(cWF) $\forall w \in W$ the set $\leq [w]$ is conversely well-founded.

Remark

A frame F satisfying WF does not imply that \leq is well-founded on W :



Class of frames

We present rich classes of frames for QFS and its fragments.

Definition

Define \mathcal{F} as the class of the following Kripke frames F closed under the disjoint union: F is **constant domain** with domain \mathcal{D} , satisfying one of the following conditions:

- 1 $|\mathcal{D}| = 1$,
- 2 $|\mathcal{D}| > 1$, \mathcal{D} is finite, and F is **linear**,
- 3 \mathcal{D} is infinite, F is **linear**, and satisfies both WF and cWF.

Class of frames

We present rich classes of frames for QFS and its fragments.

Definition

Define \mathcal{F} as the class of the following Kripke frames F closed under the disjoint union: F is **constant domain** with domain \mathcal{D} , satisfying one of the following conditions:

- 1 $|\mathcal{D}| = 1$,
- 2 $|\mathcal{D}| > 1$, \mathcal{D} is finite, and F is **linear**,
- 3 \mathcal{D} is infinite, F is **linear**, and satisfies both WF and cWF.

Define \mathcal{F}_{ED} (resp. \mathcal{F}_{SW}) as the class of Kripke frames containing \mathcal{F} and also frames of the form

- constant domain, linear, infinite domains, satisfying WF (resp. cWF).

Frame characterization

$$\text{CD} : \forall x(A(x) \vee B) \rightarrow \forall xA(x) \vee B$$

$$\text{SW} : (\forall xA(x) \rightarrow B) \rightarrow \exists x(A(x) \rightarrow B)$$

$$\text{ED} : (B \rightarrow \exists xA(x)) \rightarrow \exists x(B \rightarrow A(x))$$


Theorem (Frame characterization)

Let F be a frame.

- $F \models \text{CD}$ *if and only if* F is constant domain.
- $F \models \text{ED}$ *if and only if* $F \in \mathcal{F}_{\text{ED}}$.
- $F \models \text{SW}$ *if and only if* $F \in \mathcal{F}_{\text{SW}}$.
- $F \models \text{QFS}$ *if and only if* $F \in \mathcal{F}$.

ED : $(B \rightarrow \exists x A(x)) \rightarrow \exists x (B \rightarrow A(x))$ SW : $(\forall x A(x) \rightarrow B) \rightarrow \exists x (A(x) \rightarrow B)$

Example



- The left frame is a frame for ED:

$$w_1 \Vdash B \rightarrow \exists x A(x) \quad \text{and} \quad w_1 \Vdash \exists x (B \rightarrow A(x))$$

- The right one is a frame for QFS.

Frame incompleteness

Definition

The logic L is *sound and complete* w.r.t. the class \mathcal{C} of Kripke frames when for any formula φ

$$L \vdash \varphi \quad \text{if and only if} \quad \mathcal{C} \models \varphi.$$

Frame incompleteness

Definition

The logic L is *sound and complete* w.r.t. the class \mathcal{C} of Kripke frames when for any formula φ

$$L \vdash \varphi \quad \text{if and only if} \quad \mathcal{C} \models \varphi.$$

Definition

The logic L is called *frame-incomplete* if there is no class \mathcal{C} of Kripke frames such that L is sound and complete with respect to \mathcal{C} .

Frame incompleteness

Definition

The logic L is *sound and complete* w.r.t. the class \mathcal{C} of Kripke frames when for any formula φ

$$L \vdash \varphi \quad \text{if and only if} \quad \mathcal{C} \models \varphi.$$

Definition

The logic L is called *frame-incomplete* if there is no class \mathcal{C} of Kripke frames such that L is sound and complete with respect to \mathcal{C} .

Theorem

The following logics are frame-incomplete:

$$\text{QFS} \quad \text{IQC} + \{\text{CD}, \text{SW}\} \quad \text{IQC} + \{\text{CD}, \text{ED}\}$$

Proof sketch for QFS

Proof.

To show that QFS is frame-incomplete, we have to prove that for any class \mathcal{C} of Kripke frames for QFS, there exists a formula φ such that $\mathcal{C} \models \varphi$ but $\text{QFS} \not\models \varphi$.

Proof sketch for QFS

Proof.

To show that QFS is frame-incomplete, we have to prove that for any class \mathcal{C} of Kripke frames for QFS, there exists a formula φ such that $\mathcal{C} \models \varphi$ but $\text{QFS} \not\models \varphi$. We claim taking an instance of $\varphi = \text{Lin} \vee \text{OEP}$ works, where

$$\text{Lin} := (C \rightarrow D) \vee (D \rightarrow C) \quad \text{and} \quad \text{OEP} := \exists x A(x) \rightarrow \forall x A(x)$$

are the Linearity and One Element Principle schemata.

Proof sketch for QFS

Proof.

To show that QFS is frame-incomplete, we have to prove that for any class \mathcal{C} of Kripke frames for QFS, there exists a formula φ such that $\mathcal{C} \models \varphi$ but $\text{QFS} \not\models \varphi$. We claim taking an instance of $\varphi = \text{Lin} \vee \text{OEP}$ works, where

$$\text{Lin} := (C \rightarrow D) \vee (D \rightarrow C) \quad \text{and} \quad \text{OEP} := \exists x A(x) \rightarrow \forall x A(x)$$

are the Linearity and One Element Principle schemata. The reason is:

- 1 If a frame F validates QFS, then it is constant domain. Moreover, F is either linear or its domain is just a singleton. This means that F validates all instances of the axiom schema $\text{Lin} \vee \text{OEP}$.

Proof sketch for QFS

Proof.

To show that QFS is frame-incomplete, we have to prove that for any class \mathcal{C} of Kripke frames for QFS, there exists a formula φ such that $\mathcal{C} \models \varphi$ but $\text{QFS} \not\models \varphi$. We claim taking an instance of $\varphi = \text{Lin} \vee \text{OEP}$ works, where

$$\text{Lin} := (C \rightarrow D) \vee (D \rightarrow C) \quad \text{and} \quad \text{OEP} := \exists x A(x) \rightarrow \forall x A(x)$$

are the Linearity and One Element Principle schemata. The reason is:

- 1 If a frame F validates QFS, then it is constant domain. Moreover, F is either linear or its domain is just a singleton. This means that F validates all instances of the axiom schema $\text{Lin} \vee \text{OEP}$.
- 2 There is an instance of $\text{Lin} \vee \text{OEP}$ such that $\text{QFS} \not\models \text{Lin} \vee \text{OEP}$.

Proof sketch for QFS

Proof.

To show that QFS is frame-incomplete, we have to prove that for any class \mathcal{C} of Kripke frames for QFS, there exists a formula φ such that $\mathcal{C} \models \varphi$ but $\text{QFS} \not\models \varphi$. We claim taking an instance of $\varphi = \text{Lin} \vee \text{OEP}$ works, where

$$\text{Lin} := (C \rightarrow D) \vee (D \rightarrow C) \quad \text{and} \quad \text{OEP} := \exists x A(x) \rightarrow \forall x A(x)$$

are the Linearity and One Element Principle schemata. The reason is:

- 1 If a frame F validates QFS, then it is constant domain. Moreover, F is either linear or its domain is just a singleton. This means that F validates all instances of the axiom schema $\text{Lin} \vee \text{OEP}$.
- 2 There is an instance of $\text{Lin} \vee \text{OEP}$ such that $\text{QFS} \not\models \text{Lin} \vee \text{OEP}$.

These two points together prove that QFS is frame-incomplete. □

The following model is a model of QFS but $w \not\models$ Lin \vee OEP. Let $\mathcal{D}_w = \mathcal{D}_{v_1} = \mathcal{D}_{v_2} = \mathcal{D} = \{a, b\}$.

$$v_1 \models R(a), R(b), P \quad v_1 \not\models Q$$

$$v_2 \models R(a), R(b), Q \quad v_2 \not\models P$$

$$w \models R(a)$$



Separation of fragments

What do we know about fragments of QFS? Are they distinct?

Theorem

The following hold:

- ① $CD, SW \not\vdash ED$ (hence: $CD \not\vdash ED$ and $SW \not\vdash ED$)
- ② $CD, ED \not\vdash SW$ (hence: $CD \not\vdash SW$ and $ED \not\vdash SW$)

Therefore, QFS, IQC + {CD, SW}, and IQC + {CD, ED} are all distinct.

Separation of fragments

What do we know about fragments of QFS? Are they distinct?

Theorem

The following hold:

- ① $CD, SW \not\vdash ED$ (*hence: $CD \not\vdash ED$ and $SW \not\vdash ED$*)
- ② $CD, ED \not\vdash SW$ (*hence: $CD \not\vdash SW$ and $ED \not\vdash SW$*)

Therefore, QFS, IQC + {CD, SW}, and IQC + {CD, ED} are all distinct.

Theorem

$SW \vdash CD$ *and* $ED \not\vdash CD$

Outline

- 1 Motivation
- 2 Logic of Quantifier Shifts, QFS
- 3 QFS admits Skolemization

Revisiting Skolemization

A thought:

Problem

Let A be a formula in prenex normal form. Does $\text{QFS} \vdash A$ imply $\text{IQC} \vdash A$?

Revisiting Skolemization

A thought:

Problem

Let A be a formula in prenex normal form. Does $\text{QFS} \vdash A$ imply $\text{IQC} \vdash A$?

If yes, then the Skolemization of QFS follows from Mints' theorem that the prenex fragment of IQC admits Skolemization.

Corollary

QFS admits Skolemization.

Revisiting Skolemization

A thought:

Problem

Let A be a formula in prenex normal form. Does $\text{QFS} \vdash A$ imply $\text{IQC} \vdash A$?

If yes, then the Skolemization of QFS follows from Mints' theorem that the prenex fragment of IQC admits Skolemization.

Corollary

QFS admits Skolemization.

Unfortunately, this is not the case.

Prenex fragments of QFS and IQC

Example

QFS \vdash CD

QFS $\vdash \forall x(A(x) \vee B) \rightarrow \forall yA(y) \vee B$

QFS $\vdash \forall x(A(x) \vee B) \rightarrow \forall y(A(y) \vee B)$

QFS $\vdash \exists x((A(x) \vee B) \rightarrow \forall y(A(y) \vee B))$

QFS $\vdash \exists x \forall y((A(x) \vee B) \rightarrow (A(y) \vee B))$

Prenex fragments of QFS and IQC

Example

QFS \vdash CD

QFS $\vdash \forall x(A(x) \vee B) \rightarrow \forall yA(y) \vee B$

QFS $\vdash \forall x(A(x) \vee B) \rightarrow \forall y(A(y) \vee B)$

QFS $\vdash \exists x((A(x) \vee B) \rightarrow \forall y(A(y) \vee B))$

QFS $\vdash \exists x\forall y((A(x) \vee B) \rightarrow (A(y) \vee B))$

However, IQC $\not\vdash \exists x\forall y((A(x) \vee B) \rightarrow (A(y) \vee B))$. Take:

$\{1, 2\} \ w_2 \models A(1)$



$\{1\} \ w_1$

where $\mathcal{D}_1 = \{1\}$ and $\mathcal{D}_2 = \{1, 2\}$. Then $w_2 \models A(1) \vee B$ but $w_2 \not\models A(2) \vee B$. Hence, $w_1 \not\models \exists x\forall y((A(x) \vee B) \rightarrow (A(y) \vee B))$.

- Currently, we lack a well-structured proof system, such as a cut-free or analytic sequent calculus for QFS, and we established its frame incompleteness (with respect to Kripke frames).

- Currently, we lack a well-structured proof system, such as a cut-free or analytic sequent calculus for QFS, and we established its frame incompleteness (with respect to Kripke frames).
- So, can we prove QFS admits Skolemization in other ways?

- Currently, we lack a well-structured proof system, such as a cut-free or analytic sequent calculus for QFS, and we established its frame incompleteness (with respect to Kripke frames).
- So, can we prove QFS admits Skolemization in other ways?
- Aguilera and Baaz '19 proposed a sequent calculus for first-order (classical or intuitionistic) logic by **relaxing the usual quantifier-inference restrictions**.

- Currently, we lack a well-structured proof system, such as a cut-free or analytic sequent calculus for QFS, and we established its frame incompleteness (with respect to Kripke frames).
- So, can we prove QFS admits Skolemization in other ways?
- Aguilera and Baaz '19 proposed a sequent calculus for first-order (classical or intuitionistic) logic by **relaxing the usual quantifier-inference restrictions**.
- This calculus: allows **unsound inferences**, **lacks local correctness**, but **has global correctness**. Every lie that has been introduced during a proof, in the end will be cleared.

- Currently, we lack a well-structured proof system, such as a cut-free or analytic sequent calculus for QFS, and we established its frame incompleteness (with respect to Kripke frames).
- So, can we prove QFS admits Skolemization in other ways?
- Aguilera and Baaz '19 proposed a sequent calculus for first-order (classical or intuitionistic) logic by **relaxing the usual quantifier-inference restrictions**.
- This calculus: allows **unsound inferences**, **lacks local correctness**, but **has global correctness**. Every lie that has been introduced during a proof, in the end will be cleared.
- We will use this sequent calculus.

Definitions

- ▶ An inference is a **quantifier inference** if the principal formula has a quantifier as its outermost logical symbol. For instance

$$\frac{\Gamma \Rightarrow A(a)}{\Gamma \Rightarrow \forall x A(x)}$$

Definitions

- ▶ An inference is a **quantifier inference** if the principal formula has a quantifier as its outermost logical symbol. For instance

$$\frac{\Gamma \Rightarrow A(a)}{\Gamma \Rightarrow \forall x A(x)}$$

- ▶ The **characteristic variable** of an inference is a , if the inference yields a strongly quantified formula $\forall x A(x)$ from $A(a)$, where a is a free variable.

Definitions

- ▶ An inference is a **quantifier inference** if the principal formula has a quantifier as its outermost logical symbol. For instance

$$\frac{\Gamma \Rightarrow A(a)}{\Gamma \Rightarrow \forall x A(x)}$$

- ▶ The **characteristic variable** of an inference is a , if the inference yields a strongly quantified formula $QxA(x)$ from $A(a)$, where a is a free variable.
- ▶ Let π be a derivation. We say b is a **side variable** of a in π ($a <_{\pi} b$) if π contains a strong-quantifier inference of one of the forms:

$$\frac{\Gamma \Rightarrow A(a, b, \vec{c})}{\Gamma \Rightarrow \forall x A(x, b, \vec{c})} \quad \frac{A(a, b, \vec{c}), \Gamma \Rightarrow \Delta}{\exists x A(x, b, \vec{c}), \Gamma \Rightarrow \Delta}$$

Sequent calculi **LK** and **LJ**

First-order **LK** is the extension of the usual propositional **LK** for classical logic obtained by adding quantifier inferences:

$$\begin{array}{c} \frac{\Gamma, \varphi(t) \Rightarrow \Delta}{\Gamma, \forall x \varphi(x) \Rightarrow \Delta} \forall L \qquad \frac{\Gamma \Rightarrow \varphi(y), \Delta}{\Gamma \Rightarrow \forall x \varphi(x), \Delta} \forall R \\[1em] \frac{\Gamma, \varphi(y) \Rightarrow \Delta}{\Gamma, \exists x \varphi(x) \Rightarrow \Delta} \exists L \qquad \frac{\Gamma \Rightarrow \varphi(t), \Delta}{\Gamma \Rightarrow \exists x \varphi(x), \Delta} \exists R \end{array}$$

Constraint: in $\exists L$ and $\forall R$ the variable y is **not free** in the conclusion.

Sequent calculi **LK** and **LJ**

First-order **LK** is the extension of the usual propositional **LK** for classical logic obtained by adding quantifier inferences:

$$\frac{\Gamma, \varphi(t) \Rightarrow \Delta}{\Gamma, \forall x \varphi(x) \Rightarrow \Delta} \forall L \qquad \frac{\Gamma \Rightarrow \varphi(y), \Delta}{\Gamma \Rightarrow \forall x \varphi(x), \Delta} \forall R$$
$$\frac{\Gamma, \varphi(y) \Rightarrow \Delta}{\Gamma, \exists x \varphi(x) \Rightarrow \Delta} \exists L \qquad \frac{\Gamma \Rightarrow \varphi(t), \Delta}{\Gamma \Rightarrow \exists x \varphi(x), \Delta} \exists R$$

Constraint: in $\exists L$ and $\forall R$ the variable y is **not free** in the conclusion.

First-order **LJ** for intuitionistic logic is defined as the single-conclusion version of first-order **LK**, i.e., everywhere in the succedents there is at most one formula.

Sequent calculi **LK** and **LJ**

First-order **LK** is the extension of the usual propositional **LK** for classical logic obtained by adding quantifier inferences:

$$\begin{array}{c} \frac{\Gamma, \varphi(t) \Rightarrow \Delta}{\Gamma, \forall x \varphi(x) \Rightarrow \Delta} \forall L \qquad \frac{\Gamma \Rightarrow \varphi(y), \Delta}{\Gamma \Rightarrow \forall x \varphi(x), \Delta} \forall R \\[1em] \frac{\Gamma, \varphi(y) \Rightarrow \Delta}{\Gamma, \exists x \varphi(x) \Rightarrow \Delta} \exists L \qquad \frac{\Gamma \Rightarrow \varphi(t), \Delta}{\Gamma \Rightarrow \exists x \varphi(x), \Delta} \exists R \end{array}$$

Constraint: in $\exists L$ and $\forall R$ the variable y is **not free** in the conclusion.

First-order **LJ** for intuitionistic logic is defined as the single-conclusion version of first-order **LK**, i.e., everywhere in the succedents there is at most one formula.

The sequent calculus **QFS** is defined as **LJ** + {CD, ED, SW}.

Definition (Suitable quantifier inference)

A quantifier inference is *suitable* for a proof π if either it is a weak-quantifier inference, or the following are satisfied:

Definition (Suitable quantifier inference)

A quantifier inference is *suitable* for a proof π if either it is a weak-quantifier inference, or the following are satisfied:

- 1 (substitutability) the characteristic variable does not appear in the conclusion of π .

Definition (Suitable quantifier inference)

A quantifier inference is *suitable* for a proof π if either it is a weak-quantifier inference, or the following are satisfied:

- 1 (substitutability) the characteristic variable does not appear in the conclusion of π .
- 2 (side-variable condition) the relation $<_{\pi}$ is acyclic.

Definition (Suitable quantifier inference)

A quantifier inference is *suitable* for a proof π if either it is a weak-quantifier inference, or the following are satisfied:

- 1 (substitutability) the characteristic variable does not appear in the conclusion of π .
- 2 (side-variable condition) the relation $<_{\pi}$ is acyclic.
- 3 (very weak regularity) whenever the characteristic variable is also the characteristic variable of another strong-quantifier inference in π , then it has the same principal formula.

Definition (Suitable quantifier inference)

A quantifier inference is *suitable* for a proof π if either it is a weak-quantifier inference, or the following are satisfied:

- 1 (substitutability) the characteristic variable does not appear in the conclusion of π .
- 2 (side-variable condition) the relation $<_{\pi}$ is acyclic.
- 3 (very weak regularity) whenever the characteristic variable is also the characteristic variable of another strong-quantifier inference in π , then it has the same principal formula.

For instance, each quantifier inference of **LJ** as in the previous slide is suitable for every regular **LJ**-proof.

Justification of a suitable quantifier

Violation of each condition leads to an **undesirable** proof:

❶ violation of substitutability:
$$\frac{A(a) \Rightarrow A(a)}{A(a) \Rightarrow \forall x A(x)}$$

Justification of a suitable quantifier

Violation of each condition leads to an **undesirable** proof:

❶ violation of substitutability:
$$\frac{A(a) \Rightarrow A(a)}{A(a) \Rightarrow \forall x A(x)}$$

❷ violation of side-variable condition:

$$\frac{\frac{\frac{A(a, b) \Rightarrow A(a, b)}{\exists y A(a, y) \Rightarrow A(a, b)} *_1}{\exists y A(a, y) \Rightarrow \forall x A(x, b)} *_2}{\forall x \exists y A(x, y) \Rightarrow \forall x A(x, b)} \\ \forall x \exists y A(x, y) \Rightarrow \exists y \forall x A(x, y)$$

Note that in $*_1$ and $*_2$ we have $b <_{\pi} a$ and $a <_{\pi} b$, respectively.

Justification of a suitable quantifier

Violation of each condition leads to an **undesirable** proof:

❶ violation of substitutability:
$$\frac{A(a) \Rightarrow A(a)}{A(a) \Rightarrow \forall x A(x)}$$

❷ violation of side-variable condition:

$$\frac{\frac{\frac{A(a, b) \Rightarrow A(a, b)}{\exists y A(a, y) \Rightarrow A(a, b)} *1}{\exists y A(a, y) \Rightarrow \forall x A(x, b)} *2}{\forall x \exists y A(x, y) \Rightarrow \forall x A(x, b)} \\ \forall x \exists y A(x, y) \Rightarrow \exists y \forall x A(x, y)$$

Note that in $*_1$ and $*_2$ we have $b <_{\pi} a$ and $a <_{\pi} b$, respectively.

❸ violation of very weak regularity:

$$\frac{\frac{A(a) \Rightarrow A(a)}{A(a) \Rightarrow \forall x A(x)}}{\frac{A(a) \Rightarrow \forall x A(x) \quad \frac{A(a) \Rightarrow A(a)}{\exists x A(x) \Rightarrow A(a)}}{\exists x A(x) \Rightarrow \forall x A(x)}}$$

Definition (Aguilera, Baaz '19)

The calculus LK^{++} (resp. LJ^{++}) is defined like LK (resp. LJ), except that the constraint on $\exists L$ and $\forall R$ is removed and one adds the restriction that a proof may only contain quantifier inferences that are *suitable* for it.

Definition (Aguilera, Baaz '19)

The calculus LK^{++} (resp. LJ^{++}) is defined like LK (resp. LJ), except that the constraint on $\exists L$ and $\forall R$ is removed and one adds the restriction that a proof may only contain quantifier inferences that are *suitable* for it.

An interesting fact: all the quantifier shift axioms are provable in LJ^{++} .

Theorem (Aguilera, Baaz '19)

Let S be a sequent. We have $LJ^{++} \vdash S$ iff $QFS \vdash S$

Theorem (Aguilera and Baaz '19)

If the Skolemization of $\Gamma \Rightarrow \Delta$ is cut-free \mathbf{LJ}^{++} -derivable, then $\Gamma \Rightarrow \Delta$ is cut-free \mathbf{LJ}^{++} -derivable.

Theorem (Aguilera and Baaz '19)

If the Skolemization of $\Gamma \Rightarrow \Delta$ is cut-free \mathbf{LJ}^{++} -derivable, then $\Gamma \Rightarrow \Delta$ is cut-free \mathbf{LJ}^{++} -derivable.

Corollary

QFS admits Skolemization.

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.
- Then **QFS** $\vdash \Rightarrow A^S$.

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.
- Then **QFS** $\vdash \Rightarrow A^S$.
- Therefore, **LJ** $\vdash B_1, \dots, B_n \Rightarrow A^S$, where each B_i is an instance of CD, ED, or SW.

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.
- Then $\text{QFS} \vdash \Rightarrow A^S$.
- Therefore, $\text{LJ} \vdash B_1, \dots, B_n \Rightarrow A^S$, where each B_i is an instance of CD, ED, or SW.
- As $\text{LJ} \vdash \Rightarrow \varphi \rightarrow \varphi^S$ for any φ and $\text{LJ} \vdash \Rightarrow (B_1 \wedge \dots \wedge B_n) \rightarrow A^S$, we have $\text{LJ} \vdash \Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$.

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.
- Then $\text{QFS} \vdash \Rightarrow A^S$.
- Therefore, $\text{LJ} \vdash B_1, \dots, B_n \Rightarrow A^S$, where each B_i is an instance of CD, ED, or SW.
- As $\text{LJ} \vdash \Rightarrow \varphi \rightarrow \varphi^S$ for any φ and $\text{LJ} \vdash \Rightarrow (B_1 \wedge \dots \wedge B_n) \rightarrow A^S$, we have $\text{LJ} \vdash \Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$.
- Hence, there is a cut-free proof π of $\Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$ in **LJ**.

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.
- Then $\text{QFS} \vdash \Rightarrow A^S$.
- Therefore, $\text{LJ} \vdash B_1, \dots, B_n \Rightarrow A^S$, where each B_i is an instance of CD, ED, or SW.
- As $\text{LJ} \vdash \Rightarrow \varphi \rightarrow \varphi^S$ for any φ and $\text{LJ} \vdash \Rightarrow (B_1 \wedge \dots \wedge B_n) \rightarrow A^S$, we have $\text{LJ} \vdash \Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$.
- Hence, there is a cut-free proof π of $\Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$ in **LJ**.
- Since π is also an **LJ**⁺⁺-proof, by Theorem we obtain $\text{LJ}^{++} \vdash \Rightarrow (B_1 \wedge \dots \wedge B_n) \rightarrow A$,

Proof of Corollary

Proof.

- Suppose $\text{QFS} \vdash A^S$.
- Then $\text{QFS} \vdash \Rightarrow A^S$.
- Therefore, $\text{LJ} \vdash B_1, \dots, B_n \Rightarrow A^S$, where each B_i is an instance of CD, ED, or SW.
- As $\text{LJ} \vdash \Rightarrow \varphi \rightarrow \varphi^S$ for any φ and $\text{LJ} \vdash \Rightarrow (B_1 \wedge \dots \wedge B_n) \rightarrow A^S$, we have $\text{LJ} \vdash \Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$.
- Hence, there is a cut-free proof π of $\Rightarrow ((B_1 \wedge \dots \wedge B_n) \rightarrow A^S)^S$ in **LJ**.
- Since π is also an **LJ**⁺⁺-proof, by Theorem we obtain $\text{LJ}^{++} \vdash \Rightarrow (B_1 \wedge \dots \wedge B_n) \rightarrow A$, which yields $\text{QFS} \vdash \Rightarrow A$ hence $\text{QFS} \vdash A$.



Concluding remarks

- We introduced a logic QFS by adding the quantifier shifts to IQC.
- QFS is Kripke frame-incomplete.
- **Main result:** An intermediate logic admits Skolemization if and only if it contains all quantifier shift principles.

Concluding remarks

- We introduced a logic QFS by adding the quantifier shifts to IQC.
- QFS is Kripke frame-incomplete.
- **Main result:** An intermediate logic admits Skolemization if and only if it contains all quantifier shift principles.

Future work:

- Does the logic QFS have the disjunction property (DP)? (We know that it doesn't have the existence property (EP).)
- Investigate the full power of QFS by considering other axioms not provable in IQC and asking whether they are provable in QFS or not.

Concluding remarks

- We introduced a logic QFS by adding the quantifier shifts to IQC.
- QFS is Kripke frame-incomplete.
- **Main result:** An intermediate logic admits Skolemization if and only if it contains all quantifier shift principles.

Future work:

- Does the logic QFS have the disjunction property (DP)? (We know that it doesn't have the existence property (EP).)
- Investigate the full power of QFS by considering other axioms not provable in IQC and asking whether they are provable in QFS or not.

Thank you for your attention.