

Crossing Gap Using VQE: A comparative study

I-Chi Chen

Iowa State University
Ames, USA

ichen@iastate.edu

Nouhaila Innan[✉]

Quantum Physics and Magnetism Team,
LPMC, Faculty of Sciences Ben M'sick

Hassan II University of Casablanca, Morocco
nouhaila.innan-etu@etu.univh2c.ma

Suman Kumar Roy

Information Technology
NITK Surathkal, India

roysuman.212it031@nitk.edu.in

Jason Saroni

Iowa State University
Ames, USA

jsaroni@iastate.edu

Abstract—Within the evolving domain of quantum computational chemistry, the Variational Quantum Eigensolver (VQE) has been identified as a crucial algorithm for exploiting the capabilities of near-term quantum computers. This work investigates the proficiency of variational quantum algorithms, particularly adaptive VQE techniques, and Quantum Natural Gradient (QNG) optimization, in calculating the spectral gap of chemical molecules—a key determinant of material characteristics and chemical behavior. Our methodology introduces different approaches to estimating the k -th excited state by mapping highly excited states to the ground state via unitary transformations, broadening VQE’s applicability. Our research focuses on analyzing three molecular systems, H_2 , LiH , and BeH_2 , to demonstrate the versatility and precision of our proposed methods. A significant highlight of our study is the implementation of QNG optimization, which markedly improves the optimization process’s efficiency. It consistently minimizes the ground state energy while achieving convergence in fewer iterations than traditional gradient descent optimizers. This efficiency underscores the advantage of QNG in navigating the complex optimization landscape of quantum parameters more effectively. Additionally, our comparative analysis reveals that both subspace variational quantum eigensolver (SS-VQE) and quantum variational deflation (VQD) methods exhibit strong performance in spectral gap determination, with less than 1% relative error for LiH , and notably, VQD achieves an exceptional 99.99999% accuracy. Lastly we explore Pauli tapering and the usage of a tensor network ansatz to improve algorithm speed.

Index Terms—Variational Quantum Eigensolver, Quantum Natural Gradient

I. INTRODUCTION

Propelled by global research, quantum computing is moving closer to realization with the prospect of nearly thousand-qubit devices on the horizon. Known as NISQ (Noisy Intermediate-Scale Quantum) computers, these upcoming marvels lack fault tolerance and complete computational precision. The classic Schrödinger equation poses a significant challenge with wide-ranging implications. By solving it numerically, we can make predictions about atoms, molecules, and materials, driving progress in material design, drug discovery, and scientific exploration. Exploring excited state energies reveals insights into charge transfer in solar materials and sheds light on chemical reactions like photodissociation, where traditional methods often fall short, making quantum approaches essential for understanding the same. Quantum computers offer the potential to solve a range of problems much faster than

classical methods. However, many quantum algorithms require error correction, limiting their immediate practical application. By using hybrid quantum-classical algorithms, it can reduce gate depth, mitigate the impact of errors, and seamlessly integrate classical and quantum subroutines for more efficient computation.

The Variational Quantum Eigensolver (VQE) is a cutting-edge algorithm designed specifically for near-term quantum computers. VQE leverages the efficiency of quantum state storage over traditional methods to determine the lowest eigenvalue of a Hamiltonian (H). By utilizing parameterized quantum circuits, such as $U(\theta)$, to create trial wavefunctions and generate the state $|\psi(\theta)\rangle$, VQE optimizes the expectation value of

$$\langle H \rangle = \langle \psi(\theta) | H | \psi(\theta) \rangle \quad (1)$$

through iterative parameter optimization. The optimal parameters θ^* that minimize $\langle H \rangle$ provide an approximation of the ground state energy. With its streamlined circuit design, VQE shows great promise for advancing quantum-enhanced computation in the foreseeable future.

However, VQE is only designed for solving the lowest state and its energy. There are many VQE’s variants to solve the excited state. Here, We will test different variational quantum techniques to determine the spectral gap of molecules which is the energy difference between the ground state and the first excited state. Moreover, we extend the problem to solve the highly excited states and their spectral gap between the state and the nearest energy states by mapping the highly excited state to the ground state.

II. METHODOLOGY

A. Subspace-Search Variational Quantum Eigensolver

The Subspace-Search Variational Quantum Eigensolver (SS-VQE) is an algorithm designed to address the challenge of calculating excited states [1]. SS-VQE efficiently explores a low-energy subspace to identify the k -th excited state by utilizing orthogonal input states and leveraging unitary transformations. It is worth noting that this method involves only two parameter optimization steps and eliminates the necessity for ancilla qubits. SS-VQE further generalizes all excited states up to the k -th through a single optimization procedure. Through careful parameter optimization, this approach ensures

the orthogonality of input states and accurately maps them to energy eigenstates. This approach minimize the following loss function

$$\mathcal{L}_w(\theta) = w \langle \psi_k | U^\dagger(\theta) H U(\theta) | \psi_k \rangle \quad (2)$$

$$+ \sum_{j=0}^{k-1} \langle \psi_j | U^\dagger(\theta) H U(\theta) | \psi_j \rangle \quad (3)$$

where the cost \mathcal{L}_w gets its global optimum, the circuit $U(\theta)$ maps $|\psi_k\rangle$ to the k -th excited state $|E_k\rangle$ of the Hamiltonian(H) and others to the subspace spanned by $\sum_{j=0}^{k-1} E_j$.

B. Variational Quantum Deflation

The potential of VQE for near-term quantum computing is generating excitement for the advancement of computational capabilities. [2] study aims to enhance VQE's functionality by efficiently identifying excited states. By incorporating "overlap" terms into the optimization function and leveraging Hermitian matrices, which consist of a complete set of orthogonal eigenvectors, this work demonstrates a cost-effective approach. Utilizing VQE's ability to maintain classical parameters, low-depth quantum circuits are employed to compute these overlap terms. This methodology maintains the same qubit count as VQE for ground-state calculations, with only a minor increase in measurements. In contrast to existing methods for computing excited states in quantum computing, this approach minimizes resource overhead. The following loss function has been provided over this work.

$$\mathcal{L}(\lambda_k) = \langle \psi(\lambda_k) | H | \psi(\lambda_k) \rangle \quad (4)$$

$$+ \sum_{i=0}^{k-1} \beta_i | \langle \psi(\lambda_k) | \psi(\lambda_k) \rangle |^2. \quad (5)$$

C. Folded Spectrum VQE

Expanding on the VQE, [3] study introduces a specialized method for calculating molecular excited states. By utilizing the Folded Spectrum (FS) approach, we restructure the Hamiltonian's eigenspectrum to specifically target highly excited states. While FS has been acknowledged in the past, its quantum application was previously considered too costly due to the exponential growth of terms in the measured operator. Nevertheless, our implementation reveals a significant advancement by employing a Pauli grouping technique, which can significantly reduce the number of required measurements, making FS a cost-efficient option. This technique has a particularly notable impact on second quantized molecular Hamiltonians, thanks to their distinct structural properties. The loss function for this procedure is

$$\mathcal{L}(\theta) = \langle \psi(\theta) | (H - \omega)^2 | \psi(\theta) \rangle, \quad (6)$$

where ω is an arbitrary scalar energy value.

D. VQE with Quantum Natural Gradient

The landscape of optimization problems encountered in VQE applications is characteristically intricate, often riddled

with many local minima. This complexity underscores the necessity of employing an effective optimization strategy, pivotal for the algorithm's successful convergence to the ground state energy of the system under study.

Among various optimization techniques, the QNG optimization strategy, which stands out by using the geometric properties of the parameter space [4]. Contrary to the traditional gradient descent method, which operates under the assumption of an Euclidean metric space, QNG employs the Fubini-Study metric tensor, denoted as g , to modulate the optimization step sizes. This tensor captures the variational state space's inherent curvature, facilitating more informed and efficacious optimization steps.

The essence of the QNG approach is encapsulated in the update rule:

$$\theta_{\text{new}} = \theta - \eta g(\theta)^{-1} \nabla f(\theta), \quad (7)$$

where θ denotes the variational circuit parameters, η signifies the learning rate, $g(\theta)$ represents the Fubini-Study metric tensor, and $\nabla f(\theta)$ is the gradient of the objective function with respect to θ . The objective function f typically corresponds to the expectation value of the Hamiltonian, whose ground state energy the VQE seeks to approximate.

By accounting for the parameter space's geometry, the QNG optimizer significantly enhances the efficiency of the optimization process. It navigates the circuit's sensitivity to parameter variations, circumventing suboptimal pathways often pursued by conventional optimization methods.

Algorithm 1 delineates the procedural steps for incorporating QNG optimization within the VQE.

Algorithm 1 VQE with QNG

Require: Hamiltonian H of the quantum system, variational ansatz $U(\theta)$, initial parameters θ_0 , learning rate η

Ensure: Optimized parameters θ^* minimizing the expectation value $\langle H \rangle_\theta$

- 1: Initialize parameters $\theta \leftarrow \theta_0$
 - 2: **while** not converged **do**
 - 3: Evaluate the objective function $f(\theta) = \langle U(\theta)^\dagger | H | U(\theta) \rangle$
 - 4: Compute the gradient $\nabla f(\theta)$ with respect to θ
 - 5: Compute the Fubini-Study metric tensor $g(\theta)$
 - 6: Update the parameters using QNG: $\theta \leftarrow \theta - \eta g(\theta)^{-1} \nabla f(\theta)$
 - 7: Check for convergence (e.g., change in $f(\theta)$ is below a threshold)
 - 8: **end while**
 - 9: $\theta^* \leftarrow \theta$
 - 10: **return** θ^*
-

III. RESULTS

In our study, we meticulously analyzed the efficacy of the SS-VQE for calculating the energy states of LiH , emphasizing its performance in different spin subspaces. The results are

distilled into Figs. 1a and 1b show the energy profiles for the ground and excited states at $S_z = 0$ and $S_z = 1$ cases respectively. Fig. 1a reveals the SS-VQE’s proficiency in approximating the singlet ground state (S0) and triplet first excited state (T1) of *LiH* at $S_z = 0$. The alignment of SS-VQE results with the exact theoretical calculations is particularly noteworthy for shorter bond lengths, a region crucial for understanding chemical bonding and reactions. The observed convergence of energy values at increased atomic separations reflects the algorithm’s capability to capture the dissociative characteristics of the molecule, demonstrating its robustness in quantum state representation for non-interacting atomic scenarios. Fig. 1a extends this comparison to $S_z = 1$, offering insights into the SS-VQE’s ability to model excited spin states accurately. The close correlation between SS-VQE and exact calculations for T1 and T2 states across various atomic distances underscores the versatility of the algorithm. This precision holds even as the bond lengthens, indicating that SS-VQE can adeptly model the *LiH* molecule’s behavior under stretched conditions, which is vital for simulating molecular dynamics.

In Fig. 2a, the relative error for the S0 and T1 states at $S_z = 0$ is quantified, exhibiting SS-VQE’s remarkable precision with errors maintained below 0.8%. This high accuracy is retained across the spectrum of bond lengths, confirming the algorithm’s exceptional ability to capture quantum mechanical interactions within the molecule. Fig. 2b further corroborates the method’s precision, where the SS-VQE sustains an impressively low relative error for the T1 and T2 states at $S_z = 1$. With errors well below 0.003%, SS-VQE is an invaluable tool for precisely estimating quantum states, particularly in molecules exhibiting non-zero spin states.

These compelling findings highlight SS-VQE’s potential to transform the landscape of quantum computational chemistry. The accuracy and precision demonstrated in this study pave the way for advanced modeling of molecular systems, offering a new lens through which we can observe and predict the quantum behavior of chemical substances. The implications of this work are profound, promising advancements in fields ranging from material science to pharmaceutical development, where the understanding of quantum states is fundamental.

The VQD approach has been meticulously evaluated for its ability to predict the energy states of *LiH*, focusing on its performance across different spin subspaces, $S_z = 0$ and (b) $S_z = 1$. The analyses encompass both the accuracy of the energy estimations and the relative errors in these calculations. The energy profiles of *LiH* obtained via VQD display remarkable fidelity to the exact theoretical calculations, with a pronounced accuracy in the chemically significant region of shorter bond lengths. As shown in Figs. 3a and 3b, VQD’s calculated energies for the ground and excited states at $S_z = 0$ and $S_z = 1$, respectively, closely track the exact values across a range of atomic separations. This agreement is particularly noticeable in the molecular dissociation limit, where VQD and exact calculations predict the energy levels’ convergence toward a constant value.

The relative error analyses are shown in Figs. 4a and 4b for $S_z = 0$ and $S_z = 1$, further underscore the precision of the VQD approach. The errors remain exceptionally low, with the largest observed errors not exceeding 0.0175% for $S_z = 0$ and peaking just below 0.0010% for the excited states at $S_z = 1$. These low relative errors across the atomic units underscore the VQD’s robustness and its adaptability to quantum mechanical subtleties inherent in the excited states of the molecule.

The comprehensive analysis provided by these figures highlights the VQD’s potential as an indispensable tool for quantum chemists and material scientists. The exceptional accuracy in estimating the energy states in both $S_z = 0$ and $S_z = 1$ subspaces is critical for applications requiring precise energy level transitions, such as in the field of photochemical reactions and the development of photonic materials. The VQD approach has demonstrated its prowess as a highly reliable and accurate technique for estimating quantum states in spin-polarized systems. Its consistent performance across a spectrum of molecular configurations promises enhanced understanding and prediction of molecular properties, and it holds great potential for propelling the field of quantum computational chemistry forward.

Our investigation employed a VQE framework enhanced by the QNG optimizer to determine the ground state energies of hydrogen (*H₂*), lithium hydride (*LiH*), and beryllium hydride (*BeH₂*) molecules. The results underscore the QNG optimizer’s efficacy in navigating the optimization landscape, accelerating convergence to the ground state energy compared to traditional gradient descent methods.

For the *H₂* molecule, initial optimization efforts using traditional gradient descent exhibited gradual energy reductions, achieving a final convergence parameter of 0.00000097 Ha after 120 iterations. Notably, the introduction of QNG optimization markedly improved the efficiency, reaching a similar convergence threshold of 0.00000022 Ha in merely 16 iterations, as summarized below:


- Initial Energy: -0.09424484 Ha.
- Final GD Energy: -1.13615247 Ha after 120 iterations.
- Final QNG Energy: -1.13618926 Ha after 16 iterations.

The optimization for the *LiH* molecule demonstrated the QNG optimizer’s ability to reduce the energy across iterations consistently. The final ground state energy attained via QNG optimization was -7.60898098 Ha, showcasing a significant improvement over traditional methods:


- Initial Energy: -3.80567655 Ha.
- Final GD Energy: -7.60440889 Ha after 190 iterations.
- Final QNG Energy: -7.60898098 Ha after 190 iterations.

For *BeH₂*, both optimization approaches converged to a similar energy value, indicating the potential limitations of traditional optimization methods for more complex molecular systems. The final energies underscore the precision of QNG in fine-tuning the variational parameters:

- Initial Energy: -4.62101342 Ha.
- Final GD Energy: -13.08284703 Ha after 140 iterations.




(a)




(b)

Fig. 1: Energy estimations across various atomic distances using SS-VQE of LiH at: (a) $S_z = 0$ and (b) $S_z = 1$.




(a)




(b)

Fig. 2: Relative error analysis using SS-VQE of LiH at: (a) $S_z = 0$ and (b) $S_z = 1$.




(a)




(b)


Fig. 3: Energy estimations across various atomic distances using VQD for LiH at: (a) $S_z = 0$ and (b) $S_z = 1$.




(a)




(b)

Fig. 4: Relative error analysis using VQD of LiH at: (a) $S_z = 0$ and (b) $S_z = 1$ 

(a)



(b)



(c)

Fig. 5: Energy convergence for different molecules using QG and QNG optimizers for: (a) H_2 , (b) LiH , and (c) BeH_2 .

- Final QNG Energy: -13.08286007 Ha after 10 iterations.

As presented in Fig. 5, these results clearly illustrate the advantage of employing the QNG in the VQE algorithm, particularly in terms of efficiency and convergence speed. The marked improvement in optimization performance across different molecular systems underscores the potential of QNG to enhance quantum computational chemistry applications. Fig. 6 shows energy estimations across various atomic distances using QNG and adaptive VQE for LiH [5]. This approach includes a recipe to adaptively select gates that have a significant contribution to the desired state. We select only gates that have a predefined threshold above 10^{-5} . The results closely match the expected from exact diagonalization exemplifying QNG quantum chemistry applicability.

IV. EXPLORING ALGORITHM SPEED IMPROVEMENT

In addition to using QNG to reduce the number of iterations for convergence, we apply Pauli tapering, leveraging Z_2 molecular hamiltonian symmetries to reduce the number of qubits required for the VQE simulation. Additionally we test the use of the MERA tensor network ansatz [6] and find that it greatly improves the speed at which a SSVQE custom

algorithm finds the first excited state at the expense of accuracy at certain distance values as shown in Fig. 7.

V. CONCLUSION

Our investigation into advanced variational quantum algorithms, notably SS-VQE, VQD, Folded Spectrum VQE, and VQE with QNG, illuminates the path forward in quantum computational chemistry. These methodologies, each with its unique approach to navigating the challenges of quantum optimization, collectively underscore a significant leap toward harnessing quantum computing's potential for chemical and material science. SS-VQE and VQD have demonstrated robust capabilities in accurately determining the spectral gap of chemical molecules, a critical parameter influencing material properties. With precision that approaches near-exactness, particularly noted in the spectral gap analysis of LiH with less than 1% relative error, these strategies highlight the precision achievable with quantum computational approaches.

Folded Spectrum VQE further extends the versatility of quantum algorithms to identify highly excited states, offering a broader understanding of molecular dynamics and energetics. By facilitating a comprehensive view of the energy spectrum,




Fig. 6: Energy estimations across various atomic distances using QNG and adaptive VQE compared to exact results for LiH .




Fig. 7: First excited state energy estimations across various atomic distances. For the orange dots, we used a strongly entangling layers ansatz and an SSVQE algorithm that finds all k states in one optimization process. For the blue dots, we used a MERA [6] tensor network ansatz and a custom SSVQE algorithm that finds specific excited states and in this case the first excited state improving speed at the cost of accuracy. Additionally, we apply Pauli tapering to both algorithms. The results are compared to exact first excited state results.

this approach enriches our ability to predict and manipulate chemical behaviors at a quantum level.

Most notably, QNG stands out for its efficiency, significantly reducing the number of iterations required for convergence. By intelligently navigating the optimization landscape, QNG enhances computational efficiency and sets a new standard for precision in quantum computing optimizations. Looking ahead, the convergence of these advanced quantum computational methods opens new horizons for exploring chemical molecules in unprecedented detail. This study's remarkable accuracy and efficiency beckon a future where quantum computing plays a central role in unraveling complex chemical

mysteries, potentially revolutionizing our approach to material synthesis, drug discovery, and beyond. As we refine these algorithms and adapt them to more complex systems, the promise of quantum computational chemistry to contribute meaningful insights into the natural world becomes increasingly tangible.

REFERENCES

- [1] K. M. Nakanishi, K. Mitarai, and K. Fujii, "Subspace-search variational quantum eigensolver for excited states," *Phys. Rev. Res.*, vol. 1, p. 033062, Oct 2019. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevResearch.1.033062>
- [2] O. Higgott, D. Wang, and S. Brierley, "Variational Quantum Computation of Excited States," *Quantum*, vol. 3, p. 156, Jul. 2019. [Online]. Available: <https://doi.org/10.22331/q-2019-07-01-156>
- [3] L. C. Tazi and A. J. W. Thom, "Folded spectrum vqe : A quantum computing method for the calculation of molecular excited states," 2023.
- [4] J. Stokes, J. Izaac, N. Killoran, and G. Carleo, "Quantum natural gradient," *Quantum*, vol. 4, p. 269, May 2020.
- [5] H. R. Grimsley, S. E. Economou, E. Barnes, and N. J. Mayhall, "An adaptive variational algorithm for exact molecular simulations on a quantum computer," *Nature Communications*, vol. 10, July 2019.
- [6] G. Vidal, "Class of quantum many-body states that can be efficiently simulated," *Phys. Rev. Lett.*, vol. 101, p. 110501, Sep 2008. [Online]. Available: <https://link.aps.org/doi/10.1103/PhysRevLett.101.110501>