

# Distribution and spread of pyrethroid and DDT resistance among the *Anopheles gambiae* complex in Tanzania

B. KABULA<sup>1,2</sup>, P. TUNGU<sup>1</sup>, R. MALIMA<sup>1</sup>, M. ROWLAND<sup>3</sup>, J. MINJA<sup>4</sup>, R. WILILO<sup>5</sup>, M. RAMSAN<sup>5</sup>, P. D. MCELROY<sup>6</sup>, J. KAFUKO<sup>6</sup>, M. KULKARNI<sup>2</sup>, N. PROTOPOPOFF<sup>3</sup>, S. MAGESA<sup>1,7</sup>, F. MOSHA<sup>2</sup> and W. KISINZA<sup>1</sup>

<sup>1</sup>Amani Research Centre, National Institute for Medical Research, Ubwari, Muheza, Tanzania, <sup>2</sup>Department of Parasitology and Entomology, Kilimanjaro Christian Medical University College, Tumaini University, Moshi, Tanzania, <sup>3</sup>Department of Disease Control, London School of Hygiene and Tropical Medicine, London, U.K., <sup>4</sup>National Malaria Control Programme, Ministry of Health and Social Welfare, Dar es Salaam, Tanzania, <sup>5</sup>Global Health Division, RTI International, Dar es Salaam, Tanzania, <sup>6</sup>U.S. President's Malaria Initiative, Dar es Salaam, Tanzania and <sup>7</sup>Global Health Division, RTI International, Nairobi, Kenya

**Abstract.** The development of insecticide resistance is a threat to the control of malaria in Africa. We report the findings of a national survey carried out in Tanzania in 2011 to monitor the susceptibility of malaria vectors to pyrethroid, organophosphate, carbamate and DDT insecticides, and compare these findings with those identified in 2004 and 2010. Standard World Health Organization (WHO) methods were used to detect knock-down and mortality rates in wild female *Anopheles gambiae* s.l. (Diptera: Culicidae) collected from 14 sentinel districts. Diagnostic doses of the pyrethroids deltamethrin, lambda-cyhalothrin and permethrin, the carbamate propoxur, the organophosphate fenitrothion and the organochlorine DDT were used. *Anopheles gambiae* s.l. was resistant to permethrin in Muleba, where a mortality rate of 11% [95% confidence interval (CI) 6–19%] was recorded, Muheza (mortality rate of 75%, 95% CI 66–83%), Moshi and Arumeru (mortality rates of 74% in both). Similarly, resistance was reported to lambda-cyhalothrin in Muleba, Muheza, Moshi and Arumeru (mortality rates of 31–82%), and to deltamethrin in Muleba, Moshi and Muheza (mortality rates of 28–75%). Resistance to DDT was reported in Muleba. No resistance to the carbamate propoxur or the organophosphate fenitrothion was observed. *Anopheles gambiae* s.l. is becoming resistant to pyrethroids and DDT in several parts of Tanzania. This has coincided with the scaling up of vector control measures. Resistance may impair the effectiveness of these interventions and therefore demands close monitoring and the adoption of a resistance management strategy.

**Key words.** *Anopheles gambiae*, DDT resistance, insecticides, malaria vectors, pyrethroid resistance, Tanzania.

## Introduction

The application of insecticide-based malaria control measures in Africa is expanding with rapid increases in both the

distribution of insecticide-treated nets (ITNs) and indoor residual spraying (IRS) in many endemic countries. The longlasting insecticidal net (LLIN), the modern form of the ITN, and IRS are the cornerstones of malaria control

Correspondence: Bilali Kabula, National Institute for Medical Research, Amani Research Centre, Ubwari, P.O. Box 81, Muheza, Tanzania. Tel.: + 255 7830 21213; Fax: + 255 2726 41320; E-mail: bika72@yahoo.com

programmes (Roberts & Enserink, 2007). Vector control interventions are dependent on a limited number of insecticides from four chemical classes, namely, the organochlorines, organophosphates, carbamates and pyrethroids (World Health Organization, 2012a). The World Health Organization (WHO) has approved 12 insecticides belonging to these classes for use in IRS. The chemical arsenal for use with ITNs and LLINs is more limited and only six insecticides, all of which are pyrethroids, are available (WHO, 2012a). The pyrethroids are the only class approved for use on ITNs as a result of their low human toxicity, excito-repellent properties, rapid rate of knock-down and killing effects (Zaim *et al.*, 2000).

These four insecticide classes are also widely used for the control of agricultural pests in Africa (Santolamazza *et al.*, 2008). This can pose additional selection pressure on malaria vectors when insecticide-contaminated ground water permeates mosquito larval habitats (Ranson *et al.*, 2009). Exposure of mosquitoes to insecticides from multiple sources may combine to accelerate resistance and compromise malaria control efforts (Ranson *et al.*, 2011). Resistance to the organochlorines DDT (dichlorodiphenyltrichloroethane) and the now obsolete dieldrin was first reported in African malaria vectors in the 1950s and 1960s (Brown, 1958; Hamon *et al.*, 1968). Pyrethroid resistance was first detected in West African malaria vectors in 1993 (Elissa *et al.*, 1993), since when there has been an increasing number of reports of pyrethroid-resistance in *Anopheles gambiae* s.l. in countries in west, central, east and southern Africa (Vulule *et al.*, 1996; Chandre *et al.*, 1999; Stump *et al.*, 2004; Roberts & Enserink, 2007; Munhenga *et al.*, 2008; Protopopoff *et al.*, 2008; Ndjemai *et al.*, 2009; Hunt *et al.*, 2010; Chanda *et al.*, 2011) and in *Anopheles funestus* in Ghana, Mozambique, Malawi, Zambia and South Africa (Hargreaves *et al.*, 2000; Okoye *et al.*, 2008; Chanda *et al.*, 2011). Carbamate- and organophosphate-resistant populations of *An. gambiae* have also been reported in West Africa (Corbel *et al.*, 2007). The continuing success of IRS and LLIN interventions against the rising tide of insecticide resistance depends on the acquiring of comprehensive knowledge of the factors underlying selection in order to support the development of effective resistance management strategies. This, in turn, raises the need for regular surveys to monitor the insecticide susceptibility of vector species and to assess their impact on malaria control activities (Kelly-Hope *et al.*, 2008).

The use of ITNs in Tanzania started in the early 1990s as projects in the Muheza and Korogwe districts in Tanga, the Kilombero and Ulanga districts in Morogoro, and the Bagamoyo district in the coastal region (Schellenberg *et al.*, 1999, 2001; Abdulla *et al.*, 2001; Maxwell *et al.*, 2002; Erlanger *et al.*, 2004; Magesa *et al.*, 2005). These projects were then expanded to cover larger areas in 1998 through the Social Marketing of Insecticide-Treated Nets (SMITN) project and from 2002 onwards through the Strategic Social Marketing for Expanding the Commercial Market of ITNs (SMARNET) programme (Magesa *et al.*, 2005). This was followed by the rolling out of vouchers for ITNs, which were offered to pregnant women through the Tanzania National Voucher Scheme (TNVS) in 2004 (Magesa *et al.*, 2005; Hanson *et al.*, 2009). To accelerate coverage and address an

equity gap, Tanzania adopted the free distribution of LLINs in 2008. The ITNs were distributed through an 'Under-5 Catch-Up' campaign, which was followed by various 'universal coverage' campaigns (WHO, 2012b). All the LLINs distributed recently are Olyset™ nets (Sumitomo Chemical Co., Ltd, Tokyo, Japan), impregnated with permethrin. Indoor residual spraying was reintroduced in the country in 2007 as a way to control malaria outbreaks in selected areas of two districts (Muleba and Karagwe) in northwest Tanzania (WHO, 2012b). By March 2011, IRS operations with lambda-cyhalothrin had expanded to 18 districts in the Lake Victoria Zone and reached approximately 94% of the targeted structures in those districts (WHO, 2012b). Through these campaigns, national coverage with ITNs and LLINs rose dramatically from 38% in 2007–2008 to 91% in 2011–2012 and that of IRS reached 13% in 2011–2012 (National Bureau of Statistics, 2012). Therefore, it is imperative to regularly monitor the susceptibility status of local malaria vectors to the insecticides used in these interventions. This will enable the timely deployment of resistance mitigation tactics when resistance is recognized.

Earlier large-scale surveys in Tanzania demonstrated the efficacy of all four classes against malaria vectors (Kulkarni *et al.*, 2007) with focal points of phenotypic resistance (Kabula *et al.*, 2012). This survey was carried out as part of continued monitoring of the susceptibility status of local malaria vectors to insecticides in use in the country. The present study reports the findings of a national survey carried out in 2011 at 14 sentinel sites to monitor the susceptibility of malaria vectors to pyrethroid, organophosphate, carbamate and DDT insecticides and to compare current trends with the situations existing in 2004 and 2010. Some of the 2004 comparison data were novel and some were published (Kulkarni *et al.*, 2007). All comparison data for 2010 have been previously reported elsewhere (Kabula *et al.*, 2012).

## Materials and methods

### Study sites

The study was carried out in 14 sentinel districts selected from across the country. The selection of sentinel districts was based on WHO-recommended selection criteria, which include: a history of insecticide use by communities living in the areas (in agricultural and public health contexts); malaria endemicity (moderate to high transmission); high coverage by ITNs and LLINs; demographic setting (urban/rural), and easy site accessibility. The characteristics of these districts have been previously described (Kabula *et al.*, 2012). Additional characteristics and the distribution of the study districts are shown in Table 1 and Fig. 1, respectively. These sentinel districts and insecticide resistance monitoring sites were initially established in 1999 with funding from the Ministry of Health and Social Welfare (MoHSW). This was followed by further monitoring and the detection of insecticide resistance in the context of the scaling up of ITN distribution in 2004, at which point the survey was carried out in 11 sentinel sites. In response to the concern of the global malaria community over the threat implied by insecticide resistance, the country

**Table 1.** Geographical coordinates and characteristics of the sentinel districts.

Site	Coordinates	ITN coverage trend (at least one ITN)			Malaria prevalence‡	Other interventions using insecticides	Demographic settings
		2004*	2009†	2011‡			
Arumeru	03°08' S, 36°52' E	23.7%	32.4%	84.5%	0.1%	Floriculture, agriculture, livestock	Semi-urban
Babati	04°13' S, 35°45' E	8.1%	22.1%	88.6%	0.1%	Agriculture, livestock	Rural
Dodoma	07°00' S, 36°30' E	13.4%	28.2%	92.8%	2.6%	Agriculture, livestock	Rural
Handeni	05°43' S, 38°01' E	17.3%	38.6%	90.5%	5.5%	Agriculture, livestock	Rural
Ilala	06°49' S, 39°14' E	60.8%	70.7%	78.6%	3.6%	Horticulture	Urban
Kilombero	08°31' S, 37°22' E	32.6%	44.1%	91.2%	13.0%	Agriculture	Rural
Kyela	09°35' S, 33°55' E	13.6%	29.9%	91.4%	0.5%	Agriculture	Rural
Lushoto	04°78' S, 38°28' E	17.3%	38.6%	90.5%	5.5%	Horticulture	Rural
Magu	02°30' S, 33°30' E	28.1%	48.4%	96.1%	19.1%	IRS since 2011, agriculture, livestock	Rural
Moshi	03°21' S, 37°20' E	13.1%	29.9%	94.8%	0.1%	Agriculture	Rural
Muheza	05°10' S, 38°46' E	17.3%	38.6%	90.5%	5.5%	Agriculture	Rural
Muleba	01°45' S, 31°40' E	13.9%	29.5%	91.9%	8.5%	IRS since 2006, agriculture	Rural
Mvomero	06°30' S, 37°30' E	32.6%	44.1%	91.2%	13.0%	Agriculture, livestock	Rural
Tabora	04°55' S, 32°49' E	18.4%	39.8%	94.5%	9.2%	Agriculture, livestock	Rural

\*Tanzania Demographic and Health Survey 2004–2005.

†Tanzania HIV/AIDS and Malaria Indicator Survey 2007–2008.

‡Tanzania HIV/AIDS and Malaria Indicator Survey 2011–2012.

IRS, indoor residual spraying.

continued with the periodic monitoring of resistance. In 2008, funding from the Bill & Melinda Gates Foundation through the WHO enabled the number of surveillance sites to be expanded to the current 14.

### *Mosquito collections and identification*

At each study site, indoor-resting *Anopheles* mosquitoes were collected from human habitations, using torches, mechanical aspirators and paper cups, between 06.00 hours and 09.00 hours (WHO, 1975). The collections were made in the rainy season during May and June 2011 in all but one of the sites; this is the optimal time to capture adequate numbers of mosquitoes for susceptibility testing as per guidelines issued by the WHO (1998). An exception was the Muleba district, where collections were made during April to June and in November and December 2011 in 11 villages described by Protopopoff *et al.* (2013). The collected mosquitoes were maintained on a fresh 10% glucose solution before being tested for insecticide susceptibility. In Ilala and Dodoma districts adult collections were insufficient. Therefore larvae were collected using dippers and transferred in cool boxes to the laboratory, where they were reared on ground Tetramin<sup>®</sup> fish food at 27–30 °C. The geographical coordinates of each sampling site were determined by GPS (Trimble Geoexplorer II; Trimble Navigation Ltd, Sunnyvale, CA, U.S.A.). Adult *Anopheles* mosquitoes were carefully sorted for susceptibility tests, as recommended by the WHO (1998), and morphologically identified using a standard morphological key according to Gillies and Coetsee (1987). Technical and logistical constraints prevent the inclusion of the results of the molecular identification of *An. gambiae* complex specimens in this analysis.

### *Insecticide susceptibility/resistance bioassays*

Insecticide susceptibility/resistance bioassays were carried out using WHO insecticide susceptibility test kits (WHO, 1975) according to standard procedures (WHO, 1998) with four or five replicates of 15–25 wild-caught adult female mosquitoes per test. For convenience purposes this study used the unfed wild-caught adult female anopheles mosquitoes to test their susceptibility to insecticides in most of the sites. Therefore the age of the mosquitoes tested might be less certain. Mosquitoes were exposed to papers impregnated with the WHO-recommended discriminating concentrations (w/v) of 0.05% deltamethrin, 0.05% lambda-cyhalothrin, 0.75% permethrin, 4% DDT, 0.1% propoxur and 1% fenitrothion; the papers were prepared at University Sains Malaysia (WHO, 1998). The controls were exposed to control papers impregnated with silicone oil (pyrethroid control), risella oil (DDT control) or olive oil (organophosphate/carbamate control). During exposure, time to knock-down (KDT) rates were recorded after 10, 15, 20, 30, 40, 50 and 60 min for pyrethroid and organochlorine insecticides. Mosquitoes were then transferred to holding tubes and supplied with glucose solution. In instances of knock-down of less than 80% during the 60-min exposure period, knock-down was monitored for a further 20 min in the holding tube. Mortality was scored after a 24-h holding period (WHO, 1998).

### Statistical analyses

Mortality percentages and 95% confidence intervals (CIs) in WHO susceptibility tests were calculated using the binomial exact method in spss for Windows Version 16.0 (SPSS, Inc.,




Fig. 1. Map of Tanzania showing the distribution of insecticide resistance surveillance districts.

Chicago, IL, U.S.A.) and VassarStats, a web-based statistical package (<http://www.vassarstats.net/>). Abbott's formula was used to correct observed mortality when mortality in the control samples was between 5% and 20% (Abbott, 1925). The  $KDT_{50}$  and  $KDT_{95}$  (time required to knock down 50% and 95%, respectively, of the test mosquitoes) were estimated by probit analysis (Finney, 1971) using spss Version 16.0. Resistance ratios (RRs) at  $KDT_{50}$  were estimated by comparing the  $KDT_{50}$  of field-collected mosquitoes with that of the *An. gambiae* Kisumu susceptible strain.

## Results

### Mortality

The highest percentages of mosquitoes resistant to permethrin were observed in samples collected from Arumeru (26%), Moshi (26%) and Muheza (25%), followed by those from Kilombero (15%) and Ilala (10%). Lower levels of survival and hence resistance to permethrin (1–5%) were found in populations from Babati, Magu and Handeni districts. Complete susceptibility (mortality of 100%) following exposure to

permethrin was observed in populations collected from the five districts of Lushoto, Dodoma, Tabora, Mvomero and Kyela (Table 2).

Following exposure to WHO-recommended diagnostic doses of deltamethrin, highest survival was observed in populations from Moshi (34%), Muheza (26%), Arumeru (10%) and Handeni (7%), but low survival (<4%) was observed in mosquitoes collected from Ilala, Kilombero and Babati. None of the mosquitoes collected from Magu, Lushoto, Dodoma, Tabora, Mvomero and Kyela districts survived.

Following exposure to lambdacyhalothrin, highest survival rate was recorded in mosquitoes collected from Moshi (42%), Arumeru (30%), Muheza (18%) and Handeni (2%) (Table 2).

Conversely, following exposure to DDT, no mosquitoes from any of the sentinel districts survived, except those from Ilala and Magu, where survival rates were 12% and 5%, respectively (Table 3). Samples of mosquitoes from the 14 sentinel sites exposed to either fenitrothion or propoxur exhibited 99–100% mortality. Only 1% of tested mosquitoes from the Kilombero, Kyela and Mvomero districts survived exposure to propoxur. Similarly, 1% of mosquitoes survived exposure to fenitrothion in Kyela district (Table 4).



**Table 2.** Susceptibility status of *Anopheles gambiae* s.l. to diagnostic concentrations of permethrin, deltamethrin and lambdacyhalothrin in 2011 compared with 2004 and 2010. Mean percentage mortality in World Health Organization susceptibility tests and number of mosquitoes exposed.

Insecticide	Site	Mortality, % (mosquitoes exposed, n)		
		2004	2009/2010*	2011
Permethrin	Arumeru	91% (150)†	93% (150)	74% (125)‡
	Babati	—	98% (200)	99% (123)
	Dodoma	—	—	100% (80)
	Handeni	—	91% (80)	95% (100)
	Ilala	—	92% (64)	90.3% (75)
	Kilombero	99% (543)	96% (180)	85% (80)‡
	Kyela	99% (219)	100% (81)	100% (86)
	Lushoto	—	100% (100)	100% (100)
	Magu	100% (79)†	—	96% (80)
	Mvomero	100% (225)	91% (160)	100% (82)
	Moshi	97% (675)	77% (555)	74% (542)
	Muheza	100% (180)	100% (100)	75% (95)‡
	Muleba	—	100% (100)	11% (98)‡§
	Tabora	100% (160)	100% (75)	100% (80)
Deltamethrin	Arumeru	100% (150)†	96.7% (150)	90.4% (125)
	Babati	100% (245)	98% (200)	96% (125)
	Dodoma	—	—	100% (80)
	Handeni	—	—	93% (99)
	Ilala	—	88% (60)	97% (85)
	Kilombero	100% (100)†	90% (180)	96% (80)
	Kyela	99% (198)	100% (80)	100% (78)
	Lushoto	—	100% (100)	100% (100)
	Magu	100% (79)†	—	100% (40)
	Mvomero	100% (66)†	94% (180)	100% (80)
	Moshi	100% (453)	94% (482)	66% (533)†
	Muheza	100% (100)	100% (80)	75% (95)†
	Muleba	—	100% (100)	28% (106)*‡
Lambdacyhalothrin	Tabora	100% (80)	100% (80)	100% (80)
	Arumeru	—	87% (150)	70% (125)†
	Babati	—	98% (200)	100% (125)
	Dodoma	—	—	100% (80)
	Handeni	—	84% (80)	98% (92)
	Ilala	—	94% (94)	95% (79)
	Kilombero	—	94% (180)	100% (80)
	Kyela	—	90% (88)	100% (83)
	Lushoto	—	100% (100)	100% (100)
	Magu	—	—	100% (80)
	Mvomero	—	88% (160)	100% (83)
	Moshi	—	73% (515)	58% (531)†
	Muheza	—	95% (80)	82% (95)†
	Muleba	—	100% (100)	31% (1099)*‡
	Tabora	—	100% (76)	100% (80)

\*Kabula *et al.* (2012).†Kulkarni *et al.* (2007).

‡Significant reduction in mortality rate at 95% confidence interval.

§Protopopoff *et al.* (2013).*Mortality trends in An. gambiae* s.l. exposed to insecticides

The 2004 surveys found no clear evidence of pyrethroid resistance in any part of the country; the only exceptions were Arumeru and Moshi, where mosquitoes showed some resistance to permethrin (mortality rates of 91% and 97%,

**Table 3.** Susceptibility status of *Anopheles gambiae* s.l. to diagnostic concentrations of DDT in 2011 compared with 2004 and 2010. Mean percentage mortality in World Health Organization susceptibility tests and number of mosquitoes exposed.

Insecticide	Site	Mortality, % (mosquitoes exposed, n)		
		2004	2009/2010*	2011
DDT	Arumeru	100% (150)†	100% (150)	100% (125)
	Babati	99% (299)	99% (200)	100% (100)
	Dodoma	—	—	100% (80)
	Handeni	—	—	100% (100)
	Ilala	—	65% (88)	88% (520)
	Kilombero	93% (702)	100% (100)	100% (125)
	Kyela	100% (281)	90% (84)	100% (85)
	Lushoto	—	92% (92)	100% (100)
	Magu	98% (79)†	—	95% (80)
	Mvomero	100% (128)	100% (160)	100% (83)
	Moshi	100% (518)	99% (491)	99% (648)
	Muheza	100% (100)	99% (100)	100% (100)
	Muleba	—	100% (100)	67% (234)‡§
	Tabora	95% (20)†	100% (79)	100% (80)

\*Kabula *et al.* (2012).†Kulkarni *et al.* (2007).‡Protopopoff *et al.* (2013).

§Significant reduction in mortality rate at 95% confidence interval.

**Table 4.** Response of wild-caught *Anopheles gambiae* s.l. local populations to discriminatory dosages of 0.1% propoxur and 1% fenitrothion in treated papers in 14 sentinel districts of Tanzania in 2011. Mean percentage mortality in World Health Organization susceptibility tests and number of mosquitoes exposed.

Site	Mortality, % (mosquitoes exposed, n)	
	Propoxur	Fenitrothion
Arumeru	100% (125)	100% (125)
Babati	100% (125)	100% (100)
Dodoma	100% (80)	100% (100)
Handeni	100% (100)	100% (100)
Ilala	100% (68)	100% (60)
Kilombero	99% (80)	100% (123)
Kyela	99% (79)	99% (80)
Lushoto	100% (100)	100% (100)
Magu	100% (90)	100% (80)
Moshi	99% (338)	99% (249)
Muheza	100% (100)	100% (100)
Muleba	100% (80)	100% (60)
Mvomero	99% (80)	100% (81)
Tabora	100% (80)	100% (100)

respectively). By 2010, however, there was growing evidence of resistance in selected localities. In Moshi, for example, permethrin-induced mortality had decreased to 77% and there was some indication of permethrin resistance in Kilombero, Arumeru, Ilala and Handeni, where mortality rates ranged from 91% to 96%. This trend continued in 2011: mortality achieved by permethrin in Moshi was 74% and there was growing evidence of resistance in Arumeru (mortality: 74%), Ilala (mortality: 90%), Kilombero (mortality: 85%),



**Table 5.** Knock-down times in wild-caught *Anopheles gambiae* s.l. local populations exposed to discriminatory dosages of 0.05% deltamethrin, 0.75% permethrin, 0.05% lambdacyhalothrin and 4% DDT in 2011.

Site (sentinel district)	Permethrin		Deltamethrin		Lambdacyhalothrin		DDT	
	KDT <sub>50</sub> , min, mean (95% CI)	KDT <sub>50</sub> ratio	KDT <sub>50</sub> , min, mean (95% CI)	KDT <sub>50</sub> ratio	KDT <sub>50</sub> , min, mean (95% CI)	KDT <sub>50</sub> ratio	KDT <sub>50</sub> , min, mean (95% CI)	KDT <sub>50</sub> ratio
Kisumu strain*	12.5	–	13.8	–	15.2	–	21.5 (18.7–24.4)	–
Arumeru	84.4 (67.2–151.0)	6.8	92.2 (84.0–118.5)	6.7	114 (91.7–171.7)	7.5	29.0 (27.9–30.1)	1.3
Babati	15.5 (13.4–17.7)	1.2	34.4 (31.6–37.3)	2.5	118.0 (96.0–179.0)	7.8	38.1 (36.1–40.2)	1.8
Dodoma	17.3 (16.3–18.3)	1.4	22.2 (19.6–24.9)	1.6	21.8 (19.4–24.2)	1.4	26.4 (24.1–28.8)	1.2
Handeni	33.4 (27.3–42.2)	2.7	26.7 (25.1–28.5)	1.9	33.6 (31.8–35.4)	2.2	22.1 (12.9–31.2)	1.0
Ilala	37.9 (36.1–39.7)	3.0	32.5 (28.6–36.5)	2.4	41.5 (37.7–45.9)	2.7	31.2 (28.6–33.8)	1.5
Kilombero	31.3 (28.8–33.9)	2.5	22.3 (19.3–25.3)	1.6	31.5 (28.7–34.4)	2.1	19.2 (15.4–23.3)	0.9
Kyela	15.7 (14.5–16.8)	1.3	13.3 (11.6–14.9)	1.0	11.9 (10.9–12.9)	0.8	31.1 (28.1–34.4)	2.0
Lushoto	15.8 (13.2–18.2)	1.3	23.8 (21.4–26.4)	1.7	28.9 (22.7–35.1)	1.9	36.1 (34.7–37.4)	1.7
Magu	23.8 (19.2–27.9)	1.9	34.2 (29.3–39.4)	2.5	41.7 (35.8–49.6)	2.7	33.3 (24.7–40.2)	1.5
Moshi	47.8 (46.0–49.8)	3.8	47.6 (46.3–48.9)	3.4	59.3 (50.2–82.0)	3.9	36.2 (34.3–38.1)	1.7
Muheza	28.3 (25.8–30.8)	2.3	25.2 (23.7–26.9)	1.8	29.6 (25.9–33.5)	1.9	32.7 (29.9–35.4)	1.5
Mvomero	17.9 (16.4–19.5)	1.4	15.3 (12.9–17.4)	1.1	20.6 (18.5–22.8)	1.4	27.3 (22.7–32.7)	1.3
Tabora	23.5 (20.0–27.1)	1.9	20.9 (16.6–25.4)	1.5	26.7 (24.2–29.4)	1.8	30.5 (24.1–38.8)	1.4

\*Kisumu strain, *Anopheles gambiae* s.s. susceptible Kisumu strain was used as susceptible control.

KDT<sub>50</sub>, time required for 50% of test mosquitoes to be knocked down; KDT<sub>50</sub> ratio, KDT<sub>50</sub> of the tested population/KDT<sub>50</sub> of the susceptible Kisumu strain.

95% CI, 95% confidence interval.

has been implicated in the development of insecticide resistance in malaria vectors (Diabate *et al.*, 2002). Overall, there has been a rapid decrease in susceptibility status across Tanzania compared with that identified in the last nationwide insecticide resistance surveys (Kulkarni *et al.*, 2007; Kabula *et al.*, 2012). This may indicate the cumulative effects of increases in the use of ITNs (such as in the Under-5 Catch-Up and universal coverage campaigns), which have been ongoing since the 1990s and have shown relative increases since 2010. Similar observations have been made in Niger, Equatorial Guinea (Bioko) and Kenya (Stump *et al.*, 2004; Sharp *et al.*, 2007; Czeher *et al.*, 2008). Recent studies in Senegal and Liberia have also demonstrated increased frequencies of pyrethroid resistance after high LLIN usage (Ndiath *et al.*, 2012; Temu *et al.*, 2013). The rapid rise of pyrethroid resistance in Muleba also coincides with the intensive use of IRS in the district (since 2007) for the control of malaria transmission and was recorded a few years earlier in the neighbouring country of Uganda (Protopopoff *et al.*, 2013). Indoor residual spraying is commonly associated with the selection of pyrethroid resistance (Sharp *et al.*, 2007; WHO, 2010).

Resistance as measured in WHO susceptibility tests does not imply that ITNs/LLINs are no longer protective against local vector populations. The operational significance of resistance can be demonstrated in experimental hut trials against wild host-seeking resistant mosquitoes and through careful surveillance for increasing malaria transmission. The present level of resistance in Lower Moshi appears not to impair the effectiveness of permethrin-treated nets. Field trials in Moshi show that although such treated nets kill relatively few host-seeking *An. arabiensis* that enter local houses, the nets continue to provide personal protection through the strong excito-repellent activity of permethrin (Mosha *et al.*, 2008).

However, high pyrethroid resistance in *An. gambiae* can impair the effectiveness of ITNs/LLINs and IRS as reported in Benin (N'Guessan *et al.*, 2007; Asidi *et al.*, 2012). The operational failure of IRS with deltamethrin was noted previously in South Africa, where the malaria vector *An. funestus* was found to have developed resistance to this insecticide (Hargreaves *et al.*, 2000).

In the present context, it is of paramount importance that levels of surveillance and the monitoring of resistance are increased across the country and that the operational implications of resistance on the effectiveness of pyrethroid-based LLIN and IRS interventions are assessed. This can partially be ensured by strengthening systems for the surveillance of human malaria cases in the areas in which ongoing insecticide resistance is monitored. Without the capacity to overlay insecticide resistance data and human case surveillance data, the full implications of reductions in vector susceptibility will remain uncertain.

## Conclusions

Resistance to pyrethroids and DDT is emerging across Tanzania at unprecedented rates that have not been seen previously. This has coincided with the scaling up of malaria vector control interventions and may have serious implications for the future of the national malaria control programme, which largely depends on the use of pyrethroids in IRS and LLINs/ITNs. A resistance management strategy that develops the rotational use of insecticides, especially in areas in which IRS is used, is recommended. For example, mosquito populations remain susceptible to the organophosphate and carbamate classes, which indicates that these can be used to good effect in IRS as part of a resistance management strategy.

- © 2013 The Royal Entomological Society, *Medical and Veterinary Entomology*, **28**, 244–252



- permethrin resistance in *Anopheles arabiensis* from Lower Moshi, north-eastern Tanzania. *Malaria Journal*, **9**, 193.
- Maxwell, C.A., Msuya, E., Sudi, M., Njunwa, K.J., Carneiro, I.A. & Curtis, C.F. (2002) Effect of community-wide use of insecticide-treated nets for 3–4 years on malarial morbidity in Tanzania. *Tropical Medicine and International Health*, **7**, 1003–1008.
- Mosha, F.W., Lyimo, I.N., Oxborough, R.M. *et al.* (2008) Comparative efficacies of permethrin-, deltamethrin- and alphacypermethrin-treated nets, against *Anopheles arabiensis* and *Culex quinquefasciatus* in northern Tanzania. *Annals of Tropical Medicine and Parasitology*, **102**, 367–376.
- Munhenga, G., Masendu, H.T., Brooke, B.D., Hunt, R.H. & Koekemoer, L.K. (2008) Pyrethroid resistance in the major malaria vector *Anopheles arabiensis* from Gwave, a malaria-endemic area in Zimbabwe. *Malaria Journal*, **7**, 247.
- National Bureau of Statistics (2012) *Malaria Atlas: The 2011–2012 Tanzania HIV/AIDS and Malaria Indicator Survey*. NBS, Dar es Salaam.
- Ndiath, M.O., Sougoufara, S., Gaye, A. *et al.* (2012) Resistance to DDT and pyrethroids and increased *kdr* mutation frequency in *An. gambiae* after the implementation of permethrin-treated nets in Senegal. *PLoS One*, **7**, e31943.
- Ndjemai, H.N., Patchoke, S., Atangana, J. *et al.* (2009) The distribution of insecticide resistance in *Anopheles gambiae* s.l. populations from Cameroon: an update. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **103**, 1127–1138.
- N'Guessan, R., Corbel, V., Akogbeto, M. & Rowland, M. (2007) Reduced efficacy of insecticide-treated nets and indoor residual spraying for malaria control in pyrethroid resistance area, Benin. *Emerging Infectious Diseases*, **13**, 199–206.
- Okoye, P.N., Brooke, B.D., Koekemoer, L.L., Hunt, R.H. & Coetzee, M. (2008) Characterization of DDT, pyrethroid and carbamate resistance in *Anopheles funestus* from Obuasi, Ghana. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **102**, 591–598.
- Protopopoff, N., Van Bortel, W., Marcotty, T. *et al.* (2008) Spatial targeted vector control is able to reduce malaria prevalence in the highlands of Burundi. *American Journal of Tropical Medicine and Hygiene*, **79**, 12–18.
- Protopopoff, N., Matowo, J., Malima, R. *et al.* (2013) High level of resistance in the mosquito *Anopheles gambiae* to pyrethroid insecticides and reduced susceptibility to bendiocarb in north-western Tanzania. *Malaria Journal*, **12**, 149.
- Ranson, H., Jensen, B., Vulule, J.M., Wang, X., Hemingway, J. & Collins, F.H. (2000) Identification of a point mutation in the voltage-gated sodium channel gene of Kenyan *Anopheles gambiae* associated with resistance to DDT and pyrethroids. *Insect Molecular Biology*, **9**, 491–497.
- Ranson, H., Abdallah, H., Badolo, A. *et al.* (2009) Insecticide resistance in *Anopheles gambiae*: data from the first year of a multi-country study highlight the extent of the problem. *Malaria Journal*, **8**, 299.
- Ranson, H., N'Guessan, R., Lines, J., Moiroux, N., Nkuni, Z. & Corbel, V. (2011) Pyrethroid resistance in African anopheline mosquitoes: what are the implications for malaria control? *Trends in Parasitology*, **27**, 91–98.
- Roberts, L. & Enserink, M. (2007) Malaria. Did they really say ... eradication? *Science*, **318**, 1544–1545.
- Santolamazza, F., Calzetta, M., Etang, J. *et al.* (2008) Distribution of knock-down resistance mutations in *Anopheles gambiae* molecular forms in west and west-central Africa. *Malaria Journal*, **7**, 74.
- Schellenberg, J.R., Abdulla, S., Minja, H. *et al.* (1999) KINET: a social marketing programme of treated nets and net treatment for malaria control in Tanzania, with evaluation of child health and longterm survival. *Transactions of the Royal Society of Tropical Medicine and Hygiene*, **93**, 225–231.
- Schellenberg, J.R., Abdulla, S., Nathan, R. *et al.* (2001) Effect of large-scale social marketing of insecticide-treated nets on child survival in rural Tanzania. *Lancet*, **357**, 1241–1247.
- Sharp, B.L., Ridl, F.C., Govender, D., Kuklinski, J. & Kleinschmidt, I. (2007) Malaria vector control by indoor residual insecticide spraying on the tropical island of Bioko, Equatorial Guinea. *Malaria Journal*, **6**, 52.
- Stump, A.D., Atieli, F.K., Vulule, J.M. & Besansky, N.J. (2004) Dynamics of the pyrethroid knock-down resistance allele in western Kenyan populations of *Anopheles gambiae* in response to insecticide-treated bed net trials. *American Journal of Tropical Medicine and Hygiene*, **70**, 591–596.
- Temu, E.A., Maxwell, C., Munyekenye, G. *et al.* (2013) Pyrethroid resistance in *Anopheles gambiae*, in Bomi County, Liberia, compromises malaria vector control. *PLoS One*, **7**, e44986.
- Vulule, J.M., Beach, R.F., Atieli, F.K., Mount, D.L., Roberts, J.M. & Mwambi, R.W. (1996) Longterm use of permethrin-impregnated nets does not increase *Anopheles gambiae* permethrin tolerance. *Medical and Veterinary Entomology*, **10**, 71–79.
- World Health Organization (1975) *Manual on Practical Entomology in Malaria. Part II. Methods and Techniques*. Geneva, Switzerland.
- World Health Organization (1998) *Test Procedures for Insecticide Resistance Monitoring in Malaria Vectors, Bio-efficacy and Persistence of Insecticides on Treated Surfaces*. Geneva, Switzerland.
- World Health Organization (2010) *The Technical Basis for Coordinated Action against Insecticide Resistance: Preserving the Effectiveness of Modern Malaria Vector Control*, Technical Report. Geneva, Switzerland.
- World Health Organization (2012a) *Global Plan for Insecticide Resistance Management in Malaria Vectors (GPIRM)*. Geneva, Switzerland.
- World Health Organization (2012b) *RBM Progress and Impact Series: Country Reports: Focus on Mainland Tanzania*. WHO, Geneva.
- World Health Organization (2013) *Test Procedures for Insecticide Resistance Monitoring in Malaria Vector Mosquitoes*. Geneva, Switzerland.
- Zaim, M., Aitio, A. & Nakashima, N. (2000) Safety of pyrethroid-treated mosquito nets. *Medical and Veterinary Entomology*, **14**, 1–5.

Accepted 19 September 2013

First published online 5 November 2013