

INFORMATIONAL DISPLACEMENT & RECOIL THEORY (IDRT)

Advanced Research in Universal Information Fields

I. Abstract

The Informational Displacement & Recoil Theory (IDRT) proposes a fundamental shift in understanding high-energy physics. It suggests that the universe is governed by an underlying Informational Fabric. Any artificial creation of matter in deep-space environments results in an "Informational Debt" that generates a corrective, non-stochastic return vector toward the terrestrial origin.

"Matter is not merely mass; it is a localized configuration of the Universal Information Field (UIF). Unnatural synthesis creates a tension that necessitates a physical recoil toward its source."

II. The Universal Information Field (UIF)

Unlike standard quantum entanglement, IDRT posits that all objects and events are mapped within a UIF. When synthetic matter is generated in isolation:

- **Informational Signature:** The matter retains a 'homing' frequency tied to the laboratory's specific coordinates.
- **Field Tension:** The vacuum of deep space acts as an elastic manifold. Artificial collisions 'stretch' this manifold, creating a pressure that drives the resulting products back to the equilibrium point (The Origin).

III. Mathematical Implications

The force of the recoil is determined by the **Az Constant (α_z)**, which scales the informational complexity of the synthetic matter against the spatial displacement from the UIF origin point.

IV. Experimental Safeguards

Given the potential for 'Informational Contagion', the theory mandates that such experiments be conducted in extragalactic zones, monitored exclusively by AI systems to analyze the Recoil Vector without risking the terrestrial biosphere.

Prepared by Principal Researcher:

Abdulaziz Ali

Date: 2026-05-13

FORMAL THEORETICAL PROPOSAL

MATHEMATICAL SUPPLEMENT

Informational Dynamics of Synthetic Matter

1. The Informational Potential Field

We propose that the "Homing" effect is driven by a potential field generated by informational gradients. Let Φ_I be the informational potential of the system.

Field Potential Equation:

$$\Phi_I(x, t) = - \int [S(\rho_t) / |x - x_0|] dV$$

In this equation, $S(\rho_t)$ represents the Von Neumann entropy of the synthetic matter's state, and $|x - x_0|$ is the spatial distance from the origin x_0 . As entropy increases during the collision, a "well" is created in the informational manifold that biases the particle's probability distribution toward the origin.

2. Non-Local Connectivity (The Homing Tensor)

To describe the directional return, we utilize a second-rank tensor $T_{\mu\nu}$ which characterizes the coupling between the synthetic matter and the laboratory's frame of reference.

Homing Tensor Calculation:

$$T_{\mu\nu} = \eta_{\mu\nu} + \alpha_z (\partial_\mu I \otimes \partial_\nu I)$$

Here, α_z is the **Az Constant**, a scaling factor representing the "strength" of the informational attraction. The term $\partial \mathbf{I}$ represents the partial derivative of the shared information, defining how the field changes over space and time.

3. Probability of Informational Recoil

The probability P_r that the matter (or its informational effect) will return to the origin is given by the following path integral:

Path Integral Probability:

$$P_r = \left| \int D[\varphi] \exp\left(i/\hbar S_{\text{eff}}[\varphi] \right) \right|^2$$

Where S_{eff} is the effective action that includes the informational coupling term. This ensures that the most "stable" path for the synthetic matter is the one leading back to its point of creation.

Principal Researcher: **Abdulaziz Ali**

Date: 2026-05-13