

# Lecture Note on Terwilliger Algebra

P. Terwilliger, edited by H. Suzuki

2022-12-21



# Contents

<b>About this lecturenote</b>	<b>5</b>
Setting . . . . .	5
Another Host . . . . .	6
<b>1 Subconstituent Algebra of a Graph</b>	<b>7</b>
<b>2 Perron-Frobenius Theorem</b>	<b>13</b>
<b>3 Cayley Graphs</b>	<b>19</b>
<b>4 Examples</b>	<b>25</b>
<b>5 <math>T</math>-Modules of <math>H(D, 2)</math>, I</b>	<b>31</b>
<b>6 <math>T</math>-Modules of <math>H(D, 2)</math>, II</b>	<b>37</b>
<b>7 The Johnson Graph <math>J(D, N)</math></b>	<b>45</b>
<b>8 Thin Graphs</b>	<b>51</b>
<b>9 Thin <math>T</math>-Module, I</b>	<b>57</b>
<b>10 Thin <math>T</math>-Module, II</b>	<b>63</b>
<b>11 Examples of <math>T</math>-Module</b>	<b>69</b>
<b>12 Distance-Regular</b>	<b>73</b>
<b>13 Modules of a DRG</b>	<b>81</b>
<b>14 Parameters of Thin Modules, I</b>	<b>87</b>
<b>15 Parameters of Thin Modules, II</b>	<b>91</b>
<b>16 Thin Modoles of a DRG</b>	<b>97</b>

17 Association Schemes	105
18 Polynomial Schemes	109
19 Commutative Association Schemes	111
20 Vanishing Conditions	117
21 Norton Algebras	123
22 $Q$ -Polynomial Schemes	127
23 Representation of a Scheme	133
24 Balanced Conditions, I	139
25 Balanced Conditions, II	145
26 Representation Diagrams	151
27 $P$ -and $Q$ -Polynomial Schemes	157
28 The First Eigenspace of a $Q$ -DRG	165
29 Tridiagonal Pair $A, A^*$	169
30 $R, F, L$ Matrices	177
31 The “Inverse” of $R$	185
32 Irreducible Modules of Endpoint $i$	189
33 Algebra on First Subconstituent	197
34 Modules of Endpoint One	205
35 $\dim E_1^* T E_1^* \leq 5$	215
36 Dual Endpoint	219
37 Generalized Adjacency Matrix	225
38 An Injection from $E_{11}^*$ to $E_{22}^*$	231
39 $A^+$ and $A^-$	235
40 Title of the Chapter	237

# About this lecturenote

## Setting

sudo This note is created by `bookdown` package on RStudio.

For `bookdown` See (Xie, 2015), (Xie, 2017), (Yihui Xie, 2018).

1. Log-in to my GitHub Account
2. Go to RStudio/bookdown-demo repository: <https://github.com/rstudio/bookdown-demo>
3. Use This Template
4. Input Repository Name
5. Select Public - default
6. Create repository from template
7. From Code download ZIP
8. Move the extracted folder into a favorite directory
9. Open RStudio Project in the folder
10. Use Terminal in the bottom left pane
  - confirm that the current directory is the home directory of the project by `pwd`
11. (failed to proceed by ssh)
12. Use Console
  1. `library(usethis)`
  2. `use_git()`
  3. `use_github()` — Error
  4. `gh_token_help()`
  5. `create_github_token()`: create a token in the github page. Copy the token
  6. `gitcreds::gitcreds_set()`: paste the token, the token is to be expired in 30 days
13. Use Terminal
  1. `git remote add origin https://github.com/icu-hsuzuki/t-algebra.git`
  2. `git push -u origin main`
  3. type in the password of the computer
14. Use GIT in R Studio

## Another Host

1. create a project by version control git
2. git init
3. git remote add origin git@github.com:/.git
4. git branch -r
5. git fetch
6. git pull origin main

# Chapter 1

## Subconstituent Algebra of a Graph

Wednesday, January 20, 1993

A graph (undirected, without loops or multiple edges) is a pair  $\Gamma = (X, E)$ , where

$$X = \text{finite set (of vertices)} \quad (1.1)$$

$$E = \text{set of (distinct) 2-element subsets of } X (= \text{edges of } \Gamma). \quad (1.2)$$

vertices  $x$  and  $y \in X$  are adjacent if and only if  $xy \in E$ .

**Example 1.1.** Let  $\Gamma$  be a graph.  $X = \{a, b, c, d\}$ ,  $E = \{ab, ac, bc, bd\}$ .



Set  $n = |X|$ , the order of  $\Gamma$ .

Pick a field  $K$  ( $= \mathbb{R}$  or  $\mathbb{C}$ ). Then  $\text{Mat}_X(K)$  denotes the  $K$  algebra of all  $n \times n$  matrices with entries in  $K$ . (rows and columns are indexed by  $X$ )

*Adjacency matrix*  $A \in \text{Mat}_X(K)$  is defined by

$$A_{xy} = \begin{cases} 1 & \text{if } xy \in E \\ 0 & \text{else .} \end{cases} \quad (1.3)$$

**Example 1.2.** Let  $a, b, c, d$  be labels of rows and columns. Then

$$A = \begin{matrix} & \begin{matrix} a & b & c & d \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix}$$

The subalgebra  $M$  of  $\text{Mat}_X(K)$  generated by  $A$  is called the *Bose-Mesner algebra* of  $\Gamma$ .

Set  $V = K^n$ , the set of  $n$ -dimensional column vectors, the coordinates are indexed by  $X$ .

Let  $\langle \cdot, \cdot \rangle$  denote the Hermitean inner product:

$$\langle u, v \rangle = u^\top \cdot v \quad (u, v \in V)$$

$V$  with  $\langle \cdot, \cdot \rangle$  is the *standard module* of  $\Gamma$ .

$M$  acts on  $V$ : For every  $x \in X$ , write

$$\hat{x} = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

where 1 is at the  $x$  position.

Then

$$A\hat{x} = \sum_{y \in X, xy \in E} \hat{y}.$$

Since  $A$  is a real symmetrix matrix,

$$V = V_0 + V_1 + \cdots + V_r \quad \text{some } r \in \mathbb{Z}^{\geq 0},$$

the orthogonal direct sum of maximal  $A$ -eigenspaces.

Let  $E_i \in \text{Mat}_X(K)$  denote the orthogonal projection,

$$E_i : V \longrightarrow V_i.$$

Then  $E_0, \dots, E_r$  are the primitive idempotents of  $M$ .

$$M = \text{Span}_K(E_0, \dots, E_r),$$

$$E_i E_j = \delta_{ij} E_i \quad \text{for all } i, j, \quad E_0 + \cdots + E_r = I.$$



Let  $\theta_i$  denote the eigenvalue of  $A$  for  $V_i$  in  $\mathbb{R}$ . Without loss of generality we may assume that

$$\theta_0 > \theta_1 > \dots > \theta_r.$$

Let

$$m_i = \text{the multiplicity of } \theta_i = \dim V_i = \text{rank } E_i.$$

Set

$$\text{Spec}(\Gamma) = \begin{pmatrix} \theta_0, & \theta_1, & \dots, & \theta_r \\ m_0, & m_1, & \dots, & m_r \end{pmatrix}.$$

**Problem.** What can we say about  $\Gamma$  when  $\text{Spec}(\Gamma)$  is given?

The following Lemma 1.1, is an example of Problem.

For every  $x \in X$ ,

$$k(x) \equiv \text{valency of } x \equiv \text{degree of } x \equiv |\{y \mid y \in X, xy \in E\}|.$$

**Definition 1.1.** The graph  $\Gamma$  is regular of valency  $k$  if  $k = k(x)$  for every  $x \in X$ .

**Lemma 1.1.** *With the above notation,*

- (i)  $\theta_0 \leq \max\{k(x) \mid x \in X\} = k^{\max}$ .
- (ii) *If  $\Gamma$  is regular of valency  $k$ , then  $\theta_0 = k$ .*

*Proof.*

(i) Without loss of generality we may assume that  $\theta_0 > 0$ , else done. Let  $v := \sum_{x \in X} \alpha_x \hat{x}$  denote the eivenvector for  $\theta_0$ .

Pick  $x \in X$  with  $|\alpha_x|$  maximal. Then  $|\alpha_x| \neq 0$ .

Since  $Av = \theta_0 v$ ,

$$\theta_0 \alpha_x = \sum_{y \in X, xy \in E} \alpha_y.$$

So,

$$\theta_0 |\alpha_x| = |\theta_0 \alpha_x| \leq \sum_{y \in X, xy \in E} |\alpha_y| \leq k(x) |\alpha_x| \leq k^{\max} |\alpha_x|.$$

(ii) All 1's vector  $v = \sum_{x \in X} \hat{x}$  satisfies  $Av = kv$ .

□

### Subconstituent Algebra

Let  $x, y \in X$  and  $\ell \in \mathbb{Z}^{\geq 0}$ .

**Definition 1.2.** A path of length  $\ell$  connecting  $x, y$  is a sequence

$$x = x_0, x_1, \dots, x_\ell = y, \quad x_i \in X, \quad 0 \leq i \leq \ell$$

such that  $x_i x_{i+1} \in E$  for  $0 \leq i \leq \ell - 1$ .

**Definition 1.3.** The distance  $\partial(x, y)$  is the length of a shortest path connecting  $x$  and  $y$ .

$$\partial(x, y) \in \mathbb{Z}^{\geq 0} \cup \{\infty\}.$$

**Definition 1.4.** The graph  $\Gamma$  is connected if and only if  $\partial(x, y) < \infty$  for all  $x, y \in X$ .

From now on, assume that  $\Gamma$  is connected with  $|X| \geq 2$ .

Set

$$d_\Gamma = d = \max\{\partial(x, y) \mid x, y \in X\} \equiv \text{the diameter of } \Gamma.$$

Fix a ‘base’ vertex  $x \in X$ .

**Definition 1.5.**

$$d(x) = \text{the diameter with respect to } x = \max\{\partial(x, y) \mid y \in X\} \leq d.$$

Observe that

$$V = V_0^* + V_1^* + \cdots + V_{d(x)}^* \quad (\text{orthogonal direct sum}),$$

where

$$V_i^* = \text{Span}_K(\hat{y} \mid \partial(x, y) = i) \equiv V_i * (x)$$

and  $V_i^* = V_i^*(x)$  is called the  $i$ -th subconstituent with respect to  $x$ .

Let  $E_i^* = E_i^*(x)$  denote the orthogonal projection

$$E_i^* : V \longrightarrow V_i^*(x).$$

View  $E_i^*(x) \in \text{Mat}_X(K)$ . So,  $E_i^*(x)$  is diagonal with  $yy$  entry

$$(E_i^*(x))_{yy} = \begin{cases} 1 & \text{if } \partial(x, y) = i \\ 0 & \text{else,} \end{cases} \quad \text{for } y \in X.$$

Set

$$M^* = M^*(x) \equiv \text{Span}_K(E_0^*(x), \dots, E_{d(x)}^*(x)).$$

Then  $M^*(x)$  is a commutative subalgebra of  $\text{Mat}_X(K)$  and is called the *dual Bose-Mesner algebra with respect to  $x$* .

**Definition 1.6** (Subconstituent Algebra). Let  $\Gamma = (X, E)$ ,  $x, M, M^*(x)$  be as above. Let  $T = T(x)$  denote the subalgebra of  $\text{Mat}_X(K)$  generated by  $M$  and  $M^*(x)$ .  $T$  is the *subconstituent algebra* of  $\Gamma$  with respect to  $x$ .

**Definition 1.7.** A  $T$ -module is any subspace  $W \subset V$  such that  $aw \in W$  for all  $a \in T$  and  $w \in W$ .

$T$ -module  $W$  is *irreducible* if and only if  $W \neq 0$  and  $W$  does not properly contain a nonzero  $T$ -module.

For any  $a \in \text{Mat}_X(K)$ , let  $a^*$  denote the conjugate transpose of  $a$ .

Observe that

$$\langle au, v \rangle = \langle u, a^*v \rangle \quad \text{for all } a \in \text{Mat}_X(K), \text{ and for all } u, v \in V.$$

**Lemma 1.2.** *Let  $\Gamma = (X, E)$ ,  $x \in X$  and  $T \equiv T(x)$  be as above.*

(i) *If  $a \in T$ , then  $a^* \in T$ .*

(ii) *For any  $T$ -module  $W \subset V$ ,*

$$W^\perp := \{v \in V \mid \langle w, v \rangle = 0, \text{ for all } w \in W\}$$

*is a  $T$ -module.*

(iii)  *$V$  decomposes as an orthogonal direct sum of irreducible  $T$ -modules.*

*Proof.*

(i) It is because  $T$  is generated by symmetric real matrices

$$A, E_0^*(x), E_1^*(x), \dots, E_{d(x)(x)}^*.$$

(ii) Pick  $v \in W^\perp$  and  $a \in T$ , it suffices to show that  $av \in W^\perp$ . For all  $w \in W$ ,

$$\langle w, av \rangle = \langle a^*w, v \rangle = 0$$

as  $a^* \in T$ .

(iii) This is proved by the induction on the dimension of  $T$ -modules. If  $W$  is an irreducible  $T$ -module of  $V$ , then

$$V = W + W^\perp \quad (\text{orthogonal direct sum}).$$

□

**Problem.** What does the structure of the  $T(x)$ -module tell us about  $\Gamma$ ?

Study those  $\Gamma$  whose modules take ‘simple’ form. The  $\Gamma$ ’s involved are highly regular.

*Remark.*

1. The subconstituent algebra  $T$  is semisimple as the left regular representation of  $T$  is completely reducible. See Curtis-Reiner 25.2 (Charles W. Curtis, 2006).
2. The inner product  $\langle a, b \rangle_T = \text{tr}(a^\top \bar{b})$  is nondegenerate on  $T$ .

3. In general,

$T$ : Semisimple and Artinian  $\Leftrightarrow T$ : Artinian with  $J(T) = 0$

$\Leftrightarrow T$ : Artinian with nonzero nilpotent element

$\Leftrightarrow T \subset \text{Mat}_X(K)$  such that for all  $a \in T$  is normal.

## Chapter 2

# Perron-Frobenius Theorem

Friday, January 22, 1993

In this lecture we use the Perron Frobenius theory of nonnegative matrices to obtain information on eigenvalues of a graph.

Let  $K = \mathbb{R}$ . For  $n \in \mathbb{Z}^{>0}$ , pick a symmetric matrix  $C \in \text{Mat}_n(\mathbb{R})$ .

**Definition 2.1.** The matrix  $C$  is *reducible* if and only if there is a bipartition  $\{1, 2, \dots, n\} = X^+ \cup X^-$  (disjoint union of nonempty sets) such that  $C_{ij} = 0$  for all  $i \in X^+$ , and for all  $j \in X^-$ , and for all  $i \in X^-$ , and for all  $j \in X^+$ , i.e.,

$$C \sim \begin{pmatrix} * & O \\ O & * \end{pmatrix}.$$

**Definition 2.2.** The matrix  $C$  is *bipartite* if and only if there is a bipartition  $\{1, 2, \dots, n\} = X^+ \cup X^-$  (disjoint union of nonempty sets) such that  $C_{ij} = 0$  for all  $i, j \in X^+$ , and for all  $i, j \in X^-$ , i.e.,

$$C \sim \begin{pmatrix} O & * \\ * & O \end{pmatrix}.$$

**Note.**

1. If  $C$  is bipartite, for every eigenvalue  $\theta$  of  $C$ ,  $-\theta$  is an eigenvalue of  $C$  such that  $\text{mult}(\theta) = \text{mult}(-\theta)$ .

Indeed, let  $C = \begin{pmatrix} O & A \\ B & O \end{pmatrix}$ ,

$$\begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \theta \begin{pmatrix} x \\ y \end{pmatrix} \Leftrightarrow \begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ -y \end{pmatrix} = -\theta \begin{pmatrix} x \\ -y \end{pmatrix},$$

where  $Ay = \theta x$  and  $Bx = \theta y$ .

2. If  $C$  is bipartite,  $C^2$  is reducible.
3. The matrix  $C$  is irreducible and  $C^2$  is reducible, if  $C_{ij} \geq 0$  for all  $i, j$  and  $C$  is reducible. (Exercise)

*Remark.* Note 1. Even if  $C$  is not symmetric

$$\begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \theta \begin{pmatrix} x \\ y \end{pmatrix} \Leftrightarrow \begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ -y \end{pmatrix} = -\theta \begin{pmatrix} x \\ -y \end{pmatrix}$$

holds. So the geometrix multiplicities coincide. How about the algebraic multiplicities?

Note 3. Set  $x \sim y$  if and only if  $C_{xy} > 0$ . So the graph may have loops. Then

$$(C^2)_{xy} > 0 \Leftrightarrow \text{if there exists } z \in X \text{ such that } x \sim z \sim y.$$

Note that  $C$  is irreducible if and only if  $\Gamma(C)$  is connected. Let

$$X^+ = \{y \mid \text{there is a path of even length from } x \text{ to } y\} \quad (2.1)$$

$$X^- = \{y \mid \text{there is no path of even length from } x \text{ to } y\} \neq \emptyset. \quad (2.2)$$

If there is an edge  $y \sim z$  in  $X^+$  and  $w \in X^-$ . Then there would be a path from  $x$  to  $y$  of even length. So  $e(X^+, X^+) = e(X^-, X^-) = 0$ .

**Theorem 2.1** (Perron-Frobenius). *Given a matrix  $C$  in  $\text{Mat}_n(\mathbb{R})$  such that*

- (a)  $C$  is symmetric.
- (b)  $C$  is irreducible.
- (c)  $C_{ij} \geq 0$  for all  $i, j$ .

*Let  $\theta_0$  be the maximal eigenvalue of  $C$  with eigenspace  $V_0 \subseteq \mathbb{R}^n$ , and let  $\theta_r$  be the maximal eigenvalue of  $C$  with eigenspace  $V_r \subseteq \mathbb{R}^n$ . Then the following hold.*

$$(i) \text{ Suppose } 0 \neq v = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_n \end{pmatrix} \in V_0. \text{ Then } \alpha_0 > 0 \text{ for all } i, \text{ or } \alpha_i < 0 \text{ for all } i.$$

$$(ii) \dim V_0 = 1.$$

$$(iii) \theta_r \geq -\theta_0.$$

$$(iv) \theta_r = \theta_0 \text{ if and only if } C \text{ is bipartite.}$$

First, we prove the following lemma.

**Lemma 2.1.** *Let  $\langle \cdot, \cdot \rangle$  be the dot product in  $V = \mathbb{R}^n$ . Pick a symmetric matrix  $B \in \text{Mat}_n(\mathbb{R})$ . Suppose all eigenvalues of  $B$  are nonnegative. (i.e.,  $B$  is positive semidefinite.) Then there exist vectors  $v_1, v_2, \dots, v_n \in V$  such that  $B_{ij} = \langle v_i, v_j \rangle$  for  $(1 \leq i, j \leq n)$ .*

*Proof.* By elementary linear algebra, there exists an orthonormal basis  $w_1, w_2, \dots, w_n$  of  $V$  consisting of eigenvectors of  $B$ . Set the  $i$ -th column of  $P$  is  $w_i$  and  $D = \text{diag}(\theta_1, \dots, \theta_n)$ . Then  $P^\top P = I$  and  $BP = PD$ .

Hence,

$$B = PDP^{-1} = PDP^\top = QQ^\top,$$

where

$$Q = P \cdot \text{diag}(\sqrt{\theta_1}, \sqrt{\theta_2}, \dots, \sqrt{\theta_n}) \in \text{Mat}_n(\mathbb{R}).$$

Now, let  $v_i$  be the  $i$ -th column of  $Q^\top$ . Then

$$B_{ij} = v_i^\top \cdot v_j = \langle v_i, v_j \rangle.$$

□

Now we start the proof of Theorem 2.1.

*Proof of Theorem 2.1(i)*

Let  $\langle, \rangle$  denote the dot product on  $V = \mathbb{R}^n$ . Set

$$B = \theta I - C \tag{2.3}$$

$$= \text{symmetric matrix with eigenvalues } \theta_0 - \theta_i \geq 0 \tag{2.4}$$

$$= (\langle v_i, v_j \rangle)_{1 \leq i, j \leq n} \tag{2.5}$$

with the same  $v_1, \dots, v_n \in V$  by Lemma 2.1.

Observe:  $\sum_{i=1}^n \alpha_i v_i = 0$ .

*Pf.*

$$\left\| \sum_{i=1}^n \alpha_i v_i \right\|^2 = \left\langle \sum_{i=1}^n \alpha_i v_i, \sum_{i=1}^n \alpha_i v_i \right\rangle \tag{2.6}$$

$$= (\alpha_1 \quad \dots \quad \alpha_n) B \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \tag{2.7}$$

$$= v^\top B v \tag{2.8}$$

$$= 0, \tag{2.9}$$

since  $Bv = (\theta_0 I - C)v = 0$ .

Now set

$$s = \text{the number of indices } i, \text{ where } \alpha_i > 0.$$

Replacing  $v$  by  $-v$  if necessary, without loss of generality we may assume that  $s \geq 1$ . We want to show  $s = n$ .

Assume  $s < n$ . Without loss of generality, we may assume that  $\alpha_i > 0$  for  $1 \leq i \leq s$  and  $\alpha_i = 0$  for  $s+1 \leq i \leq n$ . Set

$$\rho = \alpha_1 v_1 + \dots + \alpha_s v_s = -\alpha_{s+1} v_{s+1} - \dots - \alpha_n v_n.$$

Then, for  $i = 1, \dots, s$ ,

$$\langle v_i, \rho \rangle = \sum_{j=s+1}^n -\alpha_j \langle v_i, v_j \rangle \quad (\langle v_i, v_j \rangle = B_{ij}, B = \theta_0 I - C) \quad (2.10)$$

$$= \sum_{j=s+1}^n (-\alpha_j)(-C_{ij}) \quad (2.11)$$

$$\leq 0. \quad (2.12)$$

Hence

$$0 \leq \langle \rho, \rho \rangle = \sum_{i=1}^s \alpha_i \langle v_i, \rho \rangle \leq 0,$$

as  $\alpha > 0$  and  $\langle v_i, \rho \rangle \leq 0$ . Thus, we have  $\langle \rho, \rho \rangle = 0$  and  $\rho = 0$ . For  $j = s+1, \dots, n$ ,

$$0 = \langle \rho, v_j \rangle = \sum_{i=1}^s \alpha_i \langle v_i, v_j \rangle \leq 0,$$

as  $\langle v_i, v_j \rangle = -C_{ij}$ .

Therefore,

$$0 = \langle v_i, v_j \rangle = -C_{ij} \text{ for } 1 \leq i \leq s, s+1 \leq j \leq n.$$

Since  $C$  is symmetric,

$$C = \begin{pmatrix} * & O \\ O & * \end{pmatrix}$$

Thus  $C$  is reducible, which is not the case. Hence  $s = n$ .

*Proof of Theorem 2.1 (ii).*

Suppose  $\dim V_0 \geq 2$ . Then,

$$\dim \left( V_0 \cap \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}^\perp \right) \geq 1.$$

So, there is a vector

$$0 \neq v = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \in V_0$$

with  $\alpha_1 = 0$ . This contradicts 1.

Now pick

$$0 \neq w = \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_n \end{pmatrix} \in V_r.$$

*Proof of Theorem 2.1 (iii).*



Suppose  $\theta_r < -\theta_0$ . Since the eigenvalues of  $C^2$  are the squares of those of  $C$ ,  $\theta_r^2$  is the maximal eigenvalue of  $C^2$ .

Also we have  $C^2 w = \theta_r^2 w$ .

Observe that  $C^2$  is irreducible. (As otherwise,  $C$  is bipartite by Note 3, and we must have  $\theta_r = -\theta_0$ .) Therefore,  $\beta_i > 0$  for all  $i$  or  $\beta_i < 0$  for all  $i$ . We have

$$\langle v, w \rangle = \sum_{i=1}^n \alpha_i \beta_i \neq 0.$$

This is a contradiction, as  $V_0 \perp V_r$ .

*Proof of Theorem 2.1 (iv)*

$\Rightarrow$ : Let  $\theta_r = -\theta_0$ . Then  $\theta = \theta_1^2 = \theta_0^2$  is the maximal eigenvalue of  $C^2$ , and  $v$  and  $w$  are linearly independent eigenvalues for  $\theta$  for  $C^2$ . Hence, for  $C^2$ ,  $\text{mult}(\theta) \geq 2$ .

Thus by 2,  $C^2$  must be reducible. Therefore,  $C$  is bipartite by Note 3.

$\Leftarrow$ : This is Note 1.  $\square$

Let  $\Gamma = (X, E)$  be any graph.

**Definition 2.3.**  $\Gamma$  is said to be *bipartite* if the adjacency matrix  $A$  is bipartite. That is,  $X$  can be written as a disjoint union of  $X^+$  and  $X^-$  such that  $X^+, X^-$  contain no edges of  $\Gamma$ .

**Corollary 2.1.** *For any (connected) graph  $\Gamma$  with*

$$\text{Spec}(\Gamma) = \begin{pmatrix} \theta_0 & \theta_1 & \cdots & \theta_r \\ m_1 & m_1 & \cdots & m_r \end{pmatrix} \quad \text{with } \theta_0 > \theta_1 > \cdots > \theta_r.$$

*Let  $V_i$  be the eigenspace of  $\theta_i$ . Then the following holds.*

1. Suppose  $0 \neq v = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \in V_0 \in \mathbb{R}^n$ . Then  $\alpha_i > 0$  for all  $i$  or  $\alpha_i < 0$  for all  $i$ .
2.  $m_0 = 1$ .
3.  $\theta_r \geq -\theta_0$  if and only if  $\Gamma$  is bipartite. In this case,

$$-\theta_i = \theta_{r-i} \text{ and } m_i = m_{r-i} \quad (0 \leq i \leq r)$$

*Proof.* This is a direct consequences of Theorem 2.1 and Note 3.  $\square$



## Chapter 3

# Cayley Graphs

Monday, January 25, 1993

Given graphs  $\Gamma = (X, E)$  and  $\Gamma' = (X', E')$ .

**Definition 3.1.** A map  $\sigma : X \rightarrow X'$  is an *isomorphism* of graphs whenever;

- i.  $\sigma$  is one-to-one and onto,
- ii.  $xy \in E$  if and only if  $\sigma x \sigma y \in E'$  for all  $x, y \in X$ .

We do not distinguish between isomorphic graphs.

**Definition 3.2.** Suppose  $\Gamma = \Gamma'$ . Above isomorphism  $\sigma$  is called an *automorphism* of  $\Gamma$ . Then set  $\text{Aut}(\Gamma)$  of all automorphisms of  $\Gamma$  becomes a finite group under composition.

**Definition 3.3.** If  $\text{Aut}(\Gamma)$  acts transitive on  $X$ ,  $\Gamma$  is called *vertex transitive*.

**Example 3.1.** A Cayley graphs:

**Definition 3.4** (Cayley Graphs). Let  $G$  be any finite group, and  $\Delta$  any generating set for  $G$  such that  $1_G \notin \Delta$  and  $g \in \Delta \rightarrow g^{-1} \in \Delta$ . Then Cayley graph  $\Gamma = \Gamma(G, \Delta)$  is defined on the vertex set  $X = G$  with the edge set  $E$  defined by the following.

$$E = \{(h_1, h_2) \mid h_1, h_2 \in G, h_1^{-1}h_2 \in \Delta\} = \{(h, hg) \mid h \in G, g \in \Delta\}$$

**Example 3.2.**  $G = \langle a \mid a^6 = 1 \rangle$ ,  $\Delta = \{a, a^{-1}\}$ .



**Example 3.3.**  $G = \langle a \mid a^6 = 1 \rangle$ ,  $\Delta = \{a, a^{-1}, a^2, a^{-2}\}$ .



**Example 3.4.**  $G = \langle a, b \mid a^6 = 1, b^2 = 1, ab = ba \rangle$ ,  $\Delta = \{a, a^{-1}, b\}$ .



*Remark.*  $\text{Aut}(\Gamma) \simeq D_6 \times \mathbb{Z}_2$  contains two regular subgroups isomorphic to  $D_6$  and  $\mathbb{Z}_5 \times \mathbb{Z}_2$  and  $\Gamma$  is obtained as Cayley graphs in two ways.

Cayley graphs are vertex transitive, indeed.

**Theorem 3.1.** *The following hold.*

(i) *For any Cayley graph  $\Gamma = \Gamma(G, \Delta)$ , the map*

$$G \rightarrow \text{Aut}(\Gamma) \quad (g \mapsto \hat{g})$$

is an injective homomorphism of groups, where

$$\hat{g}(x) = gx \quad \text{for all } g \in G \text{ and for all } x \in X (= G).$$

Also, the image  $\hat{G}$  is regular on  $X$ . i.e., the image  $\hat{G}$  acts transitively on  $X$  with trivial vertex stabilizers.

(ii) For any graph  $\Gamma = (X, E)$ , suppose there exists a subgroup  $G \subseteq \text{Aut}(\Gamma)$  that is regular on  $X$ . Pick  $x \in X$ , and let

$$\Delta = \{g \in G \mid \langle x, g(x) \rangle \in E\}.$$

Then  $1 \notin \Delta$ ,  $g \in \Delta \rightarrow g^{-1} \in \Delta$ , and  $\Delta$  generates  $G$ . Moreover,  $\Gamma \simeq \Gamma(G, \Delta)$ .

*Proof.* (i) Let  $g \in G$ . We want to show that  $\hat{g} \in \text{Aut}(\Gamma)$ . Let  $h_1, h_2 \in X = G$ . Then,

$$(h_1, h_2) \in E \rightarrow h_1^{-1}h_2 \in \Delta \quad (3.1)$$

$$\rightarrow (gh_1)^{-1}(gh_2) \in \Delta \quad (3.2)$$

$$\rightarrow (gh_1, gh_2) \in E \quad (3.3)$$

$$\rightarrow (\hat{g}(h_1), \hat{g}(h_2)) \in E. \quad (3.4)$$

Hence,  $\hat{g} \in \text{Aut}(\Gamma)$ .

Observe:  $g \mapsto \hat{g}$  is a homomorphism of groups:

$$\hat{1}_G = 1, \widehat{g_1 g_2} = \widehat{g_1} \widehat{g_2}.$$

Observe:  $g \mapsto \hat{g}$  is one-to-one:

$$\widehat{g_1} = \widehat{g_2} \rightarrow g_1 = \widehat{g_1}(1_G) = \widehat{g_2}(1_G) = g_2.$$

Observe:  $\hat{G}$  is regular on  $X$ : Clear by construction.

(ii)  $1_G \notin \Delta$ : Since  $\Gamma$  has not loops,  $(x, 1_G x) \notin E$ .

$g \in \Delta \rightarrow g^{-1} \in \Delta$ :

$$g \in \Delta \rightarrow (x, g(x)) \in E \rightarrow E \ni (g^{-1}(x), g^{-1}(g(x))) = (g^{-1}(x), x).$$

$\Delta$  generates  $G$ : Suppose  $\langle \Delta \subsetneq G$ . Let  $\hat{X} = \{g(x) \mid g \in \langle \Delta \rangle\} \subsetneq X$ . ( $\hat{X} \subsetneq X$  as  $G$  acts regularly on  $X$ .)

Since  $\Gamma$  is connected, there exists  $y \in \hat{X}$  and  $z \in X \setminus \hat{X}$  with  $yz \in E$ .

Let  $y = g(x)$ ,  $g \in \langle \Delta \rangle$ ,  $z \in h(x)$ ,  $h \in G \setminus \langle \Delta \rangle$ . Then

$$(y, z) = (g(x), h(x)) \in E \rightarrow (x, g^{-1}h(x)) \in E \rightarrow g^{-1}h \in \langle \Delta \rangle \rightarrow h \in \langle \Delta \rangle.$$

This is a contradiction. Therefore,  $\Delta$  generates  $G$ .

Let  $\Gamma' = (X', E')$  denote  $\Gamma(G, \Delta)$ . We shall show that

$$\theta : X' \rightarrow X \ (g \mapsto g(x))$$

is an isomorphism of graphs.

$\theta$  is one-to-one: For  $h_1, h_2 \in X' = G$ ,

$$\theta(h_1) = \theta(h_2) \rightarrow h_1(x) = h_2(x) \rightarrow h_1^{-1}h_2(x) = x \rightarrow h_1^{-1}h_2 \in \text{Stab}_G(x) = \{1_G\} \rightarrow h_1 = h_2.$$

( $\text{Stab}_G = \{g \in G \mid g(x) = x\}$ .)

$\theta$  is onto: Since  $G$  is transitive,

$$X = \{g(x) \mid g \in G\} = \theta(X') = \theta(G).$$

$\theta$  respects adjacency: For  $h_1, h_2 \in X' = G$ ,

$$(h_1, h_2) \in E' \leftrightarrow h_1^{-1}h_2 \in \Delta \leftrightarrow (x, h_1^{-1}h_2(x)) \in E \leftrightarrow (h_1(x), h_2(x)) \in E \leftrightarrow (\theta(h_1), \theta(h_2)) \in E.$$

Therefore  $\theta$  is an isomorphism between graphs  $\Gamma(G, \Delta)$  and  $\Gamma(X, E)$ .  $\square$

How to compute the eigenvalues of the Cayley graph of an abelian group.

Let  $G$  be any finite abelian group. Let  $\mathbb{C}^*$  be the multiplicative group on  $\mathbb{C} \setminus \{0\}$ .

**Definition 3.5.** A (linear)  $G$ -character is any group homomorphism  $\theta : G \rightarrow \mathbb{C}^*$ .

**Example 3.5.**  $G = \langle a \mid a^3 = 1 \rangle$  has three characters,  $\theta_0, \theta_1, \theta_2$ .

$$\begin{array}{c|ccc} \theta_i(a^j) & 1 & a & a^2 \\ \hline \theta_0 & 1 & 1 & 1 \\ \theta_1 & 1 & \omega & \omega^2 \\ \theta_2 & 1 & \omega^2 & \omega \end{array}, \quad \text{with } \omega = \frac{-1 + \sqrt{-3}}{2}.$$

Here  $\omega$  is a primitive cube root of  $\omega$  in  $\mathbb{C}^*$ , i.e.,  $1 + \omega + \omega^2 = 0$ .

For arbitrary group  $G$ , let  $X(G)$  be the set of all characters of  $G$ .

Observe: For  $\theta_1, \theta_2 \in X(G)$ , one can define product  $\theta_1 \theta_2$ :

$$\theta_1 \theta_2(g) = \theta_1(g) \theta_2(g) \quad \text{for all } g \in G.$$

Then  $\theta_1 \theta_2 \in X(G)$ .

Observe:  $X(G)$  with this product is an (abelian) group.

**Lemma 3.1.** *The groups  $G$  and  $X(G)$  are isomorphic for all finite abelian groups  $G$ .*

*Proof.*  $G$  is a direct sum of cyclic groups;

$$G = G_1 \oplus G_2 \oplus \cdots \oplus G_m, \quad \text{where } G_i = \langle a_i \mid a_i^{d_i} = 1 \rangle \quad (1 \leq i \leq m).$$

Pick any element  $\omega_i$  of order  $d_i$  in  $\mathbb{C}^*$ , i.e., a primitive  $d_i$ -th root of 1. Define

$$\theta_i : G \rightarrow \mathbb{C}^* \quad (a_1^{\varepsilon_1} \cdots a_m^{\varepsilon_m} \mapsto \omega_i^{\varepsilon_i} \quad \text{where } 0 \leq \varepsilon_i < d_i, 1 \leq i \leq m).$$

Then  $\theta_i \in X(G)$ . (Exercise)

Claim: There exists an isomorphism of groups  $G \rightarrow X(G)$  that sends  $a_i$  to  $\theta_i$ .

Observe:  $\theta_i^{d_i} = 1$ . For every  $g = a_1^{\varepsilon_1} \cdots a_m^{\varepsilon_m} \in G$ ,

$$\theta_i^{d_i}(g) = (\theta_i(g))^{d_i} = (\omega_i^{\varepsilon_i})^{d_i} = (\omega_i^{d_i})^{\varepsilon_i} = 1.$$

Observe: If  $\theta_1^{\varepsilon_1} \theta_2^{\varepsilon_2} \cdots \theta_m^{\varepsilon_m} = 1$  for some  $0 \leq \varepsilon_i < d_i, 1 \leq i \leq m$ . Then  $\varepsilon_1 = \varepsilon_2 = \cdots = \varepsilon_m = 0$ .

*Pf.*  $1 = \theta_1^{\varepsilon_1} \theta_2^{\varepsilon_2} \cdots \theta_m^{\varepsilon_m}(a_i) = \omega_i^{\varepsilon_i}$ , Since  $\omega_i$  is a primitive  $d_i$ -th root of 1,  $\varepsilon_i = 0$  for  $1 \leq i \leq m$ .

Observe:  $\theta_1, \dots, \theta_m$  generate  $X(G)$ . Pick  $\theta \in X(G)$ . Since  $a_i^{d_i} = 1$ ,  $1 = \theta(a_i^{d_i}) = \theta(a_i)^{d_i}$ .

Hence  $\theta(a_i) = \omega_i^{\varepsilon_i}$  for some  $\varepsilon_i$  with  $0 \leq \varepsilon_i < d_i$ .

Now  $\theta = \theta_1^{\varepsilon_1} \cdots \theta_m^{\varepsilon_m}$ , since these are both equal to  $\omega_i^{\varepsilon_i}$  at  $a_i$  for  $1 \leq i \leq m$ .

Therefore,

$$G \rightarrow X(G) \quad (a_i \mapsto \theta_i)$$

is an isomorphism of groups. □

**Note.** The correspondence above is clearly a group homomorphism.





## Chapter 4

# Examples

Wednesday, January 27, 1993

**Theorem 4.1.** *Given a Cayley graph  $\Gamma = \Gamma(G, \Delta)$ . View the standard module  $V \equiv \mathbb{C}G$  (the group algebra), so*

$$\left\langle \sum_{g \in G} \alpha_g g, \sum_{g \in G} \beta_g g \right\rangle = \sum_{g \in G} \alpha_g \overline{\beta_g}, \quad \text{with } \alpha_g, \beta_g \in \mathbb{C}.$$

For any  $\theta \in X(G)$ , write

$$\hat{\theta} = \sum_{g \in G} \theta(g^{-1})g.$$

Then the following hold.

(i)  $\langle \hat{\theta}_1, \hat{\theta}_2 \rangle = |G|$  if  $\theta_1 = \theta_2$  and 0 otherwise for  $\theta_1, \theta_2 \in X(G)$ . In particular,  $\{\hat{\theta} \mid \theta \in X(G)\}$  forms a basis for  $V$ .

(ii)  $A\hat{\theta} = \Delta_\theta \hat{\theta}$  for  $\theta \in X(G)$ , where  $A$  is the adjacency matrix and

$$\Delta_\theta = \sum_{g \in \Delta} \theta(g).$$

In particular, the eigenvalues of  $\Gamma$  are precisely

$$\Delta_\theta \mid \theta \in X(G)\}.$$

*Proof.*

(i) Claim: For every  $\theta \in X(G)$ , let

$$s := \sum_{g \in G} \theta(g^{-1}) = \begin{cases} |G| & \text{if } \theta = 1 \\ 0 & \text{if } \theta \neq 1. \end{cases}$$

*Pf.* Clear if  $\theta = 1$ .

Let  $\theta \neq 1$ . Then  $\theta(h) \neq 1$  for some  $h \in G$ .

$$s \cdot \theta(h) = \left( \sum_{g \in G} \theta(g^{-1}) \right) \theta(h) = \sum_{g \in G} \theta(g^{-1}h) = \sum_{g' \in G} \theta(g'^{-1}) = s.$$

Since  $\theta(h) \neq 1$ ,  $s = 0$ .

Claim.  $\theta(g^{-1}) = \overline{\theta(g)}$  for every  $\theta \in X(G)$  and every  $g \in G$ .

Since  $\theta(g) \in \mathbb{C}$  is a root of 1,

$$|\theta(g)|^2 = \theta(g)\overline{\theta(g)} = 1.$$

On the other hand, since  $\theta$  is a homomorphism,

$$\theta(g)\theta(g^{-1}) = \theta(1) = 1.$$

Hence  $\theta(g^{-1}) = \overline{\theta(g)}$ .

Now

$$\langle \widehat{\theta_1}, \widehat{\theta_2} \rangle = \sum_{g \in G} \theta_1(g^{-1}) \overline{\theta_2(g^{-1})} \quad (4.1)$$

$$= \sum_{g \in G} \theta_1(g^{-1}) \theta_2(g) \quad (4.2)$$

$$= \sum_{g \in G} \theta_1 \theta_2^{-1}(g^{-1}) \quad (4.3)$$

$$= \begin{cases} |G| & \text{if } \theta_1 \theta_2^{-1} = 1 \\ 0 & \text{if } \theta_1 \theta_2^{-1} \neq 1. \end{cases} \quad (4.4)$$

Since  $|G| = |X(G)|$  by Lemma 3.1, and  $\widehat{\theta_i}$ 's are orthogonal nonzero elements in  $V$ , they form a basis of  $V$ .

(ii) Let  $\Delta = \{g_1, \dots, g_r\}$ . Then

$$A\hat{\theta} = A \left( \sum_{g \in G} \theta(g^{-1}g) \right) \quad (4.5)$$

$$= \sum_{g \in G} \theta(g^{-1})(gg_1 + \cdots + gg_r) \quad (\Gamma(g) = \{gg_1, \dots, gg_r\}) \quad (4.6)$$

$$= \sum_{i=1}^r \left( \sum_{g \in G} \theta(g^{-1})(gg_i) \right) \quad (4.7)$$

$$= \sum_{i=1}^r \left( \sum_{g \in G} \theta(g_i g_i^{-1} g^{-1})(gg_i) \right) \quad (4.8)$$

$$= \sum_{i=1}^r \left( \sum_{g \in G} \theta(g_i) \theta((gg_i)^{-1}) gg_i \right) \quad (4.9)$$

$$= \sum_{i=1}^r \theta(g_i) \sum_{h \in G} \theta(h^{-1}) h \quad (4.10)$$

$$= \Delta_{\theta} \cdot \hat{\theta}. \quad (4.11)$$

Since  $\{\hat{\theta} \mid \theta \in X(G)\}$  forms a basis, the eigenvalues of  $\Gamma$  are precisely,

$$\{\Delta_{\theta} \mid \theta \in X(G)\}.$$

This completes the proof.

□

**Example 4.1.** Let  $G = \langle a \mid a^6 = 1 \rangle$ , and  $\Delta = \{a, a^{-1}\}$ . Pick a primitive 6-th root of 1,  $\omega$ . Then

$$X(G) = \{\theta^i \mid 0 \leq i \leq 5\} \quad \text{such that} \quad \theta(a) = \omega, \quad \omega + \omega^{-1} = 1.$$



$\varphi \in X(G)$	$\varphi(a)$	$\Delta_\varphi = \theta(a) + \theta(a)^{-1}$
1	1	2
$\theta$	$\omega$	$\omega + \omega^{-1} = 1$
$\theta^2$	$\omega^2$	-1
$\theta^3$	$\omega^3 = -1$	-2
$\theta^4$	$\omega^4$	-1
$\theta^5$	$\omega^5$	1

$$\text{Spec}(\Gamma) = \begin{pmatrix} 2 & 1 & -1 & -2 \\ 1 & 2 & 2 & 1 \end{pmatrix}.$$

**Example 4.2.**  $D$ -cube,  $H(D, 2)$ . Let

$$X = \{(a_1, \dots, a_D) \mid a_i \in \{1, -1\}, 1 \leq i \leq D\},$$

$$E = \{xy \mid x, y \in X, x, y: \text{different in exactly one coordinate}\}.$$

Also  $H(D, 2)$  is a Cayley graph  $\Gamma(G, \Delta)$ , where

$$G = G_1 \oplus G_2 \oplus \dots \oplus G_D,$$

$$G_i = \langle a_i \mid a_i^2 = 1 \rangle, \quad \Delta = \{a_1, \dots, a_D\}.$$

**Homework:** The spectrum of  $H(D, 2)$  is

$$\begin{pmatrix} \theta_0 & \theta_1 & \dots & \theta_D \\ m_0 & m_1 & \dots & m_D \end{pmatrix},$$

where

$$\theta_i = D - 2i \quad (0 \leq i \leq D), \quad m_i = \binom{D}{i}.$$

*Remark.* Let  $\theta \in X(G)$ . Then  $\theta : X \rightarrow \{\pm 1\}$ . If

$$\nu(\theta) = |\{i \mid \theta(a_i) = -1\}|,$$

then  $\Delta_\theta = D - 2i$ . Since there are  $\binom{D}{i}$  such  $\theta$ , we have the assertion.

We want to compute the subconstituent algebra for  $H(D, 2)$ . First, we make a few observations about arbitrary graphs.

Let  $\Gamma = (X, E)$  be any graph,  $A$ , the adjacency matrix of  $\Gamma$ , and  $V$ , the standard module over  $K = \mathbb{C}$ .

Fix a base  $x \in X$ . Write  $E_i^* = E_i^*(x)$ , and

$$T \equiv T(x) = \text{the algebra generated by } A, E_0^*, E_1^*, \dots$$

**Definition 4.1.** Let  $W$  be any irreducible  $T$ -module ( $\subseteq V$ ). Then the endpoint  $r \equiv r(W)$  satisfied

$$r = \min\{i \mid E_i^* W \neq 0\}.$$

The diameter  $d = d(W)$  satisfied

$$d = |\{i \mid E_i^* W \neq 0\}| - 1.$$

**Lemma 4.1.** With the above notation, let  $W$  be an irreducible  $T$ -module. Then

- (i)  $E_i^* A E_j^* = 0$  if  $|i - j| > 1$ ,  $E_i^* A E_j^* \neq 0$  if  $|i - j| = 1$ ,  $0 \leq i, j \leq d(x)$ .
- (ii)  $A E_j^* W \subseteq E_{j-1}^* W + E_j^* W + E_{j+1}^* W$ ,  $0 \leq j \leq d(x)$ . ( $E_i^* W = 0$  if  $i < j$  or  $i > d(x)$ .)
- (iii)  $E_j^* W \neq 0$  if  $r \leq j \leq r + d$ ,  $= 0$  if  $0 \leq j \leq r$  or  $r + d < j \leq d(x)$ .
- (iv)  $E_i^* A E_j^* W \neq 0$ , if  $|i - j| = 1$  ( $r \leq i, j \leq r + d$ ).

*Proof.*

- (i) Pick  $y \in X$  with  $\partial(x, y) = j$ . We want to find  $E_i^* A E_j^* \hat{y}$ . Note,

$$E_j^* \hat{y} = \begin{cases} 0 & \text{if } \partial(x, y) \neq j \\ \hat{y} & \text{if } \partial(x, y) = j. \end{cases}$$

$$E_i^* A E_j^* \hat{y} = E_i^* A \hat{y} \tag{4.12}$$

$$= E_i^* \sum_{z \in X, yz \in E} \hat{z} \tag{4.13}$$

$$= \sum_{z \in X, yz \in E, \partial(x, z) = i} \hat{z} \tag{4.14}$$

$$= 0 \text{ if } |i - j| > 1 \quad \text{by triangle inequality.} \tag{4.15}$$

If  $|i - j| = 1$ , there exist  $y, y' \in X$  such that  $\partial(x, y) = j$ ,  $\partial(x, y') = i$ ,  $yy' \in E$  by connectivity of  $\Gamma$ . Hence (4.14) contains  $\widehat{yy'}$  and (4.14) is not equal to zero.

- (ii) We have

$$A E_j^* W = \left( \sum_{i=0}^{d(x)} E_i^* \right) A E_j^* W \tag{4.16}$$

$$= E_{j-1}^* A E_j^* W + E_j^* A E_j^* W + E_{j+1}^* A E_j^* W \tag{4.17}$$

$$\subseteq E_{j-1}^* W + E_j^* W + E_{j+1}^* W. \tag{4.18}$$

- (iii) Suppose  $E_j^* W = 0$  for some  $j$  ( $r \leq j \leq r + d$ ). Then  $r < j$  by the definition of  $r$ . Set

$$\widetilde{W} = E_r^*W + E_{r+1}^*W + \cdots + E_{j-1}^*W.$$

Observe  $0 \subsetneq \widetilde{W} \subsetneq W$ . Also  $A\widetilde{W} \subseteq \widetilde{W}$  by (ii), and  $E_i^*\widetilde{W} \subseteq \widetilde{W}$  for every  $i$  by construction.

Thus,  $T\widetilde{W} \subseteq \widetilde{W}$ , contradicting  $W$  being irreducible.

□

## Chapter 5

# $T$ -Modules of $H(D, 2)$ , I

Friday, January 29, 1993

Let  $\Gamma = (X, E)$  be a graph,  $A$  the adjacency matrix, and  $V$  the standard module over  $K = \mathbb{C}$ .

Fix a base  $x \in X$  and write  $E_i^* \equiv E_i^*(x)$ , and  $T \equiv T(x)$ .

Let  $W$  be an irreducible  $T$ -module with endpoint  $r := \min\{i \mid E_i^*W \neq 0\}$  and diameter  $d := |\{i \mid E_i^*W \neq 0\}| - 1$ .

We have

$$E_i^*W \neq 0 \quad r \leq i \leq r + d \quad (5.1)$$

$$= 0 \quad 0 \leq i < r \text{ or } r + d < i \leq d(x). \quad (5.2)$$

Claim:  $E_i^*AE_j^*W \neq 0$  if  $|i - j| = 1$  for  $r \leq i, j \leq r + d$ . (See Lemma 4.1.)

Suppose  $E_{j+1}^*AE_j^*W = 0$  for some  $j$  with  $r \leq j < r + d$ . Observe that

$$\tilde{W} = E_r^*W + \cdots + E_j^*W$$

is  $T$ -invariant with

$$0 \subsetneq \tilde{W} \subsetneq W.$$

Because  $A\tilde{W} \subseteq \tilde{W}$  since  $AE_j^*W \subseteq E_{j-1}^*W + E_j^*W$ ,

$$E_k^*\tilde{W} \subseteq \tilde{W} \quad \text{for all } k,$$

we have  $T\tilde{W} \subseteq \tilde{W}$ .

Suppose  $E_{i-1}^*AE_i^*W = 0$  for some  $i$  with  $r \leq i < r + d$ .

Similarly,

$$\tilde{W} = E_i^*W + \cdots + E_{r+d}^*W$$

is a  $T$ -module with  $0 \subsetneq \tilde{W} \subsetneq W$ .

**Definition 5.1.** Let  $\Gamma$ ,  $E_i^*$ , and  $T$  be as above. Irreducible  $T$ -modules  $W$  and  $W'$  are isomorphic whenever there is an isomorphism  $\sigma : W \rightarrow W'$  of vector spaces such that  $a\sigma = \sigma a$  for all  $a \in T$ .

Recall that the standard module  $V$  is an orthogonal direct sum of irreducible  $T$ -modules  $W_1 \oplus W_2 \oplus \dots$ . Given  $W$  in this list, the multiplicity of  $W$  in  $V$  is

$$|\{j \mid W_j \simeq W\}|.$$

*Remark.* It is known that the multiplicity does not depend on the decomposition.

Now assume that  $\Gamma$  is the  $D$ -cube,  $H(D, 2)$  with  $D \geq 1$ . View

$$X = \{a_1 \cdots a_D \mid a_i \in \{1, -1\}, 1 \leq i \leq D\}, \quad (5.3)$$

$$E = \{xy \mid x, y \in X, x, y \text{ differ in exactly 1 coordinate}\}. \quad (5.4)$$

Find  $T$ -modules.

Claim:  $H(D, 2)$  is bipartite with a partition  $X = X^+ \cup X^-$ , where

$$X^+ = \{a_1 \cdots a_D \in X \mid \prod a_i > 0\} \quad (5.5)$$

$$X^- = \{a_1 \cdots a_D \in X \mid \prod a_i < 0\} \quad (5.6)$$

Observe: for all  $y, z \in X$ ,

$$\partial(y, z) = i \Leftrightarrow y, z \text{ differ in exactly } i \text{ coordinates with } 0 \leq i \leq D.$$

Here, the diameter of  $H(D, 2) = D = d$  for all  $x \in X$ .

**Theorem 5.1.** Let  $\Gamma = H(D, 2)$  be as above. Fix  $x \in X$ , and write  $E_i^* = E_i^*(x)$ , and  $T = T(x)$ .

Let  $W$  be an irreducible  $T$ -module with endpoint  $r$ , and diameter  $d$  with  $0 \leq r \leq r + d \leq D$ .

(i)  $W$  has a basis  $w_0, w_1, \dots, w_d$  with  $w_i \in E_{i+r}^* W$  for  $0 \leq i \leq d$ . With respect to which the matrix representing  $A$  is

$$\begin{pmatrix} 0 & d & 0 & \cdots & 0 & 0 & 0 \\ 1 & 0 & d-1 & \cdots & 0 & 0 & 0 \\ 0 & 2 & 0 & \cdots & 0 & 0 & 0 \\ 0 & 0 & 3 & \cdots & 0 & 0 & 0 \\ \cdots & \cdots & \cdots & \ddots & \ddots & \cdots & \cdots \\ 0 & 0 & 0 & \ddots & 0 & 2 & 0 \\ 0 & 0 & 0 & \cdots & d-1 & 0 & 1 \\ 0 & 0 & 0 & \cdots & 0 & d & 0 \end{pmatrix}$$



(ii)  $d = D - 2r$ . In particular,  $0 \leq r \leq D/2$ .

(iii) Let  $W'$  denote an irreducible  $T$ -module with endpoint  $r'$ . Then  $W$  and  $W'$  are isomorphic as  $T$ -modules if and only if  $r = r'$ .

(iv) The multiplicity of the irreducible  $T$ -module with endpoint  $r$  is

$$\binom{D}{r} - \binom{D}{r-1} \quad \text{if } 1 \leq r \leq D/2,$$

and 1 if  $r = 0$ .

*Proof.* Recall that  $\Gamma$  is vertex transitive. It is a Cayley graph.

Hence without loss of generality, we may assume that  $x = \overbrace{11 \cdots 1}^D$ .

Notation: Set  $\Omega = \{1, 2, \dots, D\}$ . For every subset  $S \subseteq \Omega$ , let

$$\hat{S} = a_1 \cdots a_d \in X \quad a_i = \begin{cases} -1 & \text{if } i \in S \\ 1 & \text{if } i \notin S. \end{cases}$$

In particular,  $\hat{\emptyset} = x$  and

$$|S| = i \Leftrightarrow \partial(x, \hat{S}) = i \Leftrightarrow \hat{S} \in E_i^* V.$$

For all  $S, T \subseteq \Omega$ , we say  $S$  covers  $T$  if and only if  $S \supseteq T$  and  $|S| = |T| + 1$ .

Observe that  $\hat{S}, \hat{T}$  are adjacent in  $\Gamma$  if and only if either  $T$  covers  $S$  or  $S$  covers  $T$ .

Define the ‘raising matrix’

$$R = \sum_{i=0}^D E_{i+1}^* A E_i^*.$$

Observe that

$$R E_i^* V \subseteq E_{i+1}^* V \quad \text{for } 0 \leq i \leq D, \quad \text{and } E_{D+1}^* V = 0.$$

Indeed for any  $S \subseteq \Omega$  with  $|S| = i$ ,

$$R \hat{S} = R E_i^* \hat{S} \tag{5.7}$$

$$= E_{i+1}^* A \hat{S} \tag{5.8}$$

$$= \sum_{T_1 \subseteq \Omega, S \text{ covers } T_1} E_{i+1}^* \hat{T}_1 + \sum_{T \subseteq \Omega, T \text{ covers } S} E_{i+1}^* \hat{T} \tag{5.9}$$

$$= \sum_{T \subseteq \Omega, T \text{ covers } S} E_{i+1}^* \hat{T}. \tag{5.10}$$

Define the ‘lowering matrix’

$$L = \sum_{i=0}^D E_{i-1}^* A E_i^*.$$

Observe that

$$L E_i^* V \subseteq E_{i-1}^* V \text{ for } 0 \leq i \leq D, \text{ and } E_{-1}^* V = 0.$$

Indeed for any  $S \subseteq \Omega$ ,

$$L \hat{S} = \sum_{T \subseteq \Omega, S \text{ covers } T} \hat{T}.$$

Observe that  $A = L + R$ .

For convenience, set

$$A^* = \sum_{i=0}^D (D - 2i) E_i^*.$$

Claim: The following hold.

- (a)  $LR - RL = A^*$ .
- (b)  $A^*L - LA^* = 2L$ .
- (c)  $A^*R - RA^* = -2R$ .

In particular  $\text{Span}(R, L, A^*)$  is a ‘representation of Lie algebra  $\mathfrak{sl}_2(\mathbb{C})$ .

*Remark* (Lie Algebra  $\mathfrak{sl}_2(\mathbb{C})$ ).

$$\mathfrak{sl}_2(\mathbb{C}) = \{X \mid \text{Mat}(\mathbb{C}) \mid \text{tr}(X) = 0\}.$$

For  $X, Y \in \mathfrak{sl}_2(\mathbb{C})$ , define a binary operation  $[X, Y] = XY - YX$ .

$$A^* \sim \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad L \sim \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \quad R \sim \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

Then these satisfy the relations (a) - (c) above.

*Proof of Claim.* Apply both sides to  $\hat{S}$  ( $S \subseteq \Omega$ ). Say  $|S| = i$ .

*Proof of (a):*

$$(LR - RL)\hat{S} = L \left( \sum_{\substack{T \subseteq \Omega, T \text{ covers } S \\ (D-i \text{ of them})}} \hat{T} \right) - R \left( \sum_{\substack{U \subseteq \Omega, S \text{ covers } U \\ (i \text{ of them})}} \hat{T} \right) \quad (5.11)$$

$$= (D - i)\hat{S} + \sum_{V \subseteq \Omega, |V|=i, |S \cap V|=i-1} \hat{V} - \left( i\hat{S} + \sum_{V \subseteq \Omega, |V|=i, |S \cap V|=i-1} \hat{V} \right) \quad (5.12)$$

$$= (D - 2i)\hat{S} \quad (5.13)$$

$$= A^*\hat{S}. \quad (5.14)$$

*Proof of (b):*

$$(A^*L - LA^*)\hat{S} = (D - 2(i - 1))L\hat{S} - (D - 2i)L\hat{S} \quad (\text{since } L\hat{S} \in E_{i-1}^*V) \quad (5.15)$$

$$= 2L\hat{S}. \quad (5.16)$$

*Proof of (c):*

$$(A^*R - RA^*)\hat{S} = (D - 2(i + 1))R\hat{S} - (D - 2i)R\hat{S} \quad (\text{since } R\hat{S} \in E_{i+1}^*V) \quad (5.17)$$

$$= 2R\hat{S}. \quad (5.18)$$

Let  $W$  be an irreducible  $T$ -module with endpoint  $r$  and diameter  $d$  ( $0 \leq r \leq r + d \leq D$ ).

*Proof of (i) and (ii):*

Pick  $0 \neq w \in E_r^*W$ .

Claim:  $LRw = (D - 2r)w$ .

*Pf.*

$$LRw = (A^* + RL)w \quad (\text{by Claim (a)}) \quad (5.19)$$

$$= A^*w \quad (Lw \in E_{r-1}^*W = 0) \quad (5.20)$$

$$(D - 2r)w. \quad (5.21)$$

Define

$$w_i = \frac{1}{i!}R^i w \in E_{r+i}^*W \quad (0 \leq i \leq d).$$

Then,

$$Rw_i = (i + 1)w_{i+1} \quad (0 \leq i \leq d) \quad (5.22)$$

$$Rw_d = 0 \quad (\text{by definition of } d) \quad (5.23)$$

Claim:  $Lw_0 = 0$  and

$$Lw_i = (D - 2r - i + 1)w_{i-1} \quad (1 \leq i \leq d).$$

*Pf.* We prove by induction on  $i$ . The case  $i = 0$  is trivial, and the case  $i = 1$  follows from above claim. Let  $i \geq 2$ ,

$$Lw_i = \frac{1}{i}LRw_{i-1} = \frac{1}{i}(A^* + RL)w_{i-1} \quad (\text{by Claim (a)}) \quad (5.24)$$

$$(\text{by induction hypothesis}) \quad (5.25)$$

$$= \frac{1}{i}((D - 2(r + i - 1))w_{i-1} + (D - 2r - (i - 1) + 1)Rw_{i-2}) \quad (Rw_{i-2} = (i - 1)w_{i-1}) \quad (5.26)$$

$$= \frac{1}{i}i(D - 2r - i + 1)w_{i-1} \quad (5.27)$$

$$= (D - 2r - i + 1)w_{i-1}. \quad (5.28)$$

Claim:  $w_0, \dots, w_d$  is a basis for  $W$ .

*Pf.* Let  $W' = \text{Span}\{w_0, \dots, w_d\}$ . Then  $W'$  is  $R$  and  $L$  invariant. So it is  $A = R + L$  invariant.

Also it is  $E_i^*$ -invariant for every  $i$ .

Hence  $W'$  is a  $T$ -module.

Since  $W$  is irreducible,  $W' = W$ .

As  $w_i$ 's are orthogonal, they are linearly independent. Note that  $w_i \neq 0$  by the definition of  $d$  and Lemma 4.1 (iv).

Claim:  $d = D - 2r$ .

*Pf.* By (a),

$$0 = (LR - RL - A^*)w_d \quad (5.29)$$

$$= 0 - (D - 2r - d + 1)Rw_{d-1} - (D - 2(r + d))w_d \quad (5.30)$$

$$= -d(D - 2r - d + 1)w_d - (D - 2(r + d))w_d \quad (5.31)$$

$$= (-dD + 2rd + d^2 - d - D + 2r + 2d)w_d \quad (5.32)$$

$$= (d^2 + (2r - D + 1)d + 2r - D)w_d \quad (5.33)$$

$$= (d + 2r - D)(d + 1)w_d. \quad (5.34)$$

Hence  $d = D - 2r$ .

Therefore, with respect to a basis  $w_0, w_1, \dots, w_d$ ,  $A = L + R$ ,  $w_{-1} = w_{d+1} = 0$ ,

$$Lw_i = (d - i + 1)w_{i-1}, \quad Rw_i = (i + 1)w_{i+1}.$$

$$L = \begin{pmatrix} 0 & d & 0 & \dots & 0 & 0 \\ 0 & 0 & d-1 & \dots & 0 & 0 \\ \vdots & \vdots & \ddots & \ddots & \vdots & \vdots \\ \vdots & \vdots & \dots & \dots & 0 & 1 \\ 0 & 0 & 0 & \dots & 0 & 0 \end{pmatrix}, \quad R = \begin{pmatrix} 0 & 0 & 0 & \dots & 0 & 0 \\ 1 & 0 & 0 & \dots & 0 & 0 \\ 0 & 2 & 0 & \dots & \vdots & \vdots \\ \vdots & \vdots & \ddots & \ddots & 0 & 1 \\ 0 & 0 & 0 & \dots & d & 0 \end{pmatrix}.$$

This completes the proof of (i) and (ii).  $\square$

## Chapter 6

# $T$ -Modules of $H(D, 2)$ , II

Monday, February 1, 1993

*Proof of Theorem 5.1 Continued.*

(iii) Let  $r = r'$ ,

$w_0, \dots, w_d$ : a basis for  $W$  with  $w_i \in E_i^* W$ , and

$w'_0, \dots, w'_d$ : a basis for  $W'$  with  $w'_i \in E_i^* W'$ .

Then  $d = D - 2r = D - 2r' = d'$ , and

$$\sigma : W \rightarrow W' \quad (w_i \mapsto w'_i)$$

is an isomorphism of  $T$ -modules by (i).

If  $r \neq r'$ , then

$$d = D - 2r \neq D - 2r' = d',$$

hence,  $\dim W \neq \dim W'$ .

(iv) Let  $W_i$  be an irreducible  $T$ -module with endpoint  $i$ . Then

$$\dim E_r^* V = \binom{D}{r} = \sum_{i=0}^r \text{mult}(W_i).$$

Hence, we have that

$$\text{mult}(W_r) = \binom{D}{r} - \binom{D}{r-1}$$

by induction on  $r$ .

□

**Theorem 6.1.** *Let  $\Gamma = H(D, 2)$  with  $D \geq 1$ . Fix a vertex  $x \in X$  and write*

$$E_i^* \equiv E_i^*(x), \quad T = T(x), \quad \text{and} \quad A^* \equiv \sum_{i=0}^D (D - 2i) E_i^*.$$

*Let  $W$  be an irreducible  $T$ -module with endpoint  $r$  with  $0 \leq r \leq D/2$ . Then,*

*(i)  $W$  has a basis*

$$w_0^*, w_1^*, \dots, w_d^* \quad (d = D - 2r), \quad \text{such that} \quad w_i^* \in E_{i+r} W \quad (0 \leq i \leq d)$$

*with respect to which the matrix corresponding to  $A^*$  is*

$$\begin{pmatrix} 0 & d & 0 & & & & & \\ 1 & 0 & d-1 & & & & & \\ 0 & 2 & 0 & & & & & \\ & & \ddots & \ddots & \ddots & & & \\ & & & & 0 & 2 & 0 & \\ & & & & d-1 & 0 & 1 & \\ & & & & 0 & d & 0 & \end{pmatrix}.$$

*In particular, / (ii)  $E_i A^* E_j = 0$  if  $|i - j| \neq 1$  for  $0 \leq i, j \leq D$ .*

*Proof.* We use the notation,

$$[\alpha, \beta] = \alpha\beta - \beta\alpha \quad (= -[\beta, \alpha]).$$

Recall that

$$(a) \quad [L, R] = A^*,$$

$$(b) \quad [A^*, L] = wL,$$

$$(c) \quad [A^*, R] = -2R,$$

and  $A = L + R$ .

Write (a) – (c) in terms of  $A$  and  $A^*$ , we have,

$$[A, A^*] = [L, A^*] + [R, A^*] = 2(R - L).$$

$$\begin{cases} R + L &= A \\ R - L &= [A, A^*]/2. \end{cases}.$$

Hence,

$$R = \frac{1}{4}(2A + [A, A^*]) \quad \text{and} \tag{6.1}$$

$$L = \frac{1}{4}(2A - [A, A^*]). \tag{6.2}$$

Now (a), (b) become

$$A^2A^* - 2AA^*A + A^*A^2 - 4A^* = 0 \quad (6.3)$$

$$A^{*2}A - 2A^*AA^* + AA^{*2} - 4A = 0 \quad (6.4)$$

*Pf.* By (b),

$$2A - AA^* + A^*A = 4L \quad (6.5)$$

$$= 2[A^*, L] \quad (6.6)$$

$$= A^* \frac{2A - [A, A^*]}{2} - \frac{2A - [A, A^*]}{2} A^* \quad (6.7)$$

So we have ((6.4))

$$A^{*2}A - 2A^*AA^* + AA^{*2} - 4A = 0.$$

By (a),

$$-16A^* = [2A + [A, A^*], 2A - [A, A^*]] \quad (6.8)$$

$$(2A + [A, A^*])(2A - [A, A^*]) - (2A - [A, A^*])(2A + [A, A^*]) \quad (6.9)$$

$$= [4A^2 - 2A[A, A^*] + [A, A^*](2A) - [A, A^*]^2 \quad (6.10)$$

$$- 4A^2 - 2A[A, A^*] + [A, A^*](2A) + [A, A^*]^2 \quad (6.11)$$

$$= -4A^2A^* + 4AA^*A + 4AA^*A - 4A^*A^2. \quad (6.12)$$

So,

$$A^2A^* - 2AA^*A + A^*A^2 - 4A^* = 0.$$

Claim:  $E_i^*A^*E_j = 0$  if  $|i - j| \neq 1$  for  $0 \leq i, j \leq D$ .

*Pf.* We have,

$$0 = E_i(A^2A^* - 2AA^*A + A^*A^2 - 4A^*)E_j \quad (6.13)$$

$$= E_iA^*E_j(\theta_i^2 - 2\theta_i\theta_j + \theta_j^2 - 4) \quad (6.14)$$

$$(AE_j = \theta_jE_j, E_iA = (AE_j)^\top = (\theta_iE_i)^\top = \theta_iE_i) \quad (6.15)$$

$$= E_iA^*E_j(\theta_i - \theta_j - 2)(\theta_i - \theta_j + 2) \quad (6.16)$$

$$= E_iA^*E_j(D - 2i - (D - 2j) - 2)(D - 2i - (D - 2j) + 2) \quad (6.17)$$

$$(\theta_k = D - 2k) \quad (6.18)$$

$$= E_iA^*E_j \cdot 4(i - j + 1)(i - j - 1) \quad (6.19)$$

and  $i - j + 1 \neq 0$ ,  $i - j - 1 \neq 0$ . Hence,  $E_i^*A^*E_j = 0$ .

Now define “dual raising matrix”,

$$R^* = \sum_{i=0}^D E_{i+1}A^*E_i.$$

So,

$$R^* E_i V \subseteq E_{i+1} V, \quad (0 \leq i \leq D, E_{D+1} V = 0).$$

Define “dual lowering matrix”

$$L^* = \sum_{i=0}^D E_{i-1} A^* E_i.$$

Then

$$L^* E_i V \subseteq E_{i-1} V \quad (0 \leq i \leq D, E_{-1} V = 0).$$

Observe that

$$A^* = \left( \sum_{i=0}^D E_i \right) A^* \left( \sum_{j=0}^D E_j \right) = L^* + R^*$$

by Claim 1.

Claim 2. We have

$$(a) [L^*, R^*] = A,$$

$$(b) [A, L^*] = 2L^*,$$

$$(c) [A, R^*] = -2R^*.$$

*Pf.* (b)

$$AL^* - L^* A = \sum_{i=0}^D (AE_{i-1} A^* E_i - E_{i-1} A^* E_i A) \quad (6.20)$$

$$= \sum_{i=0}^D E_{i-1} A^* E_i (\theta_{i-1} - \theta_i) \quad (6.21)$$

$$(\theta_k = D - 2k, \quad \theta_{i-1} - \theta_i = 2I - 2(i-1) = 2) \quad (6.22)$$

$$= 2L^*. \quad (6.23)$$

(c) Similar.

*Remark.*

$$AR^* - R^* A = \sum_{i=0}^D (AE_{i+1} A^* E_i - E_{i+1} A^* E_i A) \quad (6.24)$$

$$= \sum_{i=0}^D E_{i+1} A^* E_i (\theta_{i+1} - \theta_i) \quad (6.25)$$

$$= 2R^*. \quad (6.26)$$

(a) We have, by (b), (c)

$$[A, A^*] = [A, L^*] + [A, R^*] = 2(L^* - R^*). \quad (6.27)$$



Since  $A^* = L^* + R^*$ ,

$$R^* = \frac{2A^* + [A^*, A]}{4}, \quad L^* = \frac{2A^* - [A^*, A]}{4}.$$

Now (a) is seen to be equivalent to ((6.4)) upon evaluation. This proves Claim 2.

*Remark.*

$$[L^*, R^*] = \frac{1}{16}((2A^* - [A^*, A])(2A^* + [A^*, A]) - (2A^* + [A^*, A])(2A^* - [A^*, A])) \quad (6.28)$$

$$= \frac{1}{16}(4A^{*2} + 2A^*[A^*, A] - [A^*, A]2A^* - [A^*, A]^2 - 4A^{*2} + 2A^*[A^*, A] - [A^*, A]2A^* + [A^*, A]^2) \quad (6.29)$$

$$= \frac{1}{4}(A^{*2}A - 2A^*AA^* + AA^{*2}) \quad (6.30)$$

$$= A, \quad (6.31)$$

by ((6.4)).

Now apply same argument as for ((6.3)), ((6.4)) of Theorem 5.1 and observe  $A^*$  has  $D + 1$  distinct eigenvalues. So,

$$A^* = \sum_{i=0}^D (D - 2i)E_i^*$$

generates

$$M^* = \text{Span}(E_0^*, \dots, E_D^*).$$

Hence,  $E_0, \dots, E_D$ ,  $A^*$  generates  $T$ .

Take an irreducible  $T$ -module  $W$  with endpoint  $r$  with  $0 \leq r \leq D/2$ . Set  $t = \min\{i \mid E_i W\}$ .

Pick  $0 \neq w_0^* \in E_t W$ . Set

$$w_i^* = \frac{1}{i!} R^{*i} w_0^* \in E_{t+i} W \quad \text{for all } i.$$

Then,

$$R^* w_i^* = (i + 1) w_{i+1}^* \quad \text{for all } i.$$

By (a), we get by induction,  $L^* w_i^* = (D - 2t - i + 1) w_{i-1}^*$ ,

$$L^* w_i^* = \frac{1}{i} L^* R^* w_{i-1}^* \quad (6.32)$$

$$= \frac{1}{i} (A + R^* L^*) w_{i-1}^* \quad (6.33)$$

$$= \frac{1}{i} ((D - 2(t + i - 1)) w_{i-1}^* + (i - 1)(D - 2t - i + 2) w_{i-1}^*) \quad (6.34)$$

$$= (D - 2t - i + 1) w_{i-1}^*. \quad (6.35)$$

So  $\text{Span}(w_0^*, w_1^*, \dots)$  is  $L^*$ ,  $R^*$ ,  $A^*$ -invariant. Hence,  $W = (\text{Span})(w_0^*, w_1^*, \dots, w_d^*)$ ,  $w_0^*, w_1^*, \dots, w_d^* \neq 0$ ,  $w_i^* = 0$  for every  $i > d$  by dimension.

Thus  $d = D - 2t$ .

*Pf.*

$$(D - 2(t + d))w_d^* = Aw_d^* \quad (6.36)$$

$$= (L^*R^* - R^*L^*)w_d^* \quad (6.37)$$

$$= -(D - 2t - d + 1)R^*w_{d-1}^* \quad (6.38)$$

$$= -(D - 2t - d + 1)dw_d^*. \quad (6.39)$$

Hence,

$$0 = d^2 + (2t - D - 1 + 2)d - (D - 2t) = (d - D + 2t)(d + 1)$$

So  $d = D - 2t$ . □

**Definition 6.1.** For any graph  $\Gamma = (X, E)$ , pick a vertex  $x \in X$ , and set  $E_i^* \equiv E_i^*(x)$  and  $T \equiv T(x)$ .

- (i) An irreducible  $T$ -module  $W$  is thin if  $\dim E_i^*W \leq 1$  for every  $i$ .
- (ii)  $\Gamma$  is thin with respect to  $x$ , if every irreducible  $T(x)$ -module is thin,
- (iii) An irreducible  $T$ -module  $W$  is dual thin if  $\dim E_iW \leq 1$  for every  $i$ .
- (iv)  $\Gamma$  is dual thin with respect to  $x$ , if every irreducible  $T(x)$ -module is dual thin.

Observe:  $H(D, 2)$  is thin, dual thin with respect to each  $x \in X$ .

**Definition 6.2.** With above notation, write  $D \equiv D(x)$ .

- (i) An ordering  $E_0, E_1, \dots, E_R$  of primitive idempotents of  $\Gamma$  is restricted if  $E_0$  corresponds to the maximal eigenvalue.

Fix a restricted ordering,

- (ii)  $\Gamma$  is  $Q$ -polynomial with respect to  $x$ , above ordering if there exists  $A^* \equiv A^*(x)$  such that

$$(a) E_0^*V, \dots, E_D^*V \text{ are the maximal eigenspaces for } A^*.$$

$$(b) E_iA^*E_j = 0 \text{ if } |i - j| > 1 \text{ for } 0 \leq i, j \leq R.$$

Observe  $H(D, 2)$  is  $Q$ -polynomial with respect to the natural ordering of the idempotents and every vertex.

**Program.** Study graphs that are thin and  $Q$ -polynomial with respect to each vertex.

(In fact, thin with respect to  $x$  implies dual thin with respect to  $x$ .)

Get a situation like  $H(D, 2)$ , where  $T$  is generated by  $A, A^*$ . Except  $\mathfrak{sl}_2(\mathbb{C})$  is replaced by a quantum Lie algebra.



## Chapter 7

# The Johnson Graph $J(D, N)$


Wednesday, February 3, 1993

**Definition 7.1.** The Johnson graph,  $\Gamma = J(D, N)$  ( $1 \leq D \leq N - 1$ ) satisfies

$$X = \{S \mid S \subset \Omega, |S| = D\} \quad \text{where } \Omega = \{1, 2, \dots, N\} \quad (7.1)$$

$$E = \{ST \mid S, T \in X, |S \cap T| = D - 1\}. \quad (7.2)$$

**Example 7.1.**  $J(2, 4)$



**Note 1.** The symmetric group  $S_N$  acts on  $\Omega$ .  $S_N \subseteq \text{Aut}(\Gamma)$  acts vertex transitively on  $\Gamma$ .

**Note 2.**  $\Gamma = J(D, N)$  is isomorphic to  $\Gamma' = J(N - D, N)$ .

$$\Gamma = (X, E) \quad \longrightarrow \quad \Gamma' = (X', E') \quad (7.3)$$

$$X \ni S \quad \longmapsto \quad \bar{S} = \Omega \quad S \in X' \quad (7.4)$$

This correspondence induces an isomorphism of graphs.

*Pf.*

$$ST \in E \Leftrightarrow |S \cap T| = D - 1 \quad (7.5)$$

$$\Leftrightarrow |\Omega - (S \cup T)| = N - D - 1 \quad (7.6)$$

$$\Leftrightarrow |\bar{S} \cap \bar{T}| = N - D - 1 \quad (7.7)$$

$$\Leftrightarrow \bar{S}\bar{T} \in E' \quad (7.8)$$

Hence, without loss of generality, assume

$$D \leq N/2 \quad \text{for} \quad J(D, N).$$

We still need the eigenvalues of  $J(D, N)$  for certain problem later in the course. We can get these eigenvalues from our study of  $H(D, 2)$ .

**Lemma 7.1.** *The eigenvalues for  $J(D, N)$  with  $1 \leq D \leq N/2$  are given by*

$$\theta_i = (N - D - i)(D - i) - i \quad (0 \leq i \leq D), \quad (7.9)$$

$$m_i = \binom{N}{i} - \binom{N}{i-1}. \quad (7.10)$$

*Proof.* Let

$$\Gamma_J \equiv J(D, N) = (X_J, E_J) \quad (7.11)$$

$$\Gamma_H \equiv H(N, 2) = (X_H, E_H). \quad (7.12)$$

Set  $x \equiv 11 \cdots 1 \in X_H$ .

Define  $\tilde{\Gamma} \equiv (\tilde{X}, \tilde{E})$ , where

$$\tilde{X} = \{y \in X_H \mid \partial_H(x, y) = D\} \quad \partial_H : \text{distance in } \Gamma_H \quad (7.13)$$

$$\tilde{E} = \{yz \in X_H \mid \partial_H(y, z) = 2\}. \quad (7.14)$$

Observe

$$X_J \quad \rightarrow \quad \tilde{X} \quad (7.15)$$

$$S \quad \mapsto \quad \hat{S}, \quad (7.16)$$

where

$$\hat{S} = a_1 \cdots a_N, \quad a_i = \begin{cases} -1 & \text{if } i \in S \\ 1 & \text{if } i \notin S \end{cases}$$

induces an isomorphism of graphs  $\Gamma_J \rightarrow \tilde{\Gamma}$ .

*Pf.*

$$ST \in E_J \Leftrightarrow |S \cap T| = D - 1 \quad (7.17)$$

$$\Leftrightarrow \partial_H(\hat{S}, \hat{T}) = 2 \quad (7.18)$$

$$\Leftrightarrow (\hat{S}, \hat{T}) \in \tilde{E}. \quad (7.19)$$

Identify,  $\Gamma_J$  with  $\tilde{\Gamma}$ . Then the standard module  $V_J$  of  $\Gamma_J$  becomes  $\tilde{V} = E_D^* V_H$ , where  $V_H$  is the standard module of  $\Gamma_H$ , and  $E_D^* \equiv E_D^*(x)$ .

Let  $R$  be the raising matrix with respect to  $x$  in  $\Gamma_H$ , and

let  $L$  be the lowering matrix with respect to  $x$  in  $\Gamma_H$ .

Recall

$$(RL - DE_D^*)|_{\tilde{V}}$$

is the adjacency map in  $\tilde{\Gamma}$ .

To find eigenvalues of  $\tilde{A}$ , pick any irreducible  $T(x)$ -module  $W$  with the endpoint  $r \leq D$ . Then by Theorem 5.1

$$\text{diam}(W) = N - 2r.$$

Let  $w_0, w_1, \dots, w_{N-2r}$  denote a basis for  $W$  as in Theorem 5.1. Then,

$$w_{D-r} \in E_D^* W \subseteq \tilde{V}.$$

Observe:

$$\tilde{A}w_{D-r} = RLw_{D-r} - DE_D^* w_{D-r} \quad (7.20)$$

$$= R(N - 2r - D + r + 1)w_{D-r-1} - Dw_{D-r} \quad (7.21)$$

$$= ((N - D - r + 1)(D - r) - D)w_{D-r}. \quad (7.22)$$

Note that this is valid for  $D = r$  as well.

Hence,

$$\tilde{A}w_{D-r} = ((N - D - r)(D - r) - r)w_{D-r}.$$

Let

$$V_H = \sum W \quad (\text{direct sum of irreducible } T(x)\text{-modules.})$$

Then,

$$V_J = E_D^* V_H \quad (7.23)$$

$$= \sum_{W: r(W) \leq D} E_D^* W \quad (7.24)$$

$$= \text{a direct sum of 1 dimensional eigenspaces for } \tilde{A}. \quad (7.25)$$

The eigenspace for eigenvalue

$$(N - D - r)(D - r) - r \quad (\text{monotonously decreasing with respect to } r)$$

appears with multiplicity

$$\binom{N}{r} - \binom{N}{r-1}$$

in this sum by Theorem 5.1 (iv).  $\square$

**Theorem 7.1.** *Let  $\Gamma = (X, E)$  be any graph. For a fixed vertex  $x \in X$ , let*

$$E_i^* \equiv E_i^*(x), \quad T \equiv T(x), \quad D \equiv D(x), \quad \text{and } K = \mathbb{C}.$$

*Then we have the following implications of conditions:*

$$TH \Leftrightarrow C \Leftarrow S \Leftarrow G,$$

*where*

*(TH)  $\Gamma$  is thinn with respect to  $x$ .*

*(C)  $E_i^*TE_i^*$  is commutative for every  $i$ ,  $(0 \leq i \leq D)$ .*

*(S)  $E_i^*TE_i^*$  is symmetric for every  $i$ ,  $(0 \leq i \leq D)$ .*

*(G) For every  $y, z \in X$  with  $\partial(x, y) = \partial(x, z)$ , there exists  $g \in \text{Aut}(\Gamma)$  such that*

$$gx = x, \quad gy = z, \quad gz = y.$$

*Proof.*

$(TH) \Rightarrow (C)$

Fix  $i$  with  $0 \leq i \leq D$ . Let

$V = \sum W$ . The standard module written as a direct sum of irreducible  $T$ -modules.

Then,

$$E_i^*V = \sum E_i^*W. \text{ The direct sum of 1-dimensional } E_i^*TE_i^*\text{-modules.}$$

Since  $\dim E_i^*W = 1$ , for  $a, b \in E_i^*TE_i^*$ ,  $ab - ba|_{E_i^*W} = 0$ . Hence  $ab - ba = 0$ .

$(C) \Rightarrow (TH)$

Suppose  $\dim E_i^*W \geq 2$  for some irreducible  $T$ -module  $W$  with some  $i$  with  $1 \leq i \leq D$ .

Claim 1.  $E_i^*W$  is an irreducible  $E_i^*TE_i^*$ -module.

*Proof of Claim 1.* Suppose

$$0 \subsetneq U \subsetneq E_i^*W,$$



where  $U$  is an  $E_i^*TE_i^*$ -module. Then by the irreducibility,

$$TU = W.$$

So,

$$U \supseteq E_i^*TE_i^*U = E_i^*TU = E_i^*W.$$

This is a contradiction.

Claim 2. Each irreducible  $S = E_i^*TE_i^*$ -module  $U$  has dimension 1. In particular,  $\Gamma$  is thin with respect to  $x$ .

*Proof of Claim 2.* Pick

$$0 \neq a \in E_i^*TE_i^*.$$

Since  $\mathbb{C}$  is algebraically closed,  $a$  has an eigenvector  $w \in U$  with eigenvalue  $\theta$ . Then,

$$(a - \theta I)U = (a - \theta I)Sw \tag{7.26}$$

$$= S(a - \theta I)w \tag{7.27}$$

$$= 0. \tag{7.28}$$

Hence,

$$a|_U = \theta I|_U \quad \text{for all } a \in S.$$

Thus each 1 dimensional subspace of  $U$  is an  $S$ -module. We have

$$\dim U = 1.$$

By Claim 1 and Claim 2, we have (TH).

□

*Remark.* Claim 1 shows the following: *If  $W$  is an irreducible  $T$ -module, then  $E_i^*W$  is either 0 or an irreducible  $E_i^*TE_i^*$ -module.*



## Chapter 8

# Thin Graphs

**Friday, February 5, 1993**

*Proof of Theorem 7.1 continued.*

(S)  $\Rightarrow$  (C)

Fix  $i$  and pick  $a, b \in E_i^* T E_i^*$ .

Since  $a$ ,  $b$  and  $ab$  are symmetric,

$$ab = (ab)^\top = b^\top a^\top = ba.$$

Hence  $E_i^* T E_i^*$  is commutative.

(G)  $\Rightarrow$  (S)

Fix  $i$  and pick  $a \in E_i^* T E_i^*$ . Pick vertices  $y, z \in X$ .

We want to show that

$$a_{yz} = a_{zy}.$$

We may assume that

$$\partial(x, y) = \partial(x, z) = i,$$

otherwise

$$a_{yz} = a_{zy} = 0.$$

By our assumption, there exists  $g \in G$  such that

$$g(y) = z, \quad g(z) = y, \quad g(x) = x.$$

Let  $\hat{g}$  denote the permutation matrix representing  $g$ , i.e.,

$$\hat{g}\hat{y} = \widehat{g(y)} \quad \text{for all } y \in X, \quad \hat{g} = \begin{pmatrix} 0 \\ \vdots \\ 1 \\ \vdots \\ 0 \end{pmatrix} \leftarrow y.$$

If  $g \in \text{Aut}(\Gamma)$ , then

$$\hat{g}A = A\hat{g} \quad (\text{Exercise}).$$

Also, we have

$$\hat{g}E_j^* = E_j^*\hat{g} \quad (0 \leq j \leq D),$$

since

$$\partial(x, y) = \partial(g(x), g(y)) = \partial(x, g(y)).$$

Hence,  $\hat{g}$  commutes with each element of  $T$ . We have

$$a_{yz} = (\hat{g}^{-1}a\hat{g})_{yz}, \quad (\hat{g})_{yz} = \begin{cases} 1 & g(z) = y \\ 0 & \text{else.} \end{cases} \quad (8.1)$$

$$= \sum_{y', z'} (\hat{g}^{-1})_{yy'} a_{y'z'} \hat{g}_{z'z} \quad (8.2)$$

$$(\text{zero except for } g^{-1}(y') = y, g(z) = z'.) \quad (8.3)$$

$$= a_{g(y)g(z)} \quad (8.4)$$

$$= a_{zy}. \quad (8.5)$$

This proves Theorem 7.1. □

**Open Problem:** Find all the graphs that satisfy the condition (G) for every vertex  $x$ .

$H(N, 2)$  is one example, because

$$\text{Aut}\Gamma_{1\dots 1} \simeq S_\Omega, \quad x = (1\dots 1), \quad \Gamma_i(x) = \{\hat{S} \mid |S| = i\}.$$

Property (G) is clearly related to the distance-transitive property.

**Definition 8.1.** Let  $\Gamma = (X, E)$  be any graph.  $\Gamma$  with  $G \subseteq \text{Aut}(\Gamma)$  is said to be distance-transitive (or two-point homogeneous), whenever

$$\text{for all } x, x', y, y' \in X \text{ with } \partial(x, y) = \partial(x', y'),$$

there exists  $g \in G$  such that

$$g(x) = x', \quad g(y) = y'.$$

(This means  $G$  is as close to being doubly transitive as possible.)

**Lemma 8.1.** Suppose a graph  $\Gamma = (X, E)$  satisfies the property  $(G) = (G(x))$  for every  $x \in X$ . Then,

- (i) either
- (ia)  $\Gamma$  is vertex transitive; or
- (iia)  $\Gamma$  is bipartite ( $X = X^+ \cup X^-$ ) with  $X^+, X^-$  each an orbit of  $\text{Aut}(\Gamma)$ .
- (ii) if (ia) holds, then  $\Gamma$  is distance-transitive.

*Proof.* (i) Claim. Suppose  $y, z \in X$  are connected by a path of even length. Then  $y, z$  are in the same orbit of  $\text{Aut}(\Gamma)$ .

*Pf of Claim.* It suffices to assume that the path has length 2,  $y \sim w \sim z$ .

Now  $\partial(y, w) = \partial(w, z) = 1$ . So there exists  $g \in \text{Aut}(\Gamma)$  such that

$$gw = w, \quad gy = z, \quad gz = y.$$

This proves Claim.

Fix  $x \in X$ . Now suppose that  $\Gamma$  is not vertex transitive, and we shall show (ib).

Observe that  $X = X^+ \cup X^-$ , where

$$X^+ = \{y \in X \mid \text{there exists a path of even length connecting } x \text{ and } y\}, \quad (8.6)$$

$$X^- = \{y \in X \mid \text{there exists a path of odd length connecting } x \text{ and } y\}. \quad (8.7)$$

Also,  $X^+$  is contained in an orbit  $O^+$  of  $\text{Aut}(\Gamma)$ , and  $X^-$  is contained in an orbit  $O^-$  of  $\text{Aut}(\Gamma)$ .

Now  $O^+ \cap O^- = \emptyset$  (else  $O^+ = O^- = X$  and vertex transitive). So,  $X = O^+$ , and  $X^- = O^-$ .

Also  $X^+ \cup X^- = X$  is a bipartition by construction.

(ii) Fix  $x, y, x', y'$  with  $\partial(x, y) = \partial(x', y')$ .

By vertex transitivity, there exists an element

$$g_1 \in G \text{ such that } g_1x = x'.$$

Observe that

$$\partial(x', y') = \partial(x, y) = \partial(g_1x, g_1y) = \partial(x', g_1y).$$

Hence, there exists an element

$$g_2 \in G \text{ such that } g_1x' = x', g_2y' = g_1y', g_2g_1y = y'$$

by  $(G(x'))$  property.

Set  $g = g_2g_1$ . Then

$$gx = x', gy = y'$$

by construction. □

The following graphs  $\Gamma = (X, E)$  are vertex transitive, and satisfy the property  $(G(x))$  for all  $x \in X$ .

$$J(D, N), \quad H(D, r), \quad J_q(D, N),$$

where

$H(D, r)$ :

$$X = \{a_1 \cdots a_D \mid a_i \in F, 1 \leq i \leq D\} \quad (8.8)$$

$$F : \text{ any set of cardinality } r \quad (8.9)$$

$$E = \{xy \mid y, x \in X, x \text{ and } y \text{ differ in exactly one coordinate}\}. \quad (8.10)$$

$J_q(D, N)$ :

$$X = \text{the set of all } D\text{-dimensional subspaces of } N\text{-dimensional vector space over } GF(q). \quad (8.11)$$

$$F : \text{ any set of cardinality } r \quad (8.12)$$

$$E = \{xy \mid y, x \in X, \dim(x \cap y) = D - 1\}. \quad (8.13)$$

The following graph is distance-transitive but does not satisfy  $(G(x))$  for any  $x \in G$ .

$H_q(D, N)$ :

$$X = \text{the set of all } D \times N \text{ matrices with entries in } GF(q). \quad (8.14)$$

$$E = \{xy \mid y, x \in X, \text{rank}(x - y) = 1\}. \quad (8.15)$$

*Remark.*

$$H(D, r): G = S_r \text{wr} S_D, G_x = S_{r-1} \text{wr} S_D,$$

For  $x, y \in X$  with  $\partial(x, y) = \partial(x, z) = i$ ,

$$Y = \{j \in \Omega \mid x_j \neq y_j\} \leftrightarrow Z = \{j \in \Omega \mid x_j \neq z_j\} \quad (8.16)$$

$$(y_{j_1}, \dots, y_{j_i}) \leftrightarrow (z_{\ell_1}, \dots, z_{\ell_i}) \quad (8.17)$$

$$J(D, N): G = S_N, G_x = S_D \times S_{N-D}.$$

$$X \cap Y \leftrightarrow X \cap Z \quad (8.18)$$

$$(\Omega - X) \cap Y \leftrightarrow (\Omega - X) \cap Z. \quad (8.19)$$

The following graph is distance-transitive but does not satisfy  $(G(x))$  for any  $x \in G$ .

$J_q(D, N)$ :

$$X \cap Y \leftrightarrow X \cap Z.$$

The theory of single thin irreducible  $T$ -module.

Let  $\Gamma = (X, E)$  be any graph.

$$M = \text{Bose-Mesner algebra over } K/\mathbb{C} \text{ generated by the adjacency matrix } A. \quad (8.20)$$

$$= \text{Span}(E_0, \dots, E_R). \quad (8.21)$$

$M$  acts on the standard module  $V = \mathbb{C}^{|X|}$ .

Fix  $x \in X$ , let  $D \equiv D(x)$  be the  $x$ -diameter, and  $k = k(x)$  be the valency of  $x$ .





## Chapter 9

# Thin $T$ -Module, I

**Monday, February 8, 1993**

Let  $\Gamma = (X, E)$  be any graph.

$M$ : Bose-Mesner algebra over  $K/\mathbb{C}$  generated by the adjacency matrix  $A$ .

$$M = \text{Span}(E_0, \dots, E_R).$$

$M$  acts on the standard module  $V = \mathbb{C}^{|X|}$ .

Fix  $x \in X$ , let  $D \equiv D(x)$  be the  $x$ -diameter, and  $k = k(x)$  be the valency of  $x$ .

**Definition 9.1.** Pick  $x \in X$  and write  $E_i^* \equiv E_i^*(x)$  and  $T \equiv T(x)$ .

Let  $W$  be an irreducible thin  $T$ -module with endpoint  $r$ , diameter  $d$ .

Let  $a_i = a_i(W) \in \mathbb{C}$  satisfying

$$E_{r+i}^* A E_{r+i}^* |_{E_{r+i}^* W} = a_i 1 |_{E_{r+i}^* W} \quad (0 \leq i \leq d).$$

Let  $x_i = x_i(W) \in \mathbb{C}$  satisfying

$$E_{r+i-1}^* A E_{r+i}^* A E_{r+i-1}^* |_{E_{r+i}^* W} = x_i 1 |_{E_{r+i}^* W} \quad (0 \leq i \leq d).$$

**Lemma 9.1.** *With above notation, the following hold.*

(i)  $a_i \in \mathbb{R} \quad (0 \leq i \leq d)$ .

(ii)  $x_i \in \mathbb{R}^{>0} \quad (0 \leq i \leq d)$ .

(iii) Pick  $0 \neq w_0 \in E_r^* W$ . Set  $w_i = E_{r+i}^* A^i w_0$  for all  $i$ . Then

(iiia)  $w_0, w_1, \dots, w_d$  is a basis for  $W$ ,  $w_{-1} = w_{d+1} = 0$ .

(iiib)  $A w_i = w_{i+1} + a_i w_i + x_i w_{i-1} \quad (0 \leq i \leq d)$ .

(iv) Define  $p_0, p_1, \dots, p_{d+1} \in \mathbb{R}[\lambda]$  by

$$p_0 = 1, \quad \lambda p_i = p_{i+1} + a_i p_i + x_i p_{i-1} \quad (0 \leq i \leq d), \quad p_{-1} = 0.$$

(iva)  $p_i(A)w_0 = w_i$ ,  $(0 \leq i \leq d+1)$ .

(ivb)  $p_{d+1}$  is the minimal polynomial of  $A|_W$ .

*Proof.* (i)  $a_i$  is an eigenvalue of a real symmetric matrix  $E_{r+i}^* A E_{r+i}^*$ .

(ii)  $x_i$  is an eigenvalue of a real symmetric matrix  $B^\top B$ , where

$$B = E_{r+i}^* A E_{r+i-1}^*.$$

Hence,  $x_i \in \mathbb{R}$ .

Since  $B^\top B$  is positive semidefinite,

$$x_i \geq 0.$$

*Pf.* If  $B^\top Bv = \sigma v$  for some  $\sigma \in \mathbb{R}$ ,  $v \in \mathbb{R}^m \setminus \{0\}$ , then

$$0 \leq \|Bv\|^2 = v^\top B^\top Bv = \sigma v^\top v = \sigma \|v\|^2, \quad \|v\|^2 > 0.$$

Hence,  $\sigma \geq 0$ .

Moreover,  $x_i \neq 0$  by Lemma 4.1 (iv).

(iiia) Observe

$$w_i = E_{r+i}^* A E_{r+i-1}^* w_{i-1} \quad (1 \leq i \leq d).$$

So  $w_i \neq 0$   $(1 \leq i \leq d)$  by Lemma 4.1 (iv).

Hence,

$$W = \text{Span}(w_0, \dots, w_d)$$

by Lemma 4.1. (iii).

(iiib) We have that

$$Aw_i = E_{r+i+1}^* Aw_i + E_{r+i}^* Aw_i + E_{r+i-1}^* Aw_i \quad (9.1)$$

$$= w_{i+1} + E_{r+i}^* A E_{r+i}^* w_i + E_{r+i-1}^* A E_{r+i}^* A E_{r+i-1}^* w_{i-1} \quad (9.2)$$

$$= w_{i+1} + a_i w_i + x_i w_{i-1}. \quad (9.3)$$

(iva) Clear for  $i = 0$ . Assume it is valid for  $0, \dots, i$ .

$$p_{i+1}(A)w_0 = (A - a_i I)w_i - x_i w_{i-1} = w_{i+1}.$$

(ivb) By definition,

$$p_{d+1}(A)w_0 = 0.$$

Moreover,  $p_{d+1}(A)W = 0$  because of the following.

For every  $w \in W$ , write

$$w = \sum_{i=0}^d \alpha_i w_i \quad (9.4)$$

$$= \sum_{i=0}^d \alpha_i p_i(A) w_0 \quad \text{for some } \alpha_i \in \mathbb{C} \quad (9.5)$$

$$= p(A) w_0 \quad \text{for some } p \in \mathbb{C}[\lambda] \quad (9.6)$$

Hence,

$$p_{d+1}(A)w = p_{d+1}(A)p(A)w_0 \quad (9.7)$$

$$= p(A)p_{d+1}(A)w_0 \quad (9.8)$$

$$= 0. \quad (9.9)$$

Note that  $p_{d+1}$  is the minimal polynomial.

*Pf.* Suppose  $q(A)W = 0$  for some  $0 \neq q \in \mathbb{C}[\lambda]$  with  $\deg q < \deg p_{d+1} = d + 1$ . Then,

$$q = \sum_{i=0}^d \beta_i p_i \quad \text{for some } \beta_i \in \mathbb{C}.$$

We have,

$$0 = q(A)w_0 = \sum_{i=0}^d \beta_i w_i.$$

Hence  $\beta_0 = \dots = \beta_d = 0$  by (iii). Thus  $q = 0$ , and a contradiction.  $\square$

**Corollary 9.1.** *Let  $\Gamma$ ,  $W$ ,  $r$ ,  $d$  be as above. Then*

(i)  *$W$  is dual thin, that is,*

$$\dim E_i W \leq 1 \quad (1 \leq i \leq d).$$

(ii)  $d = |\{i \mid E_i W \neq 0\}| - 1$ .

*Proof.* (i) Set as in Lemma 9.1,

$$w_i = p_i(A)w_0 \in E_{r+i}^* W.$$

Then  $w_0, w_1, \dots, w_d$  is a basis for  $W$ . We have

$$W = M w_0.$$

So,

$$E_i W = E_i M w_0 = \text{Span}(E_i w_0).$$

Thus,

$$\dim E_i W = \begin{cases} 1 & \text{if } E_i w_0 \neq 0 \\ 0 & \text{if } E_i w_0 = 0. \end{cases}$$

In particular,

$$\dim E_i^* W \leq 1.$$

(ii) Immediate as

$$\dim W = d + 1.$$

This proves the lemma.  $\square$

**Lemma 9.2.** *Given an irreducible  $T(x)$ -module  $W$  with endpoint  $r = r(W)$ , diameter  $d = d(W)$ . Write*

$$x_i = x_i(W) \ (0 \leq i \leq d), \quad w_i = p_i(A)w_0 \in E_{r+i}^* W \ (0 \leq i \leq d), \quad 0 \neq w_0 \in E_r^* W.$$

Then,

$$\frac{\|w_i\|^2}{\|w_0\|^2} = x_1 x_2 \cdots x_i \quad (1 \leq i \leq d).$$

*Proof.* It suffices to show that

$$\|w_i\|^2 = x_i \|w_{i-1}\|^2 \quad (1 \leq i \leq d).$$

Recall by Lemma 9.1 (iiib) that

$$Aw_j = w_{j+1} + a_j w_j + x_j w_{j-1} \quad (0 \leq j \leq d), \quad w_{-1} = w_{d+1} = 0.$$

Now observe,

$$\langle w_{i-1}, Aw_i \rangle = \langle w_{i-1}, w_{i+1} + a_i w_i + x_i w_{i-1} \rangle \quad (9.10)$$

$$= \overline{x_i} \|w_{i-1}\|^2 \quad (9.11)$$

$$= x_i \|w_{i-1}\|^2. \quad (9.12)$$

by Lemma 9.1 (ii). Also,

$$\langle w_{i-1}, Aw_i \rangle = \langle Aw_{i-1}, w_i \rangle \quad (\text{since } \overline{A}^\top = A) \quad (9.13)$$

$$= \langle x_i + a_{i-1} w_{i-1} + x_{i-1} w_{i-2}, w_i \rangle \quad (9.14)$$

$$= \|w_i\|^2. \quad (9.15)$$

This proves the lemma.  $\square$

**Definition 9.2.** Let  $W$  be an irreducible thin  $T(x)$  module with endpoint  $r$ ,  $E_i^* \equiv E_i^*(x)$ .

The measure  $m = m_W$  is the function

$$m : \mathbb{R} \rightarrow \mathbb{R}$$

such that

$$m(\theta) = \begin{cases} \frac{\|E_i w\|^2}{\|w\|^2} & \text{where } 0 \neq w \in E_r^* W \\ & \text{if } \theta = \theta_i \text{ is an eigenvalue for } \Gamma \\ 0 & \text{if } \theta \text{ is not an eigenvalue for } \Gamma. \end{cases}$$



## Chapter 10

# Thin $T$ -Module, II

Wednesday, February 10, 1993

Let  $\Gamma = (X, E)$  be any graph.

Fix a vertex  $x \in X$ . Let  $E_i^* \equiv E_i^*(x)$ ,  $T \equiv T(x)$ , the subconstituent algebra over  $\mathbb{C}$ , and  $V = \mathbb{C}^{|X|}$  the standard module.

**Lemma 10.1.** *With above notation, let  $W$  denote a thin irreducible  $T(x)$ -module with endpoint  $r$  and diameter  $d$ . Let*

$$a_i = a_i(W) \quad (0 \leq i \leq d) \quad (10.1)$$

$$x_i = x_i(W) \quad (1 \leq i \leq d) \quad (10.2)$$

$$p_i = p_i(W) \quad (0 \leq i \leq d+1) \quad (10.3)$$

be from Lemma 9.1, and measure  $m = m_W$ . Then,

(i)  $p_0, \dots, p_{d+1}$  are orthogonal with respect to  $m$ , i.e.,

$$\sum_{\theta \in \mathbb{R}} p_i(\theta) p_j(\theta) m(\theta) = \delta_{ij} x_1 x_2 \cdots x_i \quad (0 \leq i, j \leq d+1) \text{ with } x_{d+1} = 0.$$

$$(ia) \sum_{\theta \in \mathbb{R}} p_i(\theta)^2 m(\theta) = x_1 \cdots x_i \quad (0 \leq i \leq d).$$

$$(iia) \sum_{\theta \in \mathbb{R}} m(\theta) = 1.$$

$$(iiia) \sum_{\theta \in \mathbb{R}} p_i(\theta)^2 \theta m(\theta) = x_1 \cdots x_i a_i \quad (0 \leq i \leq d).$$

*Proof.* Pick  $0 \neq w_0 \in E_r^* W$ . Set

$$w_i = p_i(A) w_0 \in E_{r+i}^* W.$$

Since  $E_i^*W$  and  $E_j^*W$  are orthogonal if  $i \neq j$ ,

$$\delta_{ij}\|w_i\|^2 = \langle w_i, w_j \rangle \quad (10.4)$$

$$= \langle p_i(A)w_0, p_j(A)w_0 \rangle \quad (10.5)$$

$$= \left\langle p_i(A) \left( \sum_{\ell=0}^R E_\ell \right) w_0, p_j(A) \left( \sum_{\ell=0}^R E_\ell \right) w_0 \right\rangle \quad (10.6)$$

$$= \left\langle \sum_{\ell=0}^R p_i(\theta_\ell) E_\ell w_0, \sum_{\ell=0}^R p_j(\theta_\ell) E_\ell w_0 \right\rangle \quad (\text{as } AE_j = \theta_j E_j) \quad (10.7)$$

$$= \sum_{\ell=0}^R p_i(\theta_\ell) \overline{p_j(\theta_\ell)} \|E_\ell w_0\|^2 \quad (10.8)$$

$$(\text{as } p_j \in \mathbb{R}[\lambda], \quad \theta_\ell \in \mathbb{R}, \quad m(\theta_i)\|w_0\|^2 = \|E_i w_0\|^2) \quad (10.9)$$

$$= \sum_{\theta \in \mathbb{R}} p_i(\theta) p_j(\theta) m(\theta) \|w_0\|^2. \quad (10.10)$$

Now we are done by Lemma 9.2 as

$$\|w_i\|^2 = \|w_0\|^2 x_1 x_2 \dots x_i.$$

For (ia), set  $i = j$ , and for (ib), set  $i = j = 0$ .

(ii) We have

$$\langle w_i, Aw_i \rangle = \langle w_i, w_{i+1} + a_i w_i + x_i w_{i-1} \rangle \quad (10.11)$$

$$= \overline{a_i} \|w_i\|^2 \quad (10.12)$$

$$= a_i x_1 \dots x_i \|w_0\|^2, \quad (10.13)$$

as  $a_i \in \mathbb{R}$  by Lemma 9.1.

Also,

$$\langle w_i, Aw_i \rangle = \langle p_i(A)w_0, Ap_i(A)w_0 \rangle \quad (10.14)$$

$$= \left\langle p_i(A) \left( \sum_{\ell=0}^R E_\ell \right) w_0, Ap_i(A) \left( \sum_{\ell=0}^R E_\ell \right) w_0 \right\rangle \quad (\text{as in (i)}) \quad (10.15)$$

$$= \sum_{\ell=0}^D p_i(\theta_\ell)^2 \theta_\ell \|E_\ell w_0\|^2 \quad (10.16)$$

$$= \sum_{\theta \in \mathbb{R}} p_i(\theta)^2 \theta m(\theta) \|w_0\|^2. \quad (10.17)$$

Thus, we have (ii).  $\square$



**Lemma 10.2.** *With above notation, let  $W$  be a thin irreducible  $T(x)$ -module with measure  $m$ . Then  $m$  determines diameter  $d(W)$ ,*

$$a_i = a_i(W) \quad (0 \leq i \leq d) \quad (10.18)$$

$$x_i = x_i(W) \quad (1 \leq i \leq d) \quad (10.19)$$

$$p_i = p_i(W) \quad (0 \leq i \leq d+1). \quad (10.20)$$

*Proof.* Note that  $d+1$  is the number of  $\theta \in \mathbb{R}$  such that  $m(\theta) \neq 0$ . Hence  $m$  determines  $d$ .

Apply (ia), (ii) of Lemma 10.1.

$$\sum_{\theta \in \mathbb{R}} m(\theta) = 1 \quad p_0 = 1. \quad (10.21)$$

$$\sum_{\theta \in \mathbb{R}} \theta m(\theta) = a_0 \quad p_1 = \lambda - a_0 \quad (10.22)$$

$$\sum_{\theta \in \mathbb{R}} p_1(\theta)^2 m(\theta) = x_1 \quad (10.23)$$

$$\sum_{\theta \in \mathbb{R}} p_1(\theta)^2 \theta m(\theta) = x_1 a \quad \rightarrow a_1 \quad (10.24)$$

$$p_2 = (\lambda - a_1)p_1 - x_1 p_0 \quad (10.25)$$

$$\sum_{\theta \in \mathbb{R}} p_2(\theta)^2 m(\theta) = x_1 x_2 \quad \rightarrow x_2 \quad (10.26)$$

$$\sum_{\theta \in \mathbb{R}} p_2(\theta)^2 \theta m(\theta) = x_1 x_2 a_2 \quad \rightarrow a_2 \quad (10.27)$$

$$p_3 = (\lambda - a_2)p_2 - x_2 p_1 \quad (10.28)$$

$$\vdots \quad (10.29)$$

$$\sum_{\theta \in \mathbb{R}} p_d(\theta)^2 m(\theta) = x_1 x_2 \cdots x_d \quad \rightarrow x_d \quad (10.30)$$

$$\sum_{\theta \in \mathbb{R}} p_d(\theta)^2 \theta m(\theta) = x_1 x_2 \cdots x_d a_d \quad \rightarrow a_d \quad (10.31)$$

$$p_{d+1} = (\lambda - a_d)p_d - x_d p_{d-1}. \quad (10.32)$$

$$(10.33)$$

This proves the assertions.  $\square$

**Corollary 10.1.** *With above notation, let  $W, W'$  denote thin irreducible  $T(x)$ -modules. The following are equivalent.*

(i)  $W, W'$  are isomorphic as  $T$ -modules.

(ii)  $r(W) = r(W')$  and  $m_W = m_{W'}$ .

(iii)  $r(W) = r(W')$ ,  $d(W) = d(W')$ ,  $a_i(W) = a_i(W')$  and  $x_i(W) = x_i(W')$  ( $0 \leq i \leq d$ ).

*Proof.* (i)  $\Rightarrow$  (iii) Write  $r \equiv r(W)$ ,  $r' \equiv r(W')$ ,  $d = d(W)$ ,  $d' = d(W')$ ,  $a_i = a_i(W)$ ,  $a'_i = a_i(W')$ ,  $x_i = x_i(W)$  and  $x'_i = x_i(W')$ .

Let  $\sigma : W \rightarrow W'$  denote an isomorphism of  $T$ -modules. (See Definition 5.1.)

For every  $i$ ,

$$\sigma E_i^* W = E_i^* \sigma W = E_i^* W'.$$

So,  $r = r'$  and  $d = d'$ .

To show  $a_i = a'_i$ , pick  $w \in E_{r+i}^* W \setminus \{0\}$ . Then,

$$E_{r+i}^* A E_{r+i}^* \sigma(W) = \sigma(E_{r+i}^* A E_{r+i}^* w) = \sigma(a_i w) = a_i \sigma(w),$$

and  $\sigma w \neq 0$ . So,

$$a_i = \text{eigenvalue of } E_{r+i}^* A E_{r+i}^* \text{ on } E_{r+i}^* W \quad (10.34)$$

$$= a'_i \quad (10.35)$$

It is similar to show  $x = x'$ .

*Remark.* Pick  $w \in E_{r+i-1}^* W \setminus \{0\}$

$$E_{r+i-1}^* A E_{r+i}^* A E_{r+i-1}^* \sigma(W) = \sigma(E_{r+i-1}^* A E_{r+i}^* A E_{r+i-1}^* w) = x_i \sigma(w).$$

Hence,  $x_i$  is the eigenvalue of  $E_{r+i-1}^* A E_{r+i}^* A E_{r+i-1}^*$  on  $E_{r+i-1}^* W = x'_i$ .

(iii)  $\Rightarrow$  (i)

Pick  $0 \neq w_0 \in E_r^* W$ ,  $0 \neq w'_0 \in E_r^* W'$ . Let  $p_i$  be in Lemma 9.1, and set

$$w_i = p_i(A)w_0 \in E_{r+i}^* W \quad (0 \leq i \leq d) \quad (10.36)$$

$$w'_i = p'_i(A)w'_0 \in E_{r+i}^* W' \quad (0 \leq i \leq d) \quad (10.37)$$

Define a linear transformation,

$$\sigma : W \rightarrow W' \quad (w_i \mapsto w'_i).$$

Since  $\{w_i\}$  and  $\{w'_i\}$  are bases with  $d = d'$ ,  $\sigma$  is an isomorphism of vector spaces.

We need to show

$$a\sigma = \sigma a \quad (\text{for all } a \in T).$$

Take  $a = E_j^*$  for some  $j$  ( $0 \leq j \leq d(x)$ ). Then for all  $i$ , we have

$$E_j^* \sigma w_i = E_j^* w'_i = \delta_{ij} w'_i,$$

$$\sigma E_j^* w_i = \delta_{ij} \sigma(w_i) = \delta_{ij} w'_i.$$

$$E_j^* \sigma w_i = \sigma E_j^* w_i?$$

Take an adjacency matrix  $A$  of  $a$ . Then,

$$A \sigma w_i = A w'_i = w'_{i+1} + a'_i w'_i + x'_i w'_{i-1} = \sigma(w_{i+1} + a_i w_i + x_i w_{i-1}) = \sigma A w_i.$$

(ii)  $\Rightarrow$  (iii) Lemma 10.2.

(iii)  $\Rightarrow$  (ii) Given  $d, a_i, x_i$ , we can compute the polynomial sequence

$$p_0, p_1, \dots, p_{d+1}$$

for  $W$ .

Show  $p_0, p_1, \dots, p_{d+1}$  determines  $m = m_W$ . Set

$$\Delta = \{\theta \in \mathbb{R} \mid p_{d+1}(\theta) = 0\}.$$

Observe:  $|\Delta| = d + 1$ . See ‘An Introcuction to Interlacing’.

$m(\theta) = 0$  if  $\theta \notin \Delta$  ( $\theta \in \mathbb{R}$ ). So it suffices to find  $m(\theta)$ ,  $\theta \in \Delta$ .

By Lemma 10.1 (i),

$$\begin{cases} \sum_{\theta \in \Delta} m(\theta) p_0(\theta) &= 1 \\ \sum_{\theta \in \Delta} m(\theta) p_1(\theta) &= 0 \\ \vdots & \\ \sum_{\theta \in \Delta} m(\theta) p_d(\theta) &= 0 \end{cases}$$

$d + 1$  linear equation with  $d + 1$  unknowns  $m(\theta)$  ( $\theta \in \Delta$ ).

But the coefficient matrix is essentially Vander Monde (since  $\deg p_i = i$ ). Hence the system is nonsingular and there are unique values for  $m(\theta)$  ( $\theta \in \Delta$ ).  $\square$

*Remark.*

$$\begin{pmatrix} \theta - a_0 & -1 & \cdots & 0 & 0 \\ -x_1 & \theta - a_1 & \cdots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & \theta - a_{d-1} & -1 \\ 0 & 0 & \cdots & -x_d & \theta - a_d \end{pmatrix} \begin{pmatrix} p_0(\theta) \\ \vdots \\ \vdots \\ \vdots \\ p_d(\theta) \end{pmatrix} = 0,$$

where  $\theta$  is an eigenvalue of a diagonalizable matrix

$$L = \begin{pmatrix} a_0 & 1 & \cdots & 0 & 0 \\ x_1 & a_1 & \cdots & 0 & 0 \\ \vdots & \ddots & \ddots & \ddots & \vdots \\ 0 & 0 & \cdots & a_{d-1} & 1 \\ 0 & 0 & \cdots & x_d & \theta a_d \end{pmatrix}$$

with multiplicity  $\dim(\text{Ker}(\theta I - L)) = 1$ .



## Chapter 11

# Examples of $T$ -Module

**Friday, February 12, 1993**

Let  $\Gamma = (X, E)$  be a connected graph.

Let  $\theta_0$  be the maximal eigenvalue of  $\Gamma$ , and  $\delta$  its corresponding eigenvector.

$$\delta = \sum_{y \in X} \delta_y \hat{y}.$$

Without loss of generality, we may assume that  $\delta_y \in \mathbb{R}^*$  for all  $y \in X$ .

**Lemma 11.1.** *Fix a vertex  $x \in X$ . Write  $T \equiv T(x)$ ,  $E_i^* \equiv E_i^*(x)$ .*

- (i)  $T\delta = T\hat{x}$  is an irreducible  $T$ -module.
- (ii) *Given any irreducible  $T$ -module  $W$ , the following are equivalent:*
  - (iia)  $W = T\delta$ .
  - (iib) *The diameter  $d(W) = d(x)$ .*
  - (iic) *The endpoint  $r(W) = 0$ .*

*Proof.* (i) Observe: there exists an irreducible  $T$ -module  $W$  that contains  $\delta$ .

Let  $V = \sum_i W_i$  be a direct sum decomposition of the standard module. Then

$$\text{Span}(\delta) = E_0 V = \sum_i E_0 W_i.$$

So,  $E_0 W_i \neq 0$  for some  $i$ . Then,

$$\delta \in E_0 W_i \subseteq W_i.$$

Observe:  $T\delta$  is an irreducible  $T$ -module.

Since  $\delta \in W$ , where  $W$  is a  $T$ -module. As  $T\delta \subseteq W$  and  $W$  is irreducible,  $T\delta = W$ .

Observe:  $T\delta = T\hat{x}$ .

Since  $\hat{x} = \delta_x^{-1} E_0^* \delta \in T\delta$ ,  $T\hat{x} \subseteq T\delta$ . Since  $T\delta$  is irreducible,  $T\hat{x} = T\delta$ .

(ii) (a)  $\rightarrow$  (b):

$$E_i^* \delta = \sum_{y \in X, \partial(x,y)=i} \delta_y \hat{y} \neq 0, \quad (0 \leq i \leq d(x)),$$

because  $\delta_y > 0$  for every  $y \in X$ .

Hence,

$$E_i^* T\delta \neq 0, \quad (0 \leq i \leq d(x)).$$

Thus,  $d(x) = d(W)$ .

(b)  $\rightarrow$  (c): Immediate.

(c)  $\rightarrow$  (a): Since  $r(W) = 0$ ,  $E_0^* W \neq 0$ . Hence,  $\hat{x} \in W$  and  $T\hat{x} \subseteq W$ .

By the irreducibility, we have  $T\hat{x} = W$ .  $\square$

**Lemma 11.2.** Assume  $\Gamma$  is bipartite ( $X = X^+ \cup X^-$ ) ( $X^+$  and  $X^-$  are nonempty). Then the following are equivalent.

(i) There exist  $\alpha^+$  and  $\alpha^- \in \mathbb{R}$  such that


$$\delta_x = \begin{cases} \alpha^+ & \text{if } x \in X^+ \\ \alpha^- & \text{if } x \in X^-. \end{cases}$$

/(ii) There exist  $k^+$  and  $k^- \in \mathbb{Z}^{>0}$  such that

$$k(x) = \begin{cases} k^+ & \text{if } x \in X^+ \\ k^- & \text{if } x \in X^-. \end{cases}$$

In this case,  $k^+ k^- = \theta_0^2$ , and  $\Gamma$  is called bi-regular.

*Proof.* (i)  $\rightarrow$  (ii)



$$A\delta = A \left( \alpha^+ \sum_{x \in X^+} \hat{x} + \alpha^- \sum_{y \in X^-} \hat{y} \right) \quad (11.1)$$

$$= \alpha^+ \sum_{y \in X^-} k(y) \hat{y} + \alpha^- \sum_{x \in X^+} k(x) \hat{x} \quad (11.2)$$

$$= \theta_0 \delta. \quad (11.3)$$

So,

$$k(x)\alpha^- = \theta_0\alpha^+, \quad k(y)\alpha^+ = \theta_0\alpha^-.$$

As  $\alpha^+ \neq 0$  and  $\alpha^- \neq 0$ ,

$$k^+ := k(x) \text{ is independent of the choice of } x \in X^+, \text{ and} \quad (11.4)$$

$$k^- := k(y) \text{ is independent of the choice of } y \in X^-. \quad (11.5)$$

Moreover,  $k^+k^- = \theta_0^2$ .

(ii)  $\rightarrow$  (i) Set

$$\delta' = \sum_{y \in X} \alpha_y \hat{y} \quad \text{where } \alpha = \begin{cases} 1/\sqrt{k^-} & \text{if } y \in X^+ \\ 1/\sqrt{k^+} & \text{if } y \in X^- \end{cases}.$$

Then one checks

$$A\delta' = A \left( \frac{1}{\sqrt{k^-}} \sum_{y \in X^+} \hat{y} + \frac{1}{\sqrt{k^+}} \sum_{y \in X^-} \hat{y} \right) \quad (11.6)$$

$$= \frac{k^-}{\sqrt{k^-}} \sum_{y \in X^-} \hat{y} + \frac{k^+}{\sqrt{k^+}} \sum_{y \in X^+} \hat{y} \quad (11.7)$$

$$= \sqrt{k^+k^-} \delta' \quad (11.8)$$

Since  $\delta' > 0$ ,  $\delta' \in \text{Span}(\delta)$ , and  $\theta_0 = \sqrt{k^+k^-}$ .  $\square$

**Definition 11.1.** For any graph  $\Gamma = (X, E)$ , fix a vertex  $x \in X$ . Set  $d = d(x)$ .

$\Gamma$  is distance-regular with respect to  $x$ , if for all  $i : (0 \leq i \leq d)$ , and all  $y \in X$  such that  $\partial(x, y) = i$ :

$$c_i(x) := |\{z \in X \mid \partial(x, z) = i-1, \partial(y, z) = 1\}| \quad (11.9)$$

$$a_i(x) := |\{z \in X \mid \partial(x, z) = i, \partial(y, z) = 1\}| \quad (11.10)$$

$$b_i(x) := |\{z \in X \mid \partial(x, z) = i+1, \partial(y, z) = 1\}| \quad (11.11)$$

depends only on  $i$ ,  $x$ , and not on  $y$ .

(In this case,  $c_0(x) = a_0(x) = b_d(x) = 0$ ,  $c_1(x) = 1$ ,  $b_0(x) = k(x)$  is the valency of  $x$ .)

We call  $c_i(x)$ ,  $a_i(x)$  and  $b_i(x)$  the intersection numbers with respect to  $x$ .

**Example 11.1.**



$$c_0 = 1 \qquad c_1 = 1 \qquad c_2 = 1 \qquad (11.12)$$

$$a_0 = 0 \qquad a_1 = 1 \qquad a_2 = 1 \qquad (11.13)$$

$$b_0 = 2 \qquad b_1 = 1 \qquad b_2 = 0 \qquad (11.14)$$



## Chapter 12

# Distance-Regular

Monday, February 15, 1993

**Lemma 12.1.** *For any connected graph  $\Gamma = (X, E)$ , the following are equivalent.*

(i) *The trivial  $T(x)$ -module is thin for all  $x \in X$ .*

(ii)  $\left\{ \sum_{y \in X, d(x,y)=i} \hat{y} \mid 0 \leq i \leq d(x) \right\}$  *is a basis for the trivial  $T(x)$ -module for every  $x \in X$ .*

(iii)  $\Gamma$  *is distance-regular with respect to  $x$  for all  $x \in X$ .*

**Note.** Let  $\Gamma = (X, E)$  be a graph, with  $X = \{x, y_1, y_2, y_3, z_1, z_2, z_3\}$ ,  $E = \{xy_1, xy_2, xy_3, y_1z_1, y_1z_2, y_2z_3, y_3z_3\}$ .



Then (i), (ii) are not equivalent for a single vertex  $x$ .

$$E_0^* T \hat{x} = \langle \hat{x} \rangle, \quad (12.1)$$

$$E_1^* T \hat{x} = \langle y_1 + y_2 + y_3 \rangle, \quad (12.2)$$

$$E_2^* T \hat{x} = \langle z_1 + z_2 + 2z_3 \rangle. \quad (12.3)$$

*Proof of Lemma 12.1.* (i)  $\rightarrow$  (ii) Let  $\delta = \sum_{y \in X} \delta_y \hat{y}$  be an eigenvector for the maximal eigenvalue  $\theta_0$ . Then,

$$\sum_{y \in X, \partial(x,y)=1} \hat{y} = A\hat{x} \in T(x)\hat{x} = T(x)\delta \ni E_1^* \delta \quad (12.4)$$

$$= \sum_{y \in X, \partial(x,y)=1} \delta_y \hat{y} \quad (12.5)$$

If the trivial  $T(x)$ -module is thin,

$$\delta_y = \delta_z \text{ for } y, z \in X, \partial(x, y) = \partial(x, z) = 1.$$

Hence,  $\delta_y = \delta_z$  if  $y$  and  $z$  in  $X$  are connected by a path of even length.

So,  $\Gamma$  is regular or bipartite biregular by Lemma 11.2.

In particular,  $\delta_y = \delta_z$  if  $\partial(x, y) = \partial(x, z)$ , as there is a path of length  $2 \cdot \partial(x, y)$ ;

$$y \sim \dots \sim x \sim \dots \sim z.$$

Hence,

$$E_i^* \delta \in \text{Span} \left( \sum_{y \in X, \partial(x,y)=i} \hat{y} \right).$$

Since  $E_0^* \delta, E_1^* \delta, \dots, E_d^* \delta$  forms a basis for  $T(x)\delta$ , we have (ii).

(ii)  $\rightarrow$  (iii) Fix  $x \in X$ , and let  $T \equiv T(x)$ ,  $E_i^* \equiv E_i^*(x)$ , and  $d \equiv d(x)$ .

$$A \sum_{y \in X, \partial(x,y)=i} \hat{y} = \sum_{z \in X} |\{y \in X \mid \partial(y, z) = 1, \partial(x, y) = i\}| \hat{z} \quad (12.6)$$

$$= \sum_{z \in X, \partial(x,y)=i-1} b_{i-1}(x, z) \hat{z} \quad (12.7)$$

$$+ \sum_{z \in X, \partial(x,y)=i} a_i(x, z) \hat{z} \quad (12.8)$$

$$+ \sum_{z \in X, \partial(x,y)=i+1} c_{i+1}(x, z) \hat{z} \quad (12.9)$$

$$\in \text{Span} \left\{ \sum_{z \in X, \partial(x,z)=j} \hat{z} \mid j = 0, 1, \dots, d \right\}. \quad (12.10)$$

Hence,  $b_{i-1}(x, z)$ ,  $a_i(x, z)$  and  $c_{i+1}(x, z)$  depend only on  $i$  and  $x$ , and not on  $z$ . Therefore,  $\Gamma$  is distance-regular with respect to  $x$ .

(iii)  $\rightarrow$  (i) Fix  $x \in X$ , and let  $T \equiv T(x)$ ,  $E_i^* \equiv E_i^*(x)$ , and  $d \equiv d(x)$ . By definition of distance-regularity, for every  $i$  ( $0 \leq i \leq d$ ),

$$A \left( \sum_{y \in X, \partial(x,y)=i} \hat{y} \right) = b_{i-1}(x) \sum_{y \in X, \partial(x,y)=i-1} \hat{y} \quad (12.11)$$

$$+ a_i(x) \sum_{y \in X, \partial(x,y)=i} \hat{y} \quad (12.12)$$

$$+ c_{i+1}(x) \sum_{y \in X, \partial(x,y)=i+1} \hat{y}. \quad (12.13)$$

Hence,

$$W = \left\{ \sum_{y \in X, \partial(x,y)=i} \hat{y} \mid 0 \leq i \leq d \right\}$$

is  $A$ -invariant and so  $T$ -invariant. Since  $\hat{x} \in W$ ,  $T\hat{x} = W$  is the trivial module and  $T\hat{x}$  is thin.  $\square$

Next, we show more is true if (i) – (iii) hold in Lemma 12.1.

In fact,  $d(x)$ ,  $a_i(x)$ ,  $c_i(x)$ , and  $b_i(x)$  are

$$\begin{cases} \text{independent of } X & \text{if } \Gamma \text{ is regular; or} \\ \text{constant over } X^+ \text{ and } X^- & \text{if } \Gamma \text{ is biregular.} \end{cases}$$

Let  $\Gamma = (X, E)$  be any (connected) graph. Pick vertices  $x, y \in X$ .

Let  $W$  be a thin, irreducible  $T(x)$ -module, and measure  $m : \mathbb{R} \rightarrow \mathbb{R}$  determined by  $W$ .

Let  $W'$  be a thin, irreducible  $T(y)$ -module, and measure  $m' : \mathbb{R} \rightarrow \mathbb{R}$  determined by  $W'$ .

Recall  $W, W'$  are orthogonal if

$$\langle w, w' \rangle = 0 \quad \text{for all } w \in W, w' \in W'.$$

We shall show if  $W$  and  $W'$  are not orthogonal, then  $m$  and  $m'$  are related:

$$m \cdot \text{poly}_1 = m' \cdot \text{poly}_2$$

for some polynomials with

$$\deg \text{poly}_1 + \deg \text{poly}_2 \leq 2 \cdot \partial(x, y).$$

**Notation.**  $V$ : standard module of  $\Gamma$ .

$H$ : any subspace of  $V$ .

$$V = H + H^\perp \quad \text{orthogonal direct sum,}$$

and for  $v = v_1 + v_2$   $\text{proj}_H : V \rightarrow H$  ( $v \mapsto v_1$ ): linear transformation.

Observe: For every  $v \in V$ ,

$$v - \text{proj}_H v \in H^\perp.$$

So,

$$\langle v - \text{proj}_H v, h \rangle = 0 \quad \text{for all } h \in H \text{ or,}$$

$$\langle v, h \rangle = \langle \text{proj}_H v, h \rangle \quad \text{for all } v \in V, \text{ and for all } h \in H.$$

**Theorem 12.1.** *Let  $\Gamma = (X, E)$  be any graph. Pick vertices  $x, y \in X$  and set  $\Delta = \partial(x, y)$ . Assume*

*$W$ : thin irreducible  $T(x)$ -module with endpoint  $r$ , diameter  $d$ , and measure  $m$ .*

*$W'$ : thin irreducible  $T(y)$ -module with endpoint  $r'$ , diameter  $d'$ , and measure  $m'$ .*

*$W$  and  $W'$  are not orthogonal.*

*Now pick*

$$0 \neq w \in E_r^*(x)W, \quad 0 \neq w' \in E_{r'}^*(x)W'.$$

*Then,*

$$(i) \quad \text{proj}_{W'} w = p(A) \frac{\|w\|}{\|w'\|} w'$$

*for some  $0 \neq p \in \mathbb{C}[\lambda]$  with  $\deg p \leq \Delta - r' + r, d'$ ,*

$$\text{proj}_W w' = p'(A) \frac{\|w'\|}{\|w\|} w$$

*for some  $0 \neq p' \in \mathbb{C}[\lambda]$  with  $\deg p' \leq \Delta - r + r', d$ .*

*(ii) For all eigenvalues  $\theta_i$  of  $\Gamma$ ,*

$$\frac{\langle E_i w, E_i w' \rangle}{\|w\| \|w'\|} = m(\theta_i) \overline{p'(\theta_i)}.$$

*(iii) For all eigenvalues  $\theta_i$  of  $\Gamma$ ,*

$$p(\theta_i) p'(\theta_i)$$

*is in a real number in interval  $[0, 1]$ .*

*Proof.* (i) Since  $W, W'$  are not orthogonal, there exist

$$v \in W, v' \in W' \text{ such that } \langle v, v' \rangle \neq 0.$$

Then there exists  $a \in M$  such that

$$v' = aw'.$$

(This is because  $w'_i = p'_i(A)w'_0$  and hence for every  $v' \in W'$ , there is a polynomial  $q \in \mathbb{C}[\lambda]$ ,  $q(A)w'_0 = v$ .)

We have

$$0 \neq \langle v', v \rangle = \langle aw', v \rangle = \langle w', a^*v \rangle$$

and  $a^*v \in W$ .

Hence,  $\text{proj}_W w' \neq 0$ .

Let  $p_0, \dots, p_d \in \mathbb{C}[\lambda]$  be from Lemma 9.1.

Then,  $w_i = p_i(A)w$  is a basis for  $E_{r+i}^*(x)W$  ( $0 \leq i \leq d$ ).

Hence,

$$\text{proj}_W w' = \alpha_0 w_0 + \dots + \alpha_d w_d \text{ for some } \alpha_j \in \mathbb{C}.$$

Set

$$p' := \frac{\|w\|}{\|w'\|} \sum_{i=0}^d \alpha_i p_i.$$

Then  $0 \neq p' \in \mathbb{C}[\lambda]$  and  $\deg p' \leq d$ .

Claim:  $\alpha_i = 0$  ( $\Delta - r + r' < i \leq d$ ).

In particular,  $\deg p' \leq \Delta - r + r'$ .

*Pf.* Observe:

$$w' \in E_{r'}^*(y)V, \quad w \in E_r^*(x)V,$$

for  $\partial(x, y) = \Delta$ .

$$E_{r'}^*(y)V \cap E_{r+i}^*(x)V = 0$$

by triangle inequality.

( $\Delta = \partial(x, y) < r + i - r'$  or  $\Delta + r' < r + i$  by our choice of  $i$ .)



Hence,

$$E_{r'}^*(y)V \perp E_{r+i}^*(x)V,$$

or

$$0 = \langle w', w_i \rangle \quad (12.14)$$

$$= \langle \text{proj}_W w', w_i \rangle \quad (12.15)$$

$$= \sum_{j=0}^d \alpha_j \langle w_j, w_i \rangle \quad (12.16)$$

$$= \alpha_i \|w_i\|^2. \quad (12.17)$$

Hence,  $\alpha_i = 0$ . Thus,

$$\text{proj}_W w' = \sum_{i=0}^{\Delta+r'-r} \alpha_i w_i \quad (12.18)$$

$$= \sum_{i=0}^{\Delta+r'-r} \alpha_i p_i(A) w_0 \quad (12.19)$$

$$= p'(A) \frac{\|w'\|}{\|w\|} w. \quad (12.20)$$

(ii) We have

$$\frac{\langle E_i w, E_i w' \rangle}{\|w\| \|w'\|} = \frac{\langle E_i w, w' \rangle}{\|w\| \|w'\|} \quad (12.21)$$

$$= \frac{\langle E_i w, \text{proj}_W w' \rangle}{\|w\| \|w'\|} \quad \text{as } \text{proj}_W w' = p'(A) \frac{\|w\|}{\|w'\|} w \quad (12.22)$$

$$= \frac{\langle E_i w, p'(A) w \rangle}{\|w\|^2} \quad (12.23)$$

$$= \frac{\langle E_i w, E_i p'(A) w \rangle}{\|w\|^2} \quad (12.24)$$

$$= \overline{p'(\theta_i)} \frac{\|E_i W\|^2}{\|w\|^2} \quad (12.25)$$

$$= \overline{p'(\theta_i)} m(\theta_i). \quad (12.26)$$

Moreover, as  $m(\theta_i), m'(\theta_i) \in \mathbb{R}$ ,

$$\frac{\langle E_i w, E_i w' \rangle}{\|w\| \|w'\|} = \frac{\overline{\langle E_i w, E_i w' \rangle}}{\|w'\| \|w\|} = \overline{p(\theta_i) m'(\theta_i)} = p(\theta_i) m'(\theta_i).$$

(iii) Since,

$$\frac{|\langle E_i w, E_i w' \rangle|^2}{\|w\|^2 \|w'\|^2} = p(\theta_i) p'(\theta_i) m(\theta_i) m'(\theta_i),$$

$$p(\theta_i)p'(\theta_i) = \frac{|\langle E_i w, E_i w' \rangle|^2}{m(\theta_i)m'(\theta_i)\|w\|^2\|w'\|^2} \in \mathbb{R} \quad (12.27)$$

$$= \frac{|\langle E_i w, E_i w' \rangle|^2}{\frac{\|E_i w\|^2}{\|w\|^2} \frac{\|E_i w'\|^2}{\|w'\|^2} \|w\|^2 \|w'\|^2}. \quad (12.28)$$

By Cauchy-Schwartz inequality,

$$(|\langle a, b \rangle| \leq \|a\| \|b\|, )$$

$$\frac{|\langle E_i w, E_i w' \rangle|^2}{\|E_i w\|^2 \|E_i w'\|^2} \leq 1.$$

Hence, we have the assertion.  $\square$





# Chapter 13

## Modules of a DRG

Wednesday, February 17, 1993

**Lemma 13.1.** *Let  $\Gamma = (X, E)$  be any graph. Pick an edge  $xy \in E$ .*

*Assume the trivial  $T(x)$ -module  $T(x)\delta$  is thin with measure  $m_x$ ,*

*and the trivial  $T(y)$ -module  $T(y)\delta$  is thin with measure  $m_y$ .*

*Then,*

$$(ia) \quad \frac{m_x(\theta)}{k_x} = \frac{m_y(\theta)}{k_y} \text{ for all } \theta \in \mathbb{R} \setminus \{0\}.$$

$$(ib) \quad \frac{m_x(0) - 1}{k_x} = \frac{m_y(0) - 1}{k_y} \text{ for all } \theta \in \mathbb{R} \setminus \{0\}.$$

$$(\delta = \sum_{y \in X} \delta_y \hat{y} \quad \text{eigenvector corresponding to the maximal eigenvalue})$$

*Proof.* Apply Theorem 12.1,

$$W = T(x)\delta \quad r = 0, \quad d = d(x) \tag{13.1}$$

$$W' = T(y)\delta \quad r' = 0, \quad d' = d(y). \tag{13.2}$$

Take  $w = \hat{x}$ ,  $w' = \hat{y}$ .

Claim.  $\text{proj}_{T(y)\delta} \hat{x} = k_y^{-1} A \hat{y}$ .

*Pf.* Since

$$\hat{y} \in T(y)\delta, \quad A\hat{y} \in T(y)\delta.$$

Show

$$(\hat{x} - k_y^{-1} A \hat{y}) \perp (T(y)\delta).$$

Recall

$$A\hat{y} = \sum_{z \in X, yz \in E} \hat{z}.$$

$$\hat{x} - k_y^{-1}Ay \in E_1^*(y)V.$$

So,

$$\hat{x} - \frac{1}{k_y}A\hat{y} \perp E_j^*(y)T(y)\delta \quad \text{if } j \neq 1 \quad (0 \leq j \leq k(y)).$$

And we have,

$$\left\langle \hat{x} - \frac{1}{k_y}A\hat{y}, A\hat{y} \right\rangle = \left\langle \hat{x}, \sum_{z \in X, yz \in E} \hat{z} \right\rangle - \frac{1}{k_y} \left\| \sum_{z \in X, yz \in E} \hat{z} \right\|^2 \quad (13.3)$$

$$= 1 - 1 \quad (13.4)$$

$$= 0 \quad (13.5)$$

This proves Claim.

Similarly,

$$\text{prof}_{T(x)\delta} \hat{y} = k_x^{-1}A\hat{x}.$$

Hence, the polynomials  $p, p' \in \mathbb{C}[\lambda]$  from Theorem 12.1 equal

$$\frac{\lambda}{k_y} \quad \text{and} \quad \frac{\lambda}{k_x}$$

respectively.

By Theorem 12.1,

$$\frac{m_x(\theta)\theta}{k_x} = m_x(\theta)\overline{p'(\theta)} = m_y(\theta)\overline{p(\theta)} = \frac{m_y(\theta)\theta}{k_y}.$$

If  $\theta \neq 0$ , we have (ia).

Also,

$$\frac{1 - m_x(0)}{k_x} = \left( \sum_{\theta \in \mathbb{R} \setminus \{0\}} m_x(0) \right) \frac{1}{k_x} \quad \text{by (ia)} \quad (13.6)$$

$$= \left( \sum_{\theta \in \mathbb{R} \setminus \{0\}} m_y(0) \right) \frac{1}{k_y} \quad (13.7)$$

$$= \frac{1 - m_y(0)}{k_y} \quad (13.8)$$

Hence, we have (ib). □

**Theorem 13.1.** *Suppose any graph  $\Gamma = (X, E)$  is distance-regular with respect to every vertex  $x \in X$ . (So  $\Gamma$  is regular or biregular by Lemma 12.1.)*

*Then,*

*Case  $\Gamma$  is regular: the diameter  $d(x)$  and the intersection numbers  $a_i(x)$ ,  $b_i(x)$ ,  $c_i(x)$  ( $0 \leq i \leq d(x)$ ) are independent of  $x \in X$ .*

*(And  $\Gamma$  is called distance-regular.)*

*Case  $\Gamma$  is biregular: ( $X = X^+ \cup X^-$ )*

*$d(x)$  and  $a_i(x)$ ,  $b_i(x)$ ,  $c_i(x)$  ( $0 \leq i \leq d(x)$ ) are constant over  $X^+$  and  $X^-$ . (And  $\Gamma$  is called distance-biregular.)*

*Proof.* We apply Lemma 13.1.

Case  $\Gamma$ : regular.

Then  $m_x = m_y$  for all  $xy \in E$ . Hence, the measure of the trivial  $T(x)$ -module is independent of  $x \in X$ .

Case  $\Gamma$  is biregular.

Then  $m_x = m_{x'}$  for all  $x, x' \in X$  with  $\partial(x, x') = 2$ .

Hence, the measure of the trivial  $T(x)$ -module is constant over  $x \in X^+$ ,  $X^-$ .

Fix  $x \in X$ . Write  $T \equiv T(x)$ ,  $E_i^* \equiv E_i^*(x)$ ,  $W = T\delta$  with measure  $m$ , diameter  $d = d(x)$ .

We know by Corollary 10.1 that  $m$  determines

$$d, \quad a_i(W) \ (0 \leq i \leq d), \quad x_i(W) \ (1 \leq i \leq d)$$

(as  $d = D(x) = d(W)$  by Lemma 11.1.)

We shall show that  $m$  determines

$$a_i(x), \ c_i(x), \ b_i(x) \quad (0 \leq i \leq d).$$

Observe:

$$a_i(W) = a_i(x) \quad (0 \leq i \leq d) \tag{13.9}$$

$$x_i(W) = b_{i-1}c_i(x) \quad (1 \leq i \leq d) \tag{13.10}$$

*Remark.*  $a_i = a_i(W)$  is an eigenvalue of

$$E_i^* A E_i^* \text{ on } E_i^* W = \langle \sum_{y \in \Gamma_i(x)} \hat{y} \rangle.$$

(See Lemma 12.1.)

$x_i = x_i(W)$  is an eigenvalue of

$$E_{i-1}^* A E_i^* A E_{i-1}^* \text{ on } E_{i-1}^* W,$$

and

$$A \sum_{y \in X, \partial(x,y)} \hat{y} = b_{i-1}(x) \sum_{y \in X, \partial(x,y)=i-1} \hat{y} \quad (13.11)$$

$$+ a_i(x) \sum_{y \in X, \partial(x,y)=i} \hat{y} \quad (13.12)$$

$$+ c_{i+1} \sum_{y \in X, \partial(x,y)=i+1} \hat{y} \quad (13.13)$$

So  $x_i = b_{i-1}(x)c_i(x)$ .

Set  $k^+ = k_x$ . Define

$$k^- = \frac{\theta_0^2}{k^+},$$

where  $\theta_0$  is the maximal eigenvalue. (See Lemma 11.1.)

(So,  $k^+ = k^-$  is the valency, if  $\Gamma$  is regular.)

For every  $i$  ( $0 \leq i \leq d$ ) and for every  $z \in X$  with  $\partial(x, z) = i$ ,

$$k_z = c_i(x) + a_i(x) + b_i(x) \quad (13.14)$$

$$= \begin{cases} k^+ & \text{if } i \text{ is even,} \\ k^- & \text{if } i \text{ is odd.} \end{cases} \quad (13.15)$$

Now  $m$  determines

$$c_0(x) = a_0(x) = 0, \quad c_1(x) = 1,$$

$$b_0(x) = b_0(x)c_1(x) = x_1(W).$$

$$k^+ = b_0(x) \quad (13.16)$$

$$k^- = \theta_0^2 / k^+ \quad (13.17)$$

$$c_i(x) = x_i(W) / b_{i-1}(x) \quad (1 \leq i \leq d) \quad (13.18)$$

$$b_i(x) = \begin{cases} k^+ - a_i(x) - c_i(x) & i: \text{ even,} \\ k^- - a_i(x) - c_i(x) & i: \text{ odd.} \end{cases} \quad (13.19)$$

This proves the assertions.  $\square$

**Proposition 13.1.** *Under the assumption of Theorem 13.1, the following hold.*

*Case  $\Gamma$ : regular.*

- (i)  $\dim E_i V = |X| m(\theta_i)$ .
- (ii)  $\Gamma$  has exactly  $d + 1$  distinct eigenvalues
- ( $d = \text{diam} \Gamma = d(x)$ , for all  $x \in X$ ).

Case  $\Gamma$ : biregular.

- (i)  $\dim E_V = |X^+| m^+(\theta_i) + |X^-| m^-(\theta_i)$ .
- (ii)  $\Gamma$  has exactly  $d^+ + 1$  distinct eigenvalues ( $d^+ \geq d^-$ ).
- (iii) If  $d^+$  is odd, the  $\Gamma$  is regular.
- (iv)  $d^+ = d^-$ , or  $d^+ = d^- + 1$  is even.
- (v)  $a_i(x) = 0$  for all  $i$  and for all  $x$ .

*Proof.* (i) Suppose  $\Gamma$  is regular.

Let  $m_x$  be the measure of the trivial  $T(x)$ -module,

$$m_x(\theta_i) = \|E_i \hat{x}\|^2, \quad \text{as } \|\hat{x}\| = 1.$$

Now,

$$|X| m_x(\theta_i) = \sum_{x \in X} m_x(\theta_i) \tag{13.20}$$

$$= \sum_{x \in X} \|E_i \hat{x}\|^2 \tag{13.21}$$

$$= \sum_{y, z \in X} |(E_i)_{yz}|^2 \tag{13.22}$$

$$= \text{trace} E_i \overline{E_i}^\top. \tag{13.23}$$

Since  $A$  is real symmetric and

$$E_i \overline{E_i}^\top = E_i^2 = E_i$$

with  $E_i$  symmetric

$$E_i \sim \begin{pmatrix} I & O \\ O & I \end{pmatrix}.$$

$$\text{trace} E_i = \text{rank} E_i = \dim E_i V.$$

Thus, we have the assertion in this case.

Suppose  $\Gamma$  is biregular.

Then, same except,

$$\sum_{x \in X} m_x(\theta_i) = |X^+| m^+(\theta_i) + |X^-| m^-(\theta_i).$$

- (ii)  $\Gamma$ : regular. Immediately, if  $\theta$  is an eigenvalue of  $\Gamma$ , then  $m(\theta) \neq 0$ .

$\Gamma$ : biregular. For each  $\theta = \theta_i \in \mathbb{R} \setminus \{0\}$ ,

$$m^-(\theta) \neq 0 \Leftrightarrow m^+(\theta) \neq 0 \quad (13.24)$$

$$\Leftrightarrow \theta \text{ is an eigenvalue of } \Gamma \quad (13.25)$$

$$\left( \frac{m^+(\theta)}{k^+} = \frac{m^-(\theta)}{k^-} \right) \quad (13.26)$$

(iv) and (v) are clear.

*Remark.* (iii) If  $d^+$  is odd,  $d^+ = d^-$  and  $\Gamma$  has even number of eigenvalues, i.e., 0 is not an eigenvalue. So  $A$  is nonsingular, and  $\Gamma$  is regular.

□

# Chapter 14

## Parameters of Thin Modules, I

Friday, February 19, 1993

Summary.

**Definition 14.1.** Assume  $\Gamma = (X, E)$  is distance-regular with respect to every vertex  $x \in X$ .

Notation: Let  $x \in X$ . The data of the trivial  $T(x)$ -module.

	Case DR	Case DBR
valency $k_x$	$k$	$\begin{cases} k^+ & \text{if } x \in X^+ \\ k^- & \text{if } x \in X^- \end{cases}$
$x$ -diameter $D_x$	$D$	$\begin{cases} D^+ & \text{if } x \in X^+ \\ D^- & \text{if } x \in X^- \end{cases}$
measure $m_x$	$m$	$\begin{cases} m^+ & \text{if } x \in X^+ \\ m^- & \text{if } x \in X^- \end{cases}$
int. number $c_i(x)$	$c_i$	$\begin{cases} c_i^+ & \text{if } x \in X^+ \\ c_i^- & \text{if } x \in X^- \end{cases}$
int. number $b_i(x)$	$b_i$	$\begin{cases} b_i^+ & \text{if } x \in X^+ \\ b_i^- & \text{if } x \in X^- \end{cases}$
int. number $a_i(x)$	$a_i$	$\begin{cases} a_i^+ & \text{if } x \in X^+ \\ a_i^- & \text{if } x \in X^- \end{cases}$

Call  $m, m^{\pm 1}$  the measure of  $\Gamma$ .

Assume  $\Gamma = (X, E)$  is distance-regular.

To what extent do  $a_i$ 's,  $b_i$ 's and  $c_i$ 's determine the structure of irreducible  $T(x)$ -modules? In general the following hold.

**Lemma 14.1.** *Assume  $\Gamma = (X, E)$  is distance-regular. Pick  $x \in X$ . Let  $X$  be a thin irreducible  $T(x)$ -module with endpoint  $r$ , diameter  $d$  and measure  $m_W$ .*

(i) *There is a unique polynomial  $f_W \in \mathbb{C}[\lambda]$  with the following properties.*

(ia)  $\deg f_W \leq D$  (diameter of  $\Gamma$ ).

(ib)  $m_W(\theta) = m(\theta)f_W(\theta)$  for every  $\theta \in \mathbb{R}$ , where  $m$  is the measure of  $\Gamma$ .

Moreover,  $f_W \in \mathbb{R}[\lambda]$ , and

(ii)  $\deg f_W \leq 2r$ .

(iii) *For all eigenvalues  $\theta_i$  of  $\Gamma$ ,  $\lambda - \theta_i$  is a factor of  $f_W$  whenever,  $E_i W = 0$ .*

In particular,  $2r - D + d \geq 0$ .

*Proof.* Let  $\theta_0, \dots, \theta_D$  denote distinct eigenvalues of  $\Gamma$ . Then  $m(\theta_i) \neq 0$  ( $0 \leq i \leq D$ ) by Proposition 13.1.

There exists a unique  $f_W \in \mathbb{C}[\lambda]$  with  $\deg f_W \leq D$  such that

$$f_W(\theta_i) = \frac{m_W(\theta_i)}{m(\theta_i)} \quad (0 \leq i \leq D)$$

by polynomial interpolation.

$f_W \in \mathbb{R}[\lambda]$  since

$$\theta_0, \dots, \theta_D \in \mathbb{R} \quad \text{and} \quad f_W(\theta_0), \dots, f_W(\theta_D) \in \mathbb{R}.$$

(ii) Without loss of generality, we may assume  $r < D/2$ , else trivial.

Pick  $0 \neq w \in E_r^*(x)W$ .

$$w = \sum_{y \in W, \partial(x,y)=r} \alpha_y \hat{y} \quad \text{some } \alpha_y \in \mathbb{C}.$$

Pick  $y \in X$  such that  $\alpha_y \neq 0$ .

Set  $W'$  be the trivial  $T(y)$ -module. ( $\langle w, \hat{y} \rangle \neq 0$ , as  $W \not\perp W'$ .)

$$r' = 0, \quad m' = m, \quad \Delta = r.$$

Apply Theorem 12.1, we have

$$\deg p \leq \Delta - r' + r = 2r, \quad p \neq 0 \tag{14.1}$$

$$\deg p' \leq \Delta - r + r' = 0, \quad p' \neq 0. \tag{14.2}$$



$$m_W(\theta)\overline{p'(\theta)} = m(\theta)p(\theta) \quad (\text{for all } \theta \in \mathbb{R}).$$

So,

$$\deg p/\bar{p}' \leq 2r,$$

and  $p/\bar{p}'$  satisfies the conditions of  $f_W$ .

$$\left( \frac{p(\theta)}{\bar{p}'(\theta)} = \frac{m_W(\theta)}{m(\theta)} \right)$$

(iii)

$$E_i W = 0 \rightarrow m_W(\theta_i) = 0 \rightarrow f_W(\theta_i) = 0.$$

that is,  $E_i W = 0$ . Hence  $\theta_i$  is a root of  $f_W(\lambda) = 0$ . So,

$$2r \geq \deg f_W \geq |\{\theta_i \mid E_i W = 0\}| = D - d.$$

Hence,

$$2r - D + d \geq 0.$$

This proves the assertions.  $\square$

**Lemma 14.2.** *Let  $\Gamma = (X, E)$  be any distance-regular graph with valency  $k$ , diameter  $D$  ( $d \geq 2$ ), measure  $m$ , and eigenvalues*

$$k = \theta_0 > \theta_1 > \dots > \theta_D.$$

*Pick  $x \in X$ . Let  $W$  be a thin irreducible  $T(x)$ -module with endpoint  $r = 1$ , diameter  $D$  and measure  $m_W = mf_W$ . Then one fo the following cases (i)–(iv) occurs.*

Case	$d$	$f_W(\lambda)$	$a_0(W)$
(i)	$D - 2$	$\frac{(\lambda-k)(\lambda-\theta_1)}{k(\theta_1+1)}$	$-\frac{b_1}{\theta_1+1} - 1$
(ii)	$D - 2$	$\frac{(\lambda-k)(\lambda-\theta_D)}{k(\theta_D+1)}$	$-\frac{b_1}{\theta_1+1} - 1$
(iii)	$D - 1$	$\frac{k-\lambda}{k}$	$-1$
(iv)	$D - 1$	$\frac{(\lambda-k)(\lambda-\beta)}{k(\beta+1)}$	$-\frac{b_1}{\beta+1} - 1$

*for some  $\beta \in \mathbb{R}$  with  $\beta \in (-\infty, \theta_D) \cup (\theta_1, \infty)$ . Moreover, the isomorphism class of  $W$  is determined by  $a_0(W)$ .*

**Note.** By (iii), the possible “shapes” of a thin irreducible  $T(x)$ -modules are:

$$r = 0 \quad d = D \tag{14.3}$$

$$r = 1 \quad d = D - 1 \tag{14.4}$$

$$r = 1 \quad d = D - 2 \tag{14.5}$$



## Chapter 15

# Parameters of Thin Modules, II

Monday, February 22, 1993

*Proof of Lemma 14.2 Continued.*

We have  $\deg f_W \leq 2$  by Lemma 14.1 (ii).

Also by Lemma 11.1,  $E_0 W = 0$ .

(As otherwise  $\langle \delta \rangle = E_0 V \subseteq W$  and  $r(W) = 0$ .)

Hence,  $\lambda - \theta_0 = \lambda - k$  is a factor of  $f_W$  by Lemma 14.1 (iii).

Let  $p_0, p_1, \dots, p_D$  denote the polynomials for the trivial  $T(x)$ -module from Lemma 9.1.

Recall,

$$\sum_{\theta \in \mathbb{R}} m(\theta) p_i(\theta) p_j(\theta) = \delta_{ij} x_1 x_2 \cdots x_i \quad (0 \leq i, j \leq D) \quad (15.1)$$

$$= \delta_{ij} b_0 b_1 \cdots b_{i-1} c_1 c_2 \cdots c_i. \quad (15.2)$$

Note that  $x_i = b_{i-1} c_i$  is in the proof of Theorem 7.1.

By construction,

$$p_0(\lambda) = 1.p_1(\lambda) \quad \quad \quad = \lambda.p_2(\lambda)\lambda^2 - a_1\lambda - k. \quad (15.3)$$

Apparently,

$$f_W = \sigma_0 p_0 + \sigma_1 p_1 + \sigma_2 p_2$$

for some  $\sigma_0, \sigma_1, \sigma_2 \in \mathbb{C}$ .

Claim:

$$\sigma_0 = 1, \quad (15.4)$$

$$\sigma_1 = \frac{a_0(W)}{k}, \quad (15.5)$$

$$\sigma_2 = \frac{1 + a_0(W)}{kb_1}. \quad (15.6)$$

*Pf of Claim.*

$$1 = \sum_{\theta \in \mathbb{R}} m_W(\theta) \quad (15.7)$$

$$= \sum_{\theta \in \mathbb{R}} m(\theta) f_W(\theta) \quad (15.8)$$

$$= \sum_{j=0}^2 \sigma_j \left( \sum_{\theta \in \mathbb{R}} m(\theta) p_j(\theta) \right) \quad (15.9)$$

$$= \sigma_0. \quad (15.10)$$

We applied Lemma 10.1 (ib), Lemma 14.1 (ib), and Lemma 10.1 (i) in this order.

Next by Lemma 10.1 (ii), and  $p_1(\theta) = \theta$ ,

$$a_0(W) = \sum_{\theta \in \mathbb{R}} m_W(\theta) \theta \quad (15.11)$$

$$= \sum_{\theta \in \mathbb{R}} f_W(\theta) \theta \quad (15.12)$$

$$= \sum_{j=0}^2 \sigma_j \sum_{\theta \in \mathbb{R}} m(\theta) p_j(\theta) p_1(\theta) \quad (15.13)$$

$$= \sigma_1 x_1(T\delta) \quad (15.14)$$

$$= \sigma_1 b_0 c_1 \quad (15.15)$$

$$= \sigma_1 k. \quad (15.16)$$

So for,

$$f_W(\lambda) = 1 + \frac{a_0(W)}{k} \lambda + \sigma_2(\lambda^2 - a_1 \lambda - k).$$

But,

$$0 = f_W(k) \quad (15.17)$$

$$= 1 + a_0(W) + \sigma_2 k(k - a_1 - 1) \quad (15.18)$$

$$1 + a_0(W) + \sigma_2 k b_1. \quad (15.19)$$

Thus,

$$\sigma_2 = -\frac{1 + a_0(W)}{k b_1}.$$

This proves Claim.

Case:  $a_0(W) = -1$ .

Here,  $\sigma_2 = 0$  and

$$f_W(\lambda) = 1 + \frac{a_0(W)\lambda}{k} = 1 - \frac{\lambda}{k}.$$

Also,

$$d + 1 = |\{\theta \mid \theta \text{ is an eigenvalue of } \Gamma, f_W(\theta) \neq 0\}| = D.$$

Case:  $a_0(W) \neq -1$ .

Here,  $\sigma_2 \neq 0$ , and  $\deg f_W = 2$ . So,

$$f_W(\lambda) = (\lambda - k)(\lambda - \beta)\alpha$$

for some  $\alpha, \beta \in \mathbb{C}$ ,  $\alpha \neq 0$ .

Comparing the coefficients in

$$(\lambda - k)(\lambda - \beta)\alpha = 1 + \frac{a_0(W)}{k}\lambda - \frac{a_0(W) + 1}{kb_1}(\lambda^2 - a_1\lambda - k),$$

we find

$$\alpha = -\frac{a_0(W) + 1}{kb_1}, \quad (15.20)$$

$$-(k + \beta)\alpha = \frac{a_0(W)}{k} + \frac{a_0(W) + 1}{kb_1}a_1, \quad (15.21)$$

$$k\beta\alpha = 1 + \frac{a_0(W) + 1}{b_1}. \quad (15.22)$$

Hence,

$$-\beta(a_0(W) + 1) = b_1 + (a_0(W) + 1).$$

Thus, we have

$$(1 + a_0(W))(1 + \beta) = -b_1. \quad (15.23)$$

In particular,  $\beta \neq -1$ , and

$$\alpha = -\frac{1 + a_0(W)}{kb_1} = \frac{1}{k(\beta + 1)}.$$

Also, by Definition 9.2,

$$0 \leq m_W(\theta) \quad (15.24)$$

$$= m(\theta)f_W(\theta) \quad (\text{for all } \theta \in \mathbb{R}). \quad (15.25)$$

But if  $\theta$  is an eigenvalue of  $\Gamma$ ,

$$0 < m(\theta).$$

So,

$$0 \leq f_W(\theta) \quad (15.26)$$

$$= \frac{(\theta - k)(\theta - \beta)}{k(\beta + 1)}. \quad (15.27)$$

Either

$$\beta + 1 > 0 \rightarrow \theta - \beta \leq 0 \text{ or } \beta \geq \theta_1,$$

or

$$\beta + 1 < 0 \rightarrow \theta - \beta \geq 0 \text{ or } \beta \leq \theta_D.$$

If  $\beta = \theta_1$ ,

$$a_0(W) = -\frac{b_1}{\beta + 1} - 1 = -\frac{b_1}{\theta_1 + 1} - 1 \quad (15.28)$$

$$f_W(\lambda) = \frac{(\lambda - k)(\lambda - \theta_1)}{k(\theta_1 + 1)}, \quad (15.29)$$

and we have (i).

If  $\beta = \theta_D$ ,

$$a_0(W) = -\frac{b_1}{\theta_D + 1} - 1 \quad (15.30)$$

$$f_W(\lambda) = \frac{(\lambda - k)(\lambda - \theta_D)}{k(\theta_D + 1)}, \quad (15.31)$$

and we have (ii).

If  $\beta \notin \{\theta_1, \theta_2\}$ ,

$$\theta \in (-\infty, \theta_D) \cup (\theta_1, \infty),$$

we have (iv).

Note using (15.23), we have (iv).

□

**Note.** Using (15.23),

$$a_0(W) \rightarrow \beta \rightarrow f_W \rightarrow m_W \rightarrow \text{isomorphism class of } W.$$

**Note on Lemma 14.2.** In fact,  $\theta_1 > -1$ ,  $\theta_D < -1$  if  $D \geq 2$ .

**Definition 15.1.** The complete graph  $K_n$  has  $n$  vertices and diameter  $D = 1$ , i.e.,  $xy \in E$  for all vertices  $x, t$ .

$K_n$  is distance-regular with valency  $k = n - 1$  and  $a_1 = n - 2$ ,  $D = 1$ . Moreover, it has two distance eigenvalues  $\theta_0, \theta_1$ .

Recall,  $\theta_0, \dots, \theta_D$  are roots of  $p_{D+1}$ , i.e.,  $D + 1$  st polynomial for the trivial module/

$$p_0 = 1 \quad (15.32)$$

$$p_1 = \lambda \quad (15.33)$$

$$p_2 = \lambda^2 - a_1\lambda - k \quad (15.34)$$

$$= \lambda^2 - (n - 2)\lambda - (n - 1) \quad (15.35)$$

$$= (\lambda - (n - 1))(\lambda + 1). \quad (15.36)$$

The roots are  $\theta_0 = n - 1 = k$  and  $\theta_1 = -1$ .

**Lemma 15.1.** *Let  $\Gamma = (X, E)$  be distance-regular of diameter  $D \geq 1$  with distinct eigenvalues*

$$k = \theta_0 > \theta_1 > \dots > \theta_D.$$

(i)  $\theta_D \leq -1$  with equality if and only if  $D = 1$ .

(ii)  $\theta_1 \geq -1$  with equality if and only if  $D = 1$ .

*Proof.* (i) Suppose  $\theta_D \geq -1$ .

Then  $I + A$  is positive semi-definite.

By Lemma 2.1, there exists vectors  $\{v_x \mid x \in X\}$  in a Euclidean space such that

$$\langle v_x, v_y \rangle = (I + A)_{xy} \quad (15.37)$$

$$= \begin{cases} 1 & \text{if } x = y \text{ or } xy \in E, \\ 0 & \text{otherwise.} \end{cases} \quad (15.38)$$

For every  $xy \in E$ ,

$$\langle v_x, v_y \rangle = \|v_x\| \|v_y\| = 1.$$

Hence,  $v_x = v_y$ , and  $v_x$  is independent of  $x \in X$ .

Shus  $\langle v_x, v_y \rangle = 1$  for all  $x, y \in X$ .

We have  $I + A = J$ , (all 1's matrix), and  $D = 1$ .

(ii) Let  $m$  be the trivial measure. Then,

$$1 = \sum_{\theta \in \mathbb{R}} m(\theta) + \sum_{\theta \in \mathbb{R}} m(\theta)\theta \quad (15.39)$$

$$= \sum_{\theta \in \mathbb{R}} m(\theta)(\theta + 1) \quad (15.40)$$

$$= m(k)(k + 1) + \sum_{\theta \neq k} m(\theta)(\theta + 1) \quad (15.41)$$

$$\leq (k + 1)|X|^{-1}. \quad (15.42)$$

Note that  $m(k) = |X|^{-1} \dim d_0 V = |X|^{-1}$ .

So  $k+1 \geq |X|$  or  $k = |X| - 1$ . Thus,  $xy \in E$  for every  $x, y \in X$ , and  $D = 1$ .  $\square$

**Note.** Lemma 15.1 does not require distance-regular assumption.



## Chapter 16

# Thin Modoles of a DRG

Wednesday, February 24, 1993

Let  $\Gamma = (X, E)$  denote any graph of diameter  $D$ .

**Definition 16.1.** For all integer  $i$ , the  $i$ -th incidence matrix  $A_i \in \text{Mat}_X(\mathbb{C})$  satisfies

$$(A_i)_{xy} = \begin{cases} 1 & \text{if } \partial(x, y) = i, \\ 0 & \text{if } \partial(x, y) \neq i, \end{cases} \quad (x, y \in X).$$

Observe,

$$A_0 = I \quad (\text{identity}) \quad (16.1)$$

$$A_1 = A \quad (\text{adjacency matrix}) \quad (16.2)$$

$$A_0 + A_1 + \cdots + A_D = J \quad (\text{all 1's matrix}). \quad (16.3)$$

In general,  $A_i$  may not belong to Bose-Mesner algebra.

**Lemma 16.1.** Assume  $\Gamma = (X, E)$  is distance-regular with diameter  $D \geq 1$  and intersection numbers  $c_i, a_i, b_i$ .

(i)

$$AA_i = c_{i+1}A_{i+1} + a_iA_i + b_{i-1}A_{i-1}, \quad (0 \leq i \leq D, A_{-1} = A_{D+1} = O).$$

(ii)  $A_i = \frac{p_i(A)}{c_1 c_2 \cdots c_i}$ ,  $(0 \leq i \leq D)$ , where  $p_0, p_1, \dots, p_D$  are polynomials for the trivial module from Lemma 9.1.

(iii)  $A_0, A_1, \dots, A_D$  form a basis for Bose-Mesner algebra  $M$ .

(iv) For all distances  $h, i, j$   $(0 \leq i, j, h \leq D)$ , and for all vertices  $x, y \in X$  with  $\partial(x, y) = h$ , the constant

$$p_{i,j}^h = |\{z \in X \mid \partial(x, z) = i, \partial(y, z) = j\}|$$

depends only on  $h, i, j$  and not on  $x, y$ .

$$(v) \ E_0 = \frac{1}{|X|} J.$$

*Proof.*

(i) Pick  $x \in X$ . Apply each side to  $\hat{x}$ , we want to show that

$$AA_i \hat{x} = c_{i+1} A_{i+1} \hat{x} + a_i A_i \hat{x} + b_{i-1} A_{i-1} \hat{x}.$$

$$\text{LHS} = A \left( \sum_{y \in X, \partial(x, y) = i} \hat{y} \right) \tag{16.4}$$

$$= c_{i+1} \left( \sum_{z \in X, \partial(x, z) = i+1} \hat{z} \right) + a_i \left( \sum_{z \in X, \partial(x, z) = i} \hat{z} \right) + b_{i-1} \left( \sum_{z \in X, \partial(x, z) = i-1} \hat{z} \right) \tag{16.5}$$

$$= \text{RHS}. \tag{16.6}$$

(ii) Recall (Lemma 9.1)

$$Ap_i(A) = p_{i+1}(A) + a_i p_i(A) + b_{i-1} c_i p_{i-1}(A) \quad (0 \leq i \leq D).$$

Dividing by  $c_1 c_2 \cdots c_i$ , we have

$$A \frac{p_i(A)}{c_1 c_2 \cdots c_i} = c_{i+1} \frac{p_{i+1}(A)}{c_1 c_2 \cdots c_{i+1}} + a_i \frac{p_i(A)}{c_1 c_2 \cdots c_i} + b_{i-1} \frac{p_{i-1}(A)}{c_1 c_2 \cdots c_i}.$$

So,  $A_i, p_i(A)/(c_1 c_2 \cdots c_i)$  satisfy the same recurrence.

Also boundary condition,

$$A_0 = p_0(A) = I.$$

Hence,

$$A_i = \frac{p_i(A)}{c_1 c_2 \cdots c_i} \quad (0 \leq i \leq D).$$

(iii) Since  $E_0, E_1, \dots, E_D$  form a basis for  $M$ ,  $\dim M = D + 1$ .

Observe  $A_0, A_1, \dots, A_D \in M$  by (ii),  $A_0, A_1, \dots, A_D$  are linearly independent, since  $p_0, p_1, \dots, p_D$  are linearly independent.

Thus,  $A_0, A_1, \dots, A_D$  form a basis for  $M$ .

(iv)  $A_0, A_1, \dots, A_D$  form a basis for an algebra  $M$ ,

$$A_i A_j = \sum_{\ell=0}^D p_{ij}^{\ell} A_{\ell} \quad \text{for some } p_{ij}^{\ell} \in \mathbb{C}. \quad (16.7)$$

Fix  $h$  ( $0 \leq h \leq D$ ). Pick  $x, y \in X$  with  $\partial(x, y) = h$ .

Compute  $x, y$  entry in (16.7),

$$(A_i A_j)_{xy} = \sum_{z \in X} (A_i)_{xz} (A_j)_{zy} \quad (16.8)$$

$$= \sum_{z \in X, \partial(x, z)=i, \partial(y, z)=j} 1 \cdot 1 \quad (16.9)$$

$$= |\{z \in X \mid \partial(x, z) = i, \partial(y, z) = j\}|. \quad (16.10)$$

On the other hand,

$$\left( \sum_{\ell=0}^D p_{ij}^{\ell} A_{\ell} \right)_{xy} = p_{ij}^h (A_h)_{xy} = p_{ij}^h.$$

(v)  $\frac{1}{|X|}J$  is the orthogonal projection onto  $\text{Span}(\delta) = E_0 V$ . Hence,

$$\frac{1}{|X|} = E_0.$$

This proves the assertions. □

**Theorem 16.1.** *Let  $\Gamma = (X, E)$  be distance-regular with diameter  $D \geq 2$  and intersection numbers  $c_i, a_i, b_i$ . Pick a vertex  $x \in X$ . Let  $W$  be a thin irreducible  $T(x)$ -module with endpoint  $r = 1$  and diameter  $d$  ( $d = D - 2$  or  $D - 1$ ). Set  $\gamma_0 = a_0(W) + 1$ .*

(i) *The scalars*

$$\gamma_i := \frac{c_2 c_3 \cdots c_{i+1} b_2 b_3 \cdots b_{i+1} \gamma_0}{x_1(W) x_2(W) \cdots x_i(W)} \quad (0 \leq i \leq d) \quad (16.11)$$

*$a_i(W), x_i(W)$  are algebraic integers in  $\mathbb{Q}[\gamma_0]$ . In particular, if  $\gamma_0 \in \mathbb{Q}$ , then  $\gamma_i, a_i(W)$  and  $x_i(W)$  are integers for all  $i$ .*

(ii) *The numbers,  $\gamma_i, a_i(W), x_i(W)$  can all be determined from  $\gamma_0$  and the intersection numbers of  $\Gamma$  in order*

$$x_1(W), \gamma_1, a_1(W), x_2(W), \gamma_2, a_2(W), \dots$$

*using (i),*

$$x_i(W) = c_i b_i + \gamma_{i-1} (a_i + c_i - c_{i+1} - a_{i-1}(W)) \quad (1 \leq i \leq D - 1), \quad (16.12)$$

and

$$a_i(W) = \gamma_i - \gamma_{i-1} + a_i + c_i - c_{i+1} \quad (1 \leq i \leq D). \quad (16.13)$$

**Note.**

$$p_i = p_1^W + \gamma_{i-1}p_{i-1}^W - c_i(p_{i-1}^W + \gamma_{i-2}^W), \quad (\gamma_{-1} = -\gamma_{-2} = 0, \quad 0 \leq i \leq d+1).$$

*Proof.* Set

$$\tilde{A}_i = A_0 + A_1 + \cdots + A_i \quad (0 \leq i \leq D).$$

$$\text{Claim 1. } A\tilde{A}_i = c_{i+1}\tilde{A}_{i+1} + (a_i - c_{i+1} + c_i)\tilde{A}_i + b_i\tilde{A}_{i-1} \quad (0 \leq i \leq D-1).$$

*Proof of Claim 1.*

$$\text{LHS} = \sum_{j=0}^i AA_j \quad (16.14)$$

$$= \sum_{j=0}^i (c_{j+1}A_{j+1} + a_jA_j + b_{j-1}A_{j-1}) \quad (16.15)$$

$$= \sum_{j=0}^{i-1} A_j(c_j + a_j + b_j) + A_i(c_i + a_i) + A_{i+1}c_{i+1} \quad (16.16)$$

$$= k(A_0 + \cdots + A_{i-1}) + (a_i + c_i)A_i + c_{i+1}A_{i+1}. \quad (16.17)$$

$$\text{RHS} = c_{i+1}(A_0 + A_1 + \cdots + A_{i-1} + A_i + A_{i+1}) \quad (16.18)$$

$$+ (a_i - c_{i+1} + c_i)(A_0 + A_1 + \cdots + A_{i-1} + A_i) \quad (16.19)$$

$$+ b_i(A_0 + A_1 + \cdots + A_{i-1}) \quad (16.20)$$

$$= k(A_0 + \cdots + A_{i-1}) + A_i(a_i + c_i) + A_{i+1}c_{i+1}. \quad (16.21)$$

This proves Claim 1.

Now pick  $0 \neq w \in E_1^*(x)W$  and let

$$w = \sum_{z \in X, \partial(x,z)=1} \alpha_z \hat{z}.$$

Pick  $y$ , where  $\alpha_y \neq 0$ .

For  $i$  ( $0 \leq i \leq D$ ), define

$$B_i = \tilde{A}_i(\hat{x} - \hat{y}) \quad (16.22)$$

$$= \sum_{z \in X, \partial(x,z) \leq i} \hat{z} - \sum_{z \in X, \partial(y,z) \leq i} \hat{z} \quad (16.23)$$

$$= \sum_{z \in X, \partial(x,z)=i, \partial(y,z)=i+1} \hat{z} - \sum_{z \in X, \partial(y,z)=i+1, \partial(y,z)=i} \hat{z}. \quad (16.24)$$

Note that  $B_D = O$ ,  $B_0 = \hat{x} - \hat{y}$ , and

$$\langle B_0, w_0 \rangle = -\alpha_y \neq 0.$$

From Claim 1,

$$AB_i = c_{i+1}B_{i+1} + (a_i - c_{i+1} + c_i)B_i + b_iB_{i-1} \quad (0 \leq i \leq D), \quad B_{-1} = O.$$

Let  $p_0^W, \dots, p_d^W$  denote polynomials for  $W$  from Lemma 9.1. So,

$$w_i = p_i^W(A)w \in E_{1+i}^*(x)W, \quad (0 \leq i \leq d).$$

Claim 2.  $\langle w_i, B_j \rangle = 0$  if  $j \notin \{i, i+1\}$ ,  $(0 \leq i \leq d, 0 \leq j \leq D)$ .

*Proof of Claim 2.*

$$w_i \in E_{1+i}^*W, \quad B_j \in E_j^*(x)W + E_{j+1}^*(x)W.$$



Vertical lines indicate possible non-orthogonality.

Compute

$$\langle Aw_i, B_j \rangle = \langle w_i, AB_j \rangle, \quad quad(0 \leq i \leq D, 0 \leq j \leq D-1). \quad (16.25)$$

$$\text{LHS} = \langle w_{i+1}, B_j \rangle + a_i(W)\langle w_i, B_j \rangle + x_i(W)\langle w_{i-1}, B_j \rangle \quad (16.26)$$

$$\text{RHD} = b_j\langle w_i, B_{j-1} \rangle + (a_j - c_{j+1} + c_j)\langle w_i, B_j \rangle + c_{j+1}\langle w_i, B_{j+1} \rangle. \quad (16.27)$$

Evaluate for  $i = j-2, j-1, j, j+1$ .

Set  $i = j-2$ .



Then (16.25) becomes

$$\langle w_{j-1}, B_j \rangle = b_j \langle w_{j-2}, B_{j-1} \rangle \quad (2 \leq j \leq D-1).$$

By induction,

$$\langle w_{j-1}, B_j \rangle = b_2 b_3 \cdots b_j \langle w_0, B_1 \rangle \quad (1 \leq j \leq D-1).$$

Define

$$\gamma_0 = \frac{\langle w_0, B_1 \rangle}{\langle w_0, B_0 \rangle}.$$

(We will show  $\gamma_0 = 1 + a_0(W)$ .)

Then,

$$\langle w_{j-1}, B_j \rangle = b_2 b_3 \cdots b_j \gamma_0 \langle w_0, B_0 \rangle. \quad (16.28)$$

Set  $i = j + 1$ . Then (16.25) becomes

$$x_{j+1}(W) \langle w_j, B_j \rangle = c_{j+1} \langle w_0, B_{j+1} \rangle \quad (0 \leq j \leq d).$$

Hence,

$$\langle w_j, B_j \rangle = \frac{x_1(W) \cdots x_j(W)}{c_1 c_2 \cdots c_j} \langle w_0, B_0 \rangle \quad (0 \leq j \leq d). \quad (16.29)$$

Set  $i = j - 1$ . Then (16.25) becomes

$$\langle w_j, B_j \rangle + a_{j-1}(W) \langle w_{j-1}, B_j \rangle = (a_j - c_{j+1} + c_j) \langle w_{j-1}, B_j \rangle + b_j \langle w_{j-1}, B_{j-1} \rangle.$$

Evaluate this using (16.28) and (16.29). ( $\langle w_0, B_0 \rangle \neq 0$ ). Then we have

$$\frac{w_1(W) \cdots x_j(W)}{c_1 \cdots c_j} + (a_{j-1}(W) - a_j + c_{j+1} - c_j) b_2 \cdots b_j \gamma_0 = b_j \frac{x_1(W) \cdots x_{j-1}(W)}{c_1 \cdots c_{j-1}},$$

$$\left( \gamma_i := \frac{c_2 c_3 \cdots c_{i+1} b_2 b_3 \cdots b_{i+1} \gamma_0}{x_0(W) x_2(W) \cdots x_i(W)} \right).$$

$$\frac{x_j(W)}{c_j} = b_j + \frac{c_1 c_3 \cdots c_{j-1} b_2 b_3 \cdots b_j \gamma_0}{x_0(W) x_2(W) \cdots x_{j-1}(W)} (a_j + c_j - c_{j+1} - a_{j-1}).$$

So,

$$x_j(W) = c_j b_j + \gamma_{j-1} (a_j + c_j - c_{j+1} - a_{j-1}(W)).$$

This proves (16.12).

Set  $i = j$ . Then (16.25) becomes

$$a_j(W) \langle w_j, B_j \rangle + x_j(W) \langle w_{j-1}, B_j \rangle = (a_j - c_{j+1} + c_j) \langle w_j, B_j \rangle + c_{j+1} \langle w_j, B_{j+1} \rangle.$$

$$(a_j(W) - (a_j - c_{j+1} + c_j)) \frac{x_1(W) \cdots x_j(W)}{c_1 \cdots c_j} x_j(W) b_2 \cdots b_j \gamma_0 - c_{j+1} b_2 \cdots b_{j+1} \gamma_0 = 0.$$

Thus,

$$a_j(W) - (a_j - c_{j+1} + c_j) + \frac{c_1 \cdots c_j b_2 \cdots b_j \gamma_0}{x_1(W) \cdots x_{j-1}(W)} - \frac{c_1 \cdots c_j c_{j+1} b_2 \cdots b_{j+1} \gamma_0}{x_1(W) \cdots x_j(W)} = 0,$$

or

$$a_j(W) = a_j + c_j - c_{j+1} - \gamma_{j-1} + \gamma_j.$$

This proves (16.13).

Also by setting  $i = j = 0$ , we have

$$a_0(W) \langle w_0, B_0 \rangle = (a_0 - c_1 + c_0) \langle w_0, B_0 \rangle + c_1 \langle w_0, B_1 \rangle \quad (16.30)$$

$$= -\langle w_0, B_0 \rangle + \gamma_0 \langle w_0, B_0 \rangle. \quad (16.31)$$

Hence,

$$\gamma_0 = 1 + a_0(W).$$

Both  $a_i(W)$  and  $x_i(W)$  are algebraic integers, since they are eigenvalues of matrices with integer entries, namely,

$$E_{i+1}^*(x) A E_{i+1}^*(x) \quad \text{and} \quad E_i^*(x) A E_{i+1}^*(x) A E_i^*(x).$$

Also  $\gamma_0 = 1 + a_0(W)$  is an algebraic integer, and  $\gamma_i - \gamma_{i-1}$  is an algebraic integer by (16.12).

Hence,  $\gamma_i$  is an algebraic integer by induction.

This completes the proof of Theorem 16.1.  $\square$

**Example 16.1** ( $D=2$ ).

$$D = 2 \Leftrightarrow \text{strongly regular.}$$

Free parameters are  $k, a_1, c_2$ . Let  $W$  be an irreducible module of endpoint 1. The matrix representation of  $A|_W$  is

$$\begin{pmatrix} a_0(W) & x_1(W) \\ 1 & a_1(W) \end{pmatrix}.$$

$a_0(W)$ : free.

$$x_1(W) = c_1 b_1 + (a_0(W) + 1)(a_1 + c_1 - c_2 - a_0(W)) \quad (16.32)$$

$$= k - a_1 - 1 + a_1 a_0(W) + a_0(W) - c_2 a_0(W) - a_0(W)^2 + a_1 + a - c_2 - a_0(W) \quad (16.33)$$

$$= a_1 a_0(W) - c_2 a_0(W) + k - c_2 - a_0(W)^2, \quad (16.34)$$

$$\gamma_1 = 0, \quad (16.35)$$

$$a_1(W) = -(a_0(W) + 1) + a_1 + c_1 - c_2 \quad (16.36)$$

$$= -a_0(W) + a_1 - c_2. \quad (16.37)$$

Then the matrix has eigenvalues  $\theta, \theta_1$ . There is one feasible condition:  $a_0(W)$  is an algebraic integer.

**Example 16.2** (D=3). Free parameters  $c_2, c_3, k, a_1, a_2$ . The matrix representation becomes

$$A|_W = \begin{pmatrix} a_0(W) & x_1(W) & 0 \\ 1 & a_1(W) & x_2(W) \\ 0 & 1 & a_2(W) \end{pmatrix}.$$

Here,  $a_0(W)$  is free ( $= \gamma - 1$ )

$$x_1(W) = k - 1 - a_1 + \gamma_0(a_1 + 1 - c_2 - a_0(W)) \quad (16.38)$$

$$= \gamma_0(a_1 - c_2 - a_0(W)) + k - a_1 + a_0(W). \quad (16.39)$$

Set

$$\gamma_1(W) = \frac{c_2 b_2 \gamma_0}{x_1(W)}.$$

$$a_1(W) = \gamma_1 - \gamma_0 + a_1 + 1 - c_2 \quad (16.40)$$

$$x_2(W) = \gamma_1(a_2 - c_3 - a_1(W)) + c_2(\gamma_0 + b_1 - a_2 + a_1(W)) \quad (16.41)$$

$$a_2(W) = -\gamma_1 + a_2 + c_2 - c_2. \quad (16.42)$$

The matrix has eigenvalues,  $\theta, \theta_2, \theta_3$ .

There are two feasibility conditions;  $\gamma_0, \gamma_1$  are algebraic integers.

For arbitrary  $D$ , there are  $D - 1$  feasibility conditions;  $\gamma_0, \gamma_1, \dots, \gamma_{D-1}$  are algebraic integers.

**Lemma 16.2.** *With the notation of Theorem 16.1, suppose*

$$f_W = \frac{k - \lambda}{k} \quad (\text{so, } a_0(W) = -1).$$

Then,

$$a_i(W) = a_i + c_i - c_{i+1} \quad (0 \leq i \leq D - 1) \quad (16.43)$$

$$x_i(W) = b_i c_i \quad (1 \leq i \leq D - 1) \quad (16.44)$$

$$\gamma_i(W) = 0. \quad (16.45)$$

*Proof.* Since  $\gamma_0 = a_0(W) = 1$ ,  $\gamma_i = 0$ . □



## Chapter 17

# Association Schemes

Monday, March 1, 1993

### Review

Let  $\Gamma = (X, E)$  be a distance-regular graph of diameter  $D \geq 2$ . Pick a vertex  $x \in X$ .

Let  $W$  be a thin irreducible  $T(x)$ -module with endpoint  $r = 1$ , diameter  $d = D - 1$  or  $D - 2$ , and  $r_0 = a(W) + 1$ .

Show

$$\gamma_i = \frac{c_2 c_3 \cdots c_{i+1} b_2 b_3 \cdots b_{i+1} \gamma_0}{x_1(W) \cdots x_i(W)},$$

$a_i(W)$  and  $x_i(W)$  are all algebraic integers in  $\mathbb{Q}[\gamma_0]$ , where

$$x_i(W) = c_i b_i + \gamma_{i-1}(a_i + c_i - c_{i+1} - a_{i-1}(W)) \quad (1 \leq i \leq d) \quad (17.1)$$

$$a_i(W) = \gamma_i - \gamma_{i-1} + a_i + c_i - c_{i+1} \quad (1 \leq i \leq d) \quad (17.2)$$

Certainly,  $x_i(W)$ ,  $\gamma_i$ , and  $a_i(W)$  are in  $\mathbb{Q}[\gamma_0]$  by the above lines and so on.

$$\gamma_0 \rightarrow a_0(W) \rightarrow x_1(W) \rightarrow \gamma_1 \rightarrow a_1(W) \rightarrow x_1(W) \rightarrow \cdots$$

Recall some  $B \in \text{Mat}_n(\mathbb{C})$  is integral whenever

$$B \in \text{Mat}_n(\mathbb{Z}).$$

In this case, the characteristic polynomial

$$\det(\lambda I - B) = \lambda^n + \alpha_{n-1} \lambda^{n-1} + \cdots + \alpha_0, \quad \text{some } \alpha_0, \dots, \alpha_{n-1} \in \mathbb{Z}.$$

Hence, eigenvalues of  $B$  are algebraic integers. But  $a_i(W)$  is an eigenvalue of an integral matrices,

$$B = E_{i+1}^*(x) A E_{i+1}^*(x).$$

Hence,  $a_i(W)$  is an algebraic integer.

Also,  $x_i(W)$  is an eigenvalue of an integral matrix

$$B = E_i^*(x)AE_{i+1}^*(x)AE_i^*(x).$$

So  $x_i(W)$  is an algebraic integer.

$$\gamma_i - \gamma_{i-1} = a_i(W) - a_i - c_i + c_{i+1}$$

is an algebraic integer.

Since  $\gamma_0 = a_0(W) + 1$  is an algebraic integer, we find  $\gamma$  is an algebraic integer for all  $i$ .

**Definition 17.1.** A (commutative) association scheme is a configuration  $Y = (X, \{R_i\}_{0 \leq i \leq D})$ , where  $X$  is a finite nonempty set (of vertices),  $R_0, R_1, \dots, R_D$  are nonempty subsets of  $X \times X$  such that

- (i)  $R_0 = \{(x, x) \mid x \in X\}$ ,
- (ii)  $R_0 \cup \dots \cup R_D = X \times X$  (disjoint union),
- (iii) for every  $i$ ,  $R_i^\top = \{(y, x) \mid xy \in R\} = R_{i'}$  some  $i' \in \{0, 1, \dots, D\}$ ,
- (iv) for every  $h, i, j$  ( $0 \leq h, i, j \leq D$ ), and every  $x, y \in X$  such that  $(x, y) \in R_h$ ,

$$p_{ij}^h = |\{z \in X \mid (x, z) \in R_i, (z, y) \in R_j\}|$$

depends only on  $h, i, j$  and not on  $x, y$ ; and

- (v)  $p_{ij}^h = p_{ji}^h$  for all  $h, i, j$ .

If  $i' = i$  for all  $i$ , we say  $Y$  is symmetric. We call  $D$  the class of scheme and  $R_i$ , the  $i$ th relation of  $Y$ . We say vertices  $x, y \in X$  are  $i$ -related, or ‘at distance  $i$ ’, whenever  $(x, y) \in R_i$ .

We always assume that a ‘scheme’ is a commutative association scheme.

Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be an association scheme.

**Definition 17.2.** The  $i$ -the association matrix  $A_i \in \text{Mat}_X(\mathbb{C})$

$$(A_i)_{xy} = \begin{cases} 1 & \text{if } (x, y) \in R_i \\ 0 & \text{if } (x, y) \notin R_i, \end{cases} \quad (x, y \in X, 0 \leq i \leq D) \quad (17.3)$$

Then,

$$(i') A_0 = I.$$

$$(ii') A_0 + A_1 + \dots + A_D = J \text{ (= all 1's matrix).}$$

$$(iii') \quad A_i^\top = A_{i'} \quad (0 \leq i \leq D).$$

$$(iv') \quad A_i A_j = \sum_{h=0}^D p_{ij}^h A_h \quad (0 \leq i, j \leq D).$$

$$(v') \quad A_i A_j = A_j A_i.$$

$M := \text{Span}_{\mathbb{C}}(A_0, \dots, A_D)$  (Bose-Mesner algebra of  $Y$ ) is a commutative  $\mathbb{C}$ -algebra of dimension  $D + 1$ .

Observe:

$$Y \text{ is symmetric} \leftrightarrow A_i^\top = A_i \text{ for all } i \leftrightarrow M \text{ is symmetric.}$$

**Example 17.1.** Let  $\Gamma = (X, E)$  be distance-regular of diameter  $D$ . Set

$$R_i = \{(x, y) \mid \partial(x, y) = i\} \quad (0 \leq i \leq D). \quad (17.4)$$

Then,

$$Y = (X, \{R_i\}_{0 \leq i \leq D})$$

is a symmetric scheme.

$$i\text{-th association matrix} = i\text{-th distance matrix} \quad \text{for all } i.$$

**Example 17.2.** Suppose a group  $G$  acts transitively on a set  $X$ . Assume  $G$  is generously transitive, i.e.,

$$\text{for all } x, y \in X, \text{ there exists } g \in G \text{ such that } gx = y, gy = x.$$

Then  $G$  acts on  $X \times X$  by rule;

$$g(x, y) = (gx, gy), \quad \text{for all } g \in G, \text{ and for all } x, y \in X.$$

Let  $R_0, \dots, R_D$  denote orbits of  $G$  on  $X \times X$ .

Observe that  $R_i^\top = R_i$  for all  $i$  by generous transitivity, and

$$Y = (X, \{R_i\}_{0 \leq i \leq D})$$

is a symmetric scheme.

**Exercise 17.1.** In Example Example 17.2, Bose-Mesner algebra

$$M = \{B \in \text{Mat}_X(\mathbb{C}) \mid Bg = gB, \text{ for all } g \in G\} \quad (17.5)$$

$$= \text{the commuting algebra of } G \text{ on } X. \quad (17.6)$$

Here, we view each  $g \in G$  as a permutation matrix in  $\text{Mat}_X(\mathbb{C})$  satisfying

$$g\hat{x} = \widehat{gx}, \quad \text{for all } x \in G.$$

**Example 17.3.** Let  $G$  be any finite group.  $G$  acts on  $X = G$  by conjugation.

$$G \times X \rightarrow X, \quad (g, x) \mapsto gxg^{-1}.$$

Let  $C_0, C_1, \dots, C_D$  denote orbits (i.e., conjugacy classes), and let  $C_0 = \{1_G\}$ . Claim that  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  is a commutative scheme (not symmetric in general).

(i)  $R_0 = \{xx \mid x \in X\}$  as  $C_0 = \{1_G\}$ .

(ii)  $R_0, \dots, R_D$  is a partition of  $X \times X$  since  $C_0, \dots, C_D$  is a partition of  $X = G$ .

(iii)  $R_i^\top = R_{i'}$ , where  $C_{i'} = \{g^{-1} \mid g \in C_i\}$ .

(iv) Set  $H = G \oplus G$ , the direct sum. Then  $H$  acts on  $X = G$ :

$$\text{for all } h = (g, gz), \text{ for all } x \in X, \quad h(x) = gx(gx)^{-1} = gxz^{-1}g^{-1}.$$

$$R_i = \{(x, y) \mid x^{-1}y \in C_i\}, \quad h_i \in C_i, \quad x^{-1}y = gh_i g^{-1}.$$

$$(x, y) = (x, xgh_i g^{-1}) \tag{17.7}$$

$$= (xgg^{-1}, xgh_i g^{-1}) \tag{17.8}$$

$$= (xg, g)(1, h_i). \tag{17.9}$$

So,  $R_0, \dots, R_D$  are the orbits of  $H$  on  $X \times X$ .

(v)  $p_{ij}^h = p_{ji}^h$ ?

Fix  $i, j, h$  and  $x, y \in X$  with  $(x, y) \in R_h$ . Set

$$S = \{z \in X \mid (x, z) \in R_i, (z, y) \in R_j\} \tag{17.10}$$

$$T = \{z \in X \mid (x, z) \in R_j, (z, y) \in R_i\}. \tag{17.11}$$

Show  $|S| = |T|$ .

For all  $z \in S$ , set  $\hat{z} = xz^{-1}y$ .

Observe,  $\hat{z} \in T$ .

$$x^{-1}z \in C_i x^{-1}\hat{z} = x^{-1}xz^{-1}y \in C_j \tag{17.12}$$

$$z^{-1}y \in C_j \hat{z}^{-1}y = y^{-1}zx^{-1}x^{-1}y = y^{-1}x(x^{-1}z)x^{-1}y \in C_i. \tag{17.13}$$

Observe

$$S \rightarrow T \quad (z \mapsto z^{-1}) \quad \text{is one-to-one and onto.}$$

# Chapter 18

## Polynomial Schemes

Wednesday, March 3, 1993

**Lemma 18.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  denote the symmetric scheme with associated matrices  $A_0, A_1, \dots, A_D$ . Then the following are equivalent.*

(i) *The graph  $\Gamma = (X, R_1)$  is distance-regular, and  $R_0, \dots, R_D$  are labelled so that*

$$R_i = \{xy \mid \partial(x, y) = i\}.$$

(ii) *There exists  $f_i \in \mathbb{C}[\lambda]$ ,  $\deg f_i = i$  such that  $f_i(A_1) = A_i$  for all  $i$  with  $0 \leq i \leq D$ .*

(iii) *The parameter  $p_{ij}^h$*

$$\begin{cases} = 0 & \text{if one of } h, i, j \text{ is larger than the sum of the other two} \\ \neq 0 & \text{if one of } h, i, j \text{ is equal to the sum of the other two.} \end{cases}$$

*Proof.*

(i)  $\Rightarrow$  (ii): Lemma 16.1.

(ii)  $\Rightarrow$  (iii): Define

$$k_i \equiv p_{ii}^0 = |\{z \mid z \text{ in } X, \partial(x, z) = i \text{ } ((x, z) \in R_i)\}|$$

for any  $x \in X$ . Then  $k_i \neq 0$  ( $0 \leq i \leq D$ ),  $k_0 = 1$ .

(By symmetry,  $(x, y) \in R_i$  if and only if  $(y, x) \in R_i$ .)

Claim.

$$k_h p_{ij}^h = k_i p_{hj}^i = k_j p_{ih}^j \quad (18.1)$$

$$= |X|^{-1} |\{xyz \in X^3 \mid \partial(x, y) = h, \partial(x, z) = i, \partial(y, z) = j\}|. \quad (18.2)$$

*Pf.* The number of  $xyz \in X^3$ ,  $\partial(x, y) = h, \partial(x, z) = i, \partial(y, z) = j$  is equal to

$$|X| k_h p_{ij}^h = |X| k_i p_{hj}^i = k_j p_{ih}^j.$$

In particular,

$$p_{ij}^h = 0 \leftrightarrow p_{hj}^i = 0 \leftrightarrow p_{ih}^j = 0.$$

Hence, it suffices to show

$$\begin{cases} p_{ij}^h = 0 & \text{if } h > i + j \\ p_{ij}^h \neq 0 & \text{if } h = i + j. \end{cases}$$

Fix  $i, j$ . Without loss of generality, we may assume that  $i + j \leq D$  as trivial otherwise.

$$f_i(A) f_j(A) = A_i A_j = \sum_{\ell=0}^D p_{ij}^{\ell} A_{\ell} = \sum_{\ell=0}^D p_{ij}^{\ell} f_{\ell}(A).$$

$$i + j = \deg \text{LHS} \quad (18.3)$$

$$= \deg \text{RHS} \quad (18.4)$$

$$= \max\{\ell \mid p_{ij}^{\ell} \neq 0\}. \quad (18.5)$$

(iii)  $\Rightarrow$  (i)

Let  $A = A_1$ , and consider a graph  $\Gamma$  with adjacency matrix  $A$ .

$$A A_j = \sum_h p_{1j}^h A_h \quad (18.6)$$

$$= p_{1j}^{j+1} A_{j+1} + p_{1j}^j A_j + p_{1j}^{j-1} A_{j-1}. \quad (18.7)$$

Then,  $p_{1j}^{j+1} \neq 0 \neq p_{1j}^{j-1}$ .

Fix a vertex  $x \in X$ , and set  $R_i(x) = \{y \mid (x, y) \in R_i\}$ .

Then each  $y \in R_i(x)$  is adjacent in  $\Gamma$  to exactly

$$p_{1,i+1}^i \neq 0 \quad \text{vertices in } R_i(x), \quad (18.8)$$

$$p_{1i}^i \quad \text{vertices in } R_{i+1}(x), \quad (18.9)$$

$$p_{1,i-1}^i \neq 0 \quad \text{vertices in } R_{i-1}(x). \quad (18.10)$$

Hence, by induction,

$$R_i(x) = \{y \mid \partial(x, y) = i \text{ in } \Gamma\} \quad (0 \leq i \leq D), \quad (18.11)$$

and  $\Gamma$  is distance regular.

□

## Chapter 19

# Commutative Association Schemes

Friday, March 5, 1993

**Lemma 19.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme with Bose-Mesner algebra  $M$ .*

*Then there exists a basis  $E_0, E_1, \dots, E_D$  for  $M$  such that*

- (i)  $E_0 = |X|^{-1}J$ .
- (ii)  $E_i E_j = E_j E_i = \delta_{ij} E_i \quad (0 \leq i, j \leq D)$ .
- (iii)  $E_0 + E_1 + \dots + E_D = I$ .
- (iv)  $E_i^\top = \overline{E_i} = E_{\hat{i}}$  for some  $\hat{i} \in \{0, 1, \dots, D\}$ .

*Proof.*  $M$  acts on Hermitean space  $V = \mathbb{C}^n$  ( $n = |X|$ ).

If  $W$  is an  $M$ -module, so is  $W^\perp$ .

Each irreducible  $M$ -module is 1 dimensional by commutativity of  $M$ . So  $V$  is orthognal direct sum of 1-dimensional  $M$ -modules.

Let  $v_1, \dots, v_n$  be an orthonormal basis for  $V$  consisiting of eigenvectors for all  $m \in M$ .

Set  $P \in \text{Mat}_X(\mathbb{C})$  so that the  $i$ -th column of  $P$  is equal to  $v_i$ . So,

$$\bar{P}^\top P = I = P \bar{P}^\top = \bar{P} P^\top,$$

and  $P$  is unitary.

Also, for all  $m \in M$ ,

$$P^{-1}mP = \text{diagonal} \quad (19.1)$$

$$= \text{diag}(\theta_1(m), \dots, \theta_n(m)). \quad (19.2)$$

for some functions

$$\theta_i : M \longrightarrow \mathbb{C}.$$

Observe: each  $\theta = \theta_i$  is a character of  $M$ , i.e.,

$$\theta : M \longrightarrow \mathbb{C}$$

is a  $\mathbb{C}$ -algebra homomorphism.

Observe: the  $\theta_1, \dots, \theta_n$  are not all distinct.

Let  $\sigma_0, \dots, \sigma_r$  denote distinct elements of

$$\theta_1, \dots, \theta_n.$$

Say  $\sigma_i$  appears  $m_i$  times. Without loss of generality, we may assume that

$$P^{-1}mP = \begin{pmatrix} \sigma_0(m)I_{m_0} & O & O & O \\ O & \sigma_1(m)I_{m_1} & O & O \\ O & O & \ddots & O \\ O & O & O & \sigma_r(m)I_{m_r} \end{pmatrix}.$$

Set

$$E_i = P \begin{pmatrix} O & O & O \\ O & I_{m_i} & O \\ O & O & O \end{pmatrix} P^{-1},$$

where  $I_{m_i}$  is in the  $i$ -th block.

Then,

$$E_i E_j = \delta_{ij} E_i \quad (0 \leq i, j \leq r),$$

$$E_0 + E_1 + \dots + E_r = I.$$

Hence for all  $m \in M$ ,

$$m = \sum_{i=0}^r \sigma_i(m) E_i \in \text{Span}(E_0, \dots, E_r).$$

So,

$$M \subseteq \text{Span}(E_0, \dots, E_r).$$

Since  $E_0, \dots, E_r$  are linearly independent,  $r \geq D$ .

Show  $E_i \in M$ .

Claim 1. For all distinct  $i, j$  ( $0 \leq i, j \leq D$ ), there exists  $m \in M$  such that  $\sigma_i(m) \neq 0$ ,  $\sigma_j(m) = 0$ .



*Pf of Claim 1.*  $\sigma_i \neq \sigma_j$  implies that there exists  $m' \in M$  such that  $\sigma_i(m') \neq \sigma_j(m')$ .

Set  $m = m' - \sigma_j(m')I$ . Then,

$$\sigma_j(m)\sigma_j(m') - \sigma_j(m') = 0, \quad (19.3)$$

$$\sigma_i(m)\sigma_i(m') - \sigma_j(m') \neq 0. \quad (19.4)$$

Claim 2.  $E_i \in M$  ( $0 \leq i \leq D$ ).

*Pf of Claim 2.* Fix a vertex  $x \in X$ . For all  $j \neq i$ , there exists  $m_j \in M$  such that  $\sigma_i(m_j) \neq 0$ ,  $\sigma_j(m_j) = 0$ ,  $i \neq j$ . Observe

$$s = \sigma_i \left( \prod_{\ell \neq i} m_\ell \right) \neq 0.$$

Set

$$m^* = \sigma_i \left( \prod_{\ell \neq i} m_\ell \right) s^{-1}.$$

Observe

$$\sigma_i(m^*) = 1, \quad \sigma_j(m^*) = 0, \quad \text{for all } j \neq i \quad (0 \leq j \leq D).$$

So

$$P^{-1}m^*P = \begin{pmatrix} O & O & O \\ O & I_{m_i} & O \\ O & O & O \end{pmatrix}.$$

We have

$$E_i = m^* \in M.$$

Now  $r = D$ ,  $M = \text{Span}(E_0, \dots, E_D)$  and  $E_0, \dots, E_D$  is a basis for  $M$ .

Observe

$$P^{-1}E_iP = \begin{pmatrix} O & O & O \\ O & I_{m_i} & O \\ O & O & O \end{pmatrix}$$

implies

$$P^{-1}\overline{E_i}^\top P = \overline{P}^\top \overline{E_i}^\top \overline{P^{-1}}^\top = \begin{pmatrix} O & O & O \\ O & I_{m_i} & O \\ O & O & O \end{pmatrix}^\top = P^{-1}E_iP.$$

Hence,

$$\overline{E_i}^\top = E_i.$$

$E_0^\top, \dots, E_D^\top$  are nonzero matrices satisfying

$$E_i^\top E_j^\top = \delta_{ij} E_i^\top,$$

$$E_0^\top + E_1^\top + \cdots + E_D^\top = I.$$

Each  $E_i^\top$  is a linear combination of  $E_0, \dots, E_D$  with coefficientss that are 0 or 1, and for no two  $E_i$ 's are coefficients of any  $E_j$  both 1's.

So,  $E_0^\top, \dots, E_D^\top$  is a permutation of  $E_0, \dots, E_D$ .

Observe  $J = A_0 + \cdots + A_D \in M$ .

The matrix  $|X|^{-1}J$  is an idempotent of rank 1.

So, without loss of generality we may assume that

$$E_0 = \frac{1}{|X|}J.$$

We have the assertions. □

Define entry-wise product  $\circ$  on  $\text{Mat}_X(\mathbb{C})$ .

$$A_i \circ A_j = \delta_{ij}A_i.$$

So,  $M$  is closed under  $\circ$ .

$$E_i \circ E_j = \frac{1}{|X|} \sum_{h=0}^D q_{ij}^h E_h.$$

The numbers  $q_{ij}^h$  is called Krein parameters of  $Y$ .

Claim.  $q_{ij}^h \in \mathbb{R}$ .

*Pf.*

$$\frac{1}{|X|} \sum_{h=0}^D \overline{q_{ij}^h} E_h = \frac{1}{|X|} \sum_{h=0}^D \overline{q_{ij}^h} \overline{E_h}^\top \quad (19.5)$$

$$= (\overline{E_i} \circ \overline{E_j})^\top \quad (19.6)$$

$$= E_i \circ E_j \quad (19.7)$$

$$= \frac{1}{|X|} \sum_{h=0}^D q_{ij}^h E_h. \quad (19.8)$$

Hence,  $q_{ij}^h = \overline{q_{ij}^h}$ .

Observe  $A_0, \dots, A_D, E_0, \dots, E_D$  are bases for  $M$ . Hence, there exist  $p_i(j), q_i(j) \in \mathbb{C}$  such that

$$A_i = \sum_{j=0}^D p_i(j) E_j \quad (19.9)$$

$$E_i = \frac{1}{|X|} \sum_{j=0}^D q_i(j) A_j. \quad (19.10)$$

Taking transpose and conjugate we find,

$$\overline{p_i(j)} = p_i(j) = p_{i'}(\hat{j}) \quad (0 \leq i, j \leq D) \quad (19.11)$$

$$\overline{q_i(j)} = q_i(j) = q_{\hat{i}}(j') \quad (0 \leq i, j \leq D). \quad (19.12)$$

Fix a vertex  $x \in X$ . Define

$$E_i^* \equiv E_i^*(x) \in \text{Mat}_X(\mathbb{C})$$

to be a diagonal matrix such that

$$(E_i^*)_{xy} = \begin{cases} 1 & \text{if } (x, y) \in R_i \\ 0 & \text{if } (x, y) \notin R_i \end{cases} \quad (0 \leq i \leq D, y \in X.)$$

Then,

$$\begin{aligned} E_i^* E_j^* &= \delta_{ij} E_i^*, \\ E_0^* + \cdots + E_D^* &= I, \\ (E_i^*)^\top &= \overline{E_i^*} = E_i^*. \end{aligned}$$

**Definition 19.1.** Dual Bose-Mesner algebra  $M^* \equiv M^*(x)$  with respect to  $x$  is

$$\text{Span}(E_0^*, \dots, E_D^*).$$

Define dual associate matrices  $A_0^*, \dots, A_D^*$ . Indeed  $A_i^* \equiv A_i^*(x) \in \text{Mat}_X(\mathbb{C})$  is a diagonal matrix with

$$(A_i^*)_{yy} = |X|(E_i)_{xy} \quad (y \in X).$$

$A_i^*$  is a diagonal matrix having the row  $x$  of  $E_i^*$  on the diagonal.

Observe

$$A_i^* = \sum_{j=0}^D q_i(j) E_j^* \quad \left( E_i = \frac{1}{|X|} \sum_{j=0}^D q_i(j) A_j \right) \quad (19.13)$$

$$E_i^* = \frac{1}{|X|} \sum_{j=0}^D p_i(j) A_j^* \quad \left( A_i = \sum_{j=0}^D p_i(j) E_j \right). \quad (19.14)$$

So,  $A_0^*, \dots, A_D^*$  form a basis for  $M^*$ .

Also,

$$A_i^* E_j^* = q_i(j) E_j^*.$$

$$\left( A_i^* E_j^* = \sum_{h=0}^D q_i(h) E_h^* E_j^* = q_i(j) E_j^* \right)$$

So,  $q_i(j)$  are dual eigenvalues of  $A_i^*$ .

Observe,

$$A_0^* = I, \quad A_0^* + \cdots + A_D^* = |X|E_0^*, \quad \overline{A_i^*} = A_i^*,$$

$$A_i^* A_j^* = \sum_{h=0}^D q_{ij}^h A_h^* \quad (0 \leq i, j \leq D).$$

*Remark. Proof.*

$$(A_0^*)_{yy} = |X|(E_0)_{xy} = (J)_{xy} = 1.$$

$$A_0^* + \cdots + A_D^* = \sum_{i=0}^D \sum_{j=0}^D q_i(j) E_j^* = |X|E_0^*.$$

Note that

$$I = E_0 + \cdots + E_D = \frac{1}{|X|} \sum_{i=0}^D \sum_{j=0}^D q_i(j) A_j.$$

$$\sum_{i=0}^D q_i(j) = \delta_{j0}|X|.$$

$$\overline{A_i^*} = \sum_{j=0}^D \overline{q_i(j) E_j^*} = \sum_{j=0}^D q_i(j) E_j^* = A_i^*.$$

$$(A_i^* A_j^*)_{yy} = |X|^2 (E_i)_{xy} (E_j)_{xy} \tag{19.15}$$

$$= |X|^2 (E_i \circ E_j)_{xy} \tag{19.16}$$

$$= |X| \sum_{h=0}^D q_{ij}^h (E_h)_{xy} \tag{19.17}$$

$$= \sum_{h=0}^D q_{ij}^h (A_h^*)_{yy}. \tag{19.18}$$

The following statements will be proved after a couple of lemmas in the next lecture.

**Lemma.** Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme. Fix a vertex  $x \in X$ , and set  $E^* \equiv E_i^*(x)$  and  $A_i^* \equiv A^*(x)$ . Then the following hold.

(i)  $E_i^* A_j E_k^* = O$  if and only if  $p_{ij}^k = 0$  for  $0 \leq i, j, k \leq D$ .

(ii)  $E_i A_j^* E_k = O$  if and only if  $q_{ij}^k = 0$  for  $0 \leq i, j, k \leq D$ .

## Chapter 20

# Vanishing Conditions

**Monday, March 15, 1993** (Monday after Spring break)

**Lemma 20.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme.*

(i)  $p_0(i) = 1$ .

(ii)  $p_i(0) = k_i$ , where

$$k_i = p_{ii'}^0 = |\{y \in X \mid (x, y) \in R_i\}|.$$

(iii)  $q_0(i) = 1$ .

(iv)  $q_i(0) = m_i$ , where

$$m_i = \text{rank} E_i.$$

*Proof.*

(i) Since  $A_0 = I$  and

$$A_0 = p_0(0)E_0 + p_0(1)E_1 + \cdots + p_0(D)E_D \quad (20.1)$$

$$I = E_0 + E_1 + \cdots + E_D, \quad (20.2)$$

$p_0(i) = 1$  for all  $i$ .

(ii) Since

$$A_i = p_i(0)E_0 + p_i(1)E_1 + \cdots + p_i(D)E_D,$$

$A_i E_0 = p_i(0) E_0$ , and

$$k_i J = A_i J = p_i(0) J$$

as there are  $k_i$  1's in each row of  $A_i$ , we have  $k_i = p_i(0)$ .

(iii) Since  $E_0 = |X|^{-1} J$  and

$$E_0 = |X|^{-1} (q_0(0) A_0 + q_0(1) A_1 + \cdots + q_0(D) A_D) \quad (20.3)$$

$$|X|^{-1} J = |X|^{-1} (A_0 + A_1 + \cdots + A_D), \quad (20.4)$$

$q_0(i) = 1$  for all  $i$ .

(iv)  $E_i = |X|^{-1} (q_i(0) A_0 + q_i(1) A_1 + \cdots + q_i(D) A_D)$ ,  $E_i^2 = E_i$ , and  $E_i$  is similar to a matrix

$$\begin{pmatrix} I_{m_i} & O \\ O & O \end{pmatrix}.$$

So,

$$m_i = \text{rank} E_i = \text{trace} E_i = \sum_{x \in X} (E_i)_{xx} = |X| |X|^{-1} q_i(0) = q_i(0).$$

Note that as

$$E_i = \frac{1}{|X|} \sum_{j=0}^D q_i(j) A_j \rightarrow (E_i)_{xx} = \frac{1}{|X|} q_i(0) (A_0)_{xx}.$$

Hence, we have all formulas.

□

**Lemma 20.2.** *With the above notation*

$$(i) \ p_{ij}^h = p_{j'i'}^{h'}.$$

$$(ii) \ k_h p_{ij}^h = k_j p_{i'h}^j = k_{hj'}^i.$$

$$(iii) \ q_{ij}^h = q_{ji}^{\hat{h}}.$$

$$(iv) \ m_h q_{ij}^h = m_j q_{ih}^j = m_i q_{hj}^i.$$

*Proof.*

(i) We have

$$\sum_{h=0}^D p_{ij}^h A_{h'} \left( \sum_{h=0}^D p_{ij}^h A_h \right)^\top \quad (20.5)$$

$$= (A_i A_j)^\top \quad (20.6)$$

$$= A_j^\top A_i^\top \quad (20.7)$$

$$= A_{j'} A_{i'} \quad (20.8)$$

$$= \sum_{h=0}^D p_{j'i'}^{h'} A_h'. \quad (20.9)$$

(ii) Count the following number,

$$|\{xyz \in X^3 \mid (x, y) \in R_h, (x, z) \in R_i, (z, y) \in R_j\}| \quad (20.10)$$

$$= |X| k_h p_{ij}^h = |X| k_j p_{i'h}^j = |X| k_{hj'}^i. \quad (20.11)$$

(iii)

$$\frac{1}{|X|} \sum_{h=0}^D q_{ij}^h E_{\hat{h}} = \left( \frac{1}{|X|} \sum_{h=0}^D q_{ij}^h E_h \right)^\top \quad (20.12)$$

$$= (E_i \circ E_j)^\top \quad (20.13)$$

$$= E_j^\top \circ E_i^\top \quad (20.14)$$

$$= E_{\hat{j}} E_{\hat{i}} \quad (20.15)$$

$$= \frac{1}{|X|} \sum_{h=0}^D q_{\hat{j}\hat{i}}^{\hat{h}} E_{\hat{h}}. \quad (20.16)$$

(iv) Let  $\tau(B)$  denote the sum of the entries in the matrix  $B$ .

Observe:  $\tau(B \circ C) = \text{trace}(BC^\top)$ .

Observe

$$\tau(E_i \circ E_j \circ E_{\hat{k}}) = \tau((E_i \circ E_j \circ E_{\hat{k}})^\top) = \tau(E_{\hat{i}} \circ E_k \circ E_{\hat{j}}) = \tau(E_k \circ E_{\hat{j}} \circ E_{\hat{i}}).$$

Compute each one.

$$\tau(E_i \circ E_j \circ E_{\hat{k}}) = \text{trace}((E_i \circ E_j)E_k) = \text{trace} \left( \left( \frac{1}{|X|} \sum_h q_{ij}^h E_h \right) E_k \right) \quad (20.17)$$

$$= \text{trace} \left( \frac{1}{|X|} q_{ij}^k E_k \right) = \frac{1}{|X|} m_k q_{ij}^k, \quad (20.18)$$

$$\tau(E_{\hat{i}} \circ E_k \circ E_{\hat{j}}) = \text{trace}((E_{\hat{i}} \circ E_k)E_{\hat{j}}) = \text{trace} \left( \left( \frac{1}{|X|} \sum_h q_{ik}^h E_h \right) E_{\hat{j}} \right) \quad (20.19)$$

$$= \text{trace} \left( \frac{1}{|X|} q_{ik}^j E_k \right) = \frac{1}{|X|} m_j q_{ik}^j, \quad (20.20)$$

$$\tau(E_k \circ E_{\hat{j}} \circ E_{\hat{i}}) = \text{trace}((E_k \circ E_{\hat{j}})E_i) = \text{trace} \left( \left( \frac{1}{|X|} \sum_h q_{kj}^h E_h \right) E_i \right) \quad (20.21)$$

$$= \text{trace} \left( \frac{1}{|X|} q_{kj}^i E_i \right) = \frac{1}{|X|} m_i q_{kj}^i. \quad (20.22)$$

Hence, we have (iv). □

**Lemma 20.3.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme. Fix a vertex  $x \in X$ , and set  $E^* \equiv E_i^*(x)$  and  $A_i^* \equiv A^*(x)$ . Then the following hold.*

(i)  $E_i^* A_j E_k^* = O$  if and only if  $p_{ij}^k = 0$  for  $0 \leq i, j, k \leq D$ .

(ii)  $E_i A_j^* E_k = O$  if and only if  $q_{ij}^k = 0$  for  $0 \leq i, j, k \leq D$ .

*Proof.*

(i) Partition rows and columns by  $R_0(x), R_1(x), \dots, R_D(x)$ . Then,

$$E_i^*(x) A_j E_h^*(x)$$

is the  $(i, h)$  block of  $A_j$ .

Hence this submatrix is zero if and only if there exists no  $y, z \in X$  such that  $(x, y) \in R_i$ ,  $(x, z) \in R_h$  and  $(y, z) \in R_j$ . This is exactly when  $p_{ij}^h = 0$ .

(ii) The sum of the squares of norms of entries in  $E_i A_j^* E_k$



$$= \tau((E_i A_j^* E_k) \circ (\overline{E_j A_j^* E_k})) \quad (20.23)$$

$$= \text{trace}(E_i A_j^* E_k (\overline{E_j A_j^* E_k})^\top) \quad (20.24)$$

$$= \text{trace}(E_i A_j^* E_k A_j^* E_i) \quad (20.25)$$

$$= \text{trace}(E_i A_j^* E_k A_j^*) \quad \text{as } \text{trace}(XY) = \text{trace}(YX) \quad (20.26)$$

$$= \sum_{y \in X} (E_i A_j^* E_k A_j^*)_{yy} \quad (20.27)$$

$$= \sum_{y \in X} \left( \sum_{z \in X} (E_i)_{yz} (A_j^*)_{zz} (E_k)_{zy} (A_j^*)_{yy} \right) \quad (20.28)$$

$$= \sum_{y \in X} \left( \sum_{z \in X} (E_i)_{zy} (|X| (E_j)_{xz}) (E_k)_{zy} (|X| (E_j)_{yx}) \right) \quad (20.29)$$

$$= |X|^2 (E_j (E_i \circ E_k))_{xx} \quad (20.30)$$

$$= |X| q_{ik}^j (E_j)_{xx} \quad (20.31)$$

$$= q_{ik}^j m_j \quad (20.32)$$

$$= m_k q_{ij}^k. \quad (20.33)$$

Note that since  $|X|E_j = q_j(0)A_0 + q_j(1)A_1 + \cdots q_j(D)A_D$ ,

$$(E_j)_{xx} = \frac{1}{|X|} q_j(0) = \frac{m_j}{|X|}.$$

Thus, we have (ii). □

**Corollary 20.1** (Krein Condition). *For any commutative scheme  $Y = (X, \{R_i\}_{0 \leq i \leq D})$ ,  $q_{ij}^h$  is a non-negative real number for  $0 \leq h, i, j \leq D$ .*

*Proof.* Since  $q_{ij}^h m_h$  is a non-negative real by the proof of Lemma 20.3 (ii).

Note that  $m_h$  is a positive integer. □

An interpretation of the Krein parameters.

Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme with standard module  $V$ .

Pick a vector  $v \in V$  with

$$v = \sum_{x \in X} \alpha_x \hat{x}.$$

View  $v$  as a function

$$v : X \longrightarrow \mathbb{C} \quad (x \mapsto \alpha_x).$$

View  $V$  as the set of all functions  $V \rightarrow \mathbb{C}$ . Then the vector space  $V$  together with product of functions is a  $\mathbb{C}$ -algebra.

For

$$v = \sum_{x \in X} \alpha_x \hat{x}, \quad w = \sum_{x \in X} \beta_x \hat{x} \in V,$$

write

$$v \circ w = \sum_{x \in X} \alpha_x \beta_x \hat{x}$$

to represent the product of  $v$  and  $w$  viewed as functions.

**Lemma 20.4.** *With the above notation,*

- (i)  $A_j^*(x)v = |X|(E_j \hat{x} \circ v)$  for all  $v \in V$  and for all  $x \in X$ .
- (ii)  $E_i V \circ E_j V \subseteq \sum_{h: q_{ij}^h \neq 0} E_h V$  for all  $0 \leq i, j \leq D$ .
- (iii)  $E_h(E_i \circ E_j V) = E_h V$  if  $q_{ij}^h \neq 0$  for all  $0 \leq h, i, j \leq D$ .

## Chapter 21

# Norton Algebras

Wednesday, March 17, 1993

*Proof of Lemma 20.4.*

(i) Suppose

$$v = \sum_{x \in X} \alpha_x \hat{x}.$$

Pick a vertex  $z \in X$  and compare  $z$ -coordinate of each side in (i).

$$(A_j^*(x)v)_z = (A_j^*(x))_{zz}v_z = |X|(E_j)_{xz}\alpha_z. \quad (21.1)$$

$$|X|(E_{\hat{j}}\hat{x} \circ v)_z = |X|(E_{\hat{j}}\hat{x})_z \cdot \alpha_z = |X|(E_j)_{xz}\alpha_z. \quad (21.2)$$

Note that  $E_{\hat{j}}\hat{x}$  is the column  $x$  of  $E_{\hat{j}}$  is the row  $x$  of  $E_j$ .

(ii) Fix  $i, j, h$  such that  $q_{ij}^h = 0$ .

Claim.  $E_h(E_i V \circ E_j V) = 0$ .

$$E_h(E_i V \circ E_j V) = E_h(\text{Span}(v \circ w \mid v \in E_i V, w \in E_j V)) \quad (21.3)$$

$$= E_h(\text{Span}(E_i \hat{y} \circ E_j \hat{z} \mid y, z \in X)) \quad (21.4)$$

$$= \text{Span}(E_h(E_j \hat{z} \circ E_i \hat{y}) \mid y, z \in X) \quad (21.5)$$

$$= \text{Span}((E_h A_j^*(z) E_i) \hat{y} \mid y, z \in X) \quad \text{by (i)} \quad (21.6)$$

But  $q_{ij}^h = 0$  implies  $q_{ji}^{\hat{h}} = 0$ .

So, by Lemma 20.3 (ii),

$$0 = (E_{\hat{i}} A_{\hat{j}}^* E_{\hat{h}})^\top = E_h A_j^* E_i.$$

Hence,  $E_h(E_i V \circ E_j V) = 0$ .

(iii) Fix  $i, j, h$  such that  $q_{ij}^h \neq 0$ . Then,

$$E_h(E_i V \circ E_j V) \subseteq E_h V$$

is clear. We show the other inclusion. Since

$$E_i \hat{y} \circ E_j \hat{y} = (\text{column } y \text{ of } E_i \circ \text{column } y \text{ of } E_j) \quad (21.7)$$

$$= \text{column } y \text{ of } E_i \circ E_j \quad (21.8)$$

$$= (E_i \circ E_j) \hat{y} \quad (21.9)$$

$$= \left( \frac{1}{|X|} \sum_{h=0}^D q_{ij}^h E_h \right) \hat{y}, \quad (21.10)$$

we have,

$$E_h(E_i V \circ E_j V) = E_h \text{Span}(E_i \hat{y} \circ E_j \hat{z} \mid y, z \in X) \quad (21.11)$$

$$\supseteq E_h \text{Span}(E_i \hat{y} \circ E_j \hat{y} \mid y \in X) \quad (21.12)$$

$$= \text{Span}(q_{ij}^h E_h \hat{y} \mid y \in X) \quad (21.13)$$

$$= \text{Span}(E_h \hat{y} \mid y \in X) \quad \text{since } q_{ij}^h \neq 0 \quad (21.14)$$

$$= E_h V. \quad (21.15)$$

This proves the assertion. □

**Lemma 21.1.** *Given a commutative scheme  $Y = (X, \{R_i\}_{0 \leq i \leq D})$ , fix  $j$  ( $0 \leq j \leq D$ ). Define binary multiplication:*

$$E_j V \times E_j V \longrightarrow E_j V \quad ((v, w) \mapsto v * w = E_j(v \circ w)).$$

Then,

(i)  $v * w = w * v$ , for all  $v, w \in E_j V$ ,

(ii)  $v * (w + w') = v * w + v * w'$  for all  $v, w, w' \in E_j V$ , and

(iii)  $(\alpha v) * w = \alpha(v * w)$  for all  $\alpha \in \mathbb{C}$ .

In particular, the vector space  $E_j V$  together with  $*$  is a commutative  $\mathbb{C}$ -algebra, (not associative in general).

$(N_j : (E_j V, *))$  is called the Norton algebra on  $E_j V$ .

(iv)  $v * w = 0$  for all  $v, w \in E_j V$  if and only if  $q_{jj}^j = 0$ .

*Proof.*

(i) – (iii) Immediate.

(iv) Immediate from Lemma 20.4 (ii), (iii).

□

Let  $Y, j, N_j$  be as in Lemma 21.1, and  $M$  Bose-Mesner algebra of  $Y$ . Let

$$\text{Aut}Y = \{\sigma \in \text{Mat}_X(\mathbb{C}) \mid \sigma : \text{permutation matrix}, \sigma \cdot m = m \cdot \sigma \text{ for all } m \in M\} \quad (21.16)$$

$$= \{\sigma \in \text{Mat}_X(\mathbb{C}) \mid \sigma : \text{permutation matrix}, \quad (21.17)$$

$$(x, y) \in R_i \rightarrow (\sigma x, \sigma y) \in R_i, \text{ for all } i, \text{ and for all } x, y \in X\} \quad (21.18)$$

$$\text{Aut}(N_j) = \{\sigma : E_j V \rightarrow E_j V \mid \sigma \text{ is } \mathbb{C}\text{-algebra isomorphisms, i.e.,} \quad (21.19)$$

$$\sigma(v * w) = \sigma(v) * \sigma(w) \text{ for all } v, w \in E_j V\}. \quad (21.20)$$

**Lemma 21.2.** *Let  $Y, j, *$  be as in Lemma 21.1.*

(i)  $E_j V$  is a module for  $\text{Aut}(Y)$ .

(ii)  $\sigma|_{E_j V} \in \text{Aut}(N_j)$  for all  $\sigma \in \text{Aut}(Y)$ .

(iii)  $\text{Aut}Y \rightarrow \text{Aut}(N_j)$ ,  $(\sigma \mapsto \sigma|_{E_j})$  is a homomorphism of groups,

(i.e., a representation of  $\text{Aut}(Y)$ ).

(iv) Suppose  $R_0, \dots, R_D$  are orbits of  $\text{Aut}(Y)$  acting on  $X \times X$ , (so, we are in Example 17.2) then above representation is irreducible.

*Proof.*

(i) Pick  $\sigma \in \text{Aut}Y$  and  $v \in V$ . Then,

$$\sigma E_j v = E_j \sigma v,$$

since  $\sigma$  commutes with each element of  $M$ .

(ii)  $\sigma|_{E_j V} : E_j V \rightarrow E_j V$  is an isomorphism of a vector space. Since  $\sigma$  is invertible, for all  $v, w \in E_j V$ ,

$$\sigma(v * w) = \sigma(E_j(E_j v \circ E_j w)) = E_j \sigma(E_j v \circ E_j w) = E_j(E_j \sigma v \circ E_j \sigma w) = \sigma(v) * \sigma(w).$$

(iii) Immediate from (i) and (ii).

(iv) Here Bose-Mesner algebra  $M$  is the full commuting algebra, i.e.,

$$M = \{m \in \text{Mat}_X(\mathbb{C}) \mid \sigma \cdot m = m \cdot \sigma, \text{ for all } \sigma \in \text{Aut}(Y)\}.$$

Suppose there is a nonzero proper subspace  $0 \neq W \subsetneq E_j V$  that is  $\text{Aut}(Y)$ -invariant.

Set

$$W^\perp = \{v \in E_j V \mid \langle w, v \rangle = 0, \text{ for all } w \in W\}.$$

Then,  $W^\perp$  is a module for  $\text{Aut}(Y)$ , since  $\text{Aut}(Y)$  is closed under transpose conjugate.

Let  $e : V \rightarrow W$  and  $f : V \rightarrow W^\perp$  be orthogonal projection such that  $e + f = E_j$ ,

$$e^e = e, f^e = f, ef = fe = 0, eE_h = 0, \text{ if } h \neq j.$$

Since  $e$  commutes with all  $\sigma \in \text{Aut}(Y)$ ,  $e \in M$  and

$$e = \sum_{i=0}^D \alpha_i E_i.$$

If  $h \neq j$ , then  $0 = eE_h$  and  $\alpha_h = 0$ . Thus,  $e = \alpha_j E_j$ , i.e.,  $e = 0$  or  $f = 0$ .

A contradiction.

□

Norton algebras were used in original construction of Monster, a finite simple group  $G$ .

Compute character table of  $G$ ,

- $p_{ij}^h, q_{ij}^h$  of group scheme on  $G$ ,
- find  $j$  where  $m_j = \dim E_j V$  is small and  $q_{jj}^j \neq 0$ ,
- guess abstract structure of  $N_j$  using the knowledge of  $p_{ij}^h$ 's and  $q_{ij}^h$ 's,
- compute  $\text{Aut}(N_j)$ ,
- $G$ .

## Chapter 22

# Q-Polynomial Schemes

Friday, March 19, 1993

**Lemma 22.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme.*

$$(i) \ p_{0j}^h = p_{j0}^h = \delta_{jh}.$$

$$(ii) \ p_{ij}^0 = \delta_{ij} k_i.$$

$$(iii) \ q_{0j}^h = q_{j0}^h = \delta_{jh}.$$

$$(iv) \ q_{ij}^0 = \delta_{ij} m_i.$$

$$(v) \ \sum_{j=0}^D p_{ij}^h = k_i.$$

$$(vi) \ \sum_{j=0}^D q_{ij}^h = m_i.$$

*Proof.*

(i), (ii) These are trivial.

(iii) We have

$$|X|^{-1} \sum_{\ell=0}^D q_{0j}^{\ell} E_{\ell} = E_0 \circ E_j = |X|^{-1} J \circ E_j = |X|^{-1} E_j.$$

(iv) Recall from Lemma 20.2

$$|X|^{-1} m_h q_{ij}^h = \tau(E_i \circ E_j \circ E_{\hat{h}}),$$

(where  $\tau(B)$  is the sum of entries in matrix  $B$ .)

$$|X|^{-1}m_0q_{ij}^0 = \tau(E_i \circ E_j \circ E_0) \quad (22.1)$$

$$= |X|^{-1}\tau(E_i \circ E_j) \quad (E_0 = |X|^{-1}J) \quad (22.2)$$

$$= |X|^{-1}\text{trace}(E_i E_j) \quad (22.3)$$

$$= |X|^{-1}\delta_{ij}\text{trace}E_i \quad (22.4)$$

$$= |X|^{-1}\delta_{ij}m_i. \quad (22.5)$$

(v) Pick  $x, y \in X$  with  $(x, y) \in R_h$ . Then,

$$\{j=0\} \wedge D \quad p \wedge h\{ij\} \quad \& \quad = \quad |\{z \in X \mid (x,z) \in R_{\perp i}, \quad ; \quad (z,y) \in R_{\perp j} \quad ; \quad \text{for some } j\}| \quad \& \quad = \\ |\{z \in X \mid (x,z) \in R_{\perp i}\}| \quad \& \quad k_{\perp i}. \quad \backslash \text{end}\{\text{align}\}$$

(vi)

$$E_i \circ E_j = |X|^{-1} \sum_{h=0}^D q_{ij}^h E_h.$$

So,

$$\sum_{j=0}^D E_i \circ E_j = |X|^{-1} \sum_{h=0}^D \left( \sum_{j=0}^D q_{ij}^h \right) E_h \quad (22.6)$$

$$= E_i \circ \sum_{j=0}^D E_j \quad (22.7)$$

$$= E_i \circ I \quad (22.8)$$

$$= |X|^{-1}(q_i(0)A_0 + q_i(1)A_1 + \cdots + q_i(D)A_D) \circ I \quad (22.9)$$

$$= |X|^{-1}q_i(0)I \quad (22.10)$$

$$= |X|^{-1}m_i(E_0 + E_1 + \cdots + E_D). \quad (22.11)$$

This proves the assertions. □

**Definition 22.1.** Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme.

$Y$  is  $Q$ -polynomial with respect to ordering  $E_0, E_1, \dots, E_D$  of primitive idempotents, if

$$q_{ij}^h \begin{cases} = 0 & \text{if one of } h, i, j \text{ is greater than the sum of the other two} \\ \neq 0 & \text{if one of } h, i, j \text{ is equal to the sum of the other two.} \end{cases}$$

In this case, set

$$c_i^* = q_{1,i-1}^i, \quad a_i^* = q_{1,i}^i, \quad b_i^* = q_{1,i+1}^i \quad (0 \leq i \leq D), \quad (c_0^* = b_D^* = 0).$$



Observe:  $Q$ -polynomial  $\rightarrow Y$  is symmetric.

Suppose  $i \neq \hat{i}$  for some  $i$ . Then, by the condition in Definition 22.1,

$$0 = q_{i\hat{i}}^0 = m_i \ (\neq 0)$$

by Lemma 22.1 (iv). This is a contradiction.

Hence,  $E_i^\top = E_{\hat{i}} = E_i$  for all  $i$ .

Therefore  $M$  is symmetric and  $Y$  is symmetric.

Observe: If  $Y$  is  $Q$ -polynomial,

$$c_i^* + a_i^* + b_i^* = m_1 \quad (0 \leq i \leq D)$$

(just as  $c_i + a_i + b_i = k$  for  $P$ -polynomial.)

By Lemma 22.1 (iv),

$$m_1 = q_{10}^i + q_{11}^i + \cdots + q_{1,i-1}^i + q_{1i}^i + q_{1,i+1}^i + \cdots$$

and  $q_{10}^i = q_{11}^i = 0$ ,  $q_{1,i-1}^i = c_i^*$ ,  $q_{1i}^i = a_i^*$ , and  $q_{1,i+1}^i = b_i^*$ .

**Lemma 22.2.** Assume  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  is a symmetric scheme. Pick  $x \in X$ , and set  $E_i^* \equiv E_i^*(x)$ ,  $A^* \equiv A^*(x)$ . Then the following are equivalent.

- (i)  $\Gamma$  is  $Q$ -polynomial with respect to  $E_0, \dots, E_D$ .
- (ii)  $\begin{aligned} & \text{\$} q_{h-j} \text{\$} \text{ if } |h-j| > 0 \text{ \& } \\ & 0 \text{ \& } \text{ if } |h-j| = 1. \end{aligned} \quad (0 \leq h, j \leq D).$
- (iii) There exists  $f_i^* \in \mathbb{C}[\lambda]$ ,  $\deg f_i^* = i$ , and

$$A_i^* = f_i^*(A_1^*) \quad (0 \leq i \leq D).$$

(iv)  $E_0^*V, \dots, E_D^*V$  are maximal eigenspaces of  $A_1^*$ , and

$$E_i A_1^* E_j = 0 \quad \text{if } |i-j| > 0, \quad (0 \leq i, j \leq D).$$

(Compare (iv) with the definition of  $Q$ -polynomial in Definition 6.2.)

*Proof.*

(i)  $\rightarrow$  (ii) Clear.

(ii)  $\rightarrow$  (iii)  $A_0^* = I$ ,

$$A_i^* A_j^* = \sum_{h=0}^D q_{ij}^h A_h^* \quad (22.12)$$

$$A_1^* A_j^* = q_{1j}^{j-1} A_{j-1}^* + q_{1j}^j A_j^* + q_{1j}^{j+1} A_{j+1}^* \quad (q_{1j}^{j+1} \neq 0, 1 \leq j \leq D-1). \quad (22.13)$$

Hence  $A_j^*$  is a polynomial of degree exactly  $j$  in  $A_1^*$  by induction on  $j$ .

$$\lambda f_j^*(\lambda) = b_{j-1}^* f_{j-1}^*(\lambda) + a_j^* f_j^*(\lambda) + c_{j+1}^* f_{j+1}^*(\lambda) \quad \text{with } c_{j+1}^* \neq 0,$$

and  $f_{-1}^* = 0$ ,  $f_0^*(\lambda) = 1$ .

(iii)  $\rightarrow$  (i) Pick  $i, j, h$  with  $0 \leq i, j, h \leq D$  and  $h \geq i + j$ . Since

$$m_h q_{ij}^h = m_j q_{ih}^j = m_i q_{hj}^i$$

by Lemma 20.2, it suffices to show that

$$q_{ij}^h \begin{cases} = 0 & \text{if } h > i + j \\ \neq 0 & \text{if } h = i + j. \end{cases}$$

$$A_i^* A_j^* = \sum_{h=0}^D q_{ij}^h A_h^* \quad (22.14)$$

$$f_i^*(A_1) f_j^*(A_1) = \sum_{h=0}^D q_{ij}^h f_h^*(A_1). \quad (22.15)$$

Hence,

$$f_i^*(\lambda) f_j^*(\lambda) = \sum_{h=0}^D q_{ij}^h f_h^*(\lambda).$$

Note that since  $A_0^*, A_1^*, \dots, A_D^*$  are linearly independent,  $f(A_1^*) = 0$  implies  $\deg f > D$ .

$$\deg \text{LHS} = i + j \rightarrow q_{ij}^{i+j} \neq 0, q_{ij}^h = 0, \text{ if } h > i + j.$$

(iii)  $\rightarrow$  (iv) Recall

$$A_1^* = q_1(0)E_0^* + q_1(1)E_1^* + \dots.$$

Each  $A_i^*$  is a polynomial in  $A_1^*$ . Then  $A_1^*$  generates the dual Bose-Mesner algebra. So,  $q_1(0), q_1(1), \dots, q_1(D)$  are distinct.

So,  $E_0^* V, \dots, E_D^* V$  are maximal eigenspaces.

Also,  $|i - j| > 1$  implies  $q_{11}^j = 0$ .

Thus,  $E_i A_1^* E_j = 0$  by Lemma 20.3 (ii).

(iv)  $\rightarrow$  (ii)  $q_{1j}^i = 0$  if  $|i - j| > 1$ . since in this case,

$E_i A_1^* E_j = 0$  implies  $q_{1j}^i = 0$  by Lemma 20.3 (ii).

Suppose  $q_{1j}^{j+1} = 0$  for some  $j$  ( $0 \leq j \leq D - 1$ ).

Without loss of generalith, choose  $j$  minimum. Then  $A_h^*$  is a polynomial of degree  $h$  in  $A_1^*$  ( $0 \leq h \leq j$ ), and

$$A_1^* A_j^* - q_{1j}^{j-1} A_{j-1}^* - q_{1j}^j A_j^* = 0.$$

the left hand side is a polynomial in  $A_1^*$  of degree  $j + 1$ .

Hence, the minimal polynomial of  $A_1^*$  has degree less than or equal to  $j + 1 \leq D$ . But  $A_1^*$  has  $D + 1$  distinct eigenvalues.

This is a contradiction.

□



## Chapter 23

# Representation of a Scheme

Monday, March 22, 1993

**Theorem 23.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a symmetric scheme. (View the standard module  $V$  as an algebra of functions from  $X$  to  $\mathbb{C}$ .) Then the following are equivalent.*

- (i)  $Y$  is  $Q$ -polynomial with respect to ordering  $E_0, E_1, \dots, E_D$  of primitive idempotents.
- (ii) For all  $i$  ( $0 \leq i \leq D$ ),

$$E_0V + E_1V + (E_1V)^2 + \dots + (E_1V)^i = E_0V + E_1V + \dots + E_iV.$$

*Proof.*

By Lemma 20.4 (ii), (iii).

$$E_h(E_iV \circ E_jV) = 0 \text{ if and only if } q_{ij}^h = 0 \quad (0 \leq i, j, h \leq D).$$

(i)  $\rightarrow$  (ii) By our assumption,

$$q_{1j}^h = 0 \text{ if } |h - j| > 1, \text{ and } q_{1j}^{j+1} \neq 0.$$

So,

$$E_1V \circ E_jV \subseteq E_{j-1}V + E_jV + E_{j+1}V \quad (0 \leq j \leq D), \quad (23.1)$$

$$E_{j+1}(E_1V \circ E_jV) = E_{j+1}V \quad (0 \leq j \leq D-1), \quad (23.2)$$

by Lemma 20.4.

Also  $E_0V \subseteq \text{Span}(\delta)$ , where  $\delta$  is all 1's vector, i.e., 1 as a function  $X \rightarrow \mathbb{C}$ . So,

$$E_0 \circ E_j V = E_j V \quad (0 \leq j \leq D). \quad (23.3)$$

Show (ii) by induction on  $i$ .

The cases  $i = 0, 1$  are trivial.

$i > 1$ :  $\subseteq$ .

$$E_0V + E_1V + (E_1V)^2 + \cdots + (E_1V)^i \quad (23.4)$$

$$= E_0V + E_1V \circ (E_0V + E_1V + \cdots + (E_1V)^{i-1}) \quad (23.5)$$

$$= E_0V + E_1V \circ (E_0V + E_1V + \cdots + E_{i-1}V) \quad (23.6)$$

$$\subseteq E_0V + E_1V + \cdots + E_iV \quad (23.7)$$

by (23.1).

$\supseteq$ .

Claim.  $E_i \subseteq E_1V \circ E_{i-1}V + E_{i-1}V + E_{i-2}V \quad (2 \leq i \leq D)$ .

*Proof of Claim.* By (23.2),

$$E_i(E_1V \circ E_{i-1}V) = E_iV.$$

For all  $v \in E_iV$ , there exists  $u \in E_1V \circ E_{i-1}V$  such that  $E_iu = v$ .

On the other hand, by (23.1),

$$E_1V \circ E_{i-1}V \subseteq E_{i-2}V + E_{i-1}V + E_{i-2}V.$$

So,  $u = w + v$ , where  $w \in E_{i-2}V + E_{i-1}V$ . We have,

$$w = u - v \in E_1V \circ E_{i-1}V + E_{i-1}V + E_{i-2}V$$

as desired.

*Remark.*

$$E_iV \circ E_jV = \text{Span}(u \circ v \mid u \in E_iV, v \in E_jV).$$

By claim,

$$E_0V + E_1V + \cdots + E_iV \quad (23.8)$$

$$\subseteq E_0V + E_1V + \cdots + E_iV + E_1V \circ E_{i-1}V \quad (23.9)$$

$$\subseteq E_0V + E_1V + \cdots + (E_1V)^{i-1} + E_1V(E_0V + E_1V + \cdots + (E_1V)^{i-1}) \quad (23.10)$$

$$\subseteq E_0V + E_1V + \cdots + (E_1V)^{i-1} + (E_1V)^i. \quad (23.11)$$

(ii)  $\rightarrow$  (i)

Claim 1. Pick  $i, j$  ( $0 \leq i, j \leq D$ ) with  $j > i + 1$ . Then  $q_{1i}^j = 0$ .

*Proof of Claim 1.*

$$E_j(E_1 \circ E_j V) \subseteq E_j(E_1 V \circ (E_0 V + E_1 V + (E_1 V)^2 + \cdots + (E_1 V)^i)) \quad (23.12)$$

$$\subseteq E_j(E_0 V + E_1 V + (E_1 V)^2 + \cdots + (E_1 V)^{i+1}) \quad (23.13)$$

$$= E_j(E_0 V + E_1 V + \cdots + E_{i+1} V) \quad (23.14)$$

$$= 0. \quad (23.15)$$

So  $q_{1i}^j = 0$  by Lemma 20.4.

Claim 2.  $q_{1i}^{i+1} \neq 0$  ( $0 \leq i < D$ ).

*Proof of Claim 2.*

$$E_0 V + E_1 V + \cdots + E_{i+1} V \quad (23.16)$$

$$= E_0 V + E_1 V + \cdots + (E_1 V)^{i+1} \quad (23.17)$$

$$= E_0 V + E_1 V \circ (E_0 V + E_1 V + \cdots + (E_1 V)^i) \quad (23.18)$$

$$= E_0 V + E_1 V \circ (E_0 V + E_1 V + \cdots + E_i V) \quad (23.19)$$

$$= E_0 V + E_1 V \circ (E_0 V + \cdots + E_i V). \quad (23.20)$$

So,

$$E_{i+1} V = E_{i+1}(E_1 V \circ (E_0 V + \cdots + E_i V)) \quad (23.21)$$

$$= E_{i+1}(E_1 V \circ E_i V) \quad (23.22)$$

by Claim 1 and Lemma 20.4.

Hence,  $q_{1i}^{i+1} \neq 0$  by Lemma 20.4.

□

Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme with standard module  $V$ .

**Definition 23.1.** A representation of  $Y$  is a pair  $(\rho, H)$ , where  $H$  is a non-zero Hermitean space (with inner product  $\langle \cdot, \cdot \rangle$ ) and  $\rho : X \rightarrow H$  is a map satisfying the following.

R1.  $H = \text{Span}(\rho(x) \mid x \in X)$ .

R2.  $\langle \rho(x), \rho(y) \rangle$  depends only on  $i$  for which  $(x, y) \in R_i$  ( $x, y \in X$ ).

R3. For every  $x \in X$  and for all  $i$  ( $0 \leq i \leq D$ ),

$$\sum_{y \in X, (y, x) \in R_i} \rho(y) \in \text{Span}(\rho(x)).$$

Above representation is nondegenerate if  $\{\rho(x) \mid x \in X\}$  are distinct.

**Example 23.1.**  $Y = H(D, 2)$ ,  $X = \{a_1 \cdots a_D \mid a_i \in \{1, -1\}, 1 \leq i \leq D\}$ . Let  $H = \mathbb{C}^D$  and  $\langle \cdot, \cdot \rangle$  usual Hermitean dot product.

For a vertex  $x = a_1 \cdots a_D \in X$ , define

$$\rho(x) = a_1 \cdots a_D \in H.$$

Then, R1 – R3 hold.

*Remark.* R1, R2 are obvious. For R3, we may assume that  $x = 1 \cdots 1$ . Restrict

$$\sum_{y \in X, (y, x) \in R_i} \rho(y)$$

on the first coordinate. Then,

$$-1 \text{ appers } \binom{D-1}{i-1} \text{ times} \quad (23.23)$$

$$1 \text{ appers } \binom{D-1}{i} \text{ times.} \quad (23.24)$$

Hence,

$$\sum_{y \in X, (y, x) \in R_i} \rho(y) = \left( \binom{D-1}{i} - \binom{D-1}{i-1} \right) \rho(x).$$

Let  $(\rho, H)$  be a representation of arbitrary commutative scheme  $Y$ . Set

$$E = (\langle \rho(x), \rho(y) \rangle)_{x, y \in X}$$

Gram matrix of the representation.

**Definition 23.2.** Representations  $(\rho, H)$ ,  $(\rho', H')$  of  $Y$  are equivalent, whenever, Gram matrices are related by

$$E' \in \text{Span} E.$$

We do not distinguish between equivalent representations.

**Note.** Suppose  $(\rho, H)$  is a representation of a symmetric scheme  $Y$ . Pick  $x, y \in X$  with  $(x, y) \in R_j$ .

Then  $(y, x) \in R_j$ . So, by R2,

$$\langle \rho(x), \rho(y) \rangle = \langle \rho(y), \rho(x) \rangle = \overline{\langle \rho(x), \rho(y) \rangle},$$

since  $\langle \cdot, \cdot \rangle$  is Hermitean.

Hence, the Gram matrix  $E$  of  $\rho$  is real symmetirc. Without loss of generality, we can view  $H$  as a real Euclidean space in this case.



**Lemma 23.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme and  $V$  a standard module.*

*Let  $E_j$  be any primitive idempotent of  $Y$ .*

*(i)  $(\rho, H)$  is a representation of  $Y$ , where  $H = E_j V$  (with inner product inherited from  $Y$ ).*

$$\rho : X \rightarrow H \quad (x \mapsto E_j \hat{x})$$

*(i.e.,  $\rho(x)$  is the  $x$ -th column of  $E_j$ .)*

*(ii)  $\langle \rho(x), \rho(y) \rangle = |X|^{-1} q_j(i)$ , if  $(x, y) \in R_i$ ,  $(x, y \in X)$ .*

*(iii) For  $0 \leq i \leq D$  and  $x, y \in X$ ,*

$$\sum_{y \in X, (y, x) \in R_i} \rho(y) = p_i(j) \rho(x).$$

*(iv)  $(\rho, H)$  is nondegenerate if and only if  $q_j(i) \neq q_j(0)$  for all  $i$ ,  $(0 \leq i \leq D)$ .*

*(v) Every representation of  $Y$  is equivalent to a representation of the above type for some  $j$  ( $0 \leq j \leq D$ ), and  $j$  is unique.*

*Proof.*

*(i) – (iii).*

R1:  $\text{Span}(\rho X)$  is the column space of  $E_j$  which is equal to  $H$ .

R2:

$$\langle \rho(x), \rho(y) \rangle = \langle E_j \hat{x}, E_j \hat{y} \rangle \quad (23.25)$$

$$= (\overline{E_j \hat{x}})^\top E_j \hat{y} \quad (23.26)$$

$$= \hat{x}^\top \overline{E_j}^\top E_j \hat{y} \quad (23.27)$$

$$= \hat{x}^\top E_j \hat{y} \quad (23.28)$$

$$(E_j)_{xy}. \quad (23.29)$$

Note that  $\overline{E_j}^\top = E_j$  by Lemma 19.1.

Recall

$$E_j = |X|^{-1} (q_j(0)A_0 + \cdots + q_j(D)A_D).$$

So,

$$(E_j)_{xy} = |X|^{-1} q_j(i), \quad \text{where } (x, y) \in R_i.$$

R2: Recall

$$A_i = p_i(0)E_0 + \cdots + p_i(D)E_D.$$

So,  $E_j A_i = p_i(j) E_j$ , and

$$p_i(j) \rho(x) = p_i(j) E_j \hat{x} = E_j A_i \hat{x} = E_j \sum_{y \in X, (y, x) \in R_i} \hat{y} = \sum_{y \in X, (y, x) \in R_i} \rho(y).$$

**Note.**

$$A_i \hat{x} = \sum_{y \in X, (x, y) \in R_{i'}} \hat{y}.$$

*Pf.*

$$z \text{ entry of LHS} = (A_i \hat{x})_z \quad (23.30)$$

$$= \sum_{w \in X} (A_i)_{zw} \hat{x}_w \quad (23.31)$$

$$= (A_i)_{zx} \quad (23.32)$$

$$= \begin{cases} 1 & \text{if } (x, z) \in R_{i'} \\ 0 & \text{else.} \end{cases} \quad (23.33)$$

$$z \text{ entry of RHS} = \sum_{y \in X, (x, y) \in R_{i'}, z=y} 1 \quad (23.34)$$

$$= \begin{cases} 1 & \text{if } (x, z) \in R_{i'} \\ 0 & \text{else.} \end{cases} \quad (23.35)$$

(iv) By (ii),

$$\|\rho(x)\|^2 = \langle \rho(x), \rho(y) \rangle \quad (23.36)$$

$$|X|^{-1} q_j(0) \quad (23.37)$$

$$|X|^{-1} m_j, \quad (23.38)$$

as  $m_j = \dim E_j V$ , and is independent of  $x \in X$ .

Pick distinct  $x, y \in X$  such that  $(x, y) \in R_i$  with  $i \neq 0$ .

Then,

$$\rho(x) = \rho(y) \Leftrightarrow \langle \rho(x), \rho(y) \rangle = \|\rho(x)\|^2 = |X|^{-1} q_j(0) \quad (23.39)$$

$$\Leftrightarrow |X|^{-1} q_j(i) = |X|^{-1} q_j(0) \quad (23.40)$$

$$\Leftrightarrow q_j(i) = q_j(0). \quad (23.41)$$

Hence, we have (iv). To be continued.

□

## Chapter 24

# Balanced Conditions, I

Wednesday, March 23, 1993

No Class on Friday (another conference).

*Proof of Lemma 23.1 continued.* Let  $E_j$  be a primitive idempotent,  $H = E_j V$  and

$$\rho : X \rightarrow H \quad (x \mapsto E_j \hat{x}).$$

(v) Every representation  $(\rho, H)$  of  $Y$  is equivalent to a representation of above type, for some  $j$  ( $0 \leq j \leq D$ ) and  $j$  is unique.

Let  $E := (\langle \rho(x), \rho(y) \rangle)_{x, y \in X}$ .

By R2,

$$E = \sum_{i=0}^D \sigma_i A_i, \quad \text{some } \sigma_0, \sigma_1, \dots, \sigma_D \in \mathbb{C}.$$

Hence,  $E$  belongs to the Bose-Mesner algebra  $\mathcal{M}$  of  $Y$ .

We want to show that  $E$  is a scalar multiple of a primitive idempotent.

Fix  $x \in X$  and fix  $i$  ( $0 \leq i \leq D$ ).

By R3,

$$\sum_{y \in X, (y, x) \in R_i} \rho(y) = \alpha \rho(x), \quad \text{some } \alpha \in \mathbb{C}. \quad (24.1)$$

So,

$$k_i \bar{\sigma}_i = \left\langle \sum_{y \in X, (y, x) \in R_i} \rho(y), \rho(x) \right\rangle = \bar{\alpha} \langle \rho(x), \rho(x) \rangle = \bar{\alpha} \sigma_0.$$

Hence,  $\alpha$  is independent of  $x$ . In maatrix form (24.1) becomes

$$EA_i \hat{x} = \alpha E \hat{x}.$$

*Remark.*

$$Eu = Ev \Leftrightarrow \langle z, Eu \rangle = \langle z, Ev \rangle \text{ for all } z \in X \Leftrightarrow (Eu)_z = (Ev)_z \text{ for all } z \in X.$$

$$(EA_i \hat{x})_z = \left\langle \rho(z), \sum_{y \in X, (y, x) \in R_i} \rho(y) \right\rangle \quad (24.2)$$

$$= \alpha \langle \rho(z), \rho(x) \rangle \quad (24.3)$$

$$= (\alpha E \hat{x})_z. \quad (24.4)$$

Hence,

$$EA_i \hat{x} = \alpha E \hat{x}.$$

Since  $x$  is arbitrary,

$$EA_i = \alpha E.$$

So,

$$EA_i \in \text{Span} E \text{ and } EM = \text{Span} E.$$

We have  $E \in E_j$  for unique  $j$  ( $0 \leq j \leq D$ ). □

*Remark.*

$$E = \tau_0 E_0 + \cdots + \tau_D E_D, \quad \tau_j \in \mathbb{C} \quad (0 \leq j \leq D).$$

And, at least one of  $\tau_j$  is nonzero, and

$$\tau_j E_j = EE_j \in \text{Span} E.$$

So,

$$\tau_j E_j = E$$

as  $E_0, \dots, E_D$  are linearly independent.

Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a symmetric scheme, and let  $E$  be a primitive idempotent.

**Definition 24.1.**  $Y$  is  $Q$ -polynomial with respect to  $E$ , if and only if  $Y$  is  $Q$ -polynomial with respect to some ordering  $E_0, E_1, \dots, E_D$  of primitive idempotents, where  $E_0 = |X|^{-1}J$ , and  $E_1 = E$ .

**Theorem 24.1.** Assume  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  is  $P$ -polynomial (i.e.,  $(X, R_1)$  is distance-regular). Let  $E$  be any primitive idempotent of  $Y$ . Let  $(\rho, H)$  be the corresponding representation.

(i) The following are equivalent.

(ia)  $Y$  is  $Q$ -polynomial with respect to  $E$ .

(ib)  $(\rho, H)$  is nondegenerate and for all  $x, y \in X$ , and for all  $i, j$  ( $0 \leq i, j \leq D$ ),

$$\sum_{z \in X, (x, z) \in R_i, (y, z) \in R_j} \rho(z) - \sum_{z' \in X, (x, z') \in R_j, (y, z') \in R_i} \rho(z') \in \text{Span}(\rho(x) - \rho(y)).$$

(ic)  $(\rho, H)$  is nondegenerate and for all  $x, y \in X$ ,

$$\sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} \rho(z) - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} \rho(z') \in \text{Span}(\rho(x) - \rho(y)).$$

(ii) Wirte

$$E = |X|^{-1} \sum_{j=0}^D \theta_j^* A_j,$$

and suppose (ia) – (ic) hold. Then the coefficient in (ib) is

$$p_{ij}^h \frac{\theta_i^* - \theta_j^*}{\theta_0^* - \theta_h^*} \quad (1 \leq h \leq D, 0 \leq i, j \leq D).$$

*Proof.*

(ia)  $\rightarrow$  (ib) Without loss of generality, assume  $E \equiv E_1$ , and  $Y$  is  $Q$ -polynomial with respect to  $E$ .

Then by Lemma 22.2,  $\theta_0^*, \dots, \theta_D^*$  are distinct. So  $\theta_h^* \neq \theta_0^*$  for all  $h \in \{1, 2, \dots, D\}$ , and  $(\rho, H)$  is nondegenerate.

Fix  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$ ,  $A_i^* \equiv A_i^*(x)$ ,  $A^* \equiv A_1^*$ .

Let  $M$  be the Bose-Mesner algebra. Set

$$L = \{mA^*n - nA^*m \mid m, n \in M\}.$$

Claim 1.  $\dim L \leq D$ .

*Proof of Claim 1.*

$$L = \text{Span}(E_i A^* E_j - E_j A^* E_i \mid 0 \leq i < j \leq D) \quad (24.5)$$

$$= \text{Span}(E_i A^* E_{i+1} - E_{i+1} A^* E_i \mid 0 \leq i \leq D-1). \quad (24.6)$$

Since  $E_i A^* E_j = O$  if  $q_{ij}^1 = 0$  by Lemma 20.2 and Lemma 20.3, and this occurs if  $|i - j| > 1$  by  $Q$ -polynomial property.

Hence,  $\dim L \leq D$ .

Claim 2. (i)  $\{A^* A_h - A_h A^* \mid 1 \leq h \leq D\}$  is a basis for  $L$ . In particular,

(ii) there exist  $r_{ij}^h \in \mathbb{C}$  ( $1 \leq h \leq D, 0 \leq i, j \leq D$ ) such that

$$A_i A^* A_j - A_j A^* A_i = \sum_{h=1}^D r_{ij}^h (A^* A_h - A_h A^*).$$

*Proof of Claim 2.*

(i) The column  $x$  of  $A^* A_h - A_h A^*$  is a nonzero scalar  $\theta_h^* - \theta_0^*$  times the column  $x$  of  $A_h$ .

*Remark.*

$$((A^* A_h - A_h A^*) \hat{x})_y = E_{xy}(A_h)_{yx} - (A_h)_{yx} E_{xx} = (\theta_h^* - \theta_0^*)(A_h)_{yz}.$$

Also the column  $x$  of  $A_0, A_1, \dots, A_D$  are linearly independent.

Hence, the matrices given are linearly independent.

They are in  $L$  by construction, so they form a basis for  $L$  by Claim 1.

(ii) This is immediate since

$$A_i A^* A_j - A_j A^* A_i \in L, \quad \text{for all } i, j.$$

Claim 3.

$$r_{ij}^\ell = p_{ij}^\ell \left( \frac{\theta^* - \theta_j^*}{\theta_0^* - \theta_\ell^*} \right) \quad (1 \leq \ell \leq D, 0 \leq i, j \leq D).$$

*Proof of Claim 3.* Fix  $i, j$ ,

$$A_i A^* A_j - A_j A^* A_i - \sum_{h=1}^D r_{ij}^h (A^* A_h - A_h A^*) = 0.$$

Pick  $\ell$  ( $1 \leq \ell \leq D$ ). Pick  $y \in X$  such that  $(x, y) \in R_\ell$ .

$$(A_i A^* A_j)_{xy} = \sum_{z \in X} (A_i)_{xz} (A^*)_{zz} (A_j)_{zy} \quad (24.7)$$

$$= \sum_{z \in X, (x, z) \in R_i, (y, z) \in R_j} (A^*)_{zz} \quad (24.8)$$

$$= |X|^{-1} p_{ij}^\ell \theta_i^*. \quad (24.9)$$

Similarly,

$$(A_j A^* A_i)_{xy} = |X|^{-1} p_{ij}^\ell \theta_j^*.$$

$$(A^*A_h - A_hA^*)_{xy} = (A_0A^*A_h - A_hA^*A_0)_{xy} \quad (24.10)$$

$$= |X|^{-1}p_{0h}^\ell(\theta_0^* - \theta_h^*) \quad (24.11)$$

$$= \begin{cases} 0 & \text{if } \ell \neq h \\ |X|^{-1}(\theta_0^* - \theta_h^*) & \text{if } \ell = h. \end{cases} \quad (24.12)$$

Hence,

$$\sum_{h=1}^D r_{ij}^h (A^*A_h - A_hA^*)_{xy} = |X|^{-1}r_{ij}^\ell(\theta_0^* - \theta_\ell^*).$$

Comparing terms, we have

$$p_{ij}^\ell(\theta_i^* - \theta_j^*) - r_{ij}^\ell(\theta_0^* - \theta_\ell^*) = 0.$$

Claim 4. For all  $h$  ( $1 \leq h \leq D$ ), for all  $i, j$  ( $0 \leq i, j \leq D$ ), for all  $w, y \in X$ ,  $(w, y) \in R_h$ ,

$$\sum_{z \in X, (w, z) \in R_i, (y, z) \in R_j} \rho(z) - \sum_{z' \in X, (w, z') \in R_j, (y, z') \in R_i} \rho(z') - r_{ij}^h(\rho(w) - \rho(y)) = 0. \quad (24.13)$$

*Proof of Claim 4.* Set  $L = \langle \text{LHS of (24.13)}, \rho(x) \rangle$ . It suffices to show that  $L = 0$ .

Note that since  $x$  is arbitrary, if LHS of (24.13) is zero.

$$L = \sum_{z \in X, (w, z) \in R_i, (y, z) \in R_j} \langle \rho(z), \rho(x) \rangle - \sum_{z' \in X, (w, z') \in R_j, (y, z') \in R_i} \langle \rho(z'), \rho(x) \rangle \quad (24.14)$$

$$- r_{ij}^h \langle \rho(w) - \rho(y), \rho(x) \rangle \quad (24.15)$$

$$= |X|^{-1}(A_iA^*A_j)_{wy} - |X|^{-1}(A_jA^*A_i)_{wy} - |X|^{-1} \sum_{\ell=1}^D r_{ij}^\ell (A^*A_\ell - A_\ell A^*)_{wy} \quad (24.16)$$

$$= |X|^{-1} \text{times } wy \text{ entry of a matrix known to be zero by Claim 2} \quad (24.17)$$

$$= 0. \quad (24.18)$$

□

*Remark.*

$$|X|^{-1} \sum_{\ell=1}^D r_{ij}^\ell (A^*A_\ell - A_\ell A^*)_{wy} = |X|^{-1} r_{ij}^h (A^*A_h - A_h A^*)_{wy} \quad (24.19)$$

$$= r_{ij}^h (\langle \rho(x), \rho(w) \rangle - \langle \rho(x), \rho(y) \rangle) \quad (24.20)$$





## Chapter 25

# Balanced Conditions, II

Monday, March 29, 1993

*Proof of Theorem 24.1 continued.*

(ib)  $\rightarrow$  (ic) Obvious.

(ic)  $\rightarrow$  (ia) Without loss of generality, we may assume  $D \geq 3$ , else trivial.

*Remark.* The case  $D = 2$  should be treated somewhere, but the assumption  $D \geq 3$  is not used.

Fix  $w \in X$ , and write  $E_i^* \equiv E_i^*(w)$ ,  $A_i^* \equiv A_i^*(w)$ ,  $A^* \equiv A_1^*$ , and  $A_i$ ,  $i$ -th distance matrix. Set

$$E \equiv E_1 = |X|^{-1} \sum_{i=0}^D \theta_i^* A_i.$$

Since  $(\rho, H)$  is nondegenerate,

$$\theta_0^* \neq \theta_h^*, \text{ for all } h \in \{1, 2, \dots, D\}$$

See Lemma 23.1 (iv).

Claim 1. Pick  $h$  ( $1 \leq h \leq D$ ), and  $x, y$  with  $(x, y) \in R_h$ . Then

$$\sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} \rho(z) - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} \rho(z') = r_{12}^h (\rho(x) - \rho(y)),$$

where

$$r_{12}^h = p_{12}^h \frac{\theta_1^* - \theta_2^*}{\theta_0^* - \theta_h^*}.$$

*Proof of Claim 1.* By our assumption,

$$\sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} \rho(z) - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} \rho(z') = \alpha (\rho(x) - \rho(y)).$$

Hence,

$$|X|^{-1}p_{12}^h(\theta_1^* - \theta_2^*) = \left\langle \sum_{z \in X, (x,z) \in R_1, (y,z) \in R_2} \rho(z) - \sum_{z' \in X, (x,z') \in R_2, (y,z') \in R_1} \rho(z'), \rho(x) \right\rangle \quad (25.1)$$

$$= \alpha \langle \rho(x) - \rho(y), \rho(x) \rangle \quad (25.2)$$

$$= \alpha |X|^{-1}(\theta_0^* - \theta_h^*). \quad (25.3)$$

We have

$$\alpha = p_{12}^h \frac{\theta_1^* - \theta_2^*}{\theta_0^* - \theta_h^*}.$$

$$\text{Claim 2. } A_1 A^* A_2 - A_2 A^* A_1 = \sum_{h=1}^D r_{12}^h (A^* A_h - A_h A^*).$$

*Proof of Claim 2.* The  $xy$  entry of the LHS – RHS is

$$|X| \left\langle \sum_{z \in X, (x,z) \in R_1, (y,z) \in R_2} \rho(z) - \sum_{z' \in X, (x,z') \in R_2, (y,z') \in R_1} \rho(z') - r_{12}^h (\rho(x) - \rho(y)), \rho(w) \right\rangle,$$

where  $(x, y) \in R_h$ ,  $h = 1, 2, \dots, D$ , and the  $xy$  entry of the LHS – RHS is 0 if  $x = y$ .

But the vector on the left in the above inner product is 0 by Claim 1, so the inner product is 0.

Thus, the  $xy$  entry of the LHS – RHS is always 0, and we have Claim 2.

$$\text{Claim 3. } A^* A_3 - A_3 A^* \in \text{Span}(AA^* A_2 - A_2 A^* A, A^* A_2 - A_2 A^*, A^* A - AA^*).$$

*Proof of Claim 3.* Since  $p_{12}^h = 0$ , if  $h > 3$ , and  $p_{12}^h \neq 0$ , if  $h = 3$ , we have  $r_{12}^h = 0$  if  $h > 0$ , and  $r_{12}^h \neq 0$ , if  $h = 3$ . Note that  $\theta_1^* \neq \theta_2^*$ .

Now we are done by Claim 2.

Claim 4. There exist  $\beta, \gamma, \delta \in \mathbb{R}$  such that

$$0 = [A, A^2 A^* - \beta A A^* A + A^* A^2 - \gamma(AA^* + A^* A) - \delta A^*] \quad (25.4)$$

$$= A^3 A^* - A^* A^3 - (\beta + 1)(A^2 A^* A - AA^* A^2) - \gamma(A^2 A^* - A^* A^2) - \delta(AA^* - A^* A). \quad (25.5)$$

*Proof of Claim 4.* There exists  $f_i \in \mathbb{R}[\lambda]$ ,  $\deg f_i = i$  such that  $A_i = f_i(A_1)$ .

Writing  $A_2, A_3$  as polynomials in  $A$  in Claim 3 and simplifying, we find

$$A^3 A^* - A^* A^3 \in \text{Span}(A^2 A^* A - AA^* A^2, A^2 A^* - A^* A^2, AA^* - A^* A).$$

*Remark.* Let  $A_3 = \beta_3 A^3 + \beta_2 A^2 + \beta_1 A + \beta_0 I$  with  $\beta_3 \neq 0$ , and  $A_2 = \gamma_2 A^2 + \gamma_1 A + \gamma_0 I$ , with  $\gamma_2 \neq 0$ . Then

$$A^* A_3 - A_3 A^* = A^*(\beta_3 A^3 + \beta_2 A^2 + \beta_1 A + \beta_0 I) - (\beta_3 A^3 + \beta_2 A^2 + \beta_1 A + \beta_0 I)A^*. \quad (25.6)$$

$$A^3 A^* - A^* A^3 \in \text{Span}(A^* A_3 - A_3 A^*, A^2 A^* - A^* A^2, AA^* - A^* A) \quad (25.7)$$

$$\subseteq \text{Span}(AA^* A_2 - A_2 A^* A, A^* A_2 - A_2 A^*, A^2 A^* - A^* A^2, AA^* - A^* A) \quad (25.8)$$

$$A^* A_2 - A_2 A^* = A^*(\gamma_2 A^2 + \gamma_1 A + \gamma_0 I) - (\gamma_2 A^2 + \gamma_1 A + \gamma_0 I)A^* \quad (25.9)$$

$$AA^* A_2 - A_2 A^* A = AA^*(\gamma_2 A^2 + \gamma_1 A + \gamma_0 I) - (\gamma_2 A^2 + \gamma_1 A + \gamma_0 I)A^* A \quad (25.10)$$

$$A^* A_2 - A_2 A^* \in \text{Span}(A^2 A^* - A^* A^2, AA^* - AA^*) \quad (25.11)$$

$$AA^* A_2 - A_2 A^* A \in \text{Span}(A^2 A^* A - AA^* A^2, AA^* - AA^*) \quad (25.12)$$

$$A^3 A^* - A^* A^3 \in \text{Span}(A^2 A^* A - AA^* A^2, A^2 A^* - A^* A^2, AA^* - A^* A). \quad (25.13)$$

Hence, we can find  $\delta, \gamma, \delta$  satisfying

$$0 = A^3 A^* - A^* A^3 - (\beta + 1)(A^2 A^* A - AA^* A^2) - \gamma(A^2 A^* - A^* A^2) - \delta(AA^* - A^* A).$$

On the other hand,

$$[A, A^2 A^* - \beta AA^* A + A^* A^2 - \gamma(AA^* + A^* A) - \delta A^*] \quad (25.14)$$

$$= A^3 A^* - A^2 A^* A - \beta A^2 A^* A + \beta AA^* A^2 + AA^* A^2 - A^* A^3 \quad (25.15)$$

$$- \gamma A^2 A^* - \gamma AA^* A + \gamma AA^* A + \gamma A^* A^2 - \delta AA^* + \delta A^* A \quad (25.16)$$

$$= A^3 A^* - A^* A^3 - (\beta + 1)(A^2 A^* A - AA^* A^2) - \gamma(A^2 A^* - A^* A^2) - \delta(AA^* - A^* A). \quad (25.17)$$

Thus we have (i) and (ii).

Define a diagram  $D_E$  on nodes  $0, 1, \dots, D$ .

Connect distinct nodes, by undirected arc if  $q_{ij}^1 \neq 0$ . (Note  $q_{ij}^1 = q_{ji}^1$ ).

Since  $q_{0j}^1 = \delta_{1j}$ , the 0-node is adjacent to the 1-node and no other node.

$Y$  is  $Q$ -polynomial with respect to  $E$  if and only if  $E_E$  is a path.

Claim 5.  $D_E$  is connected.

*Proof of Claim 5.* Suppose there exists  $\Delta \subseteq \{0, 1, \dots, D\}$  such that  $i, j$  not connected for every  $i \in \Delta$  and  $j \in \{0, 1, \dots, D\} \setminus \Delta$ .

Set

$$f = \sum_{i \in \Delta} E_i.$$

Observe

$$fA^* = \sum_{i \in \Delta} E_i A^* \left( \sum_{j=0}^D E_j \right) \quad (25.18)$$

$$= \sum_{i \in \Delta, j \in \Delta} E_i A^* E_j \quad (\text{since } E_i A^* E_j = O \text{ if } q_{ij}^1 = 0) \quad (25.19)$$

$$= fA^* f. \quad (25.20)$$

Also,  $A^* f = fA^* f$ .

Hence,  $f$  commutes with  $A^*$ .

But  $f$  is an element of the Bose-Mesner algebra

$$f = \sum_{i=0}^D \alpha_i A_i \quad \text{for some } \alpha_0, \dots, \alpha_D \in \mathbb{C}.$$

We have

$$0 = fA^* - A^* f = \sum_{i=1}^D \alpha_i (A_i A^* - A^* A_i).$$

But  $\{A_h A^* - A^* A_h \mid 1 \leq h \leq D\}$  are linearly independent. (The column  $w$  of  $A_h A^* - A^* A_h$  is  $\theta_h^* - \theta_0^*$  times the column  $w$  of  $A_h$ .)

Hence,  $\alpha_1 = \dots = \alpha_D = 0$ , and  $f = \alpha_0 I$ . Since  $f^2 = f$ ,  $\alpha_0$  is 0 or 1.

If  $\alpha_0 = 0$ ,  $f = O$  and  $\Delta = \emptyset$ .

If  $\alpha_0 = 1$ ,  $f = I$  and  $\Delta = \{0, 1, \dots, D\}$ .

This proves Claim 5. □

*Remark.* Claim 5 proves the following in general.

Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a symmetric association scheme. Fix a vertex  $x \in X$ , and let

$$E = \frac{1}{|X|} \sum_{j=0}^D \theta_j^* A_j \quad (\theta_j^* = q_1(j) \text{ if } E = E_1)$$

be a primitive idempotent and  $E_j^* \equiv E_j^*(x)$ .

$$A^* = \sum_{j=0}^D \theta_j^* E_j^*.$$

If  $\theta_0 = \theta_h^*$ ,  $h = 1, \dots, D$ , then the following hold.

- (i)  $\{A_h A^* - A^* A_h \mid 1 \leq h \leq D\}$  are linearly independent.
- (ii) The diagram  $D_E$  on nodes  $0, 1, \dots, D$  defined by

$$i \sim j \Leftrightarrow E(E_i \circ E_j) \neq O$$

is connected.

$$(iii) C_M(A^*) = \{L \in M \mid LA^* = A^*L\} = \text{Span}(I).$$

*Proof.* | (i) The column  $x$  of  $A_h A^* - A^*(A_h)$  is  $\theta_0^* - \theta_h^*$  times the column  $x$  of  $A_h$ .

$$(iii) 0 = [\sum_{h=0}^D \alpha_h A_h, A^*] = \sum_{h=1}^D \alpha_h (A_h A^* - A^* A_h). \text{ Hence, } \alpha_0 = \dots = \alpha_D = 0.$$

(ii)  $\Delta$  is a connected component. Let  $f = \sum_{i \in \Delta} E_i$ , then  $f \in C_M(A^*)$ .

Let  $Y = (X, \{R_i\}_{0 \leq i \leq 2})$  be a symmetric association scheme with  $D = 2$ . Let

$$E = \frac{1}{|X|} \sum_{j=0}^2 \theta_j^* A_j$$

be a primitive idempotent. If  $\theta_0^*, \theta_1^*, \theta_2^*$ .

Then  $Y$  is  $Q$ -polynomial with respect to  $E$ .

*Proof.* By the previous lemma,  $D_E$  is connected.

**Note.** It seems  $\theta_1^* \neq \theta_2^*$  is necessary. Clarify the condition  $\theta_1^* = \theta_2^*$ .

Terwilliger claims that  $\theta_1^* = \theta_2^*$  does not occur under the assumption (ic).  
(March 7, 1995)



## Chapter 26

# Representation Diagrams

Wednesday, March 31, 1993

*Proof of Theorem 24.1 continued.* Assume  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  is  $P$ -polynomial. Let  $E$  be a primitive idempotent of  $Y$  such that the corresponding representation  $(\rho, H)$  is nondegenerate.

Show for all  $x, y \in X$ ,

$$\sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} \rho(z) - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} \rho(z') \in \text{Span}(\rho(x) - \rho(y))$$

implies that  $Y$  is  $Q$ -polynomial with respect to  $E$ .

Define a diagram  $D_E$  on nodes  $0, 1, \dots, D$ , for  $i \neq j$ ,

$$i \frown j \leftrightarrow q_{ij}^1 \neq 0$$

by setting  $E = E_1$ .

We showed that  $0 \frown j \leftrightarrow j = 1$  ( $1 \leq j \leq D$ ) and  $D_E$  is connected.

Now it is sufficient to show the following.

Claim 6. Let  $i$  be a node in  $D_E$ . Then  $i$  is adjacent to at most 2 arcs.

*Proof of Claim 6.* Suppose the node  $j$  is adjacent to  $i$  in  $D_E$ . By claim 4,

$$0 = E_i(A^3A^* - A^*A^3 - (\beta + 1)(A^2A^*A - AA^*A^2) - \gamma(A^2A^* - A^*A) - \delta(AA^* - A^*A))E_j \quad (26.1)$$

$$= E_iA^*E_j(\theta_i^3 - \theta_j^3 - (\beta + 1)(\theta_i^2\theta_j - \theta_i\theta_j^2) - \gamma(\theta_i^2 - \theta_j^2) - \delta(\theta_i - \theta_j)) \quad (26.2)$$

$$= E_iA^*A_j(\theta_i - \theta_j)p(\theta_i, \theta_j), \quad (26.3)$$

where

$$p(s, t) = s^2 - \beta st + t^2 - \gamma(s + t) - \delta.$$

*Remark.*

$$(\theta_i - \theta_j)(\theta_i^2 - \beta\theta_i\theta_j + \theta_j^2 - \gamma(\theta_i + \theta_j) - \delta) \quad (26.4)$$

$$= \theta_i^3 - \theta_j^3 - (\beta + 1)(\theta_i^2\theta_j - \theta_i\theta_j^2) - \gamma(\theta_i^2 - \theta_j^2) - \delta(\theta_i - \theta_j) \quad (26.5)$$

Since  $i$  is adjacent to  $j$ ,  $q_{ij}^1 \neq 0$  and

$$E_i A^* E_j \neq O$$

by Lemma 20.3 (ii). Since  $Y$  is  $P$ -polynomial,

$$\theta_i \neq \theta_j \quad \text{if } i \neq j.$$

Hence  $p(\theta_i, \theta_j) = 0$ . But  $p$  is quadratic in  $t$ . So  $p(\theta_i, t) = 0$  has at most two solutions for  $\theta_j$ .

Now  $D_E$  is a pth, and  $\Gamma$  is  $Q$ -polynomial with respect to  $E$ .

This proves Theorem 24.1. □

**Corollary 26.1.** *Assume  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  is  $P$ -polynomial, and  $Q$ -polynomial with respect to a primitive idempotent*

$$E = \frac{1}{|X|} \sum_{i=0}^D \theta_i^* A_i.$$

*Then,*

$$\beta = \frac{\theta_i^* - \theta_{i+1}^* + \theta_{i+2}^* - \theta_{i+3}^*}{\theta_{i+1}^* - \theta_{i+2}^*}$$

*is independent of  $i$  ( $0 \leq i \leq D-3$ ).*

*Proof.* Fix  $i$ . Without loss of generality,  $D \geq 3$ , else vacuous.

Pick  $x, y \in X$  with  $(x, y) \in R_3$ .

Let  $(\rho, H)$  be the representation for  $E$ .

$$\sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} \rho(z) - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} \rho(z') = \frac{p_{12}^3(\theta_1^* - \theta_2^*)}{\theta_0^* - \theta_3^*} (\rho(x) - \rho(y)), \quad (26.6)$$

and  $p_{12}^3 = c_3$ .

Since  $p_{i, i+3}^3 \neq 0$ , there exists  $w \in X$  such that  $(x, w) \in R_{i+3}$ ,  $(y, w) \in R_i$ .

Take inner product of (26.6) with  $\rho(w)$ . We have

$$P_{12}^3(x, y) \subseteq P_{1, i+2}^{i+3}(x, w) \cap P_{2, i+2}^i(y, w) \quad (26.7)$$

$$P_{21}^3(x, y) \subseteq P_{2, i+1}^{i+3}(x, w) \cap P_{2, i+1}^i(y, w). \quad (26.8)$$



Hence,

$$\left\langle \sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} \rho(z) - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} \rho(z'), \rho(w) \right\rangle = |X|^{-1} c_3 (\theta_{i+2}^* - \theta_{i+1}^*),$$

$$\left\langle \frac{c_3(\theta_1^* - \theta_2^*)}{\theta_0^* - \theta_3^*} (\rho(x) - \rho(y)), \rho(w) \right\rangle = \frac{c_3(\theta_1^* - \theta_2^*)}{\theta_0^* - \theta_3^*} |X|^{-1} (\theta_{i+3}^* - \theta_{i+1}^*).$$

We have,

$$\sigma = \frac{\theta_{i+1}^* - \theta_{i+2}^*}{\theta_i^* - \theta_{i+3}^*} = \frac{\theta_1^* - \theta_2^*}{\theta_0^* - \theta_3^*}.$$

*Remark.* Note that since  $Y$  is  $P$  and  $Q$  with respect to  $A_1$  and  $E_1$ ,  $\theta_0^*, \theta_1^*, \dots, \theta_D^*$ ,  $\theta_0, \theta_1, \dots, \theta_D$  are all distinct.

So

$$\beta = \frac{1}{\sigma} - 1 = \frac{\theta_i^* - \theta_{i+1}^* + \theta_{i+2}^* - \theta_{i+3}^*}{\theta_{i+1}^* - \theta_{i+2}^*} = \frac{\theta_0^* - \theta_1^* + \theta_2^* - \theta_3^*}{\theta_1^* - \theta_2^*}.$$

We have the assertion.  $\square$

Given the intersection number of a distance-regular graph  $\Gamma$ . The following 2 lemmas give an efficient method to determine if  $\Gamma$  is  $Q$ -polynomial with respect to some primitive idempotent.

**Lemma 26.1.** *Let  $\Gamma$  be a distance-regular graph of diameter  $D \geq 1$ . Pick  $\theta, \theta_0^*, \theta_1^*, \dots, \theta_D^* \in \mathbb{R}$  such that  $\theta_0^* \neq 0$ , and set*

$$E = \frac{1}{|X|} \sum_{i=0}^D \theta_i^* A_i.$$

(i) *The following are equivalent.*

(ia)  *$\theta$  is an eigenvalue of  $\Gamma$ , and  $E$  is a corresponding primitive idempotent.*

(ib)

$$\begin{pmatrix} a_0 & b_0 & 0 & \cdots & \cdots & 0 \\ c_1 & a_1 & b_1 & 0 & \cdots & 0 \\ 0 & c_2 & a_2 & b_2 & \ddots & \vdots \\ \cdots & \ddots & \ddots & \ddots & \ddots & 0 \\ 0 & \cdots & 0 & c_{D-1} & a_{D-1} & b_{D-1} \\ 0 & \cdots & \cdots & 0 & c_D & a_D \end{pmatrix} \begin{pmatrix} \theta_0^* \\ \theta_1^* \\ \vdots \\ \vdots \\ \vdots \\ \theta_D^* \end{pmatrix} = \theta \cdots \begin{pmatrix} \theta_0^* \\ \theta_1^* \\ \vdots \\ \vdots \\ \vdots \\ \theta_D^* \end{pmatrix},$$

and  $\theta_0^* = \text{rank } E$ .

(ii) *Suppose (ia), (ib) hold. Then,*

$$\frac{\theta_1^*}{\theta_0^*}, \dots, \frac{\theta_D^*}{\theta_0^*}$$

can be computed from  $\theta$  using

$$\frac{\theta_i^*}{\theta_0^*} = \frac{p_i(\theta)}{kb_1 \cdots b_{i-1}}, \quad (1 \leq i \leq D),$$

where  $p_0 = 1$ ,  $p_1(\lambda) = \lambda$ , and

$$\lambda p_i(\lambda) = p_{i+1}(\lambda) + a_i p_i(\lambda) + b_{i-1} c_i p_{i-1}(\lambda) \quad (0 \leq i \leq D).$$

*Proof.*

(i) We have

$$(ia) \leftrightarrow (A - \theta I)E = O \text{ and } E^2 = E \quad (26.9)$$

$$\leftrightarrow 0 = \sum_{i=0}^D (A - \theta I) \theta_i^* A_i \text{ and } \text{rank} E = \text{trace} E = \theta_0^* \quad (26.10)$$

$$= \sum_{i=0}^D \theta_i^* (c_{i+1} A_{i+1} + a_i A_i + b_{i-1} A_{i-1} - \theta A_i) \quad (26.11)$$

$$= \sum_{j=0}^D A_j (c_j \theta_{j-1}^* + a_j \theta_j^* + b_j \theta_{j+1}^* - \theta \theta_j^*) \quad (26.12)$$

$$\leftrightarrow c_j \theta_{j-1}^* + a_j \theta_j^* + b_j \theta_{j+1}^* = \theta \theta_j^* \quad (0 \leq j \leq D) \text{ and } \text{rank} E = \theta_0^* \quad (26.13)$$

$$\leftrightarrow (ib). \quad (26.14)$$

*Remark.* The first  $\leftrightarrow$ .  $\rightarrow$  is clear.

$\leftarrow$ : By the first condition,  $AE = \theta E$ . So  $E$  is a scalar multiple of the primitive idempotent corresponding to  $\theta$ . Hence,  $\text{rank} E = \text{trace} E$  implies  $E$  is the primitive idempotent.

(ii) We prove by induction on  $i$ .

$i = 0$  is trivial.

$i = 1$ : Set  $j = 0$  above  $c_0 = 0, a_0 = 0, b_0 = k$ . We have

$$k\theta_1^* = \theta\theta_0^*.$$

So

$$\frac{\theta_1^*}{\theta_0^*} = \frac{\theta}{k} = \frac{p_1(\theta)}{k}.$$

$i \geq 2$ : Set  $j = i - 1$  above. We have

$$c_{i-2}\theta_{i-2}^* + a_{i-1}\theta_{i-1}^* + b_{i-1}\theta_i^* = \theta\theta_{i-1}^*.$$

So,

$$\frac{\theta_i^*}{\theta_0^*} = \frac{\theta\theta_{i-1}^* - a_{i-1}\theta_{i-1}^* - c_{i-1}\theta_{i-2}^*}{b_{i-1}\theta_0^*} \quad (26.15)$$

$$= \left( (\theta - a_{i-1}) \frac{\theta_{i-1}^*}{\theta_0^*} - c_{i-1} \frac{\theta_{i-2}^*}{\theta_0^*} \right) \frac{1}{b_{i-1}} \quad (26.16)$$

$$= \left( (\theta - a_{i-1}) \frac{p_{i-1}(\theta)}{kb_1 \cdots b_{i-2}} - c_{i-1} \frac{p_{i-2}(\theta)}{kb_1 \cdots b_{i-3}} \right) \frac{1}{b_{i-1}} \quad (26.17)$$

$$= \frac{p_i(\theta)}{kb_1 \cdots b_{i-2}b_{i-1}}, \quad (26.18)$$

as desired.

□



## Chapter 27

# $P$ -and $Q$ -Polynomial Schemes

Friday, April 2, 1993

**Theorem 27.1.** *Let  $\Gamma = (X, E)$  be a distance-regular graph of diameter  $D \geq 3$ .*

*Let  $\theta$  denote an eigenvalue of  $\Gamma$  with associated primitive idempotent*

$$E = \frac{1}{|X|} \sum_{i=0}^D \theta_i^* A_i.$$

*Then the following are equivalent.*

- (i)  $\Gamma$  is  $Q$ -polynomial with respect to  $E$ .
- (ii)  $\theta_0^* \neq \theta_h^*$  for all  $h \in \{1, 2, \dots, D\}$  and for  $i \in \{3, \dots, D\}$ ,

$$c_i \left( \theta_2^* - \theta_i^* - \frac{(\theta_1^* - \theta_{i-1}^*)^2}{\theta_0^* - \theta_i^*} \right) + b_{i-1} \left( \theta_2^* - \theta_{i-1}^* - \frac{(\theta_1^* - \theta_i^*)^2}{\theta_0^* - \theta_{i-1}^*} \right) \quad (27.1)$$

$$= (k - \theta)(\theta_1^* + \theta_2^* - \theta_{i-1}^* - \theta_i^*) - (\theta + 1)(\theta_0^* - \theta_2^*) \quad (27.2)$$

- (iii)  $\theta_0^* \neq \theta_h^*$  for all  $h \in \{1, 2, \dots, D\}$  and (27.2) holds for  $i = 3$ .

*Remark.* Note (27.2) is trivial for  $i = 1, 2$ .

$i = 1$ :

$$\text{LHS} = \left( \theta_2^* - \theta_1^* - \frac{\theta_1^* - \theta_0^*}{\theta^* - \theta_1^*} \right) + k(\theta^* - \theta_0^*) \quad (27.3)$$

$$= \theta_2^* - \theta_1^* - \theta_0^* + \theta_1^* + k(\theta_2^* - \theta_0^*) \quad (27.4)$$

$$= (k+1)(\theta_2^* - \theta_0^*) \quad (27.5)$$

$$\text{RHS} = (k-\theta)(\theta_1^* + \theta_2^* - \theta_0^* - \theta_1^*) - (\theta+1)(\theta_0^* - \theta_2^*) \quad (27.6)$$

$$= (k+1)(\theta_2^* - \theta_0^*). \quad (27.7)$$

$i = 2$ :

$$\text{LHS} = b_1 \left( \theta_2^* - \theta_1^* - \frac{\theta_1^* - \theta_0^*}{\theta^* - \theta_1^*} \right) \quad (27.8)$$

$$= b_1 \frac{(\theta_2^* - \theta_1^*)(\theta_0^* - \theta_1^* - \theta_2^* + \theta_1^*)}{\theta_0^* - \theta_1^*} \quad (27.9)$$

$$= b_1 \frac{(\theta_2^* - \theta_1^*)(\theta_0^* - \theta_2^*)}{\theta_0^* - \theta_1^*} \quad (27.10)$$

$$\text{RHS} = (\theta+1)(\theta_0^* - \theta_2^*). \quad (27.11)$$

Hence,

$$\text{LHS} = \text{RHS} \leftrightarrow b_1 \frac{\theta_2^* - \theta_1^*}{\theta_0^* - \theta_1^*} + (\theta+1) = 0 \quad (27.12)$$

$$= b_1(\theta_2^* - \theta_1^*) + (\theta+1)(\theta_0^* - \theta_1^*) = 0. \quad (27.13)$$

On the other hand,

$$b_1\theta_2^* + a_1\theta_1^* + c_1\theta_0^* = \theta\theta_1^* \quad (27.14)$$

$$b_1\theta_1^* + a_1\theta_1^* + c_1\theta_1^* = k\theta_1^*, \quad (27.15)$$

as  $\theta\theta_0^* = k\theta_1^*$  We have

$$b_1(\theta_2^* - \theta_1^*) + (\theta_0^* - \theta_1^*) = \theta(\theta_1^* - \theta_0^*).$$

*Proof.* Immediate from the proof of Theorem 2.1 in ‘A new inequality for distance-regular graphs’ (Terwilliger, 1995) and Theorem 24.1.  $\square$

**Note.** Suppose (i) – (iii) hold. In particular,  $\theta_0^*, \theta_1^*, \dots, \theta_D^*$  are distinct. Then,

$$c_i + a_i + b_i = k \quad (0 \leq i \leq D).$$

$$c_i\theta_{i-1}^* + a_i\theta_i^* + b_i\theta_{i+1}^* = \theta\theta_j^* \quad (0 \leq i \leq D).$$

$$\frac{\theta_i^* - \theta_{i+1}^* + \theta_{i+2}^* - \theta_{i-3}^*}{\theta_{i+1}^* - \theta_{i+2}^*} \quad \text{is independent of } i \quad (0 \leq i \leq D-3).$$

$$c_i \left( \theta_2^* - \theta_i^* - \frac{(\theta_1^* - \theta_{i-1}^*)^2}{\theta_0^* - \theta_i^*} \right) + b_{i-1} \left( \theta_2^* - \theta_{i-1}^* - \frac{(\theta_1^* - \theta_i^*)^2}{\theta_0^* - \theta_{i-1}^*} \right) \quad (27.16)$$

$$= (k - \theta)(\theta_1^* + \theta_2^* - \theta_{i-1}^* - \theta_i^*) - (\theta + 1)(\theta_0^* - \theta_2^*). \quad (27.17)$$

Furthermore, we can show for  $c_1, \dots, c_D, a_1, \dots, a_D, b_0, b_1, \dots, b_{D-1}$  in terms of 5 parameters.

In general, we can take the 5 parameters to be

$$D, q, s^*, r_1, r_2$$

and get

$$b_i = \frac{h(1 - q^{i-D})(1 - s^*q^{i+1})(1 - r_1q^{i+1})(1 - r_2q^{i+1})}{(1 - s^*q^{2i+1})(1 - s^*q^{2i+2})} \quad (0 \leq i \leq D), \quad (27.18)$$

$$c_i = \frac{h(1 - q^i)(1 - s^*q^{D+i+1})(r_1 - s^*q^i)(r_2 - s^*q^i)}{s^*q^D(1 - s^*q^{2i})(1 - s^*q^{2i+1})} \quad (0 \leq i \leq D), \quad (27.19)$$

$$a_i = b_0 - c_i - b_i \quad (0 \leq i \leq D), \quad (27.20)$$

where  $h$  variable is chosen so that  $c_1 = 1$ .

(We must also consider limiting cases  $h \rightarrow 0, s^* \rightarrow 0, q^* \rightarrow \pm 1$ .)

See Theorem 2.1 in “The subconstituent algebra of an association scheme, I, II, III, (Terwilliger, 1992), (Terwilliger, 1993a), (Terwilliger, 1993b).

**Definition 27.1.** Let  $\Gamma = (X, E)$  be a distance-regular graph of diameter  $D \geq 3$ . Choose  $q \in \mathbb{R} \setminus \{0, -1\}$ , set

$$\begin{bmatrix} i \\ 1 \end{bmatrix} = 1 + q + \dots + q^{i-1} = \begin{cases} \frac{q^i - 1}{q - 1} & q \neq 1 \\ i & q = 1. \end{cases}$$

**Definition 27.2.**  $\Gamma$  has classical parameters if

$$c_i = \begin{bmatrix} i \\ 1 \end{bmatrix} \left( 1 + \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \quad (27.21)$$

$$b_i = \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left( \sigma - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \quad (27.22)$$

for some  $\sigma, \alpha \in \mathbb{R}$ .

(This happens for essentially all known families of distance-regular graphs with unbounded diameter, and is essentially equivalent to  $s^* = 0$ .)

**Lemma 27.1.** *With above notation, suppose (27.21), (27.22) hold. Then,*

(i)  $\theta = \frac{b_1}{q} - 1$  is an eigenvalue of  $\Gamma$  with  $\theta \neq k$ .

(ii) Let  $E = |X|^{-1} \sum_{i=0}^D \theta_i^* A_i$  be associated primitive idempotent. Then

$$\frac{\theta_i^*}{\theta_0^*} = 1 + \left( \frac{\theta}{k} - 1 \right) \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \quad (0 \leq i \leq D).$$

In particular,  $\theta_i^* \neq \theta_0^*$  for all  $i \in \{1, 2, \dots, D\}$ .

(iii)  $\Gamma$  is  $Q$ -polynomial with respect to  $E$ .

*Proof.*

(i), (ii). Need to check

$$c_i \theta_{i-1}^* + a_i \theta_i^* + b_i \theta_{i+1}^* = \theta \theta_i^* \quad (0 \leq i \leq D),$$

where  $a_i = k - c_i - b_i \quad (0 \leq i \leq D)$ .

(equivalently: check

$$c_i(\theta_{i-1}^* - \theta_i^*) + b_i(\theta_i^* - \theta_{i+1}^*) = (\theta - k)\theta_i^* \quad (0 \leq i \leq D), \quad (27.23)$$

where  $c_i, b_i, \theta_i^*, \theta$  are as given.)

*Remark.*

$$\theta = \frac{b_1}{q} - 1, \quad \frac{\theta_i^*}{\theta_0^*} = 1 + \left( \frac{\theta}{k} - 1 \right) \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i}, \quad b_0 = \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma.$$

$i = 0$ .

$$\frac{\theta_i^*}{\theta_0^*} = \frac{\theta}{k}, \quad -k \left( 1 - \frac{\theta_1^*}{\theta_0^*} \right) = -k \left( 1 - \frac{\theta}{k} \right) = \theta - k.$$

$$\frac{\theta_{i-1}^* - \theta_i^*}{\theta_0^*} = \left( \frac{\theta}{k} - 1 \right) \left( \begin{bmatrix} i-1 \\ 1 \end{bmatrix} q^{2-i} - \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \right) = - \left( \frac{\theta}{k} - 1 \right) q^{1-i}.$$

$$\theta - k = \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - 1 \right) (\sigma - \alpha)/q - 1 \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma = \begin{bmatrix} D-1 \\ 1 \end{bmatrix} (\sigma - \alpha) - 1 - \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma.$$



$$(c_i(\theta_{i-1}^* - \theta_i^*) + b_i(\theta_i^* - \theta_{i+1}^*) - (\theta - k)\theta_i^*)/\theta_0^* \quad (27.24)$$

$$= - \begin{bmatrix} i \\ 1 \end{bmatrix} \left( 1 + \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \left( \frac{\theta}{k} - 1 \right) q^{1-i} + \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left( \sigma - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left( \frac{\theta}{k} - 1 \right) q^{-i} \quad (27.25)$$

$$- (\theta - k) \left( 1 + \left( \frac{\theta}{k} - 1 \right) \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \right) \quad (27.26)$$

$$= \left( \frac{\theta}{k} - 1 \right) \left( - \begin{bmatrix} i \\ 1 \end{bmatrix} \left( 1 + \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) q^{1-i} + \begin{bmatrix} D-i \\ 1 \end{bmatrix} \left( \sigma - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \right) \quad (27.27)$$

$$- \left( \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma + \left( \begin{bmatrix} D-1 \\ 1 \end{bmatrix} (\sigma - \alpha) - 1 - \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma \right) \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \right) \quad (27.28)$$

$$= \left( \frac{\theta}{k} - 1 \right) \left( - \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} - \alpha \left( \begin{bmatrix} i \\ 1 \end{bmatrix} \begin{bmatrix} i-1 \\ 1 \end{bmatrix} q^{1-i} + \begin{bmatrix} D-i \\ 1 \end{bmatrix} \begin{bmatrix} i \\ 1 \end{bmatrix} - \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \right) \right) \quad (27.29)$$

$$+ \sigma \left( \begin{bmatrix} D-i \\ 1 \end{bmatrix} - \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} + \begin{bmatrix} D \\ 1 \end{bmatrix} \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \right) + \begin{bmatrix} i \\ 1 \end{bmatrix} q^{1-i} \quad (27.30)$$

Check  $\theta \neq k$ . Suppose  $\theta = k$ . Then

$$\frac{b_1}{q} - 1 = k, \quad \text{and} \quad q > 0.$$

By (27.21), (27.22),

$$qc_i - b_i - q(qc_{i-1} - b_{i-1}) = (k - \theta)q \quad (1 \leq i \leq D) \quad (27.31)$$

$$= 0. \quad (27.32)$$

*Remark.* With the notation of Lemma 27.1, we have the above equality in

general.

$$qc_i - b_i - q(qc_{i-1} - b_{i-1}) \quad (27.33)$$

$$= q \begin{bmatrix} i \\ 1 \end{bmatrix} \left( 1 + \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) - \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left( \sigma - \alpha \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \quad (27.34)$$

$$- q \left( q \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \left( 1 + \alpha \begin{bmatrix} i-2 \\ 1 \end{bmatrix} \right) - \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \left( \sigma - \alpha \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \right) \quad (27.35)$$

$$= \left( q \begin{bmatrix} i \\ 1 \end{bmatrix} - q^2 \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \quad (27.36)$$

$$+ \alpha \left( q \begin{bmatrix} i \\ 1 \end{bmatrix} \begin{bmatrix} i-1 \\ 1 \end{bmatrix} + \begin{bmatrix} D \\ 1 \end{bmatrix} \begin{bmatrix} i \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \quad (27.37)$$

$$- q^2 \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \begin{bmatrix} i-2 \\ 1 \end{bmatrix} - q \begin{bmatrix} D \\ 1 \end{bmatrix} \begin{bmatrix} i-1 \\ 1 \end{bmatrix} + q \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \quad (27.38)$$

$$+ \sigma \left( - \begin{bmatrix} D \\ 1 \end{bmatrix} + \begin{bmatrix} i \\ 1 \end{bmatrix} + q \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) \quad (27.39)$$

$$= q + \alpha \left( - \begin{bmatrix} i \\ 1 \end{bmatrix} + \begin{bmatrix} D \\ 1 \end{bmatrix} + q \begin{bmatrix} i-1 \\ 1 \end{bmatrix} \right) + \sigma(q^D - 1 + 1) \quad (27.40)$$

$$= q \left( 1 + \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \alpha + q^{D-1} \sigma \right) \quad (27.41)$$

$$= q \left( \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma - \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \sigma + \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \alpha + 1 \right) \quad (27.42)$$

$$= q \left( k - \frac{\begin{bmatrix} D \\ 1 \end{bmatrix} - 1}{q} (\sigma - \alpha) + 1 \right) \quad (27.43)$$

$$= q(k - \theta). \quad (27.44)$$

Hence,

$$qc_i - b_i = q(qc_{i-1} - b_{i-1}) \quad (1 \leq i \leq D) \quad (27.45)$$

$$= q^i(qc_0 - b_0) \quad (27.46)$$

$$= -q^i k. \quad (27.47)$$

If  $i = D$ ,  $qc_D = -q^D k$ ,  $c_D = -q^{D-1} k < 0$ , a contradiction.

(iii) Check the equation (ii) of Theorem 27.1 holds for  $i = 3$ .

□

*Remark.*  $\theta_0^* \neq \theta_h^*$  for all  $h \in \{1, 2, \dots, D\}$  and

$$c_3 \left( \theta_2^* - \theta_3^* - \frac{(\theta_1^* - \theta_2^*)^2}{\theta_0^* - \theta_3^*} \right) - b_2 \frac{(\theta_1^* - \theta_3^*)^2}{\theta_0^* - \theta_2^*} = (k - \theta)(\theta_1^* - \theta_3^*) - (\theta + 1)(\theta_0^* - \theta_2^*).$$

*Pf.*

$$\frac{\text{LHS}}{\theta_0^*} = \begin{bmatrix} 3 \\ 1 \end{bmatrix} \left( 1 + \alpha \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \left( 1 - \frac{\theta}{k} \right) \left( q^{-2} - \frac{q^{-2}}{\begin{bmatrix} 3 \\ 1 \end{bmatrix} q^{-2}} \right) \quad (27.48)$$

$$- \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \left( \sigma - \alpha \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \left( 1 - \frac{\theta}{k} \right) \frac{\left( \begin{bmatrix} 3 \\ 1 \end{bmatrix} q^{1-3} - 1 \right)^2}{\begin{bmatrix} 2 \\ 1 \end{bmatrix} q^{-1}} \quad (27.49)$$

$$= \left( 1 - \frac{\theta}{k} \right) \left( \left( 1 + \alpha \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) q^{-2} \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \left( \sigma - \alpha \begin{bmatrix} 2 \\ 1 \end{bmatrix} \right) \begin{bmatrix} 2 \\ 1 \end{bmatrix} q^{-3} \right) \quad (27.50)$$

$$= \left( 1 - \frac{\theta}{k} \right) \left( q^{-2} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \alpha \left( q^{-2} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + q^{-1} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} D-2 \\ 1 \end{bmatrix} \right) \quad (27.51)$$

$$- q^{-1} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} D-2 \\ 1 \end{bmatrix} \sigma \right) \quad (27.52)$$

$$\frac{\text{RHS}}{\theta_0^*} = \left( \begin{bmatrix} D \\ 1 \end{bmatrix} \sigma - \begin{bmatrix} D-1 \\ 1 \end{bmatrix} (\sigma - \alpha) + 1 \right) \left( 1 - \frac{\theta}{k} \right) \left( \begin{bmatrix} 3 \\ 1 \end{bmatrix} q^{-2} - 1 \right) \quad (27.53)$$

$$\begin{bmatrix} D-1 \\ 1 \end{bmatrix} (\sigma - \alpha) \left( 1 - \frac{\theta}{k} \right) \begin{bmatrix} 2 \\ 1 \end{bmatrix} q^{-1} \quad (27.54)$$

$$= \left( 1 - \frac{\theta}{k} \right) \left( q^{-2} \begin{bmatrix} 2 \\ 1 \end{bmatrix} + \begin{bmatrix} 2 \\ 1 \end{bmatrix} q^{-1} \sigma \left( q^{D-2} - \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \right) \right) \quad (27.55)$$

$$+ \begin{bmatrix} 2 \\ 1 \end{bmatrix} q^{-2} \alpha \left( \begin{bmatrix} D-1 \\ 1 \end{bmatrix} + q \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \right) \right) \quad (27.56)$$

$$= \left( 1 - \frac{\theta}{k} \right) \left( q^{-2} \begin{bmatrix} 2 \\ 1 \end{bmatrix} - \sigma q^{-1} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} D-2 \\ 1 \end{bmatrix} + \alpha q^{-2} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} 2 \\ 1 \end{bmatrix} \begin{bmatrix} D-1 \\ 1 \end{bmatrix} \right) \quad (27.57)$$

**Example 27.1.**  $Q$ -polynomial distance-regular graphs with classical parameters.

$D$ -cube:  $c_i = i$ ,  $b_i = D - i$

has classical parameters:  $(q, \alpha, \sigma) = (1, 0, 1)$ .

Johnson graph  $J(D, N)$  ( $N \geq 2D$ ):

$c_i = i^2$ ,  $b_i = (D-i)(N-D-i)$  has classical parameters  $(q, \alpha, \sigma) = (1, 1, N-D)$ .

$q$ -analogue of Johnson graph  $J_q(D, N)$  ( $D \geq 2D$ ):

$$c_i = \left( \frac{q^i - 1}{q - 1} \right)^2 = \begin{bmatrix} i \\ 1 \end{bmatrix}^2, \quad b_i = \frac{q(q^D - q^i)(q^{N-D} - q^i)}{(q - 1)^2}$$

has classical parameters

$$(q, \alpha, \sigma) = \left( q, q, \left( \frac{q^{N-D+1} - 1}{q - 1} \right) - 1 \right) = \left( q, q, \begin{bmatrix} N - D + 1 \\ 1 \end{bmatrix} - 1 \right).$$

*Remark.*

$$b_i = \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left( \begin{bmatrix} N - D + 1 \\ 1 \end{bmatrix} - 1 - q \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \quad (27.58)$$

$$= \left( \begin{bmatrix} D \\ 1 \end{bmatrix} - \begin{bmatrix} i \\ 1 \end{bmatrix} \right) \left( \begin{bmatrix} N - D + 1 \\ 1 \end{bmatrix} - \begin{bmatrix} i + 1 \\ 1 \end{bmatrix} \right) \quad (27.59)$$

$$= \frac{q(q^D - q^i)(q^{N-D} - q^i)}{(q - 1)^2}. \quad (27.60)$$

## Chapter 28

# The First Eigenspace of a $Q$ -DRG

Monday, April 5, 1993

**Lemma 28.1.** *Let  $\Gamma = (X, E)$  be distance-regular of diameter  $D \geq 3$  with standard module  $V$ . Suppose  $\Gamma$  is  $Q$ -polynomial with respect to a primitive idempotent  $E_1$ . Pick a vertex  $x \in X$ . Then*

$$E_1 V = \text{Span}\{E_1 \hat{y} \mid \partial(x, y) \leq 2\}.$$

*In particular,*

$$\dim E_1 V \leq 1 + k_1 + k_2.$$

*Proof.* Let  $\Delta = \{E_1 \hat{y} \mid \partial(x, y) \leq 2\}$ .

$E_1 V \supseteq \text{Span}\Delta$ : clear.

$E_1 V \subseteq \text{Span}\Delta$ : Pick a vertex  $y \in X$ . Show that  $E_1 \hat{y} \in \text{Span}\Delta$ .

Induction on  $h = \partial(x, y)$ .

Case  $h \leq 2$ .

$E_1 \hat{y} \in \text{Span}\Delta$  follows from construction.

Case  $h \geq 3$ .

Pick a vertex  $x' \in X$  such that

$$\partial(x, x') = h - 3, \quad \partial(x', y) = 3.$$

By Theorem 24.1.

$$\sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} E_1 \hat{z} - \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} E_1 \hat{z}' = r_{12}^3 (E_1 \hat{x} - E_1 \hat{y}),$$

$$r_{12}^3 = \frac{c_3(\theta_1^* - \theta_2^*)}{\theta_0^* - \theta_3^*} \neq 0.$$

So,  $E_1 \hat{y}$  in  $\text{Span}\{f, g, E_1 \hat{x}'\}$ , where

$$f = \sum_{z \in X, (x, z) \in R_1, (y, z) \in R_2} E_1 \hat{z}, \quad g = \sum_{z' \in X, (x, z') \in R_2, (y, z') \in R_1} E_1 \hat{z}'.$$

Observe that each  $z$  in the  $f$ -sum satisfies  $\partial(x, z) = h - 2$ .

So, by induction hypothesis

$$E_1 \hat{z} \in \text{Span} \Delta, \quad \text{or } f \in \text{Span} \Delta.$$

Observe that each  $z'$  in the  $g$ -sum satisfies  $\partial(x, z') = h - 1$ .

So by induction hypothesis

$$E_1 \hat{z}' \in \text{Span} \Delta, \quad \text{or } g \in \text{Span} \Delta.$$

Also  $\partial(x, x') = h - 3$  implies  $E_1 \hat{x}' \in \text{Span} \Delta$ .

Therefore  $E_1 \hat{y} \in \text{Span} \Delta$ . □

**Note.** Let  $\Gamma$ ,  $E_1$ ,  $x$  be as in Lemma 28.1.

Assume  $D \geq 4$ .

Observe that there are many linear dependences among

$$\{E_1 \hat{y} \mid y \in \Delta\},$$

where  $\Delta = \{y \in X \mid \partial(x, y) \leq 2\}$ .

Take any  $y \in X$  such that  $\partial(x, y) \geq 4$ .

More than one choice for  $x'$  in the proof of Lemma 28.1 implies

“more than one way to put  $E_1 \hat{y} \in \text{Span} E_1 \Delta$ .”

**Open Problem:**

(i) Give a precise description of the linear dependences among

$$\{E_1 \hat{y} \mid y \in \Delta\}.$$

(ii) Find a subset  $\Delta' \subseteq \Delta$  such that

$$\{E_1 \hat{y} \mid y \in \Delta'\}$$

is a basis for  $E_1 V$ , (or find some other ‘nice’ basis for  $E_1 V$ ).

**Conjecture 28.1.** *Let  $\Gamma$ ,  $E_1$ ,  $x$  be as in Lemma 28.1. Set*

$$\widetilde{X} = \{y \in X \mid \partial(x, y) \leq 2\}, \quad (28.1)$$

$$\widetilde{\partial} = \text{the restriction of the distance function } \partial \text{ to } \widetilde{X}. \quad (28.2)$$

*Then  $\Gamma$  is determined by  $\widetilde{X}$  and  $\widetilde{\partial}$ .*

*(There should be some canonical way to reconstruct  $\Gamma$  from  $\widetilde{X}$  and  $\widetilde{\partial}$ .)*





## Chapter 29

# Tridiagonal Pair $A, A^*$

Wednesday, April 7, 1993

### Introduction to Theorem 29.1

Let  $\Gamma = (X, E)$  be distance-regular with diameter  $D \geq 3$ .

Assume  $\Gamma$  is  $Q$ -polynomial with respect to  $E_1$ .

Fix a vertex  $x \in X$ . Write  $E_i^* \equiv E_i^*(x)$ ,  $A_i^* \equiv A_i^*(x)$ ,  $A^* = A_1^*$ .

We know for  $h, i, j$  ( $0 \leq h, i, j \leq D$ ),

$$E_i^* A_h E_j^* = O \leftrightarrow p_{ij}^h = 0 \quad (29.1)$$

$$E_i A_h^* E_j = O \leftrightarrow q_{ij}^h = 0. \quad (29.2)$$

Also, for  $h, i, j$  ( $0 \leq h, i, j \leq D$ ),

$$h < |i - j| \rightarrow p_{ij}^h = 0, \quad q_{ij}^h = 0 \quad (29.3)$$

$$h = |i - j| \rightarrow p_{ij}^h \neq 0, \quad q_{ij}^h \neq 0. \quad (29.4)$$

Some  $A_h$  (resp.  $A_h^*$ ) is a polynomial of degree exactly  $h$  in  $A$  (resp.  $A^*$ ), it follows, for  $h, i, j$  ( $0 \leq h, i, j \leq D$ ),

$$E_i^* A^h E_j^*, \quad E_i A^{*h} E_j \quad \begin{cases} = 0 & \text{if } h < |i - j| \\ \neq 0 & \text{if } h = |i - j|. \end{cases}$$

We saw that there exist  $\beta, \gamma, \delta \in \mathbb{R}$  such that

$$0 = [A, A^2 A^* - \beta A A^* A + A^* A^2 - \gamma(A A^* + A^* A) - \delta A^*].$$

In fact, there exist  $\beta, \gamma^*, \delta^* \in \mathbb{R}$  such that

$$0 = [A^*, A^{*2} A - \beta A^* A A^* + A A^{*2} - \gamma^*(A^* A + A A^*) - \delta^* A]$$

as well as will now show.

Let  $K$  denote any field. Let  $V$  denote any vector space over  $K$  of finite positive dimension. Let  $\text{End}_K(V)$  denote the  $K$ -algebra of all  $K$ -linear transformations  $V \rightarrow V$ .

**Theorem 29.1.** *Given semi-simple elements  $A, A^* \in \text{End}_K(V)$ , suppose*

$$E_i(A^*)^h E_j : \begin{cases} = 0 & \text{if } h < |i - j| \\ \neq 0 & \text{if } h = |i - j|. \end{cases} \quad (0 \leq h, i, j \leq D) \quad (29.5)$$

$$E_i^* A^h E_j^* : \begin{cases} = 0 & \text{if } h < |i - j| \\ \neq 0 & \text{if } h = |i - j|. \end{cases} \quad (0 \leq h, i, j \leq R) \quad (29.6)$$

for some ordering  $E_0, E_1, \dots, E_D$  of the primitive idempotents for  $A$ , and some ordering  $E_0^*, E_1^*, \dots, E_R^*$  of primitive idempotents for  $A^*$ . Then

(i)  $R = D$ .

(ii) There exist  $\beta, \gamma, \gamma^*, \delta, \delta^* \in \mathbb{K}$  such that

$$0 = [A, A^2 A^* - \beta A A^* A + A^* A^2 - \gamma(A A^* + A^* A) - \delta A^*] \quad (29.7)$$

$$= A^3 A^* - A^* A^3 - (\beta + 1)(A^2 A^* A - A A^* A^2) \quad (29.8)$$

$$- \gamma(A^2 A^* - A^* A^2) - \delta(A A^* - A^* A) \quad (29.9)$$

$$0 = [A^*, A^{*2} A - \beta^* A^* A A^* + A A^{*2} - \gamma^*(A^* A + A A^*) - \delta^* A] \quad (29.10)$$

$$= A^{*3} A - A A^{*3} - (\beta + 1)(A^{*2} A A^* - A^* A A^{*2}) \quad (29.11)$$

$$- \gamma^*(A^{*2} A - A A^{*2}) - \delta^*(A^* A - A A^*). \quad (29.12)$$

(iii) Let  $\theta_i$  (resp.  $\theta_i^*$ ) denote the eigenvalue of  $A$  (resp.  $A^*$ ) associated with  $E_i$  (resp.  $E_i^*$ ). Then,

$$\beta = \frac{\theta_i - \theta_{i+1} + \theta_{i+2} - \theta_{i+3}}{\theta_{i+1} - \theta_{i+2}} \quad (0 \leq i \leq D-3) \quad (29.13)$$

$$= \frac{\theta_i^* - \theta_{i+1}^* + \theta_{i+2}^* - \theta_{i+3}^*}{\theta_{i+1}^* - \theta_{i+2}^*} \quad (0 \leq i \leq D-3) \quad (29.14)$$

$$\gamma = \theta_i - \beta \theta_{i+1} + \theta_{i+2} \quad (0 \leq i \leq D-2) \quad (29.15)$$

$$\gamma^* = \theta_i^* - \beta \theta_{i+1}^* + \theta_{i+2}^* \quad (0 \leq i \leq D-2) \quad (29.16)$$

$$\delta = \theta_i^2 - \beta \theta_i \theta_{i+1} + \theta_{i+1}^2 - \gamma(\theta_i + \theta_{i+1}) \quad (0 \leq i \leq D-1) \quad (29.17)$$

$$\delta^* = \theta_i^{*2} - \beta \theta_i^* \theta_{i+1}^* + \theta_{i+1}^{*2} - \gamma^*(\theta_i^* + \theta_{i+1}^*) \quad (0 \leq i \leq D-1) \quad (29.18)$$

In particular,  $\beta, \gamma, \gamma^*, \delta, \delta^*$  are uniquely determined by  $A, A^*$  and the above ordering of their primitive idempotents, whenever  $D \geq 3$ .

∴{.proof}

(i) By symmetry, it suffices to show  $D \geq R$ . Suppose  $R > D$ .

Since  $A$  is semisimple with exactly  $D + 1$  distinct eigenvalues, the minimal polynomial of  $A$  has degree  $D + 1$ .

Since  $R \geq D + 1$ ,

$$A^R \in \text{Span}\{A^j \mid 0 \leq j \leq D\}.$$

Multiplying each term on the left by  $E_R^*$  and on the right by  $E_0^*$ , we find

$$E_R^* A^R E_0^* \in \text{Span}\{E_R^* A^j E_0^* \mid 0 \leq j \leq D\}. \quad (29.19)$$

But by (29.6), the left side of (29.19) is nonzero and the right side of (29.19) is 0, a contradiction.

Hence  $D \geq R$ .

(ii), (iii)

Recalling the definitions, we have

$$A = \sum_{i=0}^D \theta_i E_i, \quad (29.20)$$

$$A^* = \sum_{i=0}^D \theta_i^* E_i^*, \quad (29.21)$$

$$A E_i = E_i A = \theta_i E_i \quad (0 \leq i \leq D), \quad (29.22)$$

$$A^* E_i^* = E_i^* A^* = \theta_i^* E_i^* \quad (0 \leq i \leq D). \quad (29.23)$$

Claim 1. For all integers  $i, j, k, \ell$  ( $0 \leq i, j, k, \ell \leq D$ ) such that  $j + k \leq i - \ell$ ,

$$E_i^* A^j A^* A^k E_\ell^* = \begin{cases} \theta_{\ell+k}^* E_i^* A^{j+k} E_\ell^* & \text{if } j + k = i - \ell, \\ 0 & \text{if } j + k < i - \ell. \end{cases} \quad (29.24)$$

*Proof of Claim 1.* The product (29.24) eqia;s

$$E_i^* A^j \left( \sum_{h=0}^D \theta_h^* E_h^* \right) A^k E_\ell^* = \sum_{h=0}^D \theta_h^* E_i^* A^j E_h^* A^k E_\ell^*.$$

Now pick any  $h$  ( $0 \leq h \leq D$ ), where

$$E_i^* A^j E_h^* A^k E_\ell^* \neq O.$$

Tehn by (29.6),  $j \geq |i - h|$ , otherwise

$$E_i^* A^j E_h^* = O$$

and by (29.5),  $k \geq |h - \ell|$  otherwise

$$E_h^* A^k E_\ell^* = O.$$

Hence

$$j + k \geq |i - h| + |h - \ell| \geq |i - \ell| \geq i - \ell.$$

Now if  $j + k < i - \ell$ , we see there is no such  $h$ , so (29.24) holds.

(Pf. Suppose  $i = j + k + \ell$  with  $0 \leq i, j, k, \ell, h \leq D$ .

Then  $i \geq j, k, \ell$ . Since  $k = |h - \ell|$ , if  $h \neq \ell + k$ ,  $h = \ell - k$  and  $j - i - h$ ,  $\ell - h + i - h = i - \ell$  implies  $h = \ell$ ,  $k = 0$  and  $h = \ell + k$ .)

This proves Claim 1.

Let  $M$  denote the subalgebra of  $\text{End}_K(V)$  generated by  $A$ . Observe that  $M$  has a basis  $E_0, \dots, E_D$  as a vector space over  $K$ . Set

$$L := \text{Span}\{mA^*m - nA^*m \mid m, n \in M\}.$$

Claim 2.  $\dim L \leq D$ .

*Proof of Claim 2.* Since  $E_0, \dots, E_D$  span  $M$ ,

$$L = \text{Span}\{E_i A^* E_j - E_j A^* E_i \mid 0 \leq i < j \leq D\} \quad (29.25)$$

$$= \text{Span}\{E_{j-1} A^* E_j - E_j A^* E_{j-1} \mid 1 \leq j \leq D\} \quad (29.26)$$

by (29.5).

In particular,  $L$  has a spanning set of order  $D$ .

So, Claim 2 holds.

Claim 3.  $\{A^i A^* - A^* A^i \mid 1 \leq i \leq D\}$  is a basis for  $L$ .

*Proof of Claim 3.* Since

$$A^i A^* - A^* A^i = A^i A^* I - i A^* A^i$$

is contained in  $L$  ( $1 \leq i \leq D$ ), and since  $\dim L \leq D$ , it suffices to show the given elements are linearly independent.

Suppose they are dependent. Then there exists an integer  $i$  ( $1 \leq i \leq D$ ) such that

$$A^i A^* - A^* A^i \in \text{Span}(A^j A^* - A^* A^j \mid 1 \leq j < i). \quad (29.27)$$

Multiplying each term in (29.27) on the left by  $E_i^*$ , and on the left by  $E_0^*$ , and simplifying using

$$E_i^*(A^\ell A^* - A^* A^\ell)E_0^* = (\theta_0^* - \theta_i^*)E_i^* A^\ell E_0^*,$$

we find

$$E_i^* A^\ell E_0^* \in \text{Span}(E_i^* A^j E_0^* \mid 1 \leq j < i). \quad (29.28)$$

But the left side of (29.28) is nonzero.

A contradiction.

Since  $a^2 A^* A - A A^* A^2$  is contained in  $L$ , we find by Claim 2,

$$A^2 A^* A - A A^* A^2 = \sum_{i=1}^D \alpha_i (A^i A^* - A^* A^i) \quad (29.29)$$

for some  $\alpha_0, \dots, \alpha_D \in K$ .

Claim 4.  $\alpha_i = 0$  ( $3 < i \leq D$ ).

*Proof of Claim 4.* Suppose not, and set

$$t = \max\{i \mid 3 < i \leq D, \alpha_i \neq 0\}.$$

Then by (29.29), and Claim 1,

$$0 = E_t^* \left( A^2 A^* A - A A^* A^2 - \sum_{i=1}^D \alpha_i (A^i A^* - A^* A^i) \right) E_0^* \quad (29.30)$$

$$= \alpha_t (\theta_t^* - \theta_0^*) E_t^* A^t E_0^* \quad (29.31)$$

$$\neq 0. \quad (29.32)$$

(Since  $\alpha_i = 0$  if  $i > t$ ,

$$E_t^* A^2 A^* A E_0^* = E_t^* A A^* A^2 E_0^* = 0 \quad (\text{as } 2 + 1 < t - 0) \quad (29.33)$$

$$E_t^* A^i A^* E_0^* = E_t^* A^* A^i E_0^* = 0 \quad (29.34)$$

$$E_t^* A^t A^* E_0^* = \theta_0^* E_t^* A^t E_0^*, \quad (29.35)$$

$$E_t^* A^* A^t E_0^* = \theta_t^* E_t^* A^* A^t E_0^*. \quad (29.36)$$

$$(29.37)$$

A contradiction. This proves Claim 4.

Claim 5. Suppose  $D \geq 3$ . Then

$$\alpha_3 = \frac{\theta_{i+1}^* - \theta_{i+2}^*}{\theta_i^* - \theta_{i+3}^*} \quad \text{for all } i, (0 \leq i \leq D-3). \quad (29.38)$$

In particular,  $\alpha \neq 0$ .

*Proof of Claim 5.* Fix an integer  $i$  ( $0 \leq i \leq D-3$ ). Then by (29.24) and (29.29),

$$0 = E_{i+3}^* \left( A^2 A^* A - A A^* A^2 - \sum_{j=1}^3 \alpha_j (A^j A^* - A^* A^j) \right) E_i^* \quad (29.39)$$

$$= (\theta_{i+1}^* - \theta_{i+2}^* - \alpha_3 (\theta_i^* - \theta_{i+3}^*)) E_{i+3}^* A^3 E_i^*. \quad (29.40)$$

But  $E_{i+3}^* A^3 E_i^* \neq O$  by (29.6), so (29.38) holds.

This proves Claim 5.

Claim 6. Lines (29.7), (29.9), (29.14) hold.

*Proof of Claim 6.* First suppose  $D \geq 3$ . Then by (29.29), Claims 4, and 5,

$$A^2 2A^* A - AA^* A^2 = \alpha_3(A^3 A^* - A^* A^3) + \alpha_2(A^2 A^* - A^* A^2) + \alpha_1(AA^* - A^* A), \quad (29.41)$$

where  $\alpha_3 \neq 0$ . Hence

$$A^3 A^* - A^* A^3 - \frac{1}{\alpha_3}(A^2 A^* A - AA^* A^2) + \frac{\alpha_2}{\alpha_3}(A^2 A^* - A^* A^2) + \frac{\alpha_1}{\alpha_3}(AA^* - A^* A) = O.$$

Now (29.9) is immediate, where

$$\beta = \frac{1}{\alpha_3} - 1, \quad (29.42)$$

$$\gamma = -\frac{\alpha_2}{\alpha_3}, \quad (29.43)$$

$$\delta = -\frac{\alpha_1}{\alpha_3}. \quad (29.44)$$

The line (29.7) follows from the definition of  $[ \ , \ ]$ .

The line (29.14) is immediate from (29.38) and (29.42).

Now suppose  $D < 3$ . Then the line (29.14) is vacuously true, so consider (29.9).

Let  $\alpha_3$  denote any nonzero element of  $K$ .

Then  $A^2 A^* - A^* A^2, AA^* - A^* A$  certainly span  $L$  by Claim 3.

So, (29.41) holds for appropriate  $\alpha_1$  and  $\alpha_2 \in K$ .

Now, (29.9) holds, where  $\beta, \gamma, \delta$  are given by (29.42), (29.43), (29.44).

Claim 7. Lines (29.13), (16.11), (29.17) hold.

*Proof of Claim 7.* Pick an integer  $i$  ( $0 \leq i \leq D-1$ ).

By (29.9), we have

$$O = E_i(A^3 A^* - A^* A^3 - (\beta + 1)(A^2 A^* A - AA^* A^2) - \gamma(A^2 A^* - A^* A^2) - \delta(AA^* - A^* A))E_{i+1} \quad (29.45)$$

$$= E_i A^* E_{i+1} (\theta_i^3 - \theta^3 - (\beta + 1)(\theta_i^* \theta_{i+1} - \theta_i \theta^2) - \gamma(\theta_i^2 - \theta_{i+1}^2) - \delta(\theta_i - \theta_{i+1})) \quad (29.46)$$

$$= E_i A^* E_{i+1} (\theta_i - \theta_{i+1})(\theta_i^2 + \theta_i \theta_{i+1} + \theta_{i+1}^2 - (\beta + 1)\theta_i \theta_{i+1} - \gamma(\theta_i + \theta_{i+1}) - \delta) \quad (29.47)$$

$$= E_i A^* E_{i+1} (\theta_i - \theta_{i+1})(\theta_i^2 - \beta \theta_i \theta_{i+1} + \theta_{i+1}^2 - \gamma(\theta_i + \theta_{i+1}) - \delta). \quad (29.48)$$

But  $E_i A^* E_{i+1} \neq O$  by (29.5), and of course,  $\theta_i \neq \theta_{i+1}$ , so

$$0 = \theta_i^2 - \beta \theta_i \theta_{i+1} + \theta_{i+1}^2 - \gamma(\theta_i + \theta_{i+1}) - \delta.$$

This proves (29.17).

To obtain (16.11), pick any integer  $i$  ( $0 \leq i \leq D-2$ ). Then by (29.17),

$$0 = \theta_i^2 - \beta \theta_i \theta_{i+1} + \theta_{i+1}^2 - \gamma(\theta_i + \theta_{i+1}) - \delta \quad (29.49)$$

$$- (\theta_{i+1}^2 - \beta \theta_{i+1} \theta_{i+2} + \theta_{i+2}^2 - \gamma(\theta_{i+1} + \theta_{i+2}) - \delta) \quad (29.50)$$

$$= \theta_i^2 - \beta \theta_i \theta_{i+1} - \gamma \theta_i + \beta \theta_{i+1} \theta_{i+2} - \theta_{i+2}^2 + \gamma \theta_{i+2} \quad (29.51)$$

$$= (\theta_i - \theta_{i+2})(\theta_i - \beta \theta_{i+1} + \theta_{i+2} - \gamma). \quad (29.52)$$

So  $0 = \theta_i - \beta \theta_{i+1} + \theta_{i+2} - \gamma$ .

This gives (16.11).

To see (29.13), pick an integer  $i$  ( $0 \leq i \leq D-3$ ).

Then by (16.11),

$$0 = (\theta_i - \beta \theta_{i+1} + \theta_{i+2} - \gamma) - (\theta_{i+1} - \beta \theta_{i+2} + \theta_{i+3} - \gamma) \quad (29.53)$$

$$= \theta_i - (\beta + 1) \theta_{i+1} + (\beta + 1) \theta_{i+2} - \theta_{i+3}. \quad (29.54)$$

We have

$$\beta = \frac{\theta_i - \theta_{i+3}}{\theta_{i+1} - \theta_{i+2}} - 1 = \frac{\theta_i - \theta_{i+1} + \theta_{i+2} - \theta_{i+3}}{\theta_{i+1} - \theta_{i+2}},$$

as desired.

This proves Claim 7.

We have now proved (29.7), (29.9), (29.13), (29.14), (29.15), (29.17).

Interchanging the roles of  $A$  and  $A^*$ , we obtain (29.10), (29.12), (29.16), (29.18).





## Chapter 30

### $R, F, L$ Matrices

**Monday, April 12, 1993** # Edit Date

Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$  with standard module  $V$ .

Assume  $\Gamma$  is  $Q$ -polynomial with respect to the ordering

$$E_0, E_1, \dots, E_D$$

of primitive idempotents. Let  $A_i$  be an  $i$ -th adjacency matrix, and  $A = A_1$ .

$$A = \sum_{i=0}^D \theta_i A_i, \quad E_i = |X|^{-1} \sum_{i=0}^D \theta_i^* A_i.$$

Fix a vertex  $x \in X$ , write

$$E_i^* \equiv E_i^*(x), \quad A^* \equiv A^*(x), \quad A^* \equiv A_1^*, \quad T \equiv T(x).$$

Then

$$A^* = \sum_{i=0}^D \theta_i^* E_i^*.$$

By Theorem 29.1, there exist  $\beta, \gamma, \gamma^*, \delta, \delta^* \in \mathbb{R}$  such that

$$0 = [A, A^2 A^* - \beta A A^* A + A^* A^2 - \gamma(A A^* + A^* A) - \delta A^*] \quad (30.1)$$

$$0 = [A^*, A^{*2} A - \beta^* A^* A A^* + A A^{*2} - \gamma^*(A^* A + A A^*) - \delta^* A] \quad (30.2)$$

Recall raising matrix

$$R = \sum_{i=0}^D E_{i+1}^* A E_i^*$$

satisfies

$$R(E_i^*V) \subseteq E_{i+1}^*V \quad (0 \leq i \leq D), \quad E_{D+1}^*V = 0,$$

lowering matrix

$$L = \sum_{i=0}^D E_{i-1}^* A E_i^*$$

satisfies

$$L(E_i^*V) \subseteq E_{i-1}^*V \quad (0 \leq i \leq D), \quad E_{-1}^*V = 0,$$

and flat matrix

$$F = \sum_{i=0}^D E_i^* A E_i^*$$

satisfies

$$F(E_i^*V) \subseteq E_i^*V \quad (0 \leq i \leq D).$$

Also,

$$A = R + F + L.$$

**Theorem 30.1.** *With the above notation and assumptions,*

(i) *For all  $i$  ( $2 \leq i \leq D$ ),*

$$g_i^- FL^2 + LFL + g_i^+ L^2 F - \gamma L^2) E_i^* = O,$$

where

$$g_i^+ = \frac{\theta_{i-2}^* - (\beta + 1)\theta_{i-1}^* + \beta\theta_i^*}{\theta_{i-2}^* - \theta_i^*} \quad (30.3)$$

$$g_i^- = \frac{\theta_{i-2}^* + (\beta + 1)\theta_{i-1}^* - \theta_i^*}{\theta_{i-2}^* - \theta_i^*}. \quad (30.4)$$

(ii) *For all  $i$  ( $0 \leq i \leq D$ ),*

$$[F, LR - h_i RL] E_i^* = O,$$

where

$$h_i = \frac{\theta_{i-1}^* - \theta_i^*}{\theta_i^* - \theta_{i+1}^*} \quad (1 \leq i \leq D-1), \quad (30.5)$$

and  $h_0, h_D$  are indeterminants.

(iii) *For all  $i$  ( $1 \leq i \leq D$ ),*

$$(e^- e_i RL^2 + (\beta + 2)LRL + e_i^+ L^2 R + LF^2 - \beta FLF + F^2 L - \gamma(LF + FL) - \delta L) E_i^* = O,$$

where

$$e_i^+ = \frac{\theta_{i-1}^* - (\beta + 2)\theta_i^* + (\beta + 1)\theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*} \quad (1 \leq i \leq D) \quad (30.6)$$

$$e_i^- = \frac{-(\beta + 1)\theta_{i-2}^* + (\beta + 2)\theta_{i-1}^* - \theta_i^*}{\theta_{i-1}^* - \theta_i^*} \quad (2 \leq i \leq D), \quad (30.7)$$

and  $e_0^+, e_1^-$  are indeterminants.

*Proof.* WE have

$$O = A^3 A^* - A^* A^3 - (\beta + 1)(A^2 A^* A - A A^* A^2) \gamma (A^2 A^* - A^* A^2) - \delta (A A^* - A^* A).$$

(i) Fix  $i$  ( $2 \leq i \leq D$ ), and multiply above on the left by  $E_{i-2}^*$ , and on the right by  $E_i^*$ . Now reduce.

For example,

$$E_{i-2}^* A^3 A^* E_i^* = \theta_i^* E_{i-2}^* A^3 E_i^*,$$

where

$$E_{i-2}^* A^3 E_i^* = E_{i-2}^* A \left( \sum_{r=0}^D E_r^* \right) A \left( \sum_{s=0}^D E_s^* \right) A E_i^* \quad (30.8)$$

$$= \sum_{r,s} E_{i-2}^* A E_r^* A E_s^* A E_i^* \quad (30.9)$$

$$= \sum_{r,s, |i-2-r| \leq 1, |r-s| \leq 1, |s-i| \leq 1} E_{i-2}^* A E_r^* A E_s^* A E_i^* \quad (30.10)$$

$$= E_{i-2}^* A E_{i-2}^* A E_{i-1}^* A E_i^* + E_{i-2}^* A E_{i-1}^* A E_{i-1}^* A E_i^* + E_{i-2}^* A E_{i-1}^* A E_i^* A E_i^* \quad (30.11)$$

$$= (FL^2 + LFL + L^2 F) E_i^*. \quad (30.12)$$

Reducing the other terms in a similar manner, and simplifying, we obtain (i).

*Remark.*

$$E_{i-2}^* A^3 A^* E_i^* = \theta_{i-2}^* E_{i-2}^* A^3 E_i^* \quad (30.13)$$

$$= \theta_{i-2}^* (FL^2 + LFL + L^2 F) E_i^* \quad (30.14)$$

$$E_{i-2}^* A^2 A^* A E_i^* = (\theta_{i-1}^* (FL^2 + LFL) + \theta_i^* L^2 F) E_i^* \quad (30.15)$$

$$E_{i-2}^* A A^* A^2 E_i^* = (\theta_{i-2}^* FL^2 + \theta_{i-1}^* (LFL + L^2 F)) E_i^* \quad (30.16)$$

$$E_{i-2}^* (A^2 A^* - A^* A^2) E_i^* = (\theta_i^* - \theta_{i-2}^*) L^2 E_i^* \quad (30.17)$$

$$E_{i-2}^* (A A^* - A^* A) E_i^* = O. \quad (30.18)$$

Then we have

$$O = ((\theta_i^* - \theta_{i-2}^*)(FL^2 + LFL + L^2F)) \quad (30.19)$$

$$- (\beta + 1)(\theta_{i-1}^*(FL^2 + LFL) + \theta^*L^2F - \theta_{i-2}^*FL^2 - \theta_{i-1}^*(LFL + L^2F)) \quad (30.20)$$

$$- \gamma(\theta_i^* - \theta_{i-2}^*)L^2)E_i^* \quad (30.21)$$

$$= ((\theta_i^* - \theta_{i-2}^* - (\beta + 1)(\theta_{i-1}^* - \theta_{i-2}^*))FL^2 + (\theta_i^* - \theta_{i-2}^*)LFL \quad (30.22)$$

$$+ (\theta_i^* - \theta_{i-2}^* - (\beta + 1)(\theta_i^* - \theta_{i-1}^*))L^2F - \gamma(\theta_i^* - \theta_{i-2}^*)L^2)E_i^* \quad (30.23)$$

$$= -(\theta_{i-2}^* - \theta_i^*) \left( \left( \frac{-\beta\theta_{i-2}^* + (\beta + 1)\theta_{i-1}^* - \theta_i^*}{\theta_{i-2}^* - \theta_i^*} \right) FL^2 + LFL \quad (30.24)$$

$$+ \left( \frac{\theta_{i-2}^* - (\beta + 1)\theta_{i-1}^* + \beta\theta_i^*}{\theta_{i-2}^* - \theta_i^*} \right) L^2F - \gamma L^2 \right) E_i^* \quad (30.25)$$

$$= (\theta_i^* - \theta_{i-2}^*)(g_i^- FL^2 + LFL + g_i^+ L^2F - \gamma L^2)E_i^*. \quad (30.26)$$

(ii), (iii) are obtained in a similar manner replacing  $i - 2$  by  $i$  (resp.  $i - 1$ ).

□

*Remark.*

(ii) We have

$$O = E_i^*(A^3A^* - A^*A^3 - (\beta + 1)(A^2A^*A - AA^*A^2) - \gamma(A^2A^* - A^*A^2) - \delta(AA^* - A^*E))E_i^*.$$

Since  $\beta + 1 \neq 0$ , by (29.42) if  $D \geq 3$ ,

$$O = E_i^*(A^2A^*A - AA^*A^2)E_i^* \quad (30.27)$$

$$= ((\theta_i^* - \theta_{i-1}^*)RLF + (\theta_i^* - \theta_{i+1}^*)LRF) + (\theta_{i-1}^* - \theta_i^*)FRL + (\theta_{i+1}^* - \theta_i^*)FLR)E_i^* \quad (30.28)$$

$$= [F, (\theta_{i-1}^* - \theta_i^*)RL - (\theta_i^* - \theta_{i+1}^*)LR]E_i^* \quad (30.29)$$

$$= (\theta_{i+1}^* - \theta_i^*) \left[ F, LR - \frac{\theta_{i-1}^* - \theta_i^*}{\theta_i^* - \theta_{i+1}^*} RL \right] E_i^* \quad (30.30)$$

$$= (\theta_{i+1}^* - \theta_i^*)[F, LR - h_i RL]E_i^*. \quad (30.31)$$

(iii) We have

$$O = E_{i-1}^*(A^3A^* - A^*A^3 - (\beta + 1)(A^2A^*A - AA^*A^2) - \gamma(A^2A^* - A^*A^2) - \delta(AA^* - A^*E))E_i^* \quad (30.32)$$

$$= ((\theta_i^* - \theta_{i-1}^*)(RL^2 + LRL + L^2R + LF^2 + FLF + F^2L)) \quad (30.33)$$

$$- (\beta + 1)((\theta_{i-1}^* - \theta_{i-2}^*)RL^2 + (\theta_{i-1}^* - \theta_i^*)LRL + (\theta_{i+1}^* - \theta_i^*)L^2R \quad (30.34)$$

$$+ (\theta_i^* - \theta_{i-1}^*)FLF \quad (30.35)$$

$$- \gamma(\theta_i^* - \theta_{i-1}^*)(LF + FL) \quad (30.36)$$

$$- \delta(\theta_i^* - \theta_{i-1}^*)E_i^* \quad (30.37)$$

$$= ((\theta_i^* - \theta_{i-1}^*) - (\beta + 1)(\theta_{i-1}^* - \theta_{i-2}^*))RL^2 \quad (30.38)$$

$$+ (\theta_i^* - \theta_{i-1}^*) - (\beta + 1)(\theta_{i-1}^* - \theta_i^*))LRL \quad (30.39)$$

$$+ (\theta_i^* - \theta_{i-1}^*) - (\beta + 1)(\theta_{i+1}^* - \theta_i^*))L^2R \quad (30.40)$$

$$+ (\theta_i^* - \theta_{i-1}^*)LF^2 + (\theta_i^* - \theta_{i-1}^*)F^2L \quad (30.41)$$

$$+ (\theta_i^* - \theta_{i-1}^* - (\beta + 1)(\theta_i^* - \theta_{i-1}^*))FLF \quad (30.42)$$

$$- \gamma(\theta_i^* - \theta_{i-1}^*)(LF + FL) \quad (30.43)$$

$$- \delta(\theta_i^* - \theta_{i-1}^*)E_i^* \quad (30.44)$$

$$= (\theta_i^* - \theta_{i-1}^*) \left( \frac{-(\beta + 1)\theta_{i-2}^* + (\beta + 2)\theta_{i-2}^* - \theta_i^*}{\theta_{i-1}^* - \theta_i^*} RL^2 + (\beta + 2)LRL \quad (30.45)$$

$$+ \frac{\theta_{i-1}^* - (\beta + 2)\theta_i^* + (\beta + 1)\theta_{i+1}^*}{\theta_{i-1}^* - \theta_i^*} L^2R + LF^2 - \beta FLF + F^2L \quad (30.46)$$

$$- \gamma(LF + FL) - \delta L \Big) E_i^* \quad (30.47)$$

$$= (e_i^- RL^2 + (\beta + 2)LRL + e_i^+ L^2R + LF^2 - \beta FLF + F^2L - \gamma(LF + FL) - \delta L)E_i^*. \quad (30.48)$$

**Lemma 30.1.** *With the notation of Theorem 30.1,*

$$e_i^+ = \frac{\theta_i^* - \theta_{i+2}^*}{\theta_i^* - \theta_{i-1}^*} \quad (1 \leq i \leq D - 2) \quad (30.49)$$

$$e_i^- = \frac{\theta_{i-1}^* - \theta_{i-3}^*}{\theta_{i-1}^* - \theta_i^*} \quad (3 \leq i \leq D) \quad (30.50)$$

$$g_i^+ = \frac{\theta_i^* - \theta_{i+1}^*}{\theta_i^* - \theta_{i-2}^*} \quad (2 \leq i \leq D - 1) \quad (30.51)$$

$$g_i^- = \frac{\theta_{i-2}^* - \theta_{i-3}^*}{\theta_{i-2}^* - \theta_i^*} \quad (3 \leq i \leq D). \quad (30.52)$$

*In particular,  $e_i^\pm, g_i^\pm$  are non-zero for the range of  $i$  given above.*

*Proof.* In each case, equate the above expression with the corresponding expression in Theorem 30.1. The resulting equation is equal to (29.13).  $\square$

*Remark.* By Corollary 26.1 and Theorem 29.1,

$$e_i^+ = \frac{\theta_{i-1}^* - (\beta + 2)\theta_i^* + (\beta + 1)\theta_{i+1}^*}{\beta_{i-1}^* - \theta_i^*},$$

and

$$\beta + 1 = \frac{\theta_{j-1}^* - \theta_j^* + \theta_{j+1}^* - \theta_{j+2}^*}{\theta_j^* - \theta_{j+1}^*} + 1 = \frac{\theta_{j-1}^* - \theta_{j+2}^*}{\theta_j^* - \theta_{j+1}^*}.$$

Hence,

$$e_i^+ = \frac{1}{\theta_{i-1}^* - \theta_i^*}(\theta_{i-1}^* - \theta_i^* - (\beta + 1)(\theta_i^* - \theta_{i+1}^*)) \quad (30.53)$$

$$= \frac{1}{\theta_{i-1}^* - \theta_i^*}(\theta_{i-1}^* - \theta_i^* - (\beta + 1)(\theta_i^* - \theta_{i+2}^*)) \quad (30.54)$$

$$= \frac{\theta_i^* - \theta_{i+2}^*}{\theta_i^* - \theta_{i-1}^*}, \quad (30.55)$$

$$e_i^- = \frac{1}{\theta_{i-1}^* - \theta_i^*}(-(\beta + 1)\theta_{i-2}^* + (\beta + 2)\theta_{i-1}^* - \theta_i^*) \quad (30.56)$$

$$= \frac{1}{\theta_{i-1}^* - \theta_i^*}(\theta_{i-1}^* - \theta_i^* - \theta_{i-3}^* + \theta_i^*) \quad (30.57)$$

$$= \frac{\theta_{i-1}^* - \theta_{i-3}^*}{\theta_{i-1}^* - \theta_i^*}, \quad (30.58)$$

$$g_i^+ = \frac{1}{\theta_{i-2}^* - \theta_i^*}(\theta_{i-2}^* - (\beta + 1)\theta_{i-1}^* + \beta\theta_i^*) \quad (30.59)$$

$$= \frac{1}{\theta_i^* - \theta_{i-2}^*}(\theta_i^* - \theta_{i-2}^* + \theta_{i-2}^* - \theta_{i+1}^*) \quad (30.60)$$

$$= \frac{\theta_i^* - \theta_{i+1}^*}{\theta_i^* - \theta_{i-3}^*}, \quad (30.61)$$

$$g_i^- = \frac{1}{\theta_{i-2}^* - \theta_i^*}(-\beta\theta_{i-2}^* + (\beta + 1)\theta_{i-1}^* - \theta_i^*) \quad (30.62)$$

$$= \frac{1}{\theta_{i-2}^* - \theta_i^*}(\theta_{i-2}^* - \theta_i^* + \theta_i^* - \theta_{i-3}^*) \quad (30.63)$$

$$= \frac{\theta_{i-2}^* - \theta_{i-3}^*}{\theta_{i-2}^* - \theta_i^*}. \quad (30.64)$$

**Corollary 30.1.** *Let  $\Gamma = (X, E)$  be distance-regular of diameter  $D \geq 3$ ,  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Fix a vertex  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$ ,  $R \equiv R(x)$ ,  $L \equiv L(x)$ ,  $F \equiv F(x)$ . Then the following hold.*

(i)  $FR^2E_j^* \in \text{Span}(RFRE_j^*, R^2FE_j^*, R^2E_j^*)$ ,  $(0 \leq j \leq D - 3)$ .

(ii)  $R^2FE_j^* \in \text{Span}(RFRE_j^*, FR^2E_j^*, R^2E_j^*)$ ,  $(1 \leq j \leq D - 2)$ .

(iii)  $LR^2E_j^* \in \text{Span}(RLRE_j^*, R^2LE_j^*, F^2RE_j^*, FRFE_j^*, RF^2E_j^*, RFE_j^*, FRE_j^*, RE_j^*)$ ,  $(0 \leq j \leq D - 3)$ .

(iv)  $R^2LE_j^* \in \text{Span}(RLRE_j^*, LR^2E_j^*, F^2RE_j^*, FRFE_j^*, RF^2E^*j, RFE_j^*, FRE_j^*, RE_j^*),$   
 $(1 \leq j \leq D).$

*Proof.* Immediate from Theorem 30.1, and Lemma 30.1.  $\square$

*Remark.* By Theorem 30.1, and Lemma 30.1, we have the following, but similarly we can obtain above.

(i)  $FL^2E_j^* \in \text{Span}(LFLE_j^*, L^2FE_j^*, L^2E_j^*), (3 \leq j \leq D).$

(ii)  $L^2FE_j^* \in \text{Span}(LFLE_j^*, FL^2E_j^*, L^2E_j^*), (2 \leq j \leq D-1).$

(iii)  $RL^2E_j^* \in \text{Span}(LRLE_j^*, L^2RE_j^*, F^2LE_j^*, FLFE_j^*, LF^2E^*j, LFE_j^*, FLE_j^*, LE_j^*),$   
 $(3 \leq j \leq D).$

(iv)  $L^2RE_j^* \in \text{Span}(LRLE_j^*, RL^2E_j^*, F^2LE_j^*, FLFE_j^*, LF^2E^*j, LFE_j^*, FLE_j^*, LE_j^*),$   
 $(2 \leq j \leq D).$





## Chapter 31

# The “Inverse” of $R$

Wednesday, April 14, 1993

Let  $\Gamma = (X, E)$  be any graph of diameter  $D \geq 2$ . Fix a vertex  $x \in X$ . Let  $E_i^* \equiv E_i^*(x)$ , and  $T \equiv T(x)$ .

Recall adjacency matrix

$$A = R + L + F \quad (31.1)$$

$$R = \sum_{i=0}^D E_{i+1}^* A E_i^*, \quad (31.2)$$

$$L = \sum_{i=0}^D E_{i-1}^* A E_i^*, \quad (31.3)$$

$$F = \sum_{i=0}^D E_i^* A E_i^*. \quad (31.4)$$

Observe  $R$  is not invertible (indeed  $RE$ .) So,  $R^{-1}$  does not exist.

Below we find a matrix “ $R^{-1}$ ”  $\in T(x)$  such that  $R^{-1}Rv = v$  for “almost all”  $v \in V$ .

**Lemma 31.1.** *Let  $\Gamma = (X, E)$  denote any graph, and the standard module  $V$  over  $\mathbb{C}$ .*

*Fix a vertex  $x \in X$ , write*

$$R \equiv R(x), \quad L \equiv L(x), \quad E_i^* \equiv E_i^*(x) \quad \text{for all } i.$$

*Then,*

*(i) There exists unique “ $R^{-1}$ ”  $\in \text{Mat}_X(\mathbb{C})$  such that;*

- (ia)  $R^{-1}v = 0$  if  $Lv = 0$  for  $v \in V$ .
- (ib)  $R^{-1}RLv = Lv$  for all  $v \in V$ .
- (ii)  $R^{-1}(E_i^*V) \subseteq E_{i-1}^*V$  ( $0 \leq i \leq D$ ),  $E_{-1}^*V = 0$ .
- (iii)  $R^{-1} \in \text{Mat}_X(\mathbb{Q})$ .
- (iv)  $R^{-1} \in T(x)$ .

*Proof.*

(i) Consider the orthogonal direct sum.

$$V = (\text{Ker}L) + (\text{Ker}L)^\perp.$$

Claim 1.  $RL(\text{Ker}L)^\perp \subseteq (\text{Ker}L)^\perp$ .

*Proof of Claim 1.* Pick  $v \in (\text{Ker}L)^\perp$ , and  $w \in \text{Ker}L$ . Show

$$\langle RLv, w \rangle = 0.$$

But

$$\bar{R}^\top = R^\top = \left( \sum_{i=0}^D E_{i+1}^* A E_i^* \right)^\top = \sum_{i=0}^D E_i^* A E_{i+1}^* = L.$$

So,

$$\langle RLv, w \rangle = \langle Lv, \bar{R}^\top w \rangle = \langle Lv, Lw \rangle = 0.$$

Claim 2.  $RL : (\text{Ker}L)^\perp \rightarrow (\text{Ker}L)^\perp$  is an isomorphism of vector spaces.

*Proof of Claim 2.* It suffices to show above map is one-to-one.

Suppose there is a vector  $v \in (\text{Ker}L)^\perp$  such that  $RLv = 0$ .

Then,

$$0 = \langle RLv, v \rangle = \langle Lv, \bar{R}^\top v \rangle = \|Lv\|^2.$$

So  $Lv = 0$ .

Hence  $v \in \text{Ker}L \cap (\text{Ker}L)^\perp = 0$ .

This proves Claim 2.

Now “ $R^{-1}$ ” denote the unique matrix in  $\text{Mat}(\mathbb{C})$  such that

$$R^{-1}v = \begin{cases} 0 & \text{if } v \in \text{Ker}L \\ L(RL)^{-1}v & \text{if } v \in (\text{Ker}L)^\perp. \end{cases} \quad (31.5)$$

Observe that  $(RL)^{-1} : (\text{Ker}L)^\perp \rightarrow (\text{Ker}L)^\perp$  exists by Claim 2.

Observe  $R^{-1}$  satisfies (ia) by (31.5).

Claim 3.  $R^{-1}$  satisfies (ib).

*Proof of Claim 3.* It suffices to check

$$R^{-1}RLv = Lv$$

for  $v \in \text{Ker}L$  and  $v \in (\text{Ker}L)^\perp$ .

The case  $v \in \text{Ker}L$  is clear. So assume  $v \in (\text{Ker}L)^\perp$  by Claim 1. So,

$$R^{-1}(RLv) = L(RL)^{-1}RLv = Lv$$

as desired.

Uniqueness: Suppose a matrix  $\hat{R}^{-1} \in \text{Mat}_X(\mathbb{C})$  satisfies (ia), (ib). Then,  $\hat{R}^{-1}$  satisfies (31.5) above.

(Pf. The first part is clear. Let  $v \in (\text{Ker}L)^\perp$ . By Claim 2, there exists  $w \in (\text{Ker}L)^\perp$  such that  $v \in RLw$ . So  $\hat{R}^{-1}v = \hat{R}^{-1}RLw = Lw = L(RL)^{-1}v$ .)

Therefore,  $\hat{R}^{-1}$  agrees with  $R^{-1}$  on a basis for  $V$ , and  $\hat{R}^{-1} = R^{-1}$ .

(ii) Pick  $v \in E_i^*V$ . Show  $R^{-1}v \in E_{i-1}^*V$ .

Without loss of generality we may assume that  $v \in \text{Ker}L$  or  $v \in (\text{Ker}L)^\perp$ .

If  $v \in \text{Ker}L$ , then  $R^{-1}v = 0 \in E_{i-1}^*V$ .

If  $v \in (\text{Ker}L)^\perp$ , then

$$R^{-1}v = L(RL)^{-1}v \in LE_i^*V \subseteq E_{i-1}^*V.$$

(iii) Observe  $R, L \in (\text{Mat})_X(\mathbb{Q})$ .

So  $V, \text{Ker}L$ , each has basis consisting of vectors in  $\mathbb{Q}^{|X|}$ .

Replacing the construction of  $R^{-1}$  with the base field replaced by  $\mathbb{Q}$ , we find a matrix  $\tilde{R}^{-1} \in \text{Mat}_X(\mathbb{Q})$  satisfying (ia), (ib).

Now  $R^{-1}$  and  $\tilde{R}^{-1}$  agree on a basis, and hence  $R^{-1} = \tilde{R}^{-1}$ .

(iv)  $RL = \bar{L}^\top L$  is a real symmetric matrix. So it is diagonalizable.

Let  $\theta$  be any eigenvalue of  $RL$ . Let  $V_\theta$  denote the corresponding maximal eigenspace in  $V$ . Then

$$V = \sum_{\theta: \text{eigenvalue for } RL} V_\theta \quad (\text{orthogonal direct sum}).$$

Let  $E_\theta : V \rightarrow V_\theta$  denote the orthogonal projection. Then  $E_\theta$  is a complex polynomial in  $RL$ .

Thus  $E_\theta \in T(x)$ .

$E_\theta$  is real. Since  $RL$  is an integral matrix, every eigenvalue of  $RL$  is an algebraic integer.

Claim 4. We have

$$R^{-1} = \sum_{\theta: \text{eigenvalue of } RL} \theta^{-1} L E_\theta. \quad (31.6)$$

In particular,  $R^{-1} \in T(x)$ .

*Proof of Claim 4.* Show two sides of (31.6) agree, when applied to arbitrary  $v \in V$ .

Without loss of generality, we may assume that  $v \in V_\theta$  for some eigenvalue  $\theta$  of  $RL$ .

Let  $\theta'$  denote any eigenvalue of  $RL$ .

$$E_{\theta'} v = \begin{cases} 0 & \text{if } \theta' \neq \theta, \\ v & \text{if } \theta' = \theta. \end{cases}$$

RHS of (31.6) applied to  $v$  equals

$$\begin{cases} 0 & \text{if } \theta = 0, \\ \theta^{-1} L v & \text{if } \theta \neq 0. \end{cases}$$

Show this equals  $R^{-1}v$ .

Case  $\theta = 0$ : Since  $RLv = 0$ ,

$$0 = \langle v, RLv \rangle = \|Lv\|^2.$$

Hence  $Lv = 0$ , or  $v \in \text{Ker } L$ . By (ia),  $R^{-1}v = 0$ .

Case  $\theta \neq 0$ : Since  $RLv = \theta v$ ,  $v = \theta^{-1}RLv$ . Hence,

$$R^{-1}v = \theta^{-1}R^{-1}RLv = \theta^{-1}Lv$$

by (ib).

□

## Chapter 32

# Irreducible Modules of Endpoint $i$

Monday, April 19, 1993

**Lemma 32.1.** *Let  $\Gamma = (X, E)$  be any graph. With the notation of Lemma 31.1, the following hold.*

(i) *Let  $W$  denote a thin irreducible  $T$ -module with endpoint  $r$ , diameter  $d$ . Pick  $i$  ( $0 \leq i \leq d$ ), and pick  $v \in E_{r+i}^* W$ . Then,*

$$R^{-1}Rv = \begin{cases} v & \text{if } i < d, \\ 0 & \text{if } i = d. \end{cases}$$

(ii) *Assume  $\Gamma$  is distance regular and thin with respect to  $x$ . Pick  $t$  ( $0 \leq t \leq D/2$ ), and pick  $v \in E_t^* V$ . Then*

$$R^{-1}R^i v = R^{i-1}v \quad (1 \leq i \leq D - 2t).$$

*In particular,  $R^{-1}Rv = v$ .*

(iii) *Assume  $\Gamma$  is distance regular and thin with respect to  $x$ . Then*

$$R : E_i^* V \rightarrow E_{i+1}^* V \quad (0 \leq i < D/2)$$

*is one-to-one.*

*Proof.*

(i) Let  $w_0, w_1, \dots, w_d$  be a basis for  $W$  and  $w_i \in E_{r+i}^* W$ ,

$$Rw_i = w_{i+1} \quad (0 \leq i < d), \quad Lw_i = x_i(W)w_{i-1} \quad (1 \leq i \leq d).$$

So,

$$RLw_i = x_i(W)w_i \quad (1 \leq i \leq d).$$

(See Lemma 9.1.)

We want to find  $R^{-1}Rw_i$ .

If  $i = d$ ,  $R^{-1}Rw_d = 0$ .

If  $0 \leq i < d$ ,

$$R^{-1}Rw_i = R^{-1}w_{i+1} \tag{32.1}$$

$$= x_{i+1}(W)^{-1}R^{-1}RLw_{i+1} \tag{32.2}$$

$$= x_{i+1}(W)^{-1}Lw_{i+1} \tag{32.3}$$

$$= x_{i+1}(W)^{-1}x_{i+1}(W)w_i \tag{32.4}$$

$$= w_i. \tag{32.5}$$

Thus, we have (i).

$$RLw_i = Rx_i(W)w_{i-1} = x_i(W)w_i, \quad LRw_i = Lw_{i+1} = x_{i+1}(W)w_i \tag{32.6}$$

$$[L, R]w_i = (x_{i+1}(W) - x_i(W))w_i, \quad (0 \leq i \leq d), \tag{32.7}$$

$$x_0(W) = 0, \quad x_{d+1}(W) = 0, \tag{32.8}$$

$$[L, R]|_W = \sum_{i=0}^d (x_{i+1} - x_i(W))E_{r+i}^*|_W. \tag{32.9}$$

(ii) Let

$$V = \sum W \quad \text{orthogonal direct sum of thin irreducible } T\text{-modules.}$$

Then,

$$E_t^*V = \sum_{r(W) \leq t} E_t^*W \quad (\text{orthogonal direct sum}).$$

Without loss of generality, we may assume

$$v \in E_t^*W$$

for some thin irreducible  $T$ -module with endpoint at most  $t$ .

Now if  $i \leq D - 2t$ , then

$$t + i \leq D - t \quad (32.10)$$

$$\leq D - r(W) \quad (32.11)$$

$$\leq r(W) + d(W) \quad (D \leq 2r + d), \quad (32.12)$$

by Lemma 14.1 (iii).

So

$$t + i - 1 \leq r(W) + d(W) - 1.$$

Hence,

$$R^{-1}R^i v = R^{-1}R(R^{i-1}v) \quad (R^{i-1}v \in E_{t+i-1}^* W) \quad (32.13)$$

$$= R^{i-1}v \quad \text{by (i)}. \quad (32.14)$$

(iii) Suppose  $Rv = 0$  for some  $v \in E_i^* V$  ( $0 \leq i < D/2$ ). Then

$$0 = R^{-1}Rv = v,$$

by (ii) with  $t = i$  and  $i = 1$ .

□

**Definition 32.1.** Let  $\Gamma = (X, E)$  denote any graph with the standard module  $V$ . Fix a vertex  $x \in X$ . Write  $E_i^* \equiv E_i^*(x)$ ,  $T \equiv T(x)$ ,  $L \equiv L(x)$ .

1. For every  $i$  ( $0 \leq i \leq D$ ), define subspace  $V_i := V_i(x) \subseteq V$  by

$$V_i = \sum W,$$

where the sum begin over irreducible  $T$ -modules  $W$  with endpoint  $i$ .

Observe:

$$V = V_0 + V_1 + \cdots + V_D \quad (\text{orthogonal direct sum.})$$

$V_0$  is the trivial  $T$ -module.

2.  $(E_i^* V)_{\text{new}} \equiv E_i^* V_i \quad (0 \leq i \leq D)$ .

In general,

$$(E_i^* V)_{\text{new}} \subseteq \text{Ker} L \cap E^* i V \subseteq \text{Ker} L \cap E_i^* V \subseteq \text{Ker}(L E_i^*).$$

If each irreducible  $T$ -module with endpoint strictly less than  $i$  is thin,

$$(E_i^* V)_{\text{new}} = \text{Ker} L \cap E_i^* V \subseteq \text{Ker}(L \cdot E_i^*).$$

We have the assertion.

*Remark.*

$$E_i^*V = \sum_{j < i} V_j + V_i.$$

For  $V_j$  part, take  $w_{i-j} \in W$  irreducible with endpoint  $j < i$ . Then,

$$Lw_{i-j} = x_{i-j}(W)w_{i-j-1} \neq 0,$$

and

$$L|_{\sum_{j < i} E_i^*V_j} : \sum_{j < i} E_i^*V_j \rightarrow V$$

is one to one.

**Lemma 32.2.** *Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$ . Fix a vertex  $x \in X$ ,  $R \equiv R(x)$ ,  $L \equiv L(x)$ ,  $F \equiv F(x)$ . Pick  $v \in (E_1^*V)_{new}$ . Then,*

- (i)  $RE_i^*A_{i-1}v = c_iE_{i+1}^*A_iv \quad (1 \leq i \leq D)$ .
- (ii)  $FE_i^*A_{i-1}v = RE_{i-1}^*A_iV + (a_{i-1} - c_i + c_{i-1})E_i^*A_{i-1}v + c_iE_i^*A_{i+1}v \quad (1 \leq i \leq D)$ .
- (iii)  $LE_i^*A_{i-1}v = FE_{i-1}^*A_iV + (a_{i-1} - c_i + c_{i-1})E_{i-1}^*A_iv + b_{i-1}E_{i-1}^*A_{i-2}v \quad (2 \leq i \leq D)$ .
- (iv)  $LE_i^*A_{i+1}v = b_iE_{i-1}^*A_iv \quad (1 \leq i \leq D-1)$ .

*Proof.*

(i) Let

$$v = \sum_{y \in X, \partial(x,y)=1} \alpha_y \hat{y} \quad \text{for some } \{\alpha_g\} \subseteq \mathbb{C}.$$

Then

$$Lv = \left( \sum_{y \in X, \partial(x,y)=1} \alpha_y \right) \hat{x} = .$$

So,

$$\sum_{y \in X, \partial(x,y)=1} \alpha_y = 0.$$

Thus,

$$v = \sum_{y \in X, \partial(x,y)=1} \alpha_y (\hat{y} - \hat{x}).$$

Let

$$\tilde{A}_i = A_0 + A_1 + \cdots + A_i \quad (0 \leq i \leq D).$$



Then

$$\tilde{A}_i v = \sum_{y \in X, \partial(x, y)=1} \alpha_y \tilde{A}_i (\hat{y} - \hat{x}) \quad (32.15)$$

$$= \sum_{y \in X, \partial(x, y)=1} \alpha_y \left( \sum_{z \in X, \partial(y, z)=i, \partial(x, z)=i+1} \hat{z} - \sum_{z' \in X, \partial(y, z')=i+1, \partial(x, z')=i} \hat{z}' \right) \quad (32.16)$$

$$= \sum_{y \in X, \partial(x, y)=1} \alpha_y (E_{i+1}^* A_i \hat{y} - E_i^* A_{i+1} \hat{y}) \quad (32.17)$$

$$= E_{i+1}^* A_i v - E_{i+1}^* A_{i+1} v. \quad (32.18)$$

Recall (Claim 1 in the proof of Theorem 16.1.)

$$A \tilde{A}_i = c_{i+1} \tilde{A}_{i+1} + (a_i - c_{i+1} + c_i) \tilde{A}_i + b_i \tilde{A}_{i-1} \quad (0 \leq i \leq D-1).$$

(This is valid for  $i = 0$  as  $A \tilde{A}_0 = AI = c_1 \tilde{A} - \tilde{A}_0 = A$  by setting  $\tilde{A}_{i-1} = O$ .)

Now (i)–(iv) are obtained by applying this to  $v$  on the right and multiplied by  $E_j^*$  ( $0 \leq j \leq D$ ) on the left.

□

*Remark.*  $A \tilde{A}_{i-1} v = A E_i^* A_{i-1} v - A E_{i-1}^* A_i v$ . For  $1 \leq i \leq D$ ,

$$(c_i \tilde{A}_i + (a_{i-1} - c_i + c_{i-1}) \tilde{A}_{i-1} + b_{i-1} \tilde{A}_{i-2}) v \quad (32.19)$$

$$= c_i E_{i+1}^* A_i v - c_i E_i^* A_{i+1} v \quad (32.20)$$

$$+ (a_{i-1} - c_i + c_{i-1}) E_i^* A_{i-1} v - (a_{i-1} - c_i + c_{i-1}) E_{i-1}^* A_i v \quad (32.21)$$

$$+ b_{i-1} E_{i-1}^* A_{i-2} v - b_{i-1} E_{i-2}^* A_{i-1} v. \quad (32.22)$$

(i)  $R E_i^* A_{i-1} v = E_{i+1}^* A E_i^* A_{i-1} v = c_i E_{i+1}^* A_i v$  ( $1 \leq i \leq D$ ).

(ii) For  $1 \leq i \leq D$ ,

$$F E_i^* A_{i-1} v = E_i^* A E_i^* A_{i-1} v \quad (32.23)$$

$$= R E_{i-1}^* A_i v - c_i E_i^* A_{i+1} v + (a_{i-1} - c_i + c_{i-1}) E_i^* A_{i-1} v. \quad (32.24)$$

(iii) For  $2 \leq i \leq D$ ,

$$L E_i^* A_{i-1} v = E_{i-1}^* A E_i^* A_{i-1} v \quad (32.25)$$

$$= F E_{i-1}^* A_i v - (a_{i-1} - c_i + c_{i-1}) E_{i-1}^* A_i v + b_{i-1} E_{i-1}^* A_{i-2} v. \quad (32.26)$$

(Even if  $i = 1$ , this is valid by setting  $A_{i-2} = O$ .)

(iv) For  $1 \leq i \leq D-1$ ,  $L E_i^* A_{i+1} v = E_{i-1}^* A E_i^* A_{i+1} v = b_i E_{i-1}^* A_i v$ .

**Lemma 32.3.** *Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$ . Fix a vertex  $x \in X$ ,  $T \equiv T(x)$ ,  $E_i^* \equiv E_i^*(x)$ ,  $R = R(x)$ ,  $F = F(x)$ ,  $L = L(x)$ .*

*For every  $v \in (E_1^*V)_{\text{new}}$ , the following are equivalent.*

- (i)  $E_i^*A_{i-1}v$ ,  $E_i^*A_{i+1}v$  are linearly dependent for every  $i$  ( $1 \leq i \leq D-1$ ).
  - (ii) *There exists a thin irreducible  $T$ -module  $W$  with endpoint 1 that contains  $v$ .*
- If (i), (ii) hold then*

$$W = \text{Span}(E_1^*, E_2^*A_1v, \dots, E_D^*A_{i-1}v).$$

*Proof.* (ii)  $\rightarrow$  (i). Clear as

$$E_i^*A_{i-1}v, E_i^*A_{i+1}v \in E_i^*W = \text{Span}(w_{i-1}).$$

(i)  $\rightarrow$  (ii) Consider the sequence

$$E_1^*A_{i-1}v, E_2^*A_1v, E_3^*A_2v, \dots, E_{D+1}^*A_Dv.$$

The first term is nonzero and the last term is 0. So there exists

$$n := \min\{i \mid 1 \leq i \leq D, E_{i+1}^*A_i v = 0\}.$$

Now

$$E_{j+1}^*A_j v = 0 \quad (n \leq j \leq D). \quad (32.27)$$

*Remark.* Use induction and Lemma 32.2 (i),

$$E_{j+1}^*A_j v \in \text{Span}(RE_j^*A_{j-1}v) \quad (j \geq 1).$$

By our assumption (i), and the definition of  $n$ ,

$$E_j^*A_{j+1}v \in \text{Span}(E_j^*A_{j-1}v) \neq 0 \quad (1 \leq j \leq n).$$

By Lemma 32.2 (i),

$$RE_j^*A_{j-1}v \in \text{Span}(E_{j+1}^*A_j v) \quad (1 \leq j \leq n).$$

By Lemma 32.2 (ii),

$$FE_j^*A_{j-1}v \in \text{Span}(RE_{j-1}^*A_j v, E_j^*A_{j-1}v, E_j^*A_{j+1}v) \quad (32.28)$$

$$\subseteq \text{Span}(RE_{j-1}^*A_{j-2}v, E_j^*A_{j-1}v) \quad (32.29)$$

$$\text{Span}(E_{j-1}^*A_{j-1}v) \quad (1 \leq j \leq n). \quad (32.30)$$

By Lemma 32.2 (iii),

$$FE_j^*A_{j-1}v \in \text{Span}(FE_{j-1}^*A_jv, E_{j-1}^*A_jv, E_{j-1}^*A_{j-2}v) \quad (32.31)$$

$$\subseteq \text{Span}(FE_{j-1}^*A_{j-2}v, E_{j-1}^*A_{j-2}v) \quad (32.32)$$

$$\text{Span}(E_{j-1}^*A_{j-2}v) \quad (2 \leq j \leq n). \quad (32.33)$$

Hence,

$$W = \text{Span}(E_1^*A_0v, E_2^*A_1v, \dots, E_n^*A_{n-1}v).$$

is  $R$ ,  $F$ ,  $L$  invariant.

Therefore  $W$  is a thin  $T$ -module with endpoint 1 that contains  $v$ . □



## Chapter 33

# Algebra on First Subconstituent

Wednesday, April 21, 1993

**Lemma 33.1.** *Let  $Y = (X, \{R_i\}_{0 \leq i \leq D})$  be a commutative scheme. Fix a vertex  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$ ,  $M^* \equiv M^*(x)$ ,  $T \equiv T(x)$ . Then the following hold.*

- (i)  $E_0^* M M^* = E_0^* M$
- (ii)  $E_0^* T = E_0^* M$ .
- (iii)  $T E_0^* T = M E_0^* M$ .
- (iv)  $E_0^* E_0 E_0^* = |X|^{-1} E_0^*$ .
- (v)  $E_0^* E_0 E_0^* = |X|^{-1} E_0^*$ .
- (vi) *Lines (i)-(iv) hold if we interchange  $(E_0, E_0^*)$ ,  $(M, M^*)$ .*

Moreover,  $M E_0^* M = M^* E_0 M^*$ .

*Proof.*

(i)  $\supseteq$ :  $1 \in M^*$  implies  $M \subseteq M M^*$ .

$\subseteq$ : Pick  $\alpha \in E_0^* M M^*$ . Show  $\alpha \in E_0^* M$ . Since  $A_0, A_1, \dots, A_D$  span  $M$ , and since  $E_0^*, E_1^*, \dots, E_D^*$  span  $M^*$ , without loss of generality we may assume that

$$\alpha = E_0^* A_i E_j^*$$

for some  $i, j \in \{0, \dots, D\}$ .

Without loss of generality we may assume that  $i = j$ , else  $\alpha = 0$  by Lemma 20.3.

$$(E_h^* A_i E_j^* \neq O \Leftrightarrow p_{hi}^j \neq 0.)$$

Now

$$\alpha = E_0^* A_i \left( \sum_{h=0}^D E_h^* \right) = E_0^* A_i \in E_0^* M.$$

(ii)  $\supseteq$ : This is clear.

$\subseteq$ :  $E_0^* T$  is the minimal right ideal of  $T$  containing  $E_0^*$ .

So, we just have to show that  $E_0^* M$  is a right ideal of  $T$  containing  $E_0^*$ .

It clearly contains  $E_0^*$  since  $I \in M$ , and is a right ideal of  $T$  by (i), and the fact that  $T$  is generated by  $M$  and  $M^*$ .

(iii) By the transpose of (ii),

$$T E_0^* = M E_0^*,$$

so,

$$T E_0^* T = (T E_0^*)(E_0^* T) = M E_0^* E_0^* M = M E_0^* M.$$

(iv) We have

$$E_0^* E_0 E_0^* = \frac{1}{|X|} E_0^* \left( \sum_{h=-}^D A_h \right) E_0^* = \frac{1}{|X|} E_0^* A_0 E_0^* = |X|^{-1} E_0^*.$$

(v) The first part is clear by using Lemma 20.3 (ii),

$$E_h A_i^* E_j \neq O \Leftrightarrow q_{hi}^j \neq 0,$$

and Lemma 22.1 (iii),

$$q_{0i}^j = \delta_{ij}.$$

Also,

$$M E_0^* M = T E_0^* T = T E_0^* E_0 E_0^* T \subseteq T E_0 T = M^* E_0 M^*,$$

and

$$M^* E_0 M^* \subseteq M E_0^* M$$

by dual argument. So,

$$M^* E_0 M^* = M E_0^* M.$$

This proves the lemma. □

**Lemma 33.2.** *Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \leq 3$ ,  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Pick a vertex  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$ ,  $M^* \equiv M^*(x)$ ,  $T \equiv T(x)$ .*

$$(i) \quad E_1^* M M^* = E^* M + E_1^* E_0 M^* + E_1^* E_1 M^*.$$

$$(ii) \quad E_1 M^* M = E_1 M^* + E_1 E_0^* M + E_1 E_1^* M.$$

*Proof.*

$$(i) \quad \text{View } E_{-1}^*, E_{D+1}^* \text{ as } O.$$

View  $\theta_{-1}^*, \theta_{D+1}^*$  as indeterminates.

Let  $\Delta$  denote RHS in (i).

$\supseteq$ :  $I \in M^*$  implies  $M \subseteq M M^*$ .

$\subseteq$ : Suppose not. Then there exists

$$\alpha \in E_1^* M M^* \setminus \Delta. \quad (33.1)$$

Since  $A_0, A_1, \dots, A_D$  span  $M$ , since  $E_0^*, E_1^*, \dots, E_D^*$  span  $M^*$ , without loss of generality we may assume that

$$\alpha = E_1^* A_i E_j^*$$

for some  $i, j \in \{0, \dots, D\}$ .

Observe  $|i - j| \leq 1$ , else  $\alpha = 0$  by Lemma 20.3.

Without loss of generality, assume  $i + j$  is minimal subject to the above constraints.

First assume

$$j = i + 1. \quad (33.2)$$

Observe

$$E_1^* A_i E_{i+1}^* + E_1^* A_i E_i^* + E_1^* A_i E_{i-1}^* \quad (33.3)$$

$$= E_1^* A_i \left( \sum_{h=0}^D E_h^* \right) \quad (33.4)$$

$$= E_1^* A_i \quad (33.5)$$

$$\in \Delta. \quad (33.6)$$

Also, observe

$$E_1^* A_i E_i^*, E_1^* A_i E_{i-1}^* \in \Delta$$

by the minimality of  $i + j$ , so

$$\alpha = E_1^* A_i E_{i+1}^* \in \Delta$$

by (33.6). Hence, (33.2) cannot occur.

Since  $|i - j| \leq 1$ ,

$$i \in \{j, j + 1\}. \quad (33.7)$$

Observe

$$E_1^* A_{j+1} E_j^* + E_1^* A_j E_j^* + E_1^* A_{j-1} E_j^* \quad (33.8)$$

$$= E_1^* \left( \sum_{h=0}^D A_h \right) E_j^* \quad (33.9)$$

$$= |X| E_1^* E_0 E_j^* \quad (33.10)$$

$$\in \Delta, \quad (33.11)$$

and

$$\theta_{j+1}^* E_1^* A_{j+1} E_j^* + \theta_j^* E_1^* A_j E_j^* + \theta_{j-1}^* E_1^* A_{j-1} E_j^* \quad (33.12)$$

$$= E_1^* \left( \sum_{h=0}^D \theta_h^* A_h \right) E_j^* \quad (33.13)$$

$$= |X| E_1^* E_1 E_j^* \quad (33.14)$$

$$\in \Delta. \quad (33.15)$$

Since  $E_1^* A_{j-1} E_j^* \in \Delta$  by the minimality of  $i + j$ , so

$$E_1 A_{j+1} E_j^* + E_1^* A_j E_j^* \in \Delta,$$

$$\theta_{j+1}^* E_1^* A_{j+1} E_{j+1}^* + \theta_j^* E_1^* A_j E_j^* \in \Delta.$$

But,  $\theta_0^*, \theta_1^*, \dots, \theta_D^*$  are distinct by Lemma 22.2 (iv), so

$$E_1^* A_{j+1} E_j^*, E_1^* A_j E_j^* \in \Delta.$$

But  $\alpha$  is one of these two matrices, so

$$\alpha \in \Delta.$$

Hence, (33.7) cannot occur either, and we have a contradiction.

(ii) Dual argument.

□

**Lemma 33.3.** *With the above notation, set*

$$\tilde{J} := E_1^* J E_1^*, \quad \tilde{A} := E_1^* A E_1^*.$$

(i)  $\tilde{J}^2 = k\tilde{J}$ . ( $k = \text{valency of } \Gamma$ )



- (ii)  $\tilde{J}\tilde{A} = \tilde{A}\tilde{J} = a_1\tilde{J}$ . ( $a_1 = p_{11}^1$  for  $\Gamma$ )
- (iii)  $E_1^*E_0E_1^* = |X|^{-1}\tilde{J}$ .
- (iv)  $E_1^*E_1E_1^* = |X|^{-1}(E_1^*(\theta_0^* - \theta_2^*) + \tilde{A}(\theta_1^* - \theta_2^*) + \tilde{J}(\theta_2^*))$ .

*Proof.*

- (i) The first subconstituent has  $k$  vertices.
- (ii) The first subconstituent is regular of valency  $a_1$ .
- (iii) Since  $E_0 = |X|^{-1}J$ ,

$$E_1^*E_0E_1^* = |X|^{-1}\tilde{J}.$$

- (iv) We have

$$E_1^*E_1E_1^* = E_1^* \left( |X|^{-1} \sum_{h=0}^D \theta_h^* A_h \right) E_1^* \quad (33.16)$$

$$= |X|^{-1}(\theta_0^* E_1^* A_0 E_1^* + \theta_1^* E_1^* A_1 E_1^* + \theta_2^* E_1^* A_2 E_1^*) \quad (33.17)$$

$$= |X|^{-1}(\theta_0^* E_1^* + \theta_1^* \tilde{A} + \theta_2^* E_1^* A_2 E_1^*). \quad (33.18)$$

Also,

$$\tilde{J} = E_1^* J E_1^* \quad (33.19)$$

$$= E^* A_0 E_1^* + E^* A_1 E_1^* + E_1^* A_2 E_1^* \quad (33.20)$$

$$= E_1^* + \tilde{A} + E_1^* A_2 E_1^*. \quad (33.21)$$

Eliminating the  $E_1^* A_2 E_1^*$  term in (33.18) using equation (33.21), we get (iv).

□

**Lemma 33.4.** *With the above notation,*

- (i)  $E_1^*T = E_1^*E_0M^* + E_1^*M + E_1^*E_1M^* + E_1^*E_1E_1^*M + \dots$ .
- (ii)  $E_1^*TE_1^* = \text{Span}(E_1^*E_0E_1^*, E_1^*, E_1^*E_1E_1^*, (E_1^*E_1E_1^*)^2, \dots)$ .
- (iii)  $E_1^*TE_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2, \dots)$ .
- (iv)  $E_1^*TE_1^*$  is symmetric (in particular, commutative).

*Proof.*

- (i)  $\supseteq$ : Clear.

$\subseteq$ :  $E_1^*T$  is the minimal right ideal of  $\Gamma$  that contains  $E_1^*$ .

RHS contains  $E_1^*$ , so show RHS is a right ideal of  $T$ .

Show RHS is closed with respect to multiplication on right by  $M$ ,  $M^*$ .

We have

$$E_1^*E_0M^*(M) = E_1^*E_0M^*, \quad E_1^*E_0M^*(M^*) = E_1^*E_0M^*$$

by dual of Lemma 33.1 (i).

By Lemma 33.2,

$$E_1^*E_1E_1^* \cdots E_1^*M(M^*) \quad (33.22)$$

$$= E_1^*E_1E_1^* \cdots E_1^*(E_1^*MM^*) \quad (33.23)$$

$$= E_1^*E_1E_1^* \cdots E_1^*(E_1^*M + E_1^*E_0M^* + E_1^*E_1M^*) \quad (33.24)$$

$$\in \text{RHS}, \quad (33.25)$$

because

$$E_1^*E_1E_1^* \cdots E_1^*E_1E_0M^* \subseteq E_1^*TE_0T = E_1^*M^*E_0M^* = E_1^*E_0M^*.$$

By Lemma 33.2,

$$E_1^*E_1E_1^* \cdots E_1^*M^*(M) \quad (33.26)$$

$$= E_1^*E_1E_1^* \cdots E_1^*(E_1^*M^*M) \quad (33.27)$$

$$= E_1^*E_1E_1^* \cdots E_1^*(E_1^*M^* + E_1^*E_0M^* + E_1^*E_1M^*) \quad (33.28)$$

$$\in \text{RHS}, \quad (33.29)$$

because by the last part of Lemma 33.1,

$$E_1^*E_1E_1^* \cdots E_1^*E_1E_0M \subseteq E_1^*TE_0^*T = E_1^*ME_0^*M = E_1^*E_0M^*.$$

(ii) Multiply (i) on the right by  $E_1^*$ , we have

$$E_1^*TE_1^* = E_1^*E_0M^*E_1^* + E_1^*ME_1^* + E_1^*E_1M^*E_1^* \quad (33.30)$$

$$+ \cdots + E_1^*E_1 \cdots E_1M^*E_1^* + E_1^*E_1 \cdots E_1^*ME_1^* \quad (33.31)$$

$$= \text{Span}(E_1^*E_0E_1^*, E_1^*, E_1^*E_1E_1^*, (E_1^*E_1E_1^*)^2, \dots). \quad (33.32)$$

*Remark.* Note that by Lemma 29.1,

$$E_1^*ME_1^* = \text{Span}(E_1^*A_0E_1^*, E_1^*A_1E_1^*, E_1^*A_2E_1^*) \quad (33.33)$$

$$= \text{Span}(E_1^*, E_1^*E_1E_1^*, E_1^*E_0E_1^*). \quad (33.34)$$

Moreover,

$$E_1^* \cdots E_1^*E_0E_1^* \subseteq E_1^*TE_0^*TE_1^* = E_1^*M^*E_0M^*E_1^* \in \text{Span}(E_1^*E_0E_1^*).$$

(iii) By (ii),  $E_1^*TE_1^*$  is generated by  $\tilde{J} = |X|E_1^*E_0E_1^*$  and  $E_1^*E_1E_1^*$ .

By Lemma 33.3 (iv),  $E_1^*TE_1^*$  is generated by  $\tilde{J}, \tilde{A}$ .

But,  $\text{Span}\tilde{J}$  is a 2-sided ideal by Lemma 33.3 (i), (ii).

Hence, we have (iii).

(iv)  $\tilde{A}, \tilde{J}$  are symmetric commuting matrices, we have the claim.

□



## Chapter 34

# Modules of Endpoint One

**Friday, April 23, 1993**

Let  $\Gamma = (X, E)$  be distance-regular of diameter  $D \geq 3$ .

Assume  $\Gamma$  is  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Write

$$\tilde{A}_i = A_0 + A_1 + \dots + A_i \quad i \in \{0, 1, \dots, D\}.$$

Fix a vertex  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$ ,  $M^* \equiv M^*(x)$ ,  $T \equiv T(x)$ .

Pick  $0 \neq v \in (E_1^*V)_{new}$ . Set  $v^* = |X|E_1v$ . We will show that

$$Tv = Mv + M^*v^*.$$

We need preliminary lemma.

**Lemma 34.1.** *With the above notation, we have the following.*

(i)  $\tilde{A}_h v = E_{h+1}^* A_h v - E_h^* A_{h+1} v$ ,  $h \in \{0, 1, \dots, D\}$ .

( $E_{D+1}^* = A_{D+1} = O$ ).

(ii)  $E_h^* v^* = (\theta_{h-1}^* - \theta_h^*) E_h^* A_{h-1} v - (\theta_h^* - \theta_{h+1}^*) E_h^* A_{h+1} v$ ,  $h \in \{0, 1, \dots, D\}$ . ( $A_{-1} = A_{D+1} = O$ ).

(iii)  $(\theta_i^* - \theta_{i+1}^*) E_{i+1}^* A_i v = \left( \sum_{h=0}^i (\theta_h^* - \theta_{i+1}^*) A_h \right) v - \left( \sum_{h=0}^i E_h^* \right) v^*$ ,  $i \in \{0, 1, \dots, D-1\}$ .

(iv)  $(\theta_i^* - \theta_{i+1}^*) E_i^* A_{i+1} v = \left( \sum_{h=0}^{i-1} (\theta_h^* - \theta_i^*) A_h \right) v - \left( \sum_{h=0}^i E_h^* \right) v^*$ ,  $i \in \{0, 1, \dots, D-1\}$ .

(v)  $Mv + M^*v^* = \text{Span}\{E_i^* A_{i-1} v, E_{i-1}^* A_i v \mid 1 \leq i \leq D\}$ .

*Proof.*

(i) It is already done in Lemma 32.2.

(ii)

$$E_h^* v^* = |X| E_h^* E_1 v \quad (34.1)$$

$$= E_h^* \left( \sum_{i=0}^D \theta_i^* A_i \right) v \quad (34.2)$$

$$= E_h^* \left( \sum_{i=0}^D \theta_i^* (\tilde{A}_i - \tilde{A}_{i-1}) \right) v \quad (34.3)$$

$$= E_h^* \left( \sum_{i=0}^{D-1} (\theta_i^* - \theta_{i+1}^*) \tilde{A}_i \right) v + E_h^* \theta_D^* \tilde{A}_D v \quad (34.4)$$

$$= E_h^* \left( \sum_{i=0}^{D-1} (\theta_i^* - \theta_{i+1}^*) (E_{i+1}^* A_i v - E_i^* A_{i+1} v) \right) \quad (34.5)$$

$$= (\theta_{h-1}^* - \theta_h^*) E_h^* A_{h-1} v - (\theta_h^* - \theta_{h+1}^*) E_h^* A_{h+1} v. \quad (34.6)$$

(iii), (iv) Call the equation in (iii),  $i^+$  and call the equation in (iv)  $i^-$ . Prove in order,

$$0^-, 0^+, 1^-, 1^+, 2^-, 2^+, \dots$$

$0^-$ : Trivial.

*Remark.*

$$\text{LHS} = (\theta_0^* - \theta_1^*) E_0^* A_1 v \quad (34.7)$$

$$= (\theta_{-1}^* - \theta_1^*) E_0^* A_{-1} v - E_h^* v^* \quad (\text{by (ii)}) \quad (34.8)$$

$$= -E_0^* v^* \quad (34.9)$$

$$= \text{RHS}. \quad (34.10)$$

$i^+$ : using (i) and  $i^-$ .

$$\text{LHS} = (\theta_i^* - \theta_{i+1}^*) E_{i+1}^* A_i v \quad (34.11)$$

$$= (\theta_i^* - \theta_{i+1}^*) E_{i+1}^* A_{i+1} v + (\theta_i^* - \theta_{i+1}^*) \tilde{A}_i v \quad (\text{by (i)}) \quad (34.12)$$

$$= \left( \sum_{h=0}^{i-1} (\theta_h^* - \theta_{i+1}^*) A_h \right) v - \left( \sum_{h=0}^i E_h^* \right) v^* + (\theta_i^* - \theta_{i+1}^*) \left( \sum_{h=0}^i A_h \right) v \quad (\text{by } i^-) \quad (34.13)$$

$$= \left( \sum_{h=0}^i (\theta_h^* - \theta_{i+1}^*) A_h \right) v - \left( \sum_{h=0}^i E_h^* \right) v^*. \quad (34.14)$$

$i^-$ : using (ii) and  $(i-1)^+$ .

$$\text{LHS} = (\theta_i^* - \theta_{i+1}^*)E_i^*A_{i+1}v \quad (34.15)$$

$$= (\theta_{i-1}^* - \theta_i^*)E_i^*A_{i-1}v - E_i^*v^* \quad (\text{by (ii)}) \quad (34.16)$$

$$= \left( \sum_{h=0}^{i-1} (\theta_h^* - \theta_i^*)A_h \right) v - \left( \sum_{h=0}^{i-1} E_h^* \right) v^* - E_i^*v^* \quad (34.17)$$

$$= \left( \sum_{h=0}^{i-1} (\theta_h^* - \theta_i^*)A_h \right) v - \left( \sum_{h=0}^i E_h^* \right) v^*. \quad (34.18)$$

(v) Immediate from (i) – (iv).

□

*Remark.*

$$Mv + M^*v^* \subseteq \text{Span}\{\tilde{A}_h v, E_h^*v^* \mid 0 \leq h \leq D\} \quad (34.19)$$

$$\subseteq \text{Span}\{E_h^*A_{h-1}v, E_{h-1}^*A_h v \mid 1 \leq h \leq D\} \quad (34.20)$$

by (i) and (ii).

On the other hand,

$$E^*hA_{h-1}v, E_{h-1}^*A_h v \in Mv + M^*v^* \quad i \in \{1, 2, \dots, D\}$$

by (iii) and (iv).

**Lemma 34.2.** *With the notation of Lemma 34.1, assume  $0 \neq v \in (E_1^*V)_{\text{new}}$  is an eigenvector for  $\tilde{A} := E_1^*AE_1^*$ . Then*

(i)  $Tv = Mv + M^*v$ , where  $v^* = |X|E_1v$ .

(ii)  $Tv = \text{Span}\{v_1^+, v_2^+, \dots, v_D^+, v_2^-, v_3^-, \dots, v_{D-1}^-\}$ , where  $v_i^+ = E_i^*A_{i-1}v$ ,  $v_i^- = E_i^*A_{i+1}v$ .

(iii)  $\dim E_1^*Tv = 1$ ,  $\dim E_i^*Tv \leq 2$  for  $i \in \{2, \dots, D-1\}$ , and  $\dim E_D^*Tv \leq 1$ .

(iv)  $Tv$  is an irreducible  $T$ -module.

*Proof.*

(i)  $\supseteq$ :  $v \in Tv$ . So  $Mv \subseteq Tv$ , and

$$v^* \in Mv \subseteq Tv.$$

Hence,  $M^*v^* \subseteq Tv$ .

$\subseteq$ : It suffices to show that  $Mv + M^*v^*$  is a  $T$ -module (since it clearly contains  $v$ ).

Show:

$$(a) M^*Mv \subseteq Mv + M^*v^*.$$

$$(b) MM^*v \subseteq Mv + M^*v^*.$$

*Proof of (a).* By the transpose of (i) in Lemma 33.2,

$$M^*ME_1^* = ME_1^* + M^*E_0E_1^* + M^*E_1E_1^*.$$

Since  $v \in E_1^*V$ ,  $E_1^*v = v$  and

$$M^*Mv = Mv + M^*E_0v + M^*E_1v.$$

But also  $E_0v = 0$  since  $v$  is orthogonal to the trivial  $T$ -module. Since  $E_1v = |X|^{-1}v^*$ ,

$$M^*Mv = Mv + M^*v^*$$

as desired.

(b) is obtained from the transpose of (ii) in Lemma 33.2.

*Remark.*

$$MM^*v = MM^*E_1v^* \quad (34.21)$$

$$= M^*E_1v^* + ME_0^*E_1v^* + ME_1^*E_1v^* \quad (34.22)$$

$$= M^*v^* + ME_0^*v^* + ME_1^*v^*. \quad (34.23)$$

$E_0^*v^* \in Tv$  and  $E_0^*Tv = 0$  as  $v \in (E_1^*V)_{new}$ . So,  $E_0^*v^* = 0$ .

$$E_1^*v^* = |X|E_1^*E_1v \quad (34.24)$$

$$= |X|E_1^*E_1E_1^*v \quad (34.25)$$

$$= ((\theta_0^* - \theta_2^*)E_1^* + (\theta_1^* - \theta_2^*)E_1^*AE_1^* + \theta_2^*|X|E_1^*E_0E_1^*)v \quad (34.26)$$

$$= (\theta_0^* - \theta_2^*)v + (\theta_1^* - \theta_2^*)E_1^*AE_1^*v + \theta_2^*|X|E_1^*E_0v \quad (34.27)$$

$$\in \text{Span}\{v\}, \quad (34.28)$$

as  $E_0v = 0$ , and  $v$  is an eigenvector of  $E_1^*AE_1^*$ .

\*  $v \in (E_1^*V)_{new}$ . If  $v$  is an eigenvector of  $E_1^*AE_1^*$ ,

$$E_1^*v^* \in \text{Span}\{v\}.$$

(ii) We have

$$Tv = Mv + M^*v^* \quad (34.29)$$

$$= \text{Span}\{E_i^*A_{i-1}v, E_{i-1}^*A_iv \mid 1 \leq i \leq D\} \quad (34.30)$$

$$= \text{Span}\{v_i^+, v_{i-1}^- \mid 1 \leq i \leq D\} \quad (34.31)$$

$$= \text{Span}\{v_1^+, v_2^+, \dots, v_D^+, v_0^-, \dots, v_{D-1}^-\} \quad (34.32)$$



by Lemma 34.1 (v).

But  $v_0^- = E_0^* A_1 v = 0$  since  $v \in (E_1^* V)_{new}$ , and  $v_1^- \in \text{Span}\{v_1^+\}$ .

Indeed,

$$v_1^- = E_1^* A_2 v = (-1 - a_0(Tv))v_1^+.$$

where  $a_0(Tv)$  is the eigenvalue of  $v$  associated with  $\tilde{A}$ .

To see this, observe

$$0 = \tilde{J}v \quad (34.33)$$

$$= E_1^* \left( \sum_{i=0}^D A_i \right) E_1^* v \quad (34.34)$$

$$= E_1^* \left( \sum_{i=0}^2 A_i \right) E_1^* v \quad (34.35)$$

$$= v + a_0(Tv)v + v_1^-. \quad (34.36)$$

Therefore,

$$Tv = \text{Span}\{v_1^+, v_2^+, \dots, v_D^*, v_0^-, \dots, v_{D-1}^-\}.$$

(iii)  $v_i^+, v_i^- \in E_i^* V$ .

(iv) Suppose  $Tv$  is reducible, i.e.,  $Tv = W_1 + W_2$ . (orthogonal direct sum of nonzero  $T$ -modules)

$$E_1^* Tv = E^* W_1 + E_1^* W_2$$

has dimension 1 by (iii). Assume  $v \in E_1^* W_1$ . Then  $Tv \subseteq W_1$ , a contradiction.  $\square$

**Lemma 34.3.** *With the notation of Lemma 34.1, assume  $0 \neq v \in (E_1^* V)_{new}$  is an eigenvector for  $\tilde{A} := E_1^* A E_1^*$ .*

(i)  $Tv$  is thin if and only if  $M^*v \subseteq Mv$ .

(ii) Let  $W$  denote any irreducible  $T$ -module with endpoint 1. Then

$$W = Tv'$$

for some  $0 \neq v' \in (E_1^* V)_{new}$  that is an eigenvector of  $\tilde{A}$ .

(iii) Denote eigenvalue of  $\tilde{A}$  associated to  $v$  (resp.  $v'$ ) by  $a_0(Tv)$  (resp.  $a_0(Tv')$ ).

Then  $Tv, Tv'$  are isomorphic  $T$ -module if and only if  $a_0(Tv) = a_1(Tv')$ .

(iv)  $E_1^* T E_1^*$  has basis

$$\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2, \dots, \tilde{A}^{\ell-1},$$

where  $\ell$  is the number of mutually nonisomorphic  $T$ -modules with endpoint 1.

*Proof.*

(i) If  $Tv$  is thin, then by Lemma 9.1,  $Tv = Mv$ . Hence  $M^*v^* \subseteq Mv$ .

*Remark.* Originally, the statement was  $Tv$  is thin if and only if  $M^*v = Mv$ . This is not the case in general. Suppose  $\Gamma$  is thin. Let  $W$  be an irreducible  $T$ -module of endpoint 1. Then, that  $W \cap E_1^*V \ni v \neq 0$  implies  $v^* \in W \cap E_1V$  gives one to one and  $k \leq m$ .

However, by ‘Distance-Regular Graphs’ (A.E. Brouwer, 1989),

$$J(v, d): v \geq 2d$$

$$b_j = (d-j)(v-d-j)c_j = j^2 \quad (34.37)$$

$$\theta_j = (d-j)(v-d-j) - jm_j = \binom{v}{j} - \binom{v}{j-1} \quad (34.38)$$

In particular,

$$k = b_0 = d(v-d) > m_1 = v-1 \quad \text{if } d \geq 2,$$

and  $J(v, d)$  is thin.

So  $|X|E_1v = v^*$  may be 0 sometimes. But as  $Tv$  is dual thin of diameter at least  $D-2$ . The dual endpoint  $r^* \leq 2$ , so in that case,  $E_2v \neq 0$ . Hence, if  $D \geq 3$ ,  $E_2v \neq 0$  always.

*Remark.* Now assume  $M^*v \subseteq Mv = Tv$ . Then

$$Mv = \{f(A)v \mid f(\lambda) \in \mathbb{C}[\lambda]\}.$$

So,

$$E_iTv = E_iMv \in \text{Span}(E_iv).$$

Hence,  $Tv$  is dual thin.

Now we can construct a basis,  $0 \neq w_0^* \in E_{r^*}W$ , where  $r^*$  is the dual endpoint, and

$$w_0^*, w_1^*, \dots, w_d^* \in W = Tv,$$

where  $w_i^* = E_{r^*+i}^*A_1^{*i}w_0^*$ .

$$A_1^*w_i^* = w_{i+1}^* + a_i^*w_i^* + x_i^*w_{i-1}^*,$$

and  $w_i^* = p_i^*(A^*)w_0^*$ .

$$E_{r^*+i}^*A_1^*E_{r^*+i}|_{E_{r^*+i}W} = a_i^* \cdot 1|_{E_{r^*+i}W},$$

$$E_{r^*+i-1}^* A_1^* E_{r^*+i} A^* E_{r^*+i-1} |_{E_{r^*+i-1} W} = x_i^* \cdot 1 |_{E_{r^*+i-1} W}.$$

See Lemma 9.1, and Lemma 22.2.

From above,  $Tv = M^* w_0^*$ . So,

$$E_i^* Tv = E_i^* M^* w_0^* \in \text{Span}\{E_i^* w_0^*\}.$$

Thus,  $Tv$  is thin.

\*Need to write down the dual at least for Lemma 9.1, Corollary 9.1.

(iii)  $E_1^* W$  is an  $\tilde{A}$ -module. So, there exists  $0 \neq v' \in E_1^* W$  that is an eigenvalue for  $\tilde{A}$ . Also  $Tv' \subseteq W$ .

Since  $W$  is irreducible,  $Tv' = W$ .

(iii) Suppose  $Tv \rightarrow Tv'$  is an isomorphism of  $T$ -modules.

Recall  $\sigma s = s\sigma$  for all  $s \in T$ .

$$\text{Span}\{\sigma v\} = \sigma E_1^* Tv = E_1^* \sigma Tv = E_1^* Tv' = \text{Span}\{v'\}.$$

Hence

$$a_0(Tv)\sigma v = \sigma(a_0(Tv)v) = \sigma \tilde{A}v = \tilde{A}\sigma v = a_0(Tv')\sigma v.$$

Since  $\sigma v \neq 0$ ,  $a_0(Tv) = a_0(Tv')$ .

Now suppose  $a_0(Tv) = a_0(Tv')$ . Show

$$\sigma : Tv \rightarrow Tv' \quad (sv \mapsto sv') \quad (s \in T)$$

is an isomorphism of  $T$ -modules.

Pick  $s \in T$ . Require  $sv = 0$  if and only if  $sv' = 0$ .

Without loss of generality,  $s \in TE_1^*$ , since  $v, v' \in E_1^* V$ .

Now  $0 = sv$  if and only if

$$0 = \|sv\|^2 = \bar{v}^\top \bar{s}^\top sv.$$

But,  $\bar{s}^\top s \in E_1^* TE_1^*$ .

Hence, by Lemma 33.4 (iii),

$$\bar{s}^\top s = \alpha \tilde{J} + p(\tilde{A})$$

for some  $\alpha \in \mathbb{C}$  and  $p(\lambda) \in \mathbb{C}[\lambda]$ .

Thus, using the fact that  $\tilde{J}v = 0$ ,

$$0 = \|sv\|^2 = \bar{v}^\top (\alpha \tilde{J} + p(\tilde{A}))v = \|v\|^2 p(a_0(Tv))$$

if and only if  $0 = p(a_0(Tv))$ .

Replacing  $v$  by  $v'$ , we have

$$0 = sv' \leftrightarrow 0 = p(a_0(Tv')) \quad (34.39)$$

$$\leftrightarrow 0 = p(a_0(Tv)) \quad (34.40)$$

$$\leftrightarrow 0 = sv \quad (34.41)$$

as desired.

(iv) The following hold.

$$\ell = \text{the number of mutually nonisomorphic } T\text{-modules with endpoint 1} \quad (34.42)$$

$$= \text{the number of distinct eigenvalues of } \tilde{A} : (E_1^*V)_{new} \rightarrow (E_1^*V)_{new} \quad (34.43)$$

$$= \text{the degree of minimal polynomial of } \tilde{A} : (E_1^*V)_{new} \rightarrow (E_1^*V)_{new}. \quad (34.44)$$

Claim 1.  $\tilde{J}.E_1^*, \tilde{A}, \dots, \tilde{A}^{\ell-1}$  are linearly independent.

*Proof of Claim 1.* Suppose not. Then

$$\alpha\tilde{J} + p(\tilde{A}) = O$$

for some  $\alpha \in \mathbb{C}$  and  $p(\lambda) \in \mathbb{C}[\lambda]$  with  $\deg p \leq \ell - 1$ .

But  $\tilde{J}|_{(E_1^*V)_{new}} = O$  implies  $p(\tilde{A})|_{(E_1^*V)_{new}} = O$ .

Since

$$\deg p < \text{the degree of minimal polynomial of } \tilde{A}|_{(E_1^*V)_{new}},$$

we find  $p$  is identically 0.

Then  $\alpha$  is identically 0 also.

Claim 2.  $\tilde{J}.E_1^*, \tilde{A}, \dots, \tilde{A}^{\ell-1}$  span  $E_i^*TE_i^*$ .

*Proof of Claim 2.* It needs to show

$$\tilde{J}.E_1^*, \tilde{A}, \dots, \tilde{A}^\ell \text{ are linearly dependent.} \quad (34.45)$$

Let  $m$  denote the minimal polynomial of  $\tilde{A}|_{(E_1^*V)_{new}}$ . So,

$$m(\tilde{A}|_{(E_1^*V)_{new}}) = 0.$$

Observe that

$$E_1^*V = (E_1^*V)_{new} + \text{Span}\{A\hat{x}\}.$$

(direct sum of  $E_1^*TE_1^*$ -modules.)

$$m(\tilde{A})A\hat{x} = f \cdot A\hat{x} \quad \text{for some } f \in \mathbb{C}.$$

On the other hand,

$$\tilde{J}A\hat{x} = kA\hat{x} \quad (k : \text{valency of } \Gamma).$$

Therefore,

$$m(\tilde{A}) - \frac{f}{k}\tilde{J} = O,$$

and (34.45) holds.

□



## Chapter 35

$$\dim E_1^* T E_1^* \leq 5$$

Monday, April 26, 1993

**Theorem 35.1.** *Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$ . Assume  $\Gamma$  is  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Fix a vertex  $x \in X$ , and write  $E_i^* \equiv E_i^*(x)$ ,  $T \equiv T(x)$ .*

(i) *Up to isomorphism, there are at most 4 thin irreducible  $T$ -modules with endpoint 1.*

(ii) *Suppose  $\Gamma$  is thin with respect to  $x$ . Then*

$$\dim E_1^* T E_1^* \leq 5.$$

*Proof.*

(ii) is immediate from (i) and part (iv) of Lemma 34.3.

(i)

Claim 1.  $E_1^* M E_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A})$ .

*Proof of Claim 1.*

$$E_1^* M E_1^* = \text{Span}\{E_1^*, E_1^* A E_1^*, E_1^* A_2 E_1^*, E_1^* A_3 E_1^*, \dots\}.$$

But  $E_1^* A_h E_1^* = O$  if  $h > 2$  (by Lemma 16.1). So,

$$E_1^* M E_1^* = \text{Span}\{E_1^*, E_1^* A E_1^*, E_1^* A_2 E_1^*\}.$$

Also,

$$\tilde{J} = E_1^* J E_1^* \quad (35.1)$$

$$= E_1^* \left( \sum_{h=0}^D A_h \right) E_1^* \quad (35.2)$$

$$= E_1^* + E_1^* A E_1^* + E_1^* A_2 E_1^*. \quad (35.3)$$

So,

$$E_1^* M E_1^* = \text{Span}\{E_1^*, E_1^* A E_1^*, \tilde{J}\}.$$

We are done, since  $\tilde{A} = E_1^* A E_1^*$ .

Claim 2.  $E_1^* M M^* M E_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2)$ .

*Proof of Claim 2.*  $\supseteq$ : Clear.

$\subseteq$ : In Lemma 33.4 (i), we say

$$E_1^* T = E_1^* E_0 M^* + E_1^* M + E_1^* E_1 M^* + E^* E_1 E_1^* M + \dots$$

In fact, the proof of that lemma gives a sequence;

$$E_1^* M M^* = E_1^* E_0 M^* + E_1^* M + E_1^* E_1 M^*, \quad (35.4)$$

$$E_1^* M M^* M = E_1^* E_0 M^* + E_1^* M + E_1^* E_1 M^* + E^* E_1 E_1^* M, \quad (35.5)$$

$$E_1^* M M^* M M^* = E_1^* E_0 M^* + E_1^* M + E_1^* E_1 M^* + E^* E_1 E_1^* M + E^* E_1 E_1^* M M^*, \quad (35.6)$$

$$\vdots \quad (35.7)$$

Multiply (35.5) through on the right by  $E_1^*$  to get

$$E_1^* M M^* M E_1^* = E_1^* M E_1^* + E^* E_1 E_1^* M E_1^* = \text{Span}\{\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2\},$$

since  $\tilde{J}^2, \tilde{A}\tilde{J} = \tilde{J}\tilde{A} \in \text{Span}\{\tilde{J}\}$ .

This proves Claim 2.

Now, let  $W$  denote any irreducible  $T$ -module with endpoint 1, and pick  $0 \neq v \in E_1^* W$ . Set

$$v_i^+ = E_i^* A_{i-1} E_1^* v, \quad v_i^- = E_i^* A_{i+1} E_1^* v, \quad i \in \{1, \dots, D\}.$$

We know by Lemma 34.2 (ii) that  $W$  is thin if and only if  $v_i^+, v_i^-$  are linearly dependent for all  $i \in \{2, \dots, D-1\}$ .

In general,

$$\Phi_i = \det \begin{pmatrix} \|v_i^+\|^2 & \langle v_i^+, v_i^- \rangle \\ \langle v_i^+, v_i^- \rangle & \|v_i^-\|^2 \end{pmatrix} \geq 0$$

with equality if and only if  $v_i^+, v_i^-$  are linearly dependent, (because  $\Phi_i$  is the determinant of a Gram matrix).



Let  $i$  be an integer in  $\{2, \dots, D-1\}$ .

Claim 3. There exists  $p^{++} \in \mathbb{C}[\lambda]$ ,  $\deg p^{++} \leq 2$  (that depends only on the intersection numbers) such that

$$\|v_i^+\|^2 = \|v\|^2 p^{++}(a_0(W)).$$

*Proof of Claim 3.*

$$\|v_i^+\|^2 = \bar{v}^\top E^* 1 A_{i-1} E_i^* E_i^* A_{i-1} E_1^* v = \bar{v}^\top E^* 1 A_{i-1} E_i^* A_{i-1} E_1^* v.$$

But,

$$E^* 1 A_{i-1} E_i^* A_{i-1} E_1^* \in E^* 1 M M^* M E_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2)$$

by Claim 2.

So, there exists  $\alpha \in \mathbb{C}$ , and  $p^{++} \in \mathbb{C}[\lambda]$  with  $\deg p^{++} \leq 2$  such that

$$E^* 1 A_{i-1} E_i^* A_{i-1} E_1^* = \alpha \tilde{J} + p^{++}(\tilde{A}), \quad (\tilde{A}^0 = E_1^*).$$

Now,

$$\|v_i^+\|^2 = \bar{v}^\top (\alpha \tilde{J} + p^{++}(\tilde{A})) v = \|v\|^2 p^{++}(a_0(W)),$$

since  $\tilde{J}v = 0$ , and  $\tilde{A}v = a_0(W)v$ .

This proves Claim 3.

Similarly, there exist  $p^{--}, p^{+-} \in \mathbb{C}[\lambda]$  with  $\deg p^{--}, \deg p^{+-} \leq 2$  such that

$$\|v_i^-\|^2 = \|v\|^2 p^{--}(a_0(W)), \quad \langle v_i^+, v_i^- \rangle = \|v\|^2 p^{+-}(a_0(W)).$$

Claim 4.  $E^* 1 A_{i-1} E_i^* A_{i+1} E_1^* = (\tilde{J} - \tilde{A} - E_1^*) p_{i-1, i+1}^2$ . In particular,

$$p^{+-}(\lambda) = -p_{i-1, i+1}^2(\lambda + 1).$$

*Proof of Claim 4.* Pick vertices  $y, z \in X$  such that  $\partial(x, y) = \partial(x, z) = 1$ .

$$(\text{LHS})_{yz} = \sum_{w \in X} (E_1^* A_{i-1} E_i^*)_{yw} (E_i^* A_{i+1} E_1^*)_{wz} \quad (35.8)$$

$$= \sum_{w \in X, \partial(y, w)=i-1, \partial(x, w)=i, \partial(w, z)=i+1} 1 \quad (35.9)$$

$$= \begin{cases} 0 & \text{if } \partial(y, z) = 0, \\ 0 & \text{if } \partial(y, z) = 1, \\ p_{i-1, i+1}^2 & \text{if } \partial(y, z) = 2, \end{cases} \quad (35.10)$$

$$= \text{RHS}_{yz} \quad (35.11)$$

Note that  $E_1^* A_2 E_1^* = \tilde{J} - \tilde{A} - E_1^*$ .

Now,

$$\langle v_i^+, v^- \rangle = \bar{v}^\top E_1^* A_{i-1} E_i^* A_{i+1} E_1^* v \quad (35.12)$$

$$= p_{i-1, i+1}^2 (\bar{v}^\top (\tilde{J} - \tilde{A} - E_1^*) v) \quad (35.13)$$

$$= (a_0(W) + 1) p_{i-1, i+1}^2 \|v\|^2. \quad (35.14)$$

Claim 5.  $\deg p^{++} = \deg p^{--} = 2$ . (only need for some  $i$ )

*Proof of Claim 5.* We need to calculate  $p^{++}$ ,  $p^{--}$ .

*Remark.* Pick vertices  $y, z \in X$  such that  $\partial(x, y) = \partial(x, z) = 1$ . Then

$$(E_1^* A_{i-1} E_i^* A_{i-1} E_1^*)_{yz} = |\gamma_{i-1}(y) \cap \Gamma_i(x) \cap \Gamma_{i-1}(z)|,$$

which is equal to  $p_{i-1, i}^1$  if  $\partial(y, z) = 0$ , but ...

$$(E_1^* A_{i+1} E_i^* A_{i+1} E_1^*)_{yz} = |\gamma_{i+1}(y) \cap \Gamma_i(x) \cap \Gamma_{i+1}(z)|,$$

which is equal to  $p_{i+1, i}^1$  if  $\partial(y, z) = 0$ , but ...

Conclusion.

$$\Phi_i = \det \begin{pmatrix} \|v_i^+\|^2 & \langle v_i^+, v_i^- \rangle \\ \langle v_i^+, v_i^- \rangle & \|v_i^-\|^2 \end{pmatrix} \geq 0 \quad (35.15)$$

$$= \|v\|^4 (p^{++}(\lambda) p^{--}(\lambda) - (p_{i-1, i+1}^2)^2 (\lambda + 1)^2) \quad (35.16)$$

$$\geq 0, \quad (35.17)$$

where  $\lambda = a_0(W)$ .

$W$  is thin if and only if  $\Phi_i(\lambda) = 0$  for all  $i \in \{2, \dots, D-1\}$ .

Each  $\Phi_i$  is degree 4, so at most 4 different thin irreducible modules  $W$  of endpoint 1 up to isomorphism.

□

**Note.** In fact  $\Phi_i(\lambda)$  is independent of  $i$  up to scalar multiple for  $i \in \{2, \dots, D-1\}$ .

If  $\Gamma$  has classical parameters  $(q, D, \alpha, \beta)$ , the roots are;

$$\beta - \alpha - 1, -1, -q - 1, dq \frac{q^{D-1} - 1}{q - 1} - 1.$$

## Chapter 36

# Dual Endpoint

**Wednesday, April 28, 1993**

Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$ ,  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Fix a vertex  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$ ,  $T \equiv T(x)$ .

Let  $W$  be an irreducible  $T$ -module of diameter  $d$ .

Recall that the endpoint

$$r(W) = \min\{i \mid 0 \leq i \leq D, E_i^* W \neq 0\}.$$

**Definition 36.1.** The dual endpoint (with respect to above ordering  $E_0, E_1, \dots, E_D$ ),

$$r^*(W) = \min\{i \mid 0 \leq i \leq D, E_i W \neq 0\}.$$

$$r(W) = 0 \leftrightarrow r^*(W) = 0 \leftrightarrow W : \text{ trivial } T\text{-module,}$$

(by Lemma 10.1).

Suppose  $W$  is thin. Then  $W$  is dual thin. (See Corollary 9.1.)

Moreover,  $\{i \mid E_i W \neq 0\}$  is a subinterval of  $\{0, 1, \dots, D\}$ . (same proof as for distance regular)

*Remark.* Dual version of Lemma 4.1.

**Lemma 4.1'.** Let  $A^* \equiv A_1^*(x)$ ,  $W$  an irreducible  $T$ -module, and  $d^* = \{i \mid E_i W \neq 0\} - 1$ .

- (i)  $E_i A^* E_j = 0$  if  $|i - j| > 1$ ,  $E_i A^* E_j \neq 0$  if  $|i - j| = 1$ ,  $0 \leq i, j \leq d^*(x)$ .
- (ii)  $A^* E_j W \subseteq E_{j-1} W + E_j W + E_{j+1} W$ ,  $0 \leq j \leq d^*(x)$ . ( $E_i W = 0$  if  $i < j$  or  $i > d^*(x)$ .)

- (iii)  $E_j W \neq 0$  if  $r \leq j \leq r + d$ ,  $= 0$  if  $0 \leq j \leq r$  or  $r + d < j \leq d^*(x)$ .  
 (iv)  $E_i A^* E_j W \neq 0$ , if  $|i - j| = 1$  ( $r^* \leq i, j \leq r^* + d^*$ ).

*Proof of 4.1' | (i)* By Lemma 20.3,

$$E_i A^* E_j = 0 \leftrightarrow q_{i1}^j = 0.$$

By Lemma 22.2,

$$\Gamma: Q\text{-polynomial} \leftrightarrow q_{i1}^j \begin{cases} = 0 & \text{if } |j - i| > 1 \\ \neq 0 & \text{if } |j - i| = 1. \end{cases} \quad (36.1)$$

$$\leftrightarrow E_i A^* E_j \begin{cases} = O & \text{if } |j - i| > 1 \\ \neq O & \text{if } |j - i| = 1. \end{cases} \quad (36.2)$$

(ii) We have

$$A^* E_j W = \left( \sum_{i=0}^D E_i \right) A^* E_j W \quad (36.3)$$

$$= E_{j-1} A^* E_j W + E_j A^* E_j W + E_{j+1} A^* E_j W \quad (36.4)$$

$$\subseteq E_{j-1} W + E_j W + E_{j+1} W. \quad (36.5)$$

(iii) Suppose  $E_j W = 0$  for some  $j \in \{r^*, \dots, r^* + d^*\}$ . Then  $r^* < j$  by the definition of  $r^*$ . Set

$$\widetilde{W} = E_{r^*} W + E_{r^*+1} W + \dots + E_{j-1} W.$$

Observe  $0 \subsetneq \widetilde{W} \subsetneq W$ . Also  $A\widetilde{W} \subseteq \widetilde{W}$  by (ii), and  $E_i^* \widetilde{W} \subseteq \widetilde{W}$  for every  $i$  by construction.

Thus,  $T\widetilde{W} \subseteq \widetilde{W}$ , contradicting  $W$  being irreducible.

(iv) Suppose  $E_{j+1} A^* E_j W = 0$  for some  $j \in \{r^*, \dots, r^* + d^* - 1\}$ . Then,

$$\widetilde{W} = E_{r^*} W + E_{r^*+1} W + \dots + E_j W$$

is  $T$ -invariant. If  $E_{j-1} A^* E_j W = 0$  for some  $j \in \{r^* + 1, \dots, r^* + d^*\}$ , then

$$\widetilde{W} = E_j W + E_{j+1} W + \dots + E_{r^*+d^*} W$$

is  $T$ -invariant. Moreover,  $0 \subsetneq \widetilde{W} \subsetneq W$  in both cases. A contradiction.

**Definition.** Let  $W$  be an irreducible dual thin  $T$ -module with dual endpoint  $r^*$  and diameter  $d^*$ .

Let  $a_i^* = a_i^*(W) \in \mathbb{C}$  satisfying

$$E_{r^*+i}A^*E_{r^*+i}|_{E_{r^*+i}W} = a_i^* \cdot 1|_{E_{r^*+i}W}.$$

Let  $x_i^* = x_i^*(W) \in \mathbb{C}$  satisfying

$$E_{r^*+i-1}A^*E_{r^*+i}A^*E_{r^*+i-1}|_{E_{r^*+i-1}W} = x_i^* \cdot 1|_{E_{r^*+i-1}W}.$$

**Lemma 9.1'.** With above notation, the following hold.

- (i)  $a_i^* \in \mathbb{R}$  for all  $i \in \{0, \dots, d^*\}$ .
- (ii)  $x_i^* \in \mathbb{R}^{>0}$  for all  $i \in \{1, \dots, d^*\}$ .
- (iii) Pick  $0 \neq w_0^* \in E_{r^*}W$ . Set  $w_i^* = E_{r^*+i}A^{*i}w_0^*$  for all  $i$ . Then
  - (iiia)  $w_0^*, w_1^*, \dots, w_{d^*}^*$  is a basis for  $W$ ,  $w_{-1}^* = w_{d^*+1}^* = 0$ .
  - (iiib)  $A^*w_i^* = w_{i+1}^* + a_i^*w_i^* + x_i^*w_{i-1}^*$  for all  $i \in \{0, \dots, d^*\}$ .
- (iv) Define  $p_0^*, p_1^*, \dots, p_{d^*+1}^* \in \mathbb{R}[\lambda]$  by

$$p_0^* = 1, \quad \lambda p_i^* = p_{i+1}^* + a_i^*p_i^* + x_i^*p_{i-1}^* \quad \text{for all } i \in \{0, \dots, d^*\}, \quad p_{-1}^* = 0.$$

$$(iva) \quad p_i^*(A^*)w_0^* = w_i^*, \text{ for all } i \in \{0, \dots, d^* + 1\}.$$

$$(ivb) \quad p_{d^*+1}^* \text{ is the minimal polynomial of } A^*|_W.$$

*Proof of Lemma 9.1' | (i)* Recall

$$A^* = \sum_{j=0}^D \theta_j^* E_j^*, \quad \theta_j^* = q_1(j) = |X|(E_1)_{xy} \in \mathbb{R}, \quad \partial(x, y) = j.$$

$a_i$  is an eigenvalue of a real symmetric matrix  $E_{r^*+i}A^*E_{r^*+i}$ .

(ii) Let  $B = E_{r^*+i}A^*E_{r^*+i-1}$ .

Then,  $x_i^*$  is an eigenvalue of a real symmetric matrix  $B^\top B$ . Let  $\text{Span}\{v_{i-1}\} = E_{r^*+i}A^*E_{r^*+i-1}W$ , and  $Bv_{i-1} \neq 0$  by Lemma 4.1' (iv) for  $i \in \{1, \dots, d^*\}$ . So,  $x_i \in \mathbb{R}^{>0}$  for all  $i \in \{1, \dots, d^*\}$ .

(iiia) Observe

$$w_i^* = E_{r^*+i}A^*E_{r^*+i-1}w_{i-1}^* \quad \text{for all } i \in \{1, \dots, d^*\}.$$

So  $w_i^* \neq 0$  for all  $i \in \{1, \dots, d^*\}$  by Lemma 4.1' (iv).

Hence,

$$W = \text{Span}(w_0^*, \dots, w_d^*)$$

by Lemma 4.1'. (iii).

(iiib) We have that

$$A^* w_i^* = E_{r^*+i+1} A^* w_i^* + E_{r^*+i} A^* w_i^* + E_{r^*+i-1} A^* w_i^* \quad (36.6)$$

$$= w_{i+1}^* + E_{r^*+i} A^* E_{r^*+i} w_i^* + E_{r^*+i-1} A^* E_{r^*+i} A^* E_{r^*+i-1} w_{i-1} \quad (36.7)$$

$$= w_{i+1}^* + a_i^* w_i^* + x_i^* w_{i-1}^*. \quad (36.8)$$

(iva) Clear for  $i = 0$ . Assume it is valid for  $0, \dots, i$ .

$$p_{i+1}^*(A^*) w_0^* = (A^* - a_i^* I) w_i^* - x_i^* w_{i-1}^* = w_{i+1}^*.$$

(ivb) By definition,

$$p_{d^*+1}^*(A^*) w_0^* = 0.$$

Since  $W = \{p(A^*) w_0^* \mid p \in \mathbb{C}[\lambda]\}$ ,  $p_{d^*+1}^*(A^*) W = 0$ , and  $p_{d^*+1}^*$  is a minimal polynomial, as  $w_0^*, w_1^*, \dots, w_{d^*}^*$  is a basis of  $W$ .

**Corollary 9.1'.** With the notation above, let  $W$  be a dualthin irreducible  $T$ -module with dual endpoint  $r^*(W)$ , and dual diameter  $d^*$ . Then,

(i)  $W$  is thin,

(ii)  $d^* = d = |\{i \mid E_i^* W \neq 0\}| - 1$ .

*Proof of Corollary 9.1'* Set as in Lemma {4.1}'.

$$w_i^* = p_i^*(A^*) w_0^* \in E_{r^*+i} W.$$

Then,  $w_0^*, w_1^*, \dots, w_{d^*}^*$  is a basis for  $W$ . We have  $W = M^* w_0^*$ .

So,  $E_i^* W = E_i^* M^* w_0^* = \text{Span}(E_i^* w_0^*)$ .

Thus,  $W$  is thin, and so, we have (ii).

Suppose  $r(W) = 1$ . Then  $d(W) = D - 2$  or  $D - 1$  by Lemma 14.1 (iii). See also Lemma 14.2.

Case  $d(W) = D - 2$ . Then

$$E_1 W = 0 \text{ implies } r^*(W) = 2.$$

$$E_1 W \neq 0 \text{ implies } r^*(W) = 1.$$

Case  $d(W) = D - 1$ . Then

$$r^*(W) = 1.$$

Up to isomorphism,

there are at most 3 thin irreducible  $T$ -modules with  $r(W) = 1$  and  $r^*(W) = 1$ ,

there are at most 1 thin irreducible  $T$ -modules with  $r(W) = 1$  and  $r^*(W) = 2$ ,

there are none thin irreducible  $T$ -modules with  $r(W) = 1$  and  $r^*(W) > 2$ .

By dual argument,

there are at most 3 thin irreducible  $T$ -modules with  $r^*(W) = 1$  and  $r(W) = 1$ ,

there are at most 1 thin irreducible  $T$ -modules with  $r^*(W) = 1$  and  $r(W) = 2$ ,

there are none thin irreducible  $T$ -modules with  $r$  and  $r(W) > 2$ .

**Conjecture 36.1.** *Let  $\Gamma = (X, E)$  be a thin distance regular graph of diameter  $D \geq 3$ . Let  $E_1$  be any primitive idempotent not equal to  $E_0$ .*

*Then the following are equivalent.*

(i) *For every vertex  $x \in X$ , there is no irreducible  $T$ -module  $W$  with  $r(W) > 2$ , and  $E_1 W \neq 0$ , there exists at most 1 irreducible  $T$ -module with  $r(W) = 2$ , and  $E_1 W \neq 0$ , and there exist at most 3 irreducible  $T$ -modules  $W$  with  $r(W) = 1$ , and  $E_1 W \neq 0$ .*

(ii)  *$\Gamma$  is  $Q$ -polynomial with respect to  $E_1$ .*

**Conjecture 36.2.** *Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$ ,  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Fix a vertex  $x \in X$ , and write  $E^* \equiv E_i^*(x)$ ,  $T \equiv T(x)$ . Let  $W$  denote an irreducible  $T$ -module with endpoint  $r$ , dual endpoint  $r^*$ , diameter  $d$  and dual diameter  $d^*$ .*

*Then the following hold.*

(i)  $d = d^*$ .

(ii) *there exists  $s \in \{r, \dots, r + d\}$  such that*

$$1 = \dim E_r^* W \leq \dim E_{r+1}^* W \leq \dots \leq \dim E_s^* W \geq \dots \geq \dim E_{r+d}^* W.$$

(iii) *there exists  $s^* \in \{r^*, \dots, r^* + d^*\}$  such that*

$$1 = \dim E_{r^*} W \leq \dim E_{r^*+1} W \leq \dots \leq \dim E_{s^*} W \geq \dots \geq \dim E_{r^*+d^*} W.$$

Let  $\Gamma = (X, E)$  be distance regular of diameter  $D \geq 3$ ,  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Fix a vertex  $x \in X$ , write  $E_i^* \equiv E_i^*(x)$  and  $T \equiv T(x)$ . Let  $W$  denote an irreducible module with endpoint 1.

**Conjecture 36.3.** *The following are equivalent.*

(i) *The sequence  $\dim E_1^*W, \dim E_2^*W, \dots, \dim E_D^*W$  equals*

$$1, 2, 2, \dots, 2, 1.$$

(ii)  *$v, Av, A_2v, \dots, A_{D_2}v, v^*, A^*v^*, A_2^*v^*, \dots, A_{D-2}^*v^*$  is a basis for  $W$ , where*

$$0 \neq v \in E_i^*W, \text{ and } v^* = |X|E_1v.$$

(iv)  *$v_1^+, v_2^+, \dots, v_D^+, v_2^-, v_3^-, \dots, v_{D-1}^-$  is a basis for  $W$ , where*

$$v_i^+ = E_i^*A_{i-1}v, \quad v_i^- = E_i^*A_{i+1}v.$$

**Problem.** Let  $B$  denote the orthogonal basis for  $W$  obtained by applying the Gram-Schmidt procedure to be basis in (iv).

Find the matrix representation  $A$  with respect to this basis.

I believe the entries are nicely factorable expressions in the basic variables,

$$q, s, s^*, r_1, r_1^*.$$

(Hint: use Theorem 35.1.)

If not, find some nice basis for  $W$ , and find the matrices representing  $A, A^*$  with respect to this basis.

Perhaps, some orthogonal basis based on (iii).

Algebraically, everything is determined by the intersection numbers and  $a_0(W)$ .

Combinatorically, certain quantities must be nonnegative integers. Does this give some new bounds, or other information on  $a_0(W)$ ?



## Chapter 37

# Generalized Adjacency Matrix

Friday, April 30, 1993

**Lemma 37.1.** *Let  $\Gamma = (X, E)$  be a distance-regular graph of diameter  $D \geq 3$ , and  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Fix a vertex  $x \in X$ , and write  $E_i^* \equiv E_i^*(x)$ , and  $T \equiv T(x)$ . Let  $W$  be an irreducible  $T$ -module of endpoint 1. If  $\dim E_2^*W = 1$ , then  $W$  is thin.*

*Proof.* Pick  $0 \neq v \in E_1^*W$ .

We want to show that

- $FR^i v \in \text{Span}(R^i v)$  for  $i \in \{0, \dots, D-1\}$ .
- $LR^i v \in \text{Span}(R^{i-1} v)$  for  $i \in \{1, \dots, D-1\}$ .

We have that

(1)  $FR^2 E_j^* \in \text{Span}(RFRE_j^*, R^2 FE_j^*, R^2 E_j^*)$  for  $i \in \{0, \dots, D-3\}$ .

(2)  $LR^2 E_j^* \in \text{Span}(RLRE_j^*, R^2 LE_j^*, F^2 E_j^*, FRFE_j^*, RF^2 E_j^*, RFE_j^*, FRE_j^*, RE_j^*)$  for  $i \in \{0, \dots, D-3\}$

by Corollary 30.1.

Claim (a)  $FR^i v \in \text{Span}(R^i v)$  for  $i \in \{0, \dots, D-2\}$ , (b)  $LR^i v \in \text{Span}(R^{i-1} v)$  for  $i \in \{1, \dots, D-2\}$ .

*Remark. Proof of Claim.* | (a) by Lemma 34.2, and an assumption

$$\dim E_1^*W = \dim E_2^*W = 1.$$

So,  $Rv \neq 0$ , and  $E_2^*W = \text{Span}(Rv)$ .

We may assume  $i \geq 2$ . Then  $R^{i-2}v \in E_{i-1}^*W$ ,

$$FR^i v = FR^2 R^{i-2}v, \quad \text{if } i \leq D-2, \quad = R(FR + RF + R)R^{i-2}v \quad (37.1)$$

$$\in R(\text{Span}(R^{i-1}v)) \quad (37.2)$$

$$= \text{Span}(R^i v), \quad (37.3)$$

by the induction hypothesis.

(b) If  $i \leq D-2$ , then  $R^{i-2}v \in E_{i-1}^*W$  with  $i-1 \leq D-3$ . Hence,

$$LR^i v = LR^2(R^{i-2}v) \quad (37.4)$$

$$= (RLR + R^2L + F^2R + FRF + RF^2 + RF + FR + R)R^{i-2}v \quad (37.5)$$

$$\in \text{Span}(R^{-1}v), \quad (37.6)$$

by induction and (a).

Suppose  $R^{D-1}v = 0$ . Then,

$$\text{Span}(v, Rv, \dots, R^{D-2}v) = \widetilde{W}.$$

is invariant under  $M$  and  $M^*$ , hence, under  $T$ .

Since  $W$  is irreducible,  $W = \widetilde{W}$ , and  $W$  is thin in this case.

Suppose  $R^{D-1}v \neq 0$ .

Observe:  $v, Av, \dots, A^{D-1}v \in \text{Span}(v, Rv, \dots, R^{D-1}v)$ .

Hence, each  $R^i v$  is a polynomial of degree  $i$  in  $A$  applied to  $v$ , and

$$\text{Span}(v, Av, \dots, A^{D-1}v) = \text{Span}(v, Rv, \dots, R^{D-1}v) = \text{Span}(v, A_1 v, \dots, A_{D-1} v).$$

Also,

$$A_D v = Jv - \left( \sum_{h=0}^{D-1} A_h \right) v \in \text{Span}(v, A_1 v, \dots, A_{D-1} v).$$

Thus,

$$Mv = \text{Span}(v, A_1 v, \dots, A_{D-1} v) = \widetilde{W}$$

is invariant under  $M$ ,  $M^*$ , and hence  $T$ . We have  $W = \widehat{W}$  and  $W$  is thin.  $\square$

**Definition 37.1.** Let  $\Gamma = (X, E)$  be any regular graph (not necessarily connected).

Let  $A$  be the adjacency matrix of  $\Gamma$ , and let  $J$  be the all 1's matrix.

Pick  $O \neq B \in \text{Mat}_X(\mathbb{C})$ .

$B$  is a generalized adjacency matrix, if

(i) for all vertices  $x, y \in X$ ,  $B_{xy} \neq 0$  implies  $A_{xy} \neq 0$  or  $x = y$ ,

(ii)  $B$  is in the subalgebra of  $\text{Mat}_X(\mathbb{C})$  generated by  $A$  and  $J$ .

**Example 37.1.** Any nonzero matrix of form

$$\alpha A + \beta I \quad (\alpha, \beta \in \mathbb{C})$$

is a generalized adjacency matrix.

If  $\Gamma$  is distance-regular, all generalized adjacency matrices are of this form.

Let  $\Gamma = (X, E)$  be a distance-regular graph of diameter  $D \geq 3$ . Assume  $\Gamma$  is thin, and  $Q$ -polynomial.

Pick a vertex  $x \in X$ , and write  $E_i^* \equiv E_i^*(x)$ ,  $T \equiv T(x)$ . Then,

$$E_1^* T E_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2, \tilde{A}^3),$$

and  $\dim E_1^* T E_1^* \leq 5$ .

We will produce a ‘nice’ spanning set

$$E_1^* T E_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A}, A^+ (= R^{-1} E_2^* A E_1^*), A^+ \tilde{A}).$$

**Lemma 37.2.** *Let  $\Gamma = (X, E)$  be a thin distance-regular graph of diameter  $D \geq 4$ .*

*Fix a vertex  $x \in X$ , and write  $E_i^* \equiv E_i^*(x)$  and  $R \equiv R(x)$ .*

*Let  $\Gamma_1$  denote the vertex subgraph induced on the first subconstituent of  $\Gamma$  relative to  $x$ . Then,*

$$\Delta = (R^{-1})^{i-1} E_i^* A_i E_1^*$$

*is a generalized adjacency matrix for  $\Gamma_1$  for all  $i \in \{1, \dots, D-3\}$ .*

*Proof.* Write  $T \equiv T(x)$ . Fix  $i \in \{1, \dots, D-3\}$ .

Recall  $R^{-1} \in T$  by Lemma 31.1 (iv).

$$\Delta \in E_1^* T E_1^* = \text{Span}(\tilde{J}, E_1^*, \tilde{A}, \tilde{A}^2, \dots)$$

by Lemma 34.3 (iv).

Hence,  $\Delta$  satisfied the condition (ii) of Definition 37.1.

To show (i), pick vertices  $y, z \in X$  such that

$$\partial(x, y) = \partial(x, z) = 1, \quad \partial(y, z) = 2.$$

We need to show

$$\Delta_{yz} = 0.$$

Suppose  $\Delta_{yz} \neq 0$ . Then,

$$\langle \Delta \hat{y}, \hat{z} \rangle \neq 0.$$

We will show this cannot occur.

Notation: Set

$$E_{ij}^* = E_i^*(x)E_j^*(y), \quad i, j \in \{0, 1, \dots, D\}.$$

Then,  $E_{ij}^*V = \text{Span}(\hat{w} \mid w \in X, \partial(x, w) = i, \partial(y, w) = j)$  for  $i, j \in \{0, 1, \dots, D\}$ .  
Let  $\delta$  denote the all 1's vector in  $V$ . Let

$$\delta_{ij} = E_{ij}^*\delta = \sum_{w \in X, \partial(x, w)=i, \partial(y, w)=j} \hat{w}.$$

Now,

$$\Delta\hat{y} \in E_1^*(x)V = E_{10}^*(x)V + E_{11}^*V + E_{12}^*V \quad (\text{orthogonal direct sum}).$$

So, there exist  $\delta_{10}^+ \in E_{10}^*(x)V$ ,  $\delta_{11}^+ \in E_{11}^*V$ , and  $\delta_{12}^+ \in E_{12}^*V$  such that

$$\Delta\hat{y} = \delta_{10}^+ + \delta_{11}^+ + \delta_{12}^+.$$

Observe:  $\hat{z} \in E_{12}^*V$  is not orthogonal to  $\Delta\hat{y}$ .

So,  $\delta_{12}^+ \neq 0$ .

Observe:

$$R^{i-1}(\delta_{10}^+ + \delta_{11}^+ + \delta_{12}^+) = R^{i-1}\Delta\hat{y} \quad (37.7)$$

$$= R^{i-1}(R^{-1})^{i-1}E_i^*A_iE_1^*\hat{y} \quad (37.8)$$

$$= E_i^*A_iE_1^*\hat{y} \quad (37.9)$$

$$= \delta_{ii} \quad (37.10)$$

$$\in E_{ii}^*V. \quad (37.11)$$

*Remark.* It is because on each irreducible hin module with standard basis  $w_r, w_{r+1}, \dots, w_{r+d}$ ,

$$R^{-1}w_i = w_{i-1}, \quad i > r, \quad R^{-1}w_r = 0,$$

and  $E_1^*V$  is an orthogonal direct sum of irreducible modules and  $r \leq 1$ .

Bu we can control  $R^{i-1}\delta_{10}^+$ ,  $R^{i-1}\delta_{11}^+$ , also.

Claim.  $RE_{jj}^*V \subseteq E_{j+1, j+1}^*V + E_{j+1, j}^*V$ ,  $j \in \{1, \dots, D-1\}$ .

*Proof of Claim.* Clear.

By Claim

$$R^{i-1}\delta_{10}^+ \in E_{i, i-1}^*V, \quad \text{and} \quad (37.12)$$

$$R^{i-1}\delta_{11}^+ \in E_{i, i-1}^*V + E_{i, i}^*V. \quad (37.13)$$

Hence, we conclude that

$$R^{i-1}\delta_{12}^+ = R^{i-1}\Delta\hat{y} - R^{i-1}\delta_{10}^+ - R^{i-1}\delta_{11}^+ \in E_{i, i-1}^*V + E_{i, i}^*V.$$

But now

$$0 = E_{i,i+1}^* R^{i-1} \delta_{12}^+ = E_{i,i+1}^* A^{i-1} E_{12}^* \delta_{12}^+ = R(y)^{i-1} \delta_{12}^+. \quad (37.14)$$

By Lemma 32.1 (ii),

$$R(y)^{i-1} : E_2^*(y)V \longrightarrow E_{i+1}^*V$$

is one-to-one, since  $\Gamma$  is thin, and  $i-1 \leq D-4$ .

So,  $\delta_{12}^+ = 0$  by (37.14).

But this contradicts (2). Hence our assumption  $\Delta_{yz} \neq 0$  is false, and the condition (i) of the definition of generalised adjacency matrices is satisfied.

This proves the lemma.  $\square$



## Chapter 38

# An Injection from $E_{11}^*$ to $E_{22}^*$

Monday, May 3, 1993

**Lemma 38.1.** *Let  $\Gamma = (X, E)$  be a thin distance-regular graph of diameter  $D \geq 5$ , and  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ . Pick vertices  $x, y \in X$  such that  $\partial(x, y) = 1$ , and write  $E_{ij}^* := E_i^*(x)E_j^*(y)$  for  $i, j \in \{0, 1, \dots, D\}$ . Then the following hold.*

- (i)  $E_{22}^*AE_{11}^* : E_{11}^*V \rightarrow E_{22}^*V$  is one-to-one.
- (ii) For every  $z \in X$  such that  $\partial(x, z) = \partial(y, z) = 1$ , there is  $w \in X$  such that

$$\partial(w, x) = \partial(w, y) = 2, \partial(w, z) = 1.$$

*Proof.*

- (i) Write  $E_i^* \equiv E_i^*(x)$ ,  $R \equiv R(x)$ ,  $F \equiv F(x)$ ,  $L \equiv L(x)$ , and  $T \equiv T(x)$ .

Suppose there exists

$$0 \neq v \in E_{11}^*V \text{ such that } E_{22}^*AE_{11}^*v = 0, \quad (38.1)$$

Claim 1.  $E_{34}^*A^2E_{12}^*AE_{11}^*v \neq 0$ .

*Proof of Claim 1.* Recall by Lemma 32.1 (ii),  $(3 \leq 5 - 2 \leq D - 2t)$ ,

$$R(y)^3 : E_1^*(y)V \rightarrow E_4^*(y)V$$

is one-to-one.

Since  $v \in E_1^*(y)V$ , we find

$$0 \neq R^4(y)v \quad (38.2)$$

$$= E_4^*(y)A^3E_1^*(y)v \quad (38.3)$$

$$= E_4^*(y)A^2E_2^*(y)AE_{11}^*v \quad (38.4)$$

$$= E_4^*(y)A^2 \left( \sum_{h=0}^D E_{h,2}^* \right) AE_{11}^*v \quad (38.5)$$

$$= E_4^*(y)A^2(E_{12}^* + E_{22}^*)AE_{11}^*v \quad (38.6)$$

$$= E_4^*(y)A^2E_{12}^*AE_{11}^*v \quad (38.7)$$

$$= E_{34}^*(y)A^2E_{12}^*AE_{11}^*v, \quad (38.8)$$

by (38.1). This proves the claim.

By Theorem 30.1 (i),

$$0 = (g_3^- R^2 F + RFR + g_3^+ FR^2 - \gamma R^2)E_1^*. \quad (38.9)$$

*Remark.* Theorem 30.1 (i) states

$$(g_i^- FL^2 + LFL + g_i^+ L^2 F - \gamma L^2)E_i^* = O \text{ for } i \in \{2, \dots, D\}.$$

For  $i = 3$ ,

$$E_1^*(g_3^- FL^2 + LFL + g_3^+ L^2 F - \gamma L^2)E_3^* = O.$$

Taking the transpose, we have

$$(g_3^- R^2 F + RFR + g_3^+ FR^2 - \gamma R^2)E_1^* = O.$$

Hence, we have (38.9).

Multiplying each term on the left by  $E_1^*(y)$ , on the right by  $E_1^*(y)$ , we find

$$O = g_3^- E_{34}^* R^2 F E_{11}^* + E_{34}^* R F R E_{11}^* + g_3^+ E_{34}^* F R^2 E_{11}^* - \gamma E_{34}^* R^2 E_{11}^* \quad (38.10)$$

$$= g_3^- E_{34}^* A^2 E_{12}^* A E_{11}^* + E_{34}^* A E_{23}^* A E_{22}^* A E_{11}^* + g_3^+ E_{34}^* A E_{33}^* A E_{22}^* A E_{11}^*. \quad (38.11)$$

Applying this to  $v$ , we find by (38.1) that

$$0 = g_3^- E_{34}^* A^2 E_{12}^* A E_{11}^* v.$$

So,  $g_3^- = 0$  by Claim 1. But by Lemma 30.1,

$$g_3^- = \frac{\theta_1^* - \theta_0^*}{\theta_1^* - \theta_3^*} \neq 0,$$

a contradiction. □



Let  $\Gamma, x, y$  be as in Lemma 38.1. We saw in Lemma 37.2,

$$R^{-1}E_2^*A_2E_1^*\hat{y} = \delta_{10}^+ + \delta_{11}^+,$$

where

$$\delta_{10}^+ \in E_{10}^*V = \text{Span}(\hat{y}), \quad \delta_{11}^+ \in E_{11}^*V.$$

**Definition 38.1.** Define  $\Psi = \Psi(x, y) \in \mathbb{C}$  by  $\delta_{10}^+ = \Psi\hat{y}$ .

We will show that  $\Psi(x, y)$  is independent of  $x, y$ .

Observe  $R^{-1}, A_i, E_i^* \in \text{Mat}_X(\mathbb{Q})$ . So  $\Psi \in \mathbb{Q}$ .

Firstly, show

$$\Psi(x, y) = \Psi(y, x).$$

**Lemma 38.2.** *With the notation of Lemma 38.1, the following hold.*

- (i)  $E_{22}^*AE_{11}^*\delta_{11}^+ = \delta_{22}$ .
- (ii)  $\$E^*\_ \{21\}AE^*\{11\}^+ \{11\} = - (x, y) \_ \{21\}$ .
- (iii)  $\langle \delta_{11}^+, \delta_{11} \rangle = \frac{a_2}{c_2} - \Psi(x, y)$ .
- (iv)  $\Psi(x, y) = \Psi(y, x)$ .
- (v)  $E_{12}^*AE_{11}^*\delta_{11}^+ = -\Psi(x, y)\delta_{12}$ .

*Proof.* Write  $\Psi \equiv \Psi(x, y)$ ,  $R \equiv R(x)$ ,  $E_i^* \equiv E_i^*(x)$ , etc.

(i) We have

$$R(\delta_{11}^+ + \Psi\hat{y}) = R(\delta_{11}^+ + \delta_{10}^+) \tag{38.12}$$

$$= R(R^{-1}(E_2^*A_2E_1^*))\hat{y} \tag{38.13}$$

$$= E_2^*A_2E_1^*\hat{y} \tag{38.14}$$

$$= \delta_{22}. \tag{38.15}$$

So,

$$\delta_{22} = R(\delta_{11}^+ + \Psi\hat{y}) \tag{38.16}$$

$$= E_2^*AE_1^*(\delta_{11}^+ + \Psi\hat{y}) \tag{38.17}$$

$$= E_{22}^*AE_{11}^*\delta_{11}^+ + \Psi E_{22}^*AE_{10}^*\hat{y}. \tag{38.18}$$

The second term is zero.

(ii) We have

$$0 = E_{21}^* \delta_{22} \quad (38.19)$$

$$= E_{21}^* R(\delta_{11}^+ + \Psi \hat{y}) \quad (38.20)$$

$$= E_{21}^* A E_{11}^* \delta_{11}^+ + \Psi E_{21}^* A E_{10}^* \hat{y} \quad (38.21)$$

$$= E_{21}^* A E_{11}^* + \Psi \delta_{21}. \quad (38.22)$$

(iii) We have

$$p_{22}^1 = \delta_{22} \|^2 \quad (38.23)$$

$$= \langle \delta_{22}, \delta_{21} + \delta_{22} + \delta_{23} \rangle \quad (38.24)$$

$$= \langle R(\delta_{11}^+ + \Psi \hat{y}), \delta_{21} + \delta_{22} + \delta_{23} \rangle \quad (38.25)$$

$$= \langle \delta_{11}^+ + \Psi \hat{y}, L(\delta_{21} + \delta_{22} + \delta_{23}) \rangle \quad (38.26)$$

$$= b_1 \langle \delta_{11}^+ + \Psi \hat{y}, \delta_{10} + \delta_{11} + \delta_{12} \rangle \quad (38.27)$$

$$= b_1 (\langle \delta_{11}^+, \delta_{11} \rangle + \Psi). \quad (38.28)$$

So,

$$\langle \delta_{11}^+, \delta_{11} \rangle = b_1^{-1} p_{22}^1 - \Psi = \frac{a_2}{c_2} - \Psi.$$

*Remark.*

$$b_1^{-1} p_{22}^1 = b_1^{-1} \frac{k_1}{k_1} p_{22}^1 = b_1^{-1} \frac{1}{k_1} k_2 p_{12}^2 = b_1^{-1} \frac{b_1}{c_2} a_2 = \frac{a_2}{c_2}.$$

(iv) Interchanging roles of  $x, y$  above, we find there exists  $\delta_{11}^{+'} \in E_{11}^* V$  such that

$$R(y)^{-1} E_2^*(y) A_2 E_1^*(y) \hat{x} = \delta_{11}^{+'} + \Psi(y, x) \hat{y}.$$

Then,

$$E_{22}^* A E_{11}^* (\delta_{11}^{+'}) = \delta_{22}.$$

So,

$$E_{22}^* A E_{11}^* (\delta_{11}^+ - \delta_{11}^{+'}) = 0.$$

Hence,  $\delta_{11}^+ = \delta_{11}^{+'}$  since

$$E_{22}^* A E_{11}^* : E_{11}^* V \rightarrow E_{22}^* V$$

is one-to-one.

Now,

$$\frac{a_2}{c_2} - \Psi(x, y) = \langle \delta_{11}^+, \delta_{11} \rangle = \langle \delta_{11}^{+'}, \delta_{11} \rangle = \frac{a_2}{c_2} - \Psi(y, x).$$

Thus,

$$\Psi(x, y) = \Psi(y, x).$$

(v) Immediate from (ii), (iv).

□

## Chapter 39

### $A^+$ and $A^-$

**Wednesday, May 5, 1993**

Assume  $\Gamma = (X, E)$  is thin, distance regular of diameter  $D \geq 5$ , and  $Q$ -polynomial with respect to  $E_0, E_1, \dots, E_D$ .

Fix a vertex  $x \in X$ , write  $E^* \equiv E_i^*(x)$ ,  $R \equiv R(x)$ ,  $T \equiv T(x)$ .

Pick  $y \in X$  with  $\partial(x, y) = 1$ . Write  $E_{i,j}^* \equiv E^*(x)E^*(y)$ ,  $\delta_{ij} = E_{ij}^*\delta$ , and  $\tilde{A} = E_1^*AE_1^*$ .

Recall that  $\delta_{11}^+ \in E_{11}^*V$  and

$$R^{-1}E_2^*A_2E_1^*\hat{y} = \delta_{11}^+ + \Psi(x, y)\hat{y}.$$

We saw  $\Psi(x, y) = \Psi(y, x)$ . We shall show below that  $\Psi(x, y)$  is independent of edge  $xy$ .

**Lemma 39.1.** *With the above notation, set  $\Psi := \Psi(x, y)$ . Then the following hold.*

$$(i) \quad \delta_{11}^- = \tilde{A}\delta_{11}^+ - \left(\frac{a_2}{c_2} - \Psi\right)\hat{y} + \Psi\delta_{12} \in E_{11}^*V.$$

$$(ii) \quad \delta_{11}^-(x, y) = \delta_{11}^{-1}(y, x).$$

*Proof.*

$$(i) \quad \delta_{12}^- \in E_{12}^*V, \delta_{11}^- \in E_{11}^*V \text{ and } \delta_{10}^- \in E_{10}^*V, \text{ and}$$

$$\tilde{A}\delta_{11}^+ = \delta_{12}^- + \delta_{11}^- + \delta_{10}^-, \tag{39.1}$$

$$\delta_{12}^- = E_{12}^*AE_{11}^*\delta_{11}^+ = -\Psi(x, y)\delta_{12}, \tag{39.2}$$

by Lemma 38.2 (v).

Also,  $\delta_{10}^- = \sigma \hat{y}$  for some  $\sigma \in \mathbb{C}$ , where

$$\sigma = \langle \tilde{A}\delta_{11}^+, \hat{y} \rangle \quad (39.3)$$

$$= \langle \delta_{11}^+, \tilde{A}\hat{y} \rangle \quad (39.4)$$

$$= \langle \delta_{11}^+, \delta_{11} \rangle \quad (39.5)$$

$$= \frac{a_2}{c_2} - \Psi. \quad (39.6)$$

Solving for  $\delta_{11}^-$  in (39.1), using (39.2) and (39.6), we have

$$\delta_{11}^- = \tilde{A}\delta_{11}^+ - \delta_{12}^- - \delta_{10}^- \quad (39.7)$$

$$= A\delta_{11}^+ + \Psi\delta_{12} - \left( \frac{a_2}{c_2} - \Psi \right) \hat{y}. \quad (39.8)$$

(ii) Since

$$\delta_{11}^- = E_{11}^* A E_{11}^* \delta_{11}^+,$$

we have  $\delta_{11}^+(x, y) = \delta_{11}^+(y, x)$ .

□

## Chapter 40

# Title of the Chapter

Wednesday, February 17, 1993 # Edit Date



# Bibliography

- A.E. Brouwer, A.M. Cohen, A. N. (1989). *Distance-Regular Graphs*. Springer-Verlag, Berlin Heidelberg. 3-540-50619-5, 0-387-50619-5.
- Charles W. Curtis, I. R. (2006). *Representation Theory of Finite Groups and Associative Algebras*. Chelsea Pub Co, uk edition. 978-1138359420.
- Terwilliger, P. (1992). The subconstituent algebra of an association scheme. i. *J. Algebraic Combin.*, 1(4):363–388.
- Terwilliger, P. (1993a). The subconstituent algebra of an association scheme. ii. *J. Algebraic Combin.*, 2(1):73–103.
- Terwilliger, P. (1993b). The subconstituent algebra of an association scheme. iii. *J. Algebraic Combin.*, 2(2):177–210.
- Terwilliger, P. (1995). A new inequality for distance-regular graphs. *Discrete Math.*, 137(1-3):319–332.
- Xie, Y. (2015). *Dynamic Documents with R and knitr*. Chapman and Hall/CRC, Boca Raton, Florida, 2nd edition. 978-0821840665.
- Xie, Y. (2017). *bookdown: Authoring Books and Technical Documents with R Markdown*. Chapman and Hall/CRC, Boca Raton, Florida, 1st edition. ISBN 978-1138469280.
- Yihui Xie, J.J Allaire, G. G. (2018). *R Markdown: The Definitive Guide*. Chapman and Hall/CRC, Boca Raton, Florida, 1st edition. 978-1138359420.

# Index

- $Q$ -polynomial, 128, 140
- association matrix, 106
- association scheme, 106
- automorphism, 19
- bipartite, 13, 17
- Cayley graph, 19
- character, 22
- complete graph, 94
- connected, 10
- diameter, 29
- distance, 10
- distance-regular, 71
- distance-transitive, 52
- dual associate matrix, 115
- dual Bose-Mesner algebra, 115
- dual endpoint, 219
- dual thin, 42
- endpoint, 29
- generalized adjacency matrix, 226
- graph, 7
- isomorphic, 32
- measure, 60
- module, 10
- multiplicity, 32
- Norton algebra, 124
- path, 9
- reducible, 13
- regular, 9
- restricted, 42
- subconstituent algebra, 10
- symmetrix, 106
- thin, 42
- valency, 9
- vertex transitive, 19