

# Lecture Note on Terwilliger Algebra

P. Terwilliger, edited by H. Suzuki

2022-11-14



# Contents

<b>About this lecturenote</b>	<b>5</b>
Setting . . . . .	5
Another Host . . . . .	6
<b>1 Lecture 1</b>	<b>7</b>
<b>2 Lecture 2</b>	<b>13</b>
<b>3 Lecture 3</b>	<b>19</b>
<b>4 Applications</b>	<b>21</b>
4.1 Example one . . . . .	21
4.2 Example two . . . . .	21
<b>5 Final Words</b>	<b>23</b>



# About this lecturenote

## Setting

This note is created by `bookdown` package on RStudio.

1. Log-in to my GitHub Account
2. Go to RStudio/bookdown-demo repository: <https://github.com/rstudio/bookdown-demo>
3. Use This Template
4. Input Repository Name
5. Select Public - default
6. Create repository from template
7. From Code download ZIP
8. Move the extracted folder into a favorite directory
9. Open RStudio Project in the folder
10. Use Terminal in the bottom left pane
  - confirm that the current directory is the home directory of the project by `pwd`
11. (failed to proceed by ssh)
12. Use Console
  1. `library(usethis)`
  2. `use_git()`
  3. `use_github()` — Error
  4. `gh_token_help()`
  5. `create_github_token()`: create a token in the github page. Copy the token
  6. `gitcreds::gitcreds_set()`: paste the token, the token is to be expired in 30 days
13. Use Terminal
  1. `git remote add origin https://github.com/icu-hsuzuki/t-algebra.git`
  2. `git push -u origin main`
  3. type in the password of the computer
14. Use GIT in R Studio

## Another Host

1. `library(usethis)`
2. `use_git()`
3. `create_github_token()`
4. `gitcreds::gitcreds_set()`: Replace these credentials

# Chapter 1

## Lecture 1

Wednesday, January 20, 1993


A graph (undirected, without loops or multiple edges) is a pair  $\Gamma = (X, E)$ , where

$$X = \text{finite set (of vertices)} \quad (1.1)$$

$$E = \text{set of (distinct) 2-element subsets of } X \text{ (= edges of ) } \Gamma. \quad (1.2)$$

vertices  $x$  and  $y \in X$  are adjacent if and only if  $xy \in E$ .

**Example 1.1.** Let  $\Gamma$  be a graph.  $X = \{a, b, c, d\}$ ,  $E = \{ab, ac, bc, bd\}$ .



Set  $n = |X|$ , the order of  $\Gamma$ .

Pick a field  $K$  ( $= \mathbb{R}$  or  $\mathbb{C}$ ). Then  $\text{Mat}_X(K)$  denotes the  $K$  algebra of all  $n \times n$  matrices with entries in  $K$ . (rows and columns are indexed by  $X$ )

*Adjacency matrix*  $A \in \text{Mat}_X(K)$  is defined by

$$A_{xy} = \begin{cases} 1 & \text{if } xy \in E \\ 0 & \text{else .} \end{cases} \quad (1.3)$$

**Example 1.2.** Let  $a, b, c, d$  be labels of rows and columns. Then

$$A = \begin{matrix} & \begin{matrix} a & b & c & d \end{matrix} \\ \begin{matrix} a \\ b \\ c \\ d \end{matrix} & \begin{pmatrix} 0 & 1 & 1 & 0 \\ 1 & 0 & 1 & 1 \\ 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 0 \end{pmatrix} \end{matrix}$$

The subalgebra  $M$  of  $\text{Mat}_X(K)$  generated by  $A$  is called the *Bose-Mesner algebra* of  $\Gamma$ .

Set  $V = K^n$ , the set of  $n$ -dimensional column vectors, the coordinates are indexed by  $X$ .

Let  $\langle \cdot, \cdot \rangle$  denote the Hermitean inner product:

$$\langle u, v \rangle = u^\top \cdot v \quad (u, v \in V)$$

$V$  with  $\langle \cdot, \cdot \rangle$  is the *standard module* of  $\Gamma$ .

$M$  acts on  $V$ : For every  $x \in X$ , write

$$\hat{x} = \begin{pmatrix} 0 \\ \vdots \\ 0 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}$$

where 1 is at the  $x$  position.

Then

$$A\hat{x} = \sum_{y \in X, xy \in E} \hat{y}.$$

Since  $A$  is a real symmetrix matrix,

$$V = V_0 + V_1 + \cdots + V_r \quad \text{some } r \in \mathbb{Z}^{\geq 0},$$

the orthogonal direct sum of maximal  $A$ -eigenspaces.

Let  $E_i \in \text{Mat}_X(K)$  denote the orthogonal projection,

$$E_i : V \longrightarrow V_i.$$

Then  $E_0, \dots, E_r$  are the primitive idempotents of  $M$ .

$$M = \text{Span}_K(E_0, \dots, E_r),$$

$$E_i E_j = \delta_{ij} E_i \quad \text{for all } i, j, \quad E_0 + \cdots + E_r = I.$$



Let  $\theta_i$  denote the eigenvalue of  $A$  for  $V_i$  in  $\mathbb{R}$ . Without loss of generality we may assume that

$$\theta_0 > \theta_1 > \dots > \theta_r.$$

Let

$$m_i = \text{the multiplicity of } \theta_i = \dim V_i = \text{rank } E_i.$$

Set

$$\text{Spec}(\Gamma) = \begin{pmatrix} \theta_0, & \theta_1, & \dots, & \theta_r \\ m_0, & m_1, & \dots, & m_r \end{pmatrix}.$$

**Problem.** What can we say about  $\Gamma$  when  $\text{Spec}(\Gamma)$  is given?

The following Lemma 1.1, is an example of Problem.

For every  $x \in X$ ,

$$k(x) \equiv \text{valency of } x \equiv \text{degree of } x \equiv |\{y \mid y \in X, xy \in E\}|.$$

**Definition 1.1.** The graph  $\Gamma$  is regular of valency  $k$  if  $k = k(x)$  for every  $x \in X$ .

**Lemma 1.1.** *With the above notation,*

1.  $\theta_0 \leq \max\{k(x) \mid x \in X\} = k^{\max}$ .
2. *If  $\Gamma$  is regular of valency  $k$ , then  $\theta_0 = k$ .*

*Proof.* 1. Without loss of generality we may assume that  $\theta_0 > 0$ , else done.

Let  $v := \sum_{x \in X} \alpha_x \hat{x}$  denote the eivenvector for  $\theta_0$ .

Pick  $x \in X$  with  $|\alpha_x|$  maximal. Then  $|\alpha_x| \neq 0$ .

Since  $Av = \theta_0 v$ ,

$$\theta_0 \alpha_x = \sum_{y \in X, xy \in E} \alpha_y.$$

So,

$$\theta_0 |\alpha_x| = |\theta_0 \alpha_x| \leq \sum_{y \in X, xy \in E} |\alpha_y| \leq k(x) |\alpha_x| \leq k^{\max} |\alpha_x|.$$

2. All 1's vector  $v = \sum_{x \in X} \hat{x}$  satisfies  $Av = kv$ .

□

### Subconstituent Algebra

Let  $x, y \in X$  and  $\ell \in \mathbb{Z}^{\geq 0}$ .

**Definition 1.2.** A path of length  $\ell$  connecting  $x, y$  is a sequence

$$x = x_0, x_1, \dots, x_\ell = y, \quad x_i \in X, \quad 0 \leq i \leq \ell$$

such that  $x_i x_{i+1} \in E$  for  $0 \leq i \leq \ell - 1$ .

**Definition 1.3.** The distance  $\partial(x, y)$  is the length of a shortest path connecting  $x$  and  $y$ .

$$\partial(x, y) \in \mathbb{Z}^{\geq 0} \cup \{\infty\}.$$

**Definition 1.4.** The graph  $\Gamma$  is connected if and only if  $\partial(x, y) < \infty$  for all  $x, y \in X$ .

From now on, assume that  $\Gamma$  is connected with  $|X| \geq 2$ .

Set

$$d_\Gamma = d = \max\{\partial(x, y) \mid x, y \in X\} \equiv \text{the diameter of } \Gamma.$$

Fix a ‘base’ vertex  $x \in X$ .

**Definition 1.5.**

$$d(x) = \text{the diameter with respect to } x = \max\{\partial(x, y) \mid y \in X\} \leq d.$$

Observe that

$$V = V_0^* + V_1^* + \cdots + V_{d(x)}^* \quad (\text{orthogonal direct sum}),$$

where

$$V_i^* = \text{Span}_K(\hat{y} \mid \partial(x, y) = i) \equiv V_i * (x)$$

and  $V_i^* = V_i^*(x)$  is called the  $i$ -th subconstituent with respect to  $x$ .

Let  $E_i^* = E_i^*(x)$  denote the orthogonal projection

$$E_i^* : V \longrightarrow V_i^*(x).$$

View  $E_i^*(x) \in \text{Mat}_X(K)$ . So,  $E_i^*(x)$  is diagonal with  $yy$  entry

$$(E_i^*(x))_{yy} = \begin{cases} 1 & \text{if } \partial(x, y) = i \\ 0 & \text{else,} \end{cases} \quad \text{for } y \in X.$$

Set

$$M^* = M^*(x) \equiv \text{Span}_K(E_0^*(x), \dots, E_{d(x)}^*(x)).$$

Then  $M^*(x)$  is a commutative subalgebra of  $\text{Mat}_X(K)$  and is called the *dual Bose-Mesner algebra with respect to  $x$* .

**Definition 1.6** (Subconstituent Algebra). Let  $\Gamma = (X, E)$ ,  $x, M, M^*(x)$  be as above. Let  $T = T(x)$  denote the subalgebra of  $\text{Mat}_X(K)$  generated by  $M$  and  $M^*(x)$ .  $T$  is the *subconstituent algebra* of  $\Gamma$  with respect to  $x$ .

**Definition 1.7.** A  $T$ -module is any subspace  $W \subset V$  such that  $aw \in W$  for all  $a \in T$  and  $w \in W$ .

$T$ -module  $W$  is *irreducible* if and only if  $W \neq 0$  and  $W$  does not properly contain a nonzero  $T$ -module.

For any  $a \in \text{Mat}_X(K)$ , let  $a^*$  denote the conjugate transpose of  $a$ .

Observe that

$$\langle au, v \rangle = \langle u, a^*v \rangle \quad \text{for all } a \in \text{Mat}_X(K), \text{ and for all } u, v \in V.$$

**Lemma 1.2.** *Let  $\Gamma = (X, E)$ ,  $x \in X$  and  $T \equiv T(x)$  be as above.*

1. *If  $a \in T$ , then  $a^* \in T$ .*
2. *For any  $T$ -module  $W \subset V$ ,*

$$W^\perp := \{v \in V \mid \langle w, v \rangle = 0, \text{ for all } w \in W\}$$

*is a  $T$ -module.*

3.  *$V$  decomposes as an orthogonal direct sum of irreducible  $T$ -modules.*

*Proof.* 1. It is because  $T$  is generated by symmetric real matrices

$$A, E_0^*(x), E_1^*(x), \dots, E_{d(x)(x)}^*.$$

2. Pick  $v \in W^\perp$  and  $a \in T$ , it suffices to show that  $av \in W^\perp$ . For all  $w \in W$ ,

$$\langle w, av \rangle = \langle a^*w, v \rangle = 0$$

as  $a^* \in T$ .

3. This is proved by the induction on the dimension of  $T$ -modules. If  $W$  is an irreducible  $T$ -module of  $V$ , then

$$V = W + W^\perp \quad (\text{orthogonal direct sum}).$$

□

**Problem.** What does the structure of the  $T(x)$ -module tell us about  $\Gamma$ ?

Study those  $\Gamma$  whose modules take ‘simple’ form. The  $\Gamma$ ’s involved are highly regular.

- Remark.*
1. The subconstituent algebra  $T$  is semisimple as the left regular representation of  $T$  is completely reducible. See Curtis-Reiner 25.2.
  2. The inner product  $\langle a, b \rangle_T = \text{tr}(a^\top \bar{b})$  is nondegenerate on  $T$ .
  3. In general,

$$T: \text{Semisimple and Artinian} \Leftrightarrow T: \text{Artinian with } J(T) = 0$$

$$\Leftrightarrow T: \text{Artinian with nonzero nilpotent element}$$

$$\Leftrightarrow T \subset \text{Mat}_X(K) \text{ such that for all } a \in T \text{ is normal.}$$



## Chapter 2

## Lecture 2

Friday, January 22, 1993

In this lecture we use the Perron Frobenius theory of nonnegative matrices to obtain information on eigenvalues of a graph.

Let  $K = \mathbb{R}$ . For  $n \in \mathbb{Z}_{>0}$ , pick a symmetric matrix  $C \in \text{Mat}_n(\mathbb{R})$ .

**Definition 2.1.** The matrix  $C$  is *reducible* if and only if there is a bipartition  $\{1, 2, \dots, n\} = X^+ \cup X^-$  (disjoint union of nonempty sets) such that  $C_{ij} = 0$  for all  $i \in X^+$ , and for all  $j \in X^-$ , and for all  $i \in X^-$ , and for all  $j \in X^+$ , i.e.,

$$C \sim \begin{pmatrix} * & O \\ O & * \end{pmatrix}.$$

**Definition 2.2.** The matrix  $C$  is *bipartite* if and only if there is a bipartition  $\{1, 2, \dots, n\} = X^+ \cup X^-$  (disjoint union of nonempty sets) such that  $C_{ij} = 0$  for all  $i, j \in X^+$ , and for all  $i, j \in X^-$ , i.e.,

$$C \sim \begin{pmatrix} O & * \\ * & O \end{pmatrix}.$$

**Note.**

1. If  $C$  is bipartite, for every eigenvalue  $\theta$  of  $C$ ,  $-\theta$  is an eigenvalue of  $C$  such that  $\text{mult}(\theta) = \text{mult}(-\theta)$ .

Indeed, let  $C = \begin{pmatrix} O & A \\ B & O \end{pmatrix}$ ,

$$\begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \theta \begin{pmatrix} x \\ y \end{pmatrix} \Leftrightarrow \begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ -y \end{pmatrix} = -\theta \begin{pmatrix} x \\ -y \end{pmatrix},$$

where  $Ay = \theta x$  and  $Bx = \theta y$ .

2. If  $C$  is bipartite,  $C^2$  is reducible.
3. The matrix  $C$  is irreducible and  $C^2$  is reducible, if  $C_{ij} \geq 0$  for all  $i, j$  and  $C$  is reducible. (Exercise)

*Remark.* Note 1. Even if  $C$  is not symmetric

$$\begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \theta \begin{pmatrix} x \\ y \end{pmatrix} \Leftrightarrow \begin{pmatrix} O & A \\ B & O \end{pmatrix} \begin{pmatrix} x \\ -y \end{pmatrix} = -\theta \begin{pmatrix} x \\ -y \end{pmatrix}$$

holds. So the geometrix multiplicities coincide. How about the algebraic multiplicities?

Note 3. Set  $x \sim y$  if and only if  $C_{xy} > 0$ . So the graph may have loops. Then

$$(C^2)_{xy} > 0 \Leftrightarrow \text{if there exists } z \in X \text{ such that } x \sim z \sim y.$$

Note that  $C$  is irreducible if and only if  $\Gamma(C)$  is connected. Let

$$X^+ = \{y \mid \text{there is a path of even length from } x \text{ to } y\} \quad (2.1)$$

$$X^- = \{y \mid \text{there is no path of even length from } x \text{ to } y\} \neq \emptyset. \quad (2.2)$$

If there is an edge  $y \sim z$  in  $X^+$  and  $w \in X^-$ . Then there would be a path from  $x$  to  $y$  of even length. So  $e(X^+, X^+) = e(X^-, X^-) = 0$ .

**Theorem 2.1** (Perron-Frobenius). *Given a matrix  $C$  in  $\text{Mat}_n(\mathbb{R})$  such that*

- a.  $C$  is symmetric.
- b.  $C$  is irreducible.
- c.  $C_{ij} \geq 0$  for all  $i, j$ . Let  $\theta_0$  be the maximal eigenvalue of  $C$  with eigenspace  $V_0 \subseteq \mathbb{R}^n$ , and let  $\theta_r$  be the maximal eigenvalue of  $C$  with eigenspace  $V_r \subseteq \mathbb{R}^n$ . Then the following hold.

$$1. \text{ Suppose } 0 \neq v = \begin{pmatrix} \alpha_1 \\ \alpha_2 \\ \vdots \\ \alpha_n \end{pmatrix} \in V_0. \text{ Then } \alpha_0 > 0 \text{ for all } i, \text{ or } \alpha_i < 0 \text{ for all } i.$$

$$2. \dim V_0 = 1.$$

$$3. \theta_r \geq -\theta_0.$$

$$4. \theta_r = \theta_0 \text{ if and only if } C \text{ is bipartite.}$$

First, we prove the following lemma.

**Lemma 2.1.** *Let  $\langle, \rangle$  be the dot product in  $V = \mathbb{R}^n$ . Pick a symmetric matrix  $B \in \text{Mat}_n(\mathbb{R})$ . Suppose all eigenvalues of  $B$  are nonnegative. (i.e.,  $B$  is positive semidefinite.) Then there exist vectors  $v_1, v_2, \dots, v_n \in V$  such that  $B_{ij} = \langle v_i, v_j \rangle$  for  $(1 \leq i, j \leq n)$ .*

By elementary linear algebra, there exists an orthonormal basis  $w_1, w_2, \dots, w_n$  of  $V$  consisting of eigenvectors of  $B$ . Set the  $i$ -th column of  $P$  is  $w_i$  and  $D = \text{diag}(\theta_1, \dots, \theta_n)$ . Then  $P^\top P = I$  and  $BP = PD$ .

Hence,

$$B = PDP^{-1} = PDP^\top = QQ^\top,$$

where

$$Q = P \cdot \text{diag}(\sqrt{\theta_1}, \sqrt{\theta_2}, \dots, \sqrt{\theta_n}) \in \text{Mat}_n(\mathbb{R}).$$

Now, let  $v_i$  be the  $i$ -th column of  $Q^\top$ . Then

$$B_{ij} = v_i^\top \cdot v_j = \langle v_i, v_j \rangle.$$

Now we start the proof of Theorem 2.1.

*Proof of Theorem 2.1*

1. Let  $\langle, \rangle$  denote the dot product on  $V = \mathbb{R}^n$ . Set

$$B = \theta I - C \tag{2.3}$$

$$= \text{symmetric matrix with eigenvalues } \theta_0 - \theta_i \geq 0 \tag{2.4}$$

$$= (\langle v_i, v_j \rangle)_{1 \leq i, j \leq n} \tag{2.5}$$

with the same  $v_1, \dots, v_n \in V$  by Lemma 2.1.

Observe:  $\sum_{i=1}^n \alpha_i v_i = 0$ .

*Pf.*

$$\left\| \sum_{i=1}^n \alpha_i v_i \right\|^2 = \left\langle \sum_{i=1}^n \alpha_i v_i, \sum_{i=1}^n \alpha_i v_i \right\rangle \tag{2.6}$$

$$= (\alpha_1 \quad \dots \quad \alpha_n) B \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \tag{2.7}$$

$$= v^\top B v \tag{2.8}$$

$$= 0, \tag{2.9}$$

since  $Bv = (\theta_0 I - C)v = 0$ .

Now set

$s =$  the number of indices  $i$ , where  $\alpha_i > 0$ .

Replacing  $v$  by  $-v$  if necessary, without loss of generality we may assume that  $s \geq 1$ . We want to show  $s = n$ .

Assume  $s < n$ . Without loss of generality, we may assume that  $\alpha_i > 0$  for  $1 \leq i \leq s$  and  $\alpha_i = 0$  for  $s+1 \leq i \leq n$ . Set

$$\rho = \alpha_1 v_1 + \dots + \alpha_s v_s = -\alpha_{s+1} v_{s+1} - \dots - \alpha_n v_n.$$

Then, for  $i = 1, \dots, s$ ,

$$\langle v_i, \rho \rangle = \sum_{j=s+1}^n -\alpha_j \langle v_i, v_j \rangle \quad (\langle v_i, v_j \rangle = B_{ij}, B = \theta_0 I - C) \quad (2.10)$$

$$= \sum_{j=s+1}^n (-\alpha_{ij})(-C_{ij}) \quad (2.11)$$

$$\leq 0. \quad (2.12)$$

Hence

$$0 \leq \langle \rho, \rho \rangle = \sum_{i=1}^s \alpha_i \langle v_i, \rho \rangle \leq 0,$$

as  $\alpha > 0$  and  $\langle v_i, \rho \rangle \leq 0$ . Thus, we have  $\langle \rho, \rho \rangle = 0$  and  $\rho = 0$ . For  $j = s+1, \dots, n$ ,

$$0 = \langle \rho, v_j \rangle = \sum_{i=1}^s \alpha_i \langle v_i, v_j \rangle \leq 0,$$

as  $\langle v_i, v_j \rangle = -C_{ij}$ .

Therefore,

$$0 = \langle v_i, v_j \rangle = -C_{ij} \text{ for } 1 \leq i \leq s, s+1 \leq j \leq n.$$

Since  $C$  is symmetric,

$$C = \begin{pmatrix} * & O \\ O & * \end{pmatrix}$$

Thus  $C$  is reducible, which is not the case. Hence  $s = n$ .

*Proof of Theorem 2.1 2.*

Suppose  $\dim V_0 \geq 2$ . Then,

$$\dim \left( V_0 \cap \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}^\perp \right) \geq 1.$$

So, there is a vector

$$0 \neq v = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \in V_0$$

with  $\alpha_1 = 0$ . This contradicts 1.

Now pick

$$0 \neq w = \begin{pmatrix} \beta_1 \\ \vdots \\ \beta_n \end{pmatrix} \in V_r.$$

*Proof of Theorem 2.1 3.*



Suppose  $\theta_r < -\theta_0$ . Since the eigenvalues of  $C^2$  are the squares of those of  $C$ ,  $\theta_r^2$  is the maximal eigenvalue of  $C^2$ .

Also we have  $C^2 w = \theta_r^2 w$ .

Observe that  $C^2$  is irreducible. (As otherwise,  $C$  is bipartite by Note 3, and we must have  $\theta_r = -\theta_0$ .) Therefore,  $\beta_i > 0$  for all  $i$  or  $\beta_i < 0$  for all  $i$ . We have

$$\langle v, w \rangle = \sum_{i=1}^n \alpha_i \beta_i \neq 0.$$

This is a contradiction, as  $V_0 \perp V_r$ .

*Proof of Theorem 2.1 4.*

$\Rightarrow$ : Let  $\theta_r = -\theta_0$ . Then  $\theta = \theta_1^2 = \theta_0^2$  is the maximal eigenvalue of  $C^2$ , and  $v$  and  $w$  are linearly independent eigenvalues for  $\theta$  for  $C^2$ . Hence, for  $C^2$ ,  $\text{mult}(\theta) \geq 2$ .

Thus by 2,  $C^2$  must be reducible. Therefore,  $C$  is bipartite by Note 3.

$\Leftarrow$ : This is Note 1.  $\square$

Let  $\Gamma = (X, E)$  be any graph.

**Definition 2.3.**  $\Gamma$  is said to be *bipartite* if the adjacency matrix  $A$  is bipartite. That is,  $X$  can be written as a disjoint union of  $X^+$  and  $X^-$  such that  $X^+, X^-$  contain no edges of  $\Gamma$ .

**Corollary 2.1.** *For any (connected) graph  $\Gamma$  with*

$$\text{Spec}(\Gamma) = \begin{pmatrix} \theta_0 & \theta_1 & \cdots & \theta_r \\ m_1 & m_1 & \cdots & m_r \end{pmatrix} \quad \text{with } \theta_0 > \theta_1 > \cdots > \theta_r.$$

*Let  $V_i$  be the eigenspace of  $\theta_i$ . Then the following holds.*

1. Suppose  $0 \neq v = \begin{pmatrix} \alpha_1 \\ \vdots \\ \alpha_n \end{pmatrix} \in V_0 \in \mathbb{R}^n$ . Then  $\alpha_i > 0$  for all  $i$  or  $\alpha_i < 0$  for all  $i$ .
2.  $m_0 = 1$ .
3.  $\theta_r \geq -\theta_0$  if and only if  $\Gamma$  is bipartite. In this case,

$$-\theta_i = \theta_{r-i} \text{ and } m_i = m_{r-i} \quad (0 \leq i \leq r)$$

*Proof.* This is a direct consequences of Theorem 2.1 and Note 3.  $\square$




# Chapter 3

## Lecture 3


Monday, January 25, 1993

Given graphs  $\Gamma = (X, E)$  and  $\Gamma' = (X', E')$ .


**Example 3.1.**  $\Delta = \{a, a^{-1}\}$



**Example 3.2.**  $\Delta = \{a, a^{-1}, a^2, a^{-2}\}$



**Example 3.3.**  $G = \langle a, b \mid a^6 = 1, b^2 = 1, ab = ba \rangle$ ,  $\Delta = \{a, a^{-1}, b\}$



## Chapter 4

# Applications

Some *significant* applications are demonstrated in this chapter.

### 4.1 Example one

### 4.2 Example two



## Chapter 5

# Final Words

We have finished a nice book.