

IL-6 Linear Regression Model

Variable	Standardized Coefficient	P	R ²
Model 1			0.134
PPT. Lower leg	-0.013	0.930	
HADS depression	0.211	0.270	
hsa-miR-30d-5p	-0.299	0.219	
hsa-miR-27b-3p	-0.316	0.125	
hsa-miR-34a-5p	-0.390	0.140	
hsa-miR-4479	0.099	0.572	
hsa-miR-487b-3p	-0.091	0.573	
hsa-miR-324-5p	0.661	0.030	
VEGF	-0.043	0.811	
hsa-miR-1304-3p	-0.306	0.116	
hsa-miR-433-3p	-0.295	0.099	
hsa-miR-143-3p	0.070	0.699	
hsa-miR-217-5p	-0.026	0.899	
IL-13	-0.140	0.597	
IL-15	0.080	0.677	
IL-8	0.026	0.890	
IL-6	0.448	0.213	
Model 2			0.276
HADS depression	0.253	0.099	
hsa-miR-27b-3p	-0.348	0.034	
hsa-miR-34a-5p	-0.371	0.084	
hsa-miR-324-5p	0.448	0.017	
hsa-miR-1304-3p	-0.389	0.012	
hsa-miR-433-3p	-0.286	0.068	
IL-6	0.370	0.072	

Table 3: Linear regression model of 17 QST, pain, questionnaire factors, miRNA, Inflammatory markers, and IL-6, selected to explain the variance in BDNF in patients with knee osteoarthritis 5 years after TKR. Model 1 consists of all variables. Model 2 uses backward elimination to identify independent factors explaining BDNF levels. The adjusted R^2 value suggests the combined predictive power of these values.