

## Computing the Lower Bound Efficiently

We have

$$\|y - Rx\|_2^2 = \|y_{k:m} - R_{k:m,k:m}x_{k:m}\|_2^2 + \|y_{1:k-1} - R_{1:k-1,1:k-1}x_{1:k-1} - R_{1:k-1,k:m}x_{k:m}\|_2^2.$$

Suppose that  $U = R^{-1}$  has been computed, then the lower bound on the second term of the above equation is:

$$\sigma_{\min}(R_{1:k-1,1:k-1}) \min_{x_{1:k-1}} \|x_{1:k-1} - U_{1:k-1,1:k-1}(y_{1:k-1} - R_{1:k-1,k:m}x_{k:m})\|_2^2.$$

We need to efficiently compute:

$$f^{k-1} \equiv U_{1:k-1,1:k-1}(y_{1:k-1} - R_{1:k-1,k:m}x_{k:m}). \quad (1)$$

For  $k = m$  we compute  $f^{k-1}$  directly by matrix-vector multiplication. For  $k < m$ , a recursion is derived to relate  $f^{k-2}$  to the already known  $f^{k-1}$ .

First split  $f^{k-1}$  into 2 parts,  $f_{1:k-2}^{k-1}$  and  $f_{k-1:m}^{k-1}$  as follows:

$$\begin{aligned} f^{k-1} &= U_{1:k-1,1:k-1}(y_{1:k-1} - R_{1:k-1,k:m}x_{k:m}) \\ &= \begin{bmatrix} U_{1:k-2,1:k-2} & U_{1:k-2,k-1} \\ 0 & u_{k-1,k-1} \end{bmatrix} \begin{bmatrix} y_{1:k-2} \\ y_{k-1} \end{bmatrix} - \begin{bmatrix} U_{1:k-2,1:k-2} & U_{1:k-2,k-1} \\ 0 & u_{k-1,k-1} \end{bmatrix} \begin{bmatrix} R_{1:k-2,k:m} \\ R_{k-1,k:m} \end{bmatrix} x_{k:m}. \\ &= \begin{bmatrix} U_{1:k-2,1:k-2}y_{1:k-2} + U_{1:k-2,k-1}y_{k-1} \\ u_{k-1,1:k-1}y_{k-1} \end{bmatrix} - \begin{bmatrix} U_{1:k-2,1:k-2}R_{1:k-2,k:m}x_{k:m} + U_{1:k-2,k-1}R_{k-1,k:m}x_{k:m} \\ u_{k-1,k-1}R_{k-1,k:m}x_{k:m} \end{bmatrix}. \end{aligned}$$

So,

$$f_{1:k-2}^{k-1} = U_{1:k-2,1:k-2}y_{1:k-2} + U_{1:k-2,k-1}y_{k-1} - U_{1:k-2,1:k-2}R_{1:k-2,k:m}x_{k:m} - U_{1:k-2,k-1}R_{k-1,k:m}x_{k:m}. \quad (2)$$

We also know from (1) that

$$\begin{aligned} f^{k-2} &= U_{1:k-2,1:k-2}y_{1:k-2} - U_{1:k-2,1:k-2}R_{1:k-2,k-1:m}x_{k-1:m} \\ &= U_{1:k-2,1:k-2}y_{1:k-2} - U_{1:k-2,1:k-2} \begin{bmatrix} R_{1:k-2,k-1} & R_{1:k-2,k:m} \end{bmatrix} \begin{bmatrix} x_{k-1} \\ x_{k:m} \end{bmatrix} \\ &= U_{1:k-2,1:k-2}y_{1:k-2} - U_{1:k-2,1:k-2}R_{1:k-2,k:m}x_{k:m} - U_{1:k-2,1:k-2}R_{1:k-2,k-1}x_{k-1}. \end{aligned} \quad (3)$$

Comparing (2) and (3), we notice that the first 2 terms in (3) also appear in (2) as the first and third term. This gives us the following equation to relate  $f^{k-1}$  and  $f^{k-2}$ :

$$f^{k-2} = f_{1:k-2}^{k-1} - U_{1:k-2,k-1}y_{k-1} + U_{1:k-2,k-1}R_{k-1,k:m}x_{k:m} - U_{1:k-2,1:k-2}R_{1:k-2,k-1}x_{k-1}. \quad (4)$$

The last term in the above equality involves the matrix-vector multiplication  $U_{1:k-2,1:k-2}R_{1:k-2,k-1}$ , which can be simplified. In fact, since  $UR = I$ , it is easy to verify that

$$\begin{bmatrix} U_{1:k-2,1:k-2} & U_{1:k-2,k-1} \end{bmatrix} \begin{bmatrix} R_{1:k-2,k-1} \\ r_{k-1,k-1} \end{bmatrix} = 0.$$

Thus  $U_{1:k-2,1:k-2}R_{1:k-2,k-1} = -U_{1:k-2,k-1}r_{k-1,k-1}$ . Then from (4) we have

$$\begin{aligned} f^{k-2} &= f_{1:k-2}^{k-1} - U_{1:k-2,k-1}y_{k-1} + U_{1:k-2,k-1}R_{k-1,k:m}x_{k:m} + U_{1:k-2,k-1}r_{k-1,k-1}x_{k-1} \\ &= f_{1:k-2}^{k-1} - U_{1:k-2,k-1}(y_{k-1} - R_{k-1,k-1:m}x_{k-1:m}). \end{aligned}$$

Notice that  $y_{k-1} - R_{k-1,k-1:m}x_{k-1:m}$  is computed in the search process. So if  $f^{k-1}$  is known, the cost for computing  $f^{k-2}$  is about  $2k$  flops. Note that when  $x_{k:m}$  is determined, the lower bound is determined. During the search process, when move from a higher lever to a lower level, we need to compute the lower bound and store the value; when move a lower level to a higher level, we just use the stored values of the lower bounds.