

Class10: Structural Bioinformatics pt.1

Idara Ibekwe: A16865157

##What is in the PDB database

The main repository of the biomolecular structure info is the PDB <www.rcsb.org>.

Let's see what the database contains:

```
stats=read.csv("Data Export Summary.csv", row.names=1)
stats
```

	X.ray	EM	NMR	Multiple.methods	Neutron	Other
Protein (only)	163,468	13,582	12,390	204	74	32
Protein/Oligosaccharide	9,437	2,287	34	8	2	0
Protein/NA	8,482	4,181	286	7	0	0
Nucleic acid (only)	2,800	132	1,488	14	3	1
Other	164	9	33	0	0	0
Oligosaccharide (only)	11	0	6	1	0	4
Total						
Protein (only)	189,750					
Protein/Oligosaccharide	11,768					
Protein/NA	12,956					
Nucleic acid (only)	4,438					
Other	206					
Oligosaccharide (only)	22					

Q1. What percentage of structures in the PDB are solved by X-Ray and Electron Microscopy.

```
as.numeric(stats$X.ray)
```

Warning: NAs introduced by coercion

```
[1] NA NA NA NA 164 11
```

We got to get rid of the commas. Can you find a function to get rid of the commas

```
x <- stats$X.ray
sum(as.numeric(gsub(",", "", x)))
```

```
[1] 184362
```

I am going to turn this into a function and then use `apply()` to work on the entire table of data.

```
sumcomma <- function(x) {
  sum(as.numeric(gsub(",", "", x)))
}

sumcomma(stats$X.ray)
```

```
[1] 184362
```

```
n.total <- sumcomma(stats$Total)
n.total
```

```
[1] 219140
```

```
sumcomma(stats$EM)
```

```
[1] 20191
```

```
apply(stats, 2, sumcomma)
```

X.ray	EM	NMR	Multiple.methods
184362	20191	14237	234
Neutron	Other	Total	
79	37	219140	

```
apply(stats, 2, sumcomma)/sumcomma(stats$Total)
```

X.ray	EM	NMR	Multiple.methods
0.8412978005	0.0921374464	0.0649676006	0.0010678105
Neutron	Other	Total	
0.0003605001	0.0001688418	1.0000000000	

84% of the structures were X-ray and 9.2% were EM

Q2: What proportion of structures in the PDB are protein?

Q3: Type HIV in the PDB website search box on the home page and determine how many HIV-1 protease structures are in the current PDB?

186898/248805733 * 100

[1] 0.07511804

##Visualizing the HIV-1 protease structure

Mol*(“mol-star”) viewer is now everywhere.


I want to insert my image from Mol* here.

Q4: Water molecules normally have 3 atoms. Why do we see just one atom per water molecule in this structure?

Q5: There is a critical “conserved” water molecule in the binding site. Can you identify this water molecule? What residue number does this water molecule have



Figure 1: My first molecular image



##Working with the bio3d package

```
library(bio3d)
```

```
pdb <- read.pdb("1hsg")
```

Note: Accessing on-line PDB file

```
pdb
```

```
Call: read.pdb(file = "1hsg")
```

```
Total Models#: 1
```

```
Total Atoms#: 1686, XYZs#: 5058 Chains#: 2 (values: A B)
```

```
Protein Atoms#: 1514 (residues/Calpha atoms#: 198)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 172 (residues: 128)
Non-protein/nucleic resid values: [ HOH (127), MK1 (1) ]
```

Protein sequence:

```
PQITLWQRPLVTIKIGGQLKEALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYD
QILIEICGHKAIGTVLVGPTPVNIIGRNLLTQIGCTLNFPQITLWQRPLVTIKIGGQLKE
ALLDTGADDTVLEEMSLPGRWKPKMIGGIGGFIKVRQYDQILIEICGHKAIGTVLVGPTP
VNIIGRNLLTQIGCTLNF
```

```
+ attr: atom, xyz, seqres, helix, sheet,
      calpha, remark, call
```

```
head(pdb$atom)
```

	type	eleno	elety	alt	resid	chain	resno	insert	x	y	z	o	b
1	ATOM	1	N	<NA>	PRO	A	1	<NA>	29.361	39.686	5.862	1	38.10
2	ATOM	2	CA	<NA>	PRO	A	1	<NA>	30.307	38.663	5.319	1	40.62
3	ATOM	3	C	<NA>	PRO	A	1	<NA>	29.760	38.071	4.022	1	42.64
4	ATOM	4	O	<NA>	PRO	A	1	<NA>	28.600	38.302	3.676	1	43.40
5	ATOM	5	CB	<NA>	PRO	A	1	<NA>	30.508	37.541	6.342	1	37.87
6	ATOM	6	CG	<NA>	PRO	A	1	<NA>	29.296	37.591	7.162	1	38.40

	segid	elesy	charge
1	<NA>	N	<NA>
2	<NA>	C	<NA>
3	<NA>	C	<NA>
4	<NA>	O	<NA>
5	<NA>	C	<NA>
6	<NA>	C	<NA>

```
pdbseq(pdb)[25]
```

```
25
"D"
```

##Predicting functional motions of a single structure

We can do a bioinformatics predictions of functional motions (i.e. flexibility/dynamics)

```
pdb <- read.pdb("6s36")
```

Note: Accessing on-line PDB file
PDB has ALT records, taking A only, rm.alt=TRUE

```
pdb
```

```
Call: read.pdb(file = "6s36")
```

```
Total Models#: 1
```

```
Total Atoms#: 1898, XYZs#: 5694 Chains#: 1 (values: A)
```

```
Protein Atoms#: 1654 (residues/Calpha atoms#: 214)
```

```
Nucleic acid Atoms#: 0 (residues/phosphate atoms#: 0)
```

```
Non-protein/nucleic Atoms#: 244 (residues: 244)
```

```
Non-protein/nucleic resid values: [ CL (3), HOH (238), MG (2), NA (1) ]
```

```
Protein sequence:
```

```
MRIILLGAPGAGKGTQAQFIMEKYGIPQISTGDMRLRAAVKSGSELGKQAKDIMDAGKLV  
DELVIALVKERIAQEDCRNGFLLDGFPRTIPQADAMKEAGINVDYVLEFDVPDELIVDKI  
VGRRVHAPSGRVYHVKFNPPKVEGKDDVTGEELTTRKDDQEETVRKRLVEYHQMTAPLIG  
YYSKEAEAGNTKYAKVDGTPVAEVRADLEKILG
```


```
+ attr: atom, xyz, seqres, helix, sheet,  
      calpha, remark, call
```

```
m <- nma(pdb)
```

```
Building Hessian... Done in 0.05 seconds.
```

```
Diagonalizing Hessian... Done in 0.31 seconds.
```

```
plot(m)
```



```
mktrj(m,file="adk_m7.pdb")
```