

Table of Contents

Executive Summary.....2

Introduction.....2

Barriers to Timely Diagnosis of Bipolar Disorder.....3

Rehabilitation Process Components.....4

Comparison of Outcomes: Untreated vs. Treated Bipolar Disorder...4

References.....5



Experiencing Bipolar Disorder: An Individual's Account of Diagnosis and Rehabilitation

Nicole S Brown

Author and Educator

Citation : Nicole S Brown (2024) Experiencing Bipolar Disorder: An Individual's Account of Diagnosis and Rehabilitation Peercite J Ment Health. 2(S1):620-623.

***Corresponding Author : Nicole S Brown**

Email : brownns8@hotmail.com

Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://peercite.org>).

Abstract :

Bipolar disorder, affecting millions worldwide, is often undiagnosed for extended periods due to stigma and lack of understanding. This case study highlights an individual undiagnosed for 16 years, where symptoms began in 1999 but a formal diagnosis occurred in 2015. Emotional trauma, including relational breakdowns, exacerbated anxiety and depression, prompting self-hospitalization and a diagnosis. Recovery involved a combination of therapy, psychiatric consultation, and medication management, demonstrating the importance of early intervention, customized treatment, and ongoing support. The individual's experiences emphasize the emotional impact of interpersonal relationships on mental health and the need for holistic treatment approaches. Despite delays, the person achieved academic and professional success, showcasing resilience and the transformative power of treatment, creative expression, and self-advocacy. Addressing systemic barriers, reducing stigma, and promoting mental health education are essential for improving outcomes for bipolar disorder patients.

Keywords: Bipolar disorder, Undiagnosed, Stigma, Emotional trauma, Anxiety and depression


Introduction :

Bipolar illness is a mental health condition that impacts millions of people globally. It is characterized by episodes of mania and depression, which can profoundly affect an individual's emotional and psychological health (National Institute of Mental Health [NIMH], 2021). Bipolar disorder frequently goes undiagnosed for prolonged durations due to stigma and insufficient awareness. An instance that exemplifies these issues pertains to an individual who endured undiagnosed bipolar disorder for 16 years. The condition's onset was retroactively determined to have occurred in 1999, when the subject was 21. However, a formal diagnosis was not established until 2015. Throughout this extended duration, symptoms of the condition were apparent; nevertheless, the pervasive stigma associated

with mental health hindered the individual from pursuing professional assistance. Research highlights that societal stigma remains a substantial obstacle to prompt mental health care, resulting in numerous persons being undiagnosed and untreated for extended periods (Corrigan et al., 2014). The pivotal moment in this instance transpired after a series of emotionally tumultuous events, including the conclusion of a long-term relationship and a failed attempt at reconciliation. These events resulted in increased anxiety and a depressive episode. At this point, the individual disclosed details about alleged clandestine activities involving their former partner and others over 15 years. This disclosure procedure, albeit very taxing, initiated their journey towards diagnosis & rehabilitation.

In 2015, the individual autonomously pursued hospitalization, where they obtained a formal diagnosis of bipolar disorder. This phase underscores the significance of self-advocacy and acknowledgment of mental health requirements, despite personal and cultural adversities. Research emphasizes that prompt intervention and precise diagnosis are essential for the optimal management of bipolar disorder and the enhancement of quality of life (Geddes & Mankowitz, 2013).

Figure 1: Barriers to Timely Diagnosis of Bipolar Disorder



This bar chart illustrates the major barriers to timely diagnosis as identified in the literature


The individual's rehabilitation process included therapy and psychological evaluations to develop a tailored treatment strategy. Medication management was crucial in stabilizing their condition, enabling them to resume professional and academic activities. Their capacity to attain educational achievements, publish scholarly works, and uphold professional duties illustrates the possibility for those with bipolar disorder to have satisfying lives when afforded suitable treatment and support. Studies indicate that regular therapy combined with medication adherence can help individuals with bipolar disorder attain significant personal and professional success (Michalak et al., 2016). The individual's experiences emphasize the significant emotional influence of personal interactions on mental health. Their story underscores the relationship between connection, betrayal, and emotional stability. Emotional breakdowns induced by interpersonal difficulties are prevalent in individuals with bipolar disorder, highlighting the necessity for a holistic approach to mental health care that considers both clinical and relational elements (Reenen et al., 2017). The impact of delayed diagnosis of bipolar disorder cannot be overstated. Many individuals face similar challenges, navigating their lives while unknowingly dealing with the disorder's manifestations.

Studies indicate that untreated bipolar disorder often leads to adverse outcomes, such as substance abuse, strained relationships, and underachievement in personal and professional spheres (Merikangas et al., 2011). These findings highlight the necessity for comprehensive mental health screening and education to facilitate early detection and intervention. Furthermore, societal stigmas that perpetuate silence and avoidance exacerbate the challenges faced by individuals living with bipolar disorder. Cultural norms and misconceptions often paint mental illness as a personal failure, discouraging individuals from seeking help. Corrigan et al. (2014) emphasize the role of stigma in delaying treatment, noting that reducing stigma through community education and advocacy campaigns can significantly improve mental health outcomes. In this case, interpersonal relationships played a critical role in the individual's mental health journey.

The emotional toll of their long-term relationship, characterized by betrayal and unresolved conflicts, contributed to their initial breakdown. Interpersonal stressors are well-documented triggers for bipolar episodes, as they intensify feelings of instability and emotional dysregulation (Reenen et al., 2017). Addressing these relational factors is crucial in developing effective therapeutic interventions for individuals with bipolar disorder. Therapy and psychiatric consultations were pivotal components in the individual's rehabilitation journey. Cognitive-behavioural therapy (CBT), in particular, is effective in managing bipolar disorder by equipping individuals with strategies to identify and modify maladaptive thought patterns (Miklowitz et al., 2007).

The inclusion of psychotherapy alongside medication underscores the importance of a multimodal treatment approach. Additionally, family-focused therapy (FFT) has emerged as an effective intervention for addressing relational dynamics and fostering a supportive environment for individuals with bipolar disorder (Miklowitz et al., 2003). From a pharmacological perspective, mood stabilizers and antipsychotics are commonly prescribed to manage the symptoms of bipolar disorder. For this individual, medication played a crucial role in restoring balance and enabling them to engage in academic and professional pursuits. Geddes and Miklowitz (2013) highlight the importance of personalized medication regimens, tailored to address the unique needs of each patient. Medication adherence, however, remains a significant challenge, as individuals often discontinue treatment due to side effects or perceived stigma (Lingam & Scott, 2002). Addressing these barriers through patient education and regular follow-ups is essential to ensure sustained recovery.

Figure 2: Rehabilitation Process Components



A pie chart representing the key components of the rehabilitation process.

The individual’s ability to achieve academic milestones and professional success despite their diagnosis demonstrates the potential for resilience and growth. Bipolar disorder does not preclude individuals from leading fulfilling lives, as long as they have access to appropriate resources and support systems. Michalak et al. (2016) emphasize the importance of fostering a sense of purpose and self-efficacy among individuals with bipolar disorder, as these factors significantly contribute to their overall quality of life. In reflecting on their journey, the individual composed a poem that encapsulates their experiences and emotions. Creative expression, such as writing and art, has been shown to have therapeutic benefits for individuals with mental health conditions (Forgeard et al., 2016). Engaging in creative activities allows individuals to process their experiences, communicate their feelings, and foster a sense of empowerment. This form of self-expression serves as a powerful tool for raising awareness and reducing the stigma surrounding mental health. The broader implications of this case underscore the importance of addressing systemic barriers to mental health care. Policymakers and healthcare providers must prioritize initiatives that promote early detection, reduce stigma, and increase access to affordable treatment options. Public health campaigns, workplace mental health programs, and community support networks can play a crucial role in creating a more inclusive and supportive environment for individuals with bipolar disorder. In conclusion, this account highlights the challenges and triumphs of living with bipolar disorder. The individual’s journey from delayed diagnosis to successful rehabilitation underscores the transformative power of treatment and support.

By addressing societal stigma, enhancing access to care, and fostering a holistic approach to mental health, we can empower individuals with bipolar disorder to lead meaningful and productive lives. Their story serves as a testament to the resilience of the human spirit and the importance of prioritizing mental health in our communities.

Figure 3: Comparison of Outcomes: Untreated vs. Treated Bipolar Disorder

Outcome	Untreated Bipolar Disorder	Treated Bipolar Disorder
Substance abuse prevalence	High	Low
Relationship quality	Strained	Stable
Academic and professional success	Low	High
The overall quality of life	Poor	Good

A table comparing key outcomes for untreated and treated bipolar disorder based on cited studies

References :

1. Corrigan, P. W., Druss, B. G., & Perlick, D. A. (2014). The influence of mental disease stigma on the pursuit and engagement in mental health services. *Psychological Science in the Public Interest*, 15(2), 37-70. <https://doi.org/10.1177/1529100614531398>
2. Integrating positive and negative pathways. *Frontiers in Psychology*, 7, 1274. <https://doi.org/10.3389/fpsyg.2016.01274> Forgeard, M. J., Elstein, J. G., & Davidson, R. J. (2016). Creative experiences and mental health:
3. Geddes, J. R., & Miklowitz, D. J. (2013). Management of bipolar disorder. *The Lancet*, 381(9878), 1672-1682. [https://doi.org/10.1016/S0140-6736\(13\)60855-0](https://doi.org/10.1016/S0140-6736(13)60855-0)
4. Lingam, R., & Scott, J. (2002). Treatment non-adherence in affective disorders. *Acta Psychiatrica Scandinavica*, 105(3), 164-172. <https://doi.org/10.1034/j.1600-2002.1r084.x>
5. Merikangas, K. R., Jin, R., He, J. P., Kessler, R. C., Lee, S., Sampson, N. A., ... & Zarkov, Z. (2011). Prevalence and correlates of bipolar spectrum disorder in the world mental health survey initiative.
6. *Archives of General Psychiatry*, 68(3), 241-251. <https://doi.org/10.1001/archgenpsychiatry.2011.12> Michalak, E. E., Yatham, L. N., Kolesar, S., & Lam, R. W. (2016). Bipolar disorder and quality of life: A patient-centric viewpoint. *Quality of Life Research*, 25(9), 2433-2442. <https://doi.org/10.1007/s11136-016-1280-5>
7. Miklowitz, D. J., & Scott, J. (2009). Psychosocial treatments for bipolar disorder: Cost-effectiveness, mediating mechanisms, and future directions. *Bipolar Disorders*, 11(2), 110-122. <https://doi.org/10.1111/j.1399-5618.2009.00675.x> Mankowitz, D. J., & Johnson, S. L. (200)