Received: January 19,, 2024 | Accepted: February 9, 2024 | Published: February 16, 2024

ORIGINAL RESEARCH

Peercite - Journal of Agriculture and Food Research

A REVIEW: ON THE HEALTH BENEFITS OF ORGANIC FOOD AND ITS DISTINCTIONS FROM CONVENTIONAL FOOD

ENAS SHEHAB AHMED[1] , RAWAA ADNAN KHALAF[2]

1 Department of Biomedical Engineering, College of Engineering, Al-Nahrain University, Jadriya, Baghdad, Iraq

2 Department of Biotechnology Research Center/Al-Nahrain University, Iraq

★***Corresponding Autor :***
ENAS SHEHAB AHMED
Email :

Citation: Galanakis CM (2024) A review: on the health benefits of organic food and its distinctions from conventional food. Peercite J Agric Food Res. 1: pages.

Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// peercite.org).

Abstract :

As public awareness of the benefits of organic food for health, social well-being, the environment, and sustainable development has increased, so has knowledge about its benefits. As this sector continues to grow, it is important to understand consumer opinion. The current study was systematically designed to assess the strength of data supporting the health benefits of feeding organic foods. We use only natural fertilizers as organic food. Regulations surrounding organic livestock farming are extensive and affect various aspects such as nutrition, reproduction, housing, medical care, and treatment. Additionally, these foods are not produced using genetically modified pesticides or insecticides. Animals raised on organic farms are given more living space than animals raised in conventional systems and should also receive organic feed, straw bedding and roughage. The fatty acid composition of milk, eggs, and meat is greatly influenced by animal nutrition. Breed selection is related to the animal's health and well-being. Proper nutrition, husbandry, and management all contribute to disease prevention and improved animal welfare. Antibiotic expiry dates are a global health concern today, and many international health organizations recognize that they pose a threat to current health care systems. Therefore, the market demand for organic products and their ease of use has increased in recent years, contributing significantly to the economy. Many people choose organic foods over conventional foods because they are healthier and contain fewer chemicals. Therefore, increasing consumer knowledge about the value of organic products and supporting their production is crucial for the growth of organic agriculture and pure water production. The organic food market is growing rapidly because consumers believe that organic foods are healthier and have a better nutritional profile than conventional foods.

Keywords : Conventional food, Prepacked foods, Organic food, Health concern

Introduction :

The importance of organic food production is at odds with conventional agriculture, human life, food safety animal welfare considerations, and environmental concerns. These concerns and growing attitudes of organic users have led in part to the emergence of different organic user groups called food phoebes, environmentalists, healthy consumers, welfare advocates, and humanists [1]. His interest in organic farming has led to his different studies comparing the properties of conventional and organically produced foods.When meeting such consumer expectations, it is important to carefully consider the issue of feed levels for conventional and organic foods to refute or support this distinction. Customers buy organic food because of their trust in the high quality of products. Generality of determination of the items evaluated showed significant differences in total carbohydrates, vitamin C dry matte and polyphenolic compounds between conventionally and organically produced fruits and vegetables [2].

Effects of fertilizers:

The main difference between organic and conventional foods is that they are free of pesticides, heavy metal residues, and fertilizers due to regulated production practices. The majority of the literature studies measure these chemicals using organic foods to demonstrate limitations. Pesticide use affects nitrogen fertilization and its deficiency, leading to the production of plant metabolites and bioactive substances [3].

The results are categorized according to the content of minerals, polyunsaturated fatty acids, polyphenols, and essential amino acids. He has already published, all of which come to different conclusions. Specifically, there was no variation in food prices between the two alternatives, demonstrating that natural products were more satisfying than others in these areas.

The opposite result was essentially characteristic of a lack of balance in the focus on execution and planning. In fact, studies have repeatedly shown that whole foods are better than conventional foods [4]. As, a rigorous test was recently carried out to analyze the formation of synthetic mixtures in different food sources, focusing on process and application and checking for diversity. The dietary sources considered were beta-carotene, all other cancer prevention drugs, rascorbic acid, potassium, vitamin E, phosphorus, complete phenols, pure protein, nitrates, and polyphenols.

Customers are willing to charge significantly higher prices for natural milk because they believe it is better than regular milk. However, not many researchers have conducted logical studies to ensure the contrast in food value between natural milk and regular milk. Additionally, there are no or minimal logical studies examining the specific characteristics and potential contrasts of the two food sources [5].

A recent study compared the food composition of consumer-purchased processed milk with commercially available milks [2]. Further convergence of protein studies reveals contrast in polyunsaturated and forming linoleic acid (CLA) and unsaturated fat profiles between retail produced natural milk and conventional retail milk. Of milk from different production systems. Additionally, folic acid (an important B nutrient during pregnancy) was less concentrated in natural milk, as very high temperatures (UHT) are more prevalent when natural milk is administered [4].

Results:

After analyzing the results of various previous articles that investigated the differences between the two types of foods, we found the following: Cereals: Cereals play an important role in the human diet due to the great regulation of food intake (table 1).

Substance	result
Phenolic compounds	organic is higher than conventional food

Table 1: Examination of chosen nature of organic and ordinary grains.

Based on pathological data, conventional feeding of bran products and whole grains may contribute to the prevention of cardiovascular diseases, especially diabetes, certain types of cancer, and the gastrointestinal system. Precious happiness [6]. Efficacy is usually determined by the presence of micronutrients, bioactive substances, and fiber co-metabolites in the diet (table 2).

Markers	Results
Protein	Lower
Dry matter	Higher
Fatty acids	Higher level of CLA
Total fat	Higher

Table 2: Chosen quality markers of organic milk in correlation with traditional milk.

Phenol corrosion concentrations in normal and natural winter wheat and mature spring wheat. A study by Dimberg and colleagues found a surprising naturally produced winter and spring wheats with higher total phenolic caustic fixation than normal wheat, although the difference was not large [7]. Showed that, contrary to standards, natural oatmeal and regular oatmeal do not differ in terms of phenolic compounds (table 3) [3].

Fatty acids	Higher level of n-3 with Lower level of n-6
Total fat	Lower
Carcass mass	Lower

Table 3: quality markers of organic beef meat in correlation with conventional beef meat.

Estimated the amount of flavonoids in open buckwheat grains. Results showed that differences in agricultural practices do not affect all plants and secondary metabolites in the same way. Concentrations of two of the four flavonoids were slightly higher in organic products, while the other concentrations remained the same [1,8]. The difference depends on the range (table 4).

Total fat	Higher
Fatty acids	Higher level of PUFA
mass of Carcass	Lower

Table 4: Quality markers of organic pork meat in correlation with conventional pork meat.

Conclusion:

Natural cultivation is a horticultural system developed by livestock breeders since ancient times that avoids the use of pesticides, growth regulators, and artificial fertilizers. Livestock diets eliminate most of the products used by the animals. The application of natural His cultivation has an extraordinary impact on overall health, reducing the risk of overweight/obesity in adults and irritability in adolescents, and reducing the development of certain Her diseases in consumers promotes it [9,10]. Overall, many people are choosing to purchase natural foods such as traditional His foods. This is because they have health benefits and fewer complex ingredients. Despite the fact that natural agriculture plays the most important role in human well-being and biodiversity security, there are still numerous limitations, such as high production costs, that hinder its development stage [11]. In summary, private and public organic farm owners need training, support, and encouragement in organic farm management. The public sector and related foundations need to draw attention to and support the value of natural products for customers through deductive educational research [12].

Conflict of interest:

Author declares no conflict of interest.

References

1. Wilkins JL and Hillers VN (1994) Influences of pesticide residue and environmental concerns on organic food preference among food cooperative members and non-members in Washington State. J of Nutrit Edu. 26:26-33.

2. Al-Karawi AS, Abid FM, Mustafa A et al. (2024) Revealing the Urinary Microbiota in Prostate Cancer: A Comprehensive Review Unveiling Insights into Pathogenesis and Clinical Application. Al-Salam J for Med Sci. 3:45-54.

3. Davies A, Titterington AJ, Cochrane C (1995) Who buys organic food? A profile of the purchasers of organic in Northern Ireland. British Food J. 97(10):17-23.

4. Ellis KA, Innocent G, Grove-White D et al. (2006) Comparing the fatty acid composition of organic and conventional milk. J Dairy Sci. 89:1938-1950.

5. Bourn D, Prescott J (2002) A Comparison of the Nutritional Value, Sensory Qualities, and Food Safety of Organically and Conventionally Produced Foods. Crit Rev Food Sci Nutr. 42:1.

6. Dangour D, Dodhia S, Hayter A et al. (2009) Nutritional quality of organic foods: a systematic review. Am J Clin Nutr. 90:680.

7. Brandt K, Leifert C, Sanderson R et al. (2011) Agroecosystem management and nutritional quality of plant foods: the case of organic fruits and vegetables. Crit Rev Plant Sci. 30:177-197.

8. Hansen LL, Claudi-Magnussen C, Jensen SK et al. (2006) Effect of organic pig production systems on performance and meat quality. Meat Science. 74:605-615.

9. Woodward BW, Fernandez MI (1999) Comparison of conventional and organic beef production systems II. Carcass characteristics. Livestock Prod Sci. 61:225-231.

10. Meyer KA, Kushi LH, Jacobs DR Jr et al. (2000) Carbohydrates, dietary fiber, and incident type 2 diabetes in older women. Am J Clin Nutr. 71:921-930.

11. Kalinova J, Vrchotova N (2011) The influence of organic and conventional crop management, variety and year on the yield and flavonoid level in common buckwheat groats. Food Chem. 127:602-508.

12. Dimberg LH, Gissen C, Nilsson J (2005) Phenolic compounds in oat grains (Avena sativa L.) grown in conventional and organic systems. Ambio. 34:331-337.