Received: January 02, 2023 | Accepted:January 29, 2023 | Published: February 06, 2023

Crossref PEERCITE PUBLISHERS Open Access

COMMENTARY

Peercite Journal of Cell Biology

POMEGRANATE EXTRACT IN MCF-7 BREAST CANCER CELLS EFFECTS OF ANTIPROLIFERATIVE

Prasad Kovvuru

Department of Biomedical Sciences, Discovery Dr, Rensselaer, NY. Discovery Dr is in Rensselaer, NY, USA

Corresponding Author : Prasad Kovvuru

Email : dr.prasadkovvuru@gmail.com

Citation: Kovvuru p(2023) Pomegranate Extract in MCF-7 Breast Cancer Cells Effects of Antiproliferative. Peercite J Cell Biol. 1(2):4507-4509.

Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// peercite.org).

Abstract :

Pomegranate extract (PE) inhibits the proliferation of breast most cancers cells and stimulates apoptosis in MCF-7 breast most cancers cells. Whilst PE is a mighty antioxidant, the prevailing studies were carried out to have a look at the mechanisms of action of PE past antioxidation via reading cellular and molecular mechanisms underlying breast tumorigenesis. PE inhibited cell increase by using inducing mobile cycle arrest in G2/M observed by means of the induction of apoptosis. In comparison, antioxidants N-acetylcysteine and Trolox did not affect mobile growth at doses containing equivalent antioxidant capacity as PE, suggesting that boom inhibition by way of PE can't entirely be attributed to its excessive antioxidant potential. DNA microarray evaluation revealed that PE downregulated genes related to mitosis, chromosome organization, RNA processing, DNA replication and DNA restore, and upregulated genes concerned in law of apoptosis and mobile proliferation. HR represents a unique goal for cancer remedy, downregulation of HR by means of PE can be exploited for sensitization of tumors to anticancer capsules.

Keywords : DNA repair, Homologous recombination

Introduction :

The antioxidant capacity of pomegranate ellagitannins has been proposed to account for his or her pharmacological moves, despite the fact that the hyperlink between antioxidant potential and tumor growth inhibition has now not been hooked up. The genomewide transcriptome evaluation of PE results in breast most cancers cells have now not been pronounced to date. Inside the present examine, an integrative technique changed into implemented using Affymetrix gene arrays to identify modifications in gene expression following treatment with PE in MCF-7 breast most cancers cells after which validating relative mRNA expression by using quantitative RT-PCR (qRT-PCR) [1].

In addition, similarly analysis of the expression of predicted microRNAs (miRNAs) for substantially modulated genes became carried out.

To assess whether or not the antioxidant effects of PE could give an explanation for the found increase inhibition of breast cancer cells, cells were exposed to traditional antioxidants, N-acetylcysteine (NAC) and Trolox.

Commentary:

General RNA became extracted with the aid of Trizol technique according to the manufacturer instructions (existence technologies, Ambion, Carlsbad, CA) from cells dealt with with 50 mg/mL of PE for 72 h and their respective controls, accrued from three impartial biological replicates at three one of a kind passage numbers. The pleasant of the RNA samples become assessed by NanoDrop 1000 Spectrophotometer and samples having 260/280 ratio 2.0 and 260/230 ratio 2.0 have been decided on.

RNA integrity number (RIN) was measured for the treatment and manage samples the usage of Bioanalyzer and appropriate exceptional samples (RIN > 8) were further processed for hybridization into whole human genome Affymetrix Human Gene-1.0 ST array platform on the center for functional Genomics. All statistics were baseline transformed to the median of the manipulate facts [2].

The facts were filed to cast off probesets with expression degrees within the backside twentieth percentile throughout all samples. The entities had been in addition processed via 1.5-fold filter out settings to consist of those entities that have been differentially expressed between PE treatments and controls.

Differentially expressed entities had been then surpassed to a t-take a look at ($P < 0.05$) with a Benjamini-Hochberg fake discovery fee correction to attain the statistically large and differentially expressed genes. DAVID useful annotation evaluation become finished to focus on the maximum relevant gene ontology (go)-term enrichment related in the gene list. The downregulated/upregulated gene entities had been then filtered by using the default settings of EASE rating threshold to calculate the Fisher genuine P-fee for gene enrichment evaluation (smaller the P-value, the more potent the enrichment in annotation categories).

Validation of the microarray facts of decided on genes became performed in MCF-7 cells treated with 50 μg/mL of PE biological experiments unbiased from the ones used for microarray analysis. Total RNA was remoted the usage of Trizol and purified the usage of NucleoSpin package (Qiagen, Germantown, MD) [3].

The purity of the RNA was assessed through NanoDrop a thousand Spectrophotometer. cDNA become synthesized from 1 mg of overall RNA using Multiscribe reverse transcriptase enzyme and qRT-PCR changed into executed the use of SYBR green (implemented Biosciences) assays in ABI7900 actual-time PCR device. GAPDH turned into used as housework gene for normalizations. Three independent organic replicates had been analyzed in reproduction. All primers were subjected to a satisfactory control to assure that each primer set generates a specific PCR product.

Three exceptional bioinformatics algorithms, miRanda, TargetScan, and PicTar, have been used to pick out miRNAs expected to goal a fixed of genes that were notably downregulated by means of PE

The application of a single miRNA target prediction program results in high stages of false nice and false poor discovery charges but, an aggregate of two or greater packages will increase the price of known goal prediction as much as 72.6% or a hundred% for evolutionary conserved targets. The miRNA application predicts miRNA targets with the aid of taking into consideration most weightage of collection complementarity to the bases comprising of seed area. It also calculates hybridization strength between miRNA-mRNA using RNAFold [4]. The TargetScan software unearths the simple 7-nucleotide best seed sequence in shape between the mRNA target area and 50 give up of miRNA.

TargetScan further extends the series match via allowing G:U wobble pairing and uses hybridization electricity (RNAFold) to calculate thermodynamic loose energy for binding the closing base pairing element. The PicTar application fits ideal 7-mer starting at position 1 or 2 at the 5' give up of the miRNA calculates hybridization strength and discards volatile duplexes.

General RNA was extracted using Trizol reagent in line with the producer's hints (life technology, Ambion). The full RNA arrangements that include miRNAs were analyzed for purity using NanoDrop a thousand Spectrophotometer (Thermo Fisher medical). miRNA cDNA turned into synthesized from 1 mg of purified general RNA using qScript miRNA cDNA synthesis package (Quanta Biosciences, Gaithersburg, MD) and qRT-PCR changed into achieved in 20 mL reaction extent containing 4 mL template cDNA and 16 mL of response blend containing PerfeCTa SYBR inexperienced grasp blend, 10 mM PerfeCTa ordinary PCR primer and 10 mM PerfeCTa miRNA assay primer in ABI7900 real-time PCR device. The samples have been to start with heated at 958C for 2 min followed with the aid of 40 cycles of 95°C for 5 s and 60°C for 30 s. The outcomes were normalized to RNU6B. Three independent biological replicates have been analyzed in replica [5].

γ-H2AX foci form rapidly after DSB formation and solve following restore and as a result represent a touchy measure of DSB formation and repair. PE remedy resulted in an improved frequency of γ-H2AX foci with the highest increase (2-fold) at 72 h [3]. Binding of short noncoding RNA, miRNA, to the goal mRNA can result in mRNA degradation or translational repression. Dietary compounds had been shown to regulate miRNAs involved in a selection of biological approaches however whether or not PE can regulate miRNAs concerned in DNA restore is unknown. We used three exclusive bioinformatics algorithms, miRanda.

This finding may be of medical relevance and shows that intermittent exposure to PE nutritional supplements is probably effective without compromising anticancer activities, which can be deemed as a more viable dietary approach. Comparative increase inhibition assessment of PE, NAC, and Trolox suggests that the boom inhibition by means of PE isn't entirely attributed to its high antioxidant capacity. This changed into a sturdy purpose for research of other feasible mechanisms of motion, for which we hired Affymetrix microarrays.

Microarray analysis confirmed that PE treatmentsignificantly modulated over 900 genes. The maximum enriched move-phrases among downregulated genes protected mitosis, chromosomal corporation, RNA processing, DNA replication, DNA damage reaction, and DNA restore, whereas the most enriched move-terms amongst upregulated genes covered law of apoptosis and law of cellular proliferation.

Even though a lot of these mobile methods might also manage mobile boom and dying, this observe links for the primary time PE publicity and downregulation of DNA restore genes. Of unique interest, maximum of the DNA restores genes that were downregulated by means of PE are concerned in HR-mediated restore [2,4].

As an instance, the MRN complex collectively enables 5'-3' resection of DSB ends required for initiation of HR. BRCA1 interacts with each MRN may be worried in stop resection, despite the fact that BRCA1 has pleiotropic capabilities and its exact function in HR is hard to dissect. At the contrary, BRCA2 is predominantly involved in HR. The 30-unmarried-stranded DNA overhang generated at some stage in stop resection is bound to replication protein A (RPA), which is required for next recruitment of RAD51. BRCA2 displaces RPA and enables loading of RAD51 onto single-stranded DNA.

Whether or not PE solely downregulates DNA repair genes in cancer cells is unknown. Regardless of this, due to the fact the MRN complex is dispensable for the maintenance and viability of postmitotic tissues and HR is handiest energetic in replicating cells, it's miles probably that downregulation of HR in ordinary tissues could have handiest a minor impact, provided the NHEJ pathway is undamaged [6].

Thus, our look at showing that PE downregulates HR genes warrants in addition investigations regarding using PE as a nontoxic HR inhibitor available in the form of a whole fruit, fruit juice, or a nutritional complement tablet. At gift, it is unknown whether PE downregulates HR in vivo [3].

In summary, our take a look at shows that PE objectives a particular DNA restore pathway this is crucial for most cancers cell survival both spontaneously and after drug venture [4]. This gives a strong motive for similarly trying out of PE as a safe adjunct to the prevention of breast most cancers recurrence via its movements on breast cancer cells. But, further research is wanted to demonstrate these consequences in animal research and in tissue samples from girls with breast cancer.

Conflict of interest:

Author declares there is no conflict of interest.

References:

1. Yuan J, Chen J (2010) MRE11-RAD50-NBS1 complex dictates DNA repair independent of H2AX. J Biol Chem. 285:1097-1104.

2. Kusano C, Takao S, Noma H, et al. (2000) N-acetyl cysteine inhibits cell cycle progression in pancreatic carcinoma cells. Hum Cell. 13:213-220.

3. Lee YJ, Lee DM, Lee CH, et al. (2011) Suppression of human prostate cancer PC-3 cell growth by N-acetylcysteine involves over-expression of Cyr61. Toxicol In Vitro. 25:199-205.

4. Helleday T, Lo J, van Gent DC et al. (2007) DNA doublestrand break repair: From mechanistic understanding to cancer treatment. DNA Repair (Amst). 6:923-935.

5. Roy R, Chun J, Powell SN (2012) BRCA1 and BRCA2: Different roles in a common pathway of genome protection. Nat Rev Cancer. 12:68-78.

6.Chen X, Arciero CA, Wang C et al. (2006) BRCC36 is essential for ionizing radiation-induced BRCA1 phosphorylation and nuclear foci formation. Cancer Res. 66:5039-5046.