Received: June 29, 2023 | Accepted: July 27, 2023 | Published: August 03, 2023

Crossref PEERCITE PUBLISHERS Open Access

Peercite Journal of Pharmaceutical Sciences

REVIEW

The Rising Presence of Multidrug-Resistant Bacteria and Fungi in the Arabian Peninsula

Sayed AbdulAzeez

Department of Genetic Research, Institute for Research and Medical Consultations (IRMC), Imam Abdulrahman Bin Faisal University, Dammam, Saudi Arabia

***Corresponding Author : Sayed AbdulAzeez**

Email : d asayed@iau.edu.sa

Citation : AbdulAzeez S (2023) The Rising Presence of Multidrug-Resistant Bacteria and Fungi in the Arabian Peninsula. Peercite J Pharm Sci. 1(3):20-24.

http://doi.org/10.61641/pjps.2023.1.3.20-24

Copyright: © 2023 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (https:// peercite.org).

Abstract :

We aimed to pick out the superiority and emerging reputation of multidrug-resistant bacteria and fungi and their related mortality in nine countries in the Arabian Peninsula. Authentic studies articles and case research regarding multidrug-resistant bacteria and fungi within the Arabian Peninsula, published during the last 10 years, had been retrieved from PubMed and Scopus. A total of 382 studies have been blanketed as in step with the inclusion and exclusion standards, in addition to the PRISMA pointers, from an intensive screening of 1705 articles, so as to analyse the rising reputation and mortality. The emerging nature of one hundred twenty multidrug-resistant (MDR) micro-organism and fungi within the Arabian Peninsula is a severe subject that requires continuous tracking and immediate preventive measures. Extra than 50% (n = 453) of multidrug-resistant, microbe-related mortality (n = 871) within the Arabian Peninsula become due to MDR *Acinetobacter baumannii,* Mycobacterium tuberculosis and Staphylococcus aureus contamination. usual, a 17.51% mortality became suggested among MDR-inflamed patients inside the Arabian Peninsula from the 382 articles of this registered systematic evaluation. MDR *A. baumannii* (5600 isolates) prevailed in all of the nine nations of the Arabian Peninsula and changed into one of the quickest rising MDR bacteria with the best mortality (n = 210). A complete of 13,087 Mycobacterium tuberculosis isolates were mentioned within the region. *Candida auris* (580 strains) is the maximum standard some of the MDR fungal pathogen in the Arabian Peninsula, having triggered 44 mortalities. Active surveillance, consistent monitoring, the development of a candidate vaccine, an early diagnosis of MDR infection, the removal of multidrug resistance modulators and uninterrupted preventive measures with more advantageous data sharing are mandatory to govern MDR infection and related sicknesses of the Arabian Peninsula. Correct and speedy detection strategies are needed to differentiate MDR pressure from different strains of the species. This overview summarises the logical relation, incidence, emerging repute and related mortality of MDR microbes within the Arabian Peninsula.

Keywords : Multidrug-resistant bacteria, *Acinetobacter baumannii*, Mycobacterium tuberculosis, Mortality

Introduction:

Due to its large vicinity and population, the kingdom of Saudi Arabia has the most tremendous wide variety of scientific research close to MDR organisms a few of the nations of the Arabian Peninsula. Consequently, it has numerous obstacles that could sell the emergence and transmission of MDR micro-organisms.

Sooner or later, a total of 79 particular MDR microbes have been said. The problems of this were overcome due to a success efforts of different sectors to restriction the boom and established order of MDR micro-organisms within the USA. The lively monitoring of the onset and spread of MDR micro-organisms could be very crucial.

Many applicable articles were posted in studies and case research from Saudi Arabia on MDR bacteria and fungi [1]. The inclusion standards have been carried out carefully to all the seek outcomes (for articles posted after 2010) from Scopus and PubMed. A complete of 198 research, as in keeping with PRISMA 2009, were eligible for inclusion.

Three guides said MDR micro-organisms in multiple USA of the Arabian Peninsula: the report through Sonnevend and associates in Bahrain, Saudi Arabia and United Arab Emirates; an observe by way of Zowawi and co-workers in Saudi Arabia, United Arab Emirates, Oman, Qatar, Bahrain, and Kuwait and a record with the aid of Sonnev end and colleagues in Saudi Arabia, Kuwait, Oman and United Arab Emirates.

Those reports were considered one after the other for person counties and also considered as most effective 3 studies in widespread. A hundred and ninety-8 research stated diverse MDR organisms in Saudi Arabia, in which there had been appreciably extra MDR organisms compared to the other international locations in the Arabian Peninsula [2].

Saudi Arabia:

Saudi Arabia had the very best quantity of MDR traces stated compared to other nations 10972 *Escherichia coli*, 9829 *Mycobacterium tuberculosis*, 4092 *Klebsiella pneumonia,* 3787 *Acinetobacter baumannii,* 2594 *Pseudomonas aeruginosa*, 2014 *Acinetobacter,* 986 *Staphylococcus aureus,* 927 *Klebsiella* spp., 812 *Enterobacter*, 722 emphProteus vulgaris, 715 *Enterobacteriaceae*, 695 *Methicillin-resistant Staphylococcus aureus* (MRSA) and 553 Enterococcus. Recently, the emergence of 18 *Clostridioides difficile* and 19 *Candida auris* had been mentioned among 2010 and 2021, the top 5 research papers suggested MDR micro-organisms in Saudi Arabia [3].

The first file protected *A. baumannii* 59 isolates from human samples: urine, blood lifestyle respiratory specimens, pores and skin and tender tissue specimens, blood, sterile frame fluids Biofilm-Formation in Clonally Unrelated Multidrug-Resistant *Acinetobacter baumannii* Isolates), wound swabs, rectal swabs, and sputum. The second file blanketed E. coli isolates from human and animal samples, the isolates from human samples originated from tissue swabs (the genomic variety and antimicrobial resistance genes in multidrug-resistant CTX-M-fantastic isolates of Escherichia coli at a healthcare facility in Jeddah), urine, wound pus, vaginal swabs, blood culture, stools and ear swabs.

The isolates from animal samples originated from: chicken camel and minced meat samples. The 1/3 document covered 36 *k. pneumonia* isolates from human samples: blood, urine, wound swabs, sputum rectal swabs and suction swabs. The fourth file covered 36 *P. aeruginosa* isolates from human samples: respiration, urine, surgical, blood subculture, genital swabs, eye swabs, ear swabs, burns, pores and skin tissue, intra-stomach fluid, bone tissue, wound swabs, sputum, and tracheal swabs. The fifth document blanketed 31 *M. tuberculosis* isolates from human samples: sputum, tissues and biopsies, abscesses, belly fluids, breast fluid, cerebrospinal fluid, urine, knee fluid and synovial fluid, wound swabs, ENT swabs, blood, tracheal swabs [4].

Saudi Arabia had the highest mortality price, with 365 sufferers who died due to MDR microbes. The top three MDR organisms with the highest mortality prices had been *A. baumannii, M. tuberculosis* and *P. aeruginosa*. Six MDR organisms have been associated with mortality: *S. epidermidis, A. baumannii, M. tuberculosis, C. auris, P. aeruginosa* and *Enterobacteriaceae* there had been 365 MDR lines related to dying tuberculosis prompted the very best stage of mortality.

The effect of the migrant workers from different nations cannot be understated; in 2017, a Filipino resident of Saudi Arabia became recognized with MDR *Mycobacterium leprae*. The etiologic agent become found the use of metagenomic sequencing a biopsy pattern of the affected person's skin [5]. In one of the studies, one hundred and five adult sufferers admitted to the in depth care unit (ICU) at Aseer relevant sanatorium in 2014 and 2015 have been reviewed retrospectively. The species of *Acinetobacter* were isolated using particular phenotypes and proven by way of the automatic gadget, Vitek 2. Of the one hundred and five stains, genus *Acinetobacter* accounted for *A. baumannii* 49, *A. baumannii* complex 19, *A. baumannii/haemolyticus* 32, *A. haemolyticus* 4, *A. lwoffii* 1. There are properly-mounted studies on electron microscopic structures, which stated both lifeless and alive multidrug-resistant micro-organisms in Saudi Arabia, the use of confocal laser scanning micrography.

The kingdom of Bahrain had the bottom variety of studies articles, in evaluation to the relaxation of the Arabian Peninsula's countries, due to its small place and populace. The papers in this evaluation, posted between 2010 and 2021 and retrieved from PubMed and Scopus, were very well screened for their eligibility, as per the inclusion and exclusion criteria.

Best 3 studies published on MDR bacteria have been protected in the final qualitative evaluation, as according to PRISMA 2009. Two studies reported MDR micro-organisms in multiple countries: a have a look at by way of Sonnevend and co-workers in Bahrain, Saudi Arabia, and the United Arab Emirates and a examine via Zowawi and associates in Saudi Arabia, United Arab Emirates, Oman, Qatar, Bahrain, and Kuwait particular MDR microbes had been stated: *Acinetobacter baumannii* and *Escherichia coli* as compared to the other nations mentioned on this overview, Bahrain had the bottom said MDR stress (n = 11): eight *A. baumannii* and three *E. coli* (Supplementary desk S2). three *E. coli* isolates from human specimens had been acquired, such as urine and tissue swabs and 8 *A. baumannii* isolates, from which human samples of blood, sputum, swab, and urine were amassed in contrast to the mortality charges of the other nations, Bahrain said no MDR microbial contamination-associated deaths.

In one of the clinical articles, researchers screened for the mcr-1 gene of colistin resistant *Enterobacteriaceae* in Bahrain; the *mcr-1* genes had been wonderful and their antibiotic resistance turned into decided. The *mcr-1* gene changed into located in *E. coli* isolates [6]. In the study by Thani, on a urinary tract infection (UTI) cefotaxime resistant *E. coli* stress, a new DNA fragment was observed.

The DNA collection changed into located to be related to the pathogenicity island III536 locus. Researchers used the unmarried Genome unique Primer-PCR (SGSP-PCR) to look at the genomic nation of the newly identified locus, which become discovered to gain the antibiotic resistance gene blaCTX m. Bahrain changed into covered inside the studies on the molecular epidemiology and resistance mechanisms of carbapenem-resistant acinophilic microorganism (CRAB) for the Arabian Peninsula countries; eight *A. baumannii* isolates have been recognized in Bahrain hospitals, and OXA-51 was found in all of the isolates. Antibiotic resistance genes have been detected the use of PCR, and clonality turned into determined the use of repetitive series-primarily based PCR (rep-PCR). The majority (84%) of the isolates on this investigation had been related to health publicity. The increased cognizance of multidrug-resistant organisms in the Gulf Cooperation Council (GCC) states has great implications for enhancing infection manage approaches, developing antimicrobial stewardship packages and highlighting the significance of regional energetic tracking structures [7].

The research, published among 2010 and 2021, from decided on databases had been evaluated for eligibility, as in line with the inclusion and exclusion criteria, consistent with PRISMA 2009. Thirty-8 papers on multidrug-resistant (MDR) micro-organisms were protected in the very last qualitative evaluation. A total of 22 precise MDR micro-organisms had been recognized from Kuwait such as the studies that suggested MDR micro-organisms in numerous countries. There were seven publications that mentioned >100 MDR lines inclusive of: 7138 methicillin-resistant *Staphylococcus aureus* (MRSA), 1176 *Mycobacterium tuberculosis,* 561 *Candida auris*, 410 *Acinetobacter baumannii*, 302 *Staphylococcus aureus*, 222 *Klebsiella pneumonia* and 149 Enterobacteriaceae. Considerably, five studies in Kuwait from 2010 to 2021 discovered MDR microorganisms. The first have a look at found out eleven *M. tuberculosis*, from human samples together with, sputum, pus, tissue biopsy, lymph node and endotracheal secretion, and cerebrospinal samples [8]. The second examine discovered eight isolates of *A. baumannii* obtained from human urine and swab specimens, sputum, blood.

The third have a look at located six *C. auris* human isolates from axilla, groin, anterior nares, vascularline go out web sites, the oropharynx, respiration and urinary tract. The isolates were also taken from environmental samples: rooms, units occupied via all of the inflamed *C. auris*, colonised patients, medical gadgets, linen, partitions, floors, furnishings, doorknobs, bed railings, bedside drawers, toilet taps, and flush handles [9]. The fourth study located five isolates of *Enterobacteriaceae* received from blood, urine, wound, and tissue swabs in the end, the 5th observe analysed four *E. coli* isolates of human samples derived from blood subculture, urine, wounds, and crucial venous strain. After Saudi Arabia, Qatar, Jordan, and Iraq, Kuwait has the fifth highest mortality charge, with a complete variety of 78 non-survivors associated with MDR microbial infection. Seven articles pronounced MDR microbial infection associated with mortality. the highest mortality in Kuwait became associated with the MDR organism, *C. auris*, with 37 non-survivor sufferers, whereas *C. auris* were the most prominent MDRO in terms of the variety of deaths [10].

Healthcare-associated pneumonia as a result of MDR pathogens poses a sizeable deal with ment task. The frequency of antibiotic medication results in inefficient antimicrobial therapy. Jamal and associates pronounced particular MDR organisms: two isolates of *Haemophilus influenzae*, isolates of *Legionella pneumophila*, and one isolate of *Moraxella catarrhalis*.

Severely sick sufferers with clinically identified respiration tract infections, hospitalised at Mubarak Al Kabeer medical institution between January and April 2013, had been decided on for his or her take a look at [11]. Currently, a fast-multiplex, PCR-based totally Unyvero pneumonia utility (UPA) assay that aids in early decision making became on hand. The respiration samples from sufferers with nosocomial pneumonia covered the sputum, bronchoalveolar lavage fluid, and endotracheal secretions. From January to December 2016, a take a look at carried out via Hamza and associates discovered a unique pathogen of an MDR micro-organism. All person patients who had the specified GI strategies, used to estimate surgical website infection (SSI) quotes among gastrointestinal surgical procedures in all Kuwait governmental hospitals, had been tested on this take a look at. In the end, a single isolate of *Aeromonas hydrophila* become received from small bowel (SB) surgery [12].

Conclusion:

The active surveillance and consistent preventive efforts to control MDR infectionsand associated sicknesses can enhance healthcare facilities and the socio-economic fame of patients. Specifically, MDR *E. coli* is predominantly associated with urinary tract infections in being pregnant. The emergence of multi drug-resistant *Candida albicansis* commonplace amongst pregnant and non-pregnant women with vaginitis. *Listeria monocytogenes* from patients with spontaneous abortions and multi drug resistant medical *Enterococcus faecalis* from pregnant ladies spotlight the want for extra precautions and constant monitoring, with the right identification of MDR and susceptibility assessments prior to medication. The improvement of a candidate vaccine and the early analysis of MDR infection will be a higher alternative for protection towards MDR contamination. The maximum familiar multidrug-resistant *Escherichia coli, Mycobacterium tuberculosis, Acinetobacter baumannii, Klebsiella pneumoniae, Pseudomonas aeruginosa, Staphylococcus aureus, Acinetobacter, Enterobacteriaceae, Providencia* spp., *Streptococcus pneumoniae,* methicillin-resistant *Staphylococcus aureus* (MRSA), *Klebsiella* spp., *Enterobacter, Proteus vulgaris, Candida auris* and *Enterococcus* within the Arabian Peninsula should be recognized as it should be and rapidly the use of advanced molecular strategies with the aid of developing region-specific diagnostic equipment. The manipulate of infections by using the usage of specificity detecting structures is a critical protection method against MDRO.

One examine develops an instantaneous detecting machine for *Candida auris* with the aid of the usage of a singular multiplex actual-time quantitative PCR that has a high degree of specificity in detecting distinct species-level of Candida. The present cumulative observations will be informative for effective planning, designing preventive strategies, and prioritizing research goals to achieve MDRO-free zones.

Conflict of interest:

The authors declare no conflict of interest.

References:

1. Zowawi HM, Sartor AL, Sidjabat HE et al. (2015) Molecular Epidemiology of Carbapenem-Resistant Acinetobacter Baumannii Isolates in the Gulf Cooperation Council States: Dominance of OXA-23-Type Producers. J Clin Microbiol. 53:896-903.

2. Rolain JM, Loucif L, Al-Maslamani M et al. (2016) Emergence of Multidrug-Resistant Acinetobacter Baumannii Producing OXA-23 Carbapenemase in Qatar. New Microbes New Infect. 11:47-51.

3. Khan FY, Abukhattab M, AbuKamar M et al. (2014) Adult K Lebsiella Pneumoniae Meningitis in Qatar: Clinical Pattern of Ten Cases. Asian Pac J Trop Biomed. 4:669-672.

4. Khan FY, Elshafie SS, Almaslamani M et al. (2010) Epidemiology of Bacteraemia in Hamad General Hospital, Qatar: A One Year Hospital-Based Study. Travel Med Infect. Dis. 8:377-387.

5. Jamal W, Rotimi VO, Albert MJ et al. (2013) High Prevalence of VIM-4 and NDM-1 Metallo-β-Lactamase among Carbapenem-Resistant *Enterobacteriaceae*. J Med Microbiol. 62:1239-1244.

6. Jamal W, Al Roomi E, AbdulAziz LR et al. (2014) Evaluation of Curetis Unyvero, a Multiplex PCR-Based Testing System, for Rapid Detection of Bacteria and Antibiotic Resistance and Impact of the Assay on Management of Severe Nosocomial Pneumonia. J Clin Microbiol. 52:2487-2492.

7. Jamal WY, Albert MJ, Khodakhast F et al. (2015) Emergence of New Sequence Type OXA-48 Carbapenemase Producing *Enterobacteriaceae* in Kuwait. Microb Drug Resist. 21:329-334.

8. Alhussain FA, Yenugadhati N, Al Eidan FA et al. (2021) Risk Factors, Antimicrobial Susceptibility Pattern and Patient Outcomes of Pseudomonas Aeruginosa Infection: A Matched Case-Control Study. J Infect Public Health. 14:152-157.

9. Almaghrabi RS, Albalawi R, Mutabagani M et al. (2020) Molecular Characterisation and Clinical Outcomes of *Candida Auris* Infection: Single-centre Experience in Saudi Arabia. Mycoses. 63:452-460.

10. Abdalhamid B, Almaghrabi R, Althawadi S et al. (2018) First Report of *Candida Auris* Infections from Saudi Arabia. J Infect Public Health. 11:598-599.

11. Abed AR, Khudhair AM, Hussein IM (2020) Effects of Misuse of Antibiotics on the Resistance of Escherichia coli Isolated from the Intestines of Broiler Chickens. Int J Drug Deliv Technol. 10:190-194.

12. Al Wutayd O, Al Nafeesah A, Adam I et al. (2018) The Antibiotic Susceptibility Patterns of Uropathogens Isolated in Qassim, Saudi Arabia. J Infect Dev Ctries. 12:946-952.